



Partner-specific induction of *Spodoptera frugiperda* immune genes in response to the entomopathogenic nematobacterial complex *Steinernema carpocapsae*-*Xenorhabdus nematophila*

Louise Huot, Audrey Bigourdan, Sylvie Pages, Jean-Claude Ogier,
Pierre-Alain Girard, Nicolas Nègre, Bernard Duvic

► To cite this version:

Louise Huot, Audrey Bigourdan, Sylvie Pages, Jean-Claude Ogier, Pierre-Alain Girard, et al.. Partner-specific induction of *Spodoptera frugiperda* immune genes in response to the entomopathogenic nematobacterial complex *Steinernema carpocapsae*-*Xenorhabdus nematophila*. *Developmental and Comparative Immunology*, 2020, 108, 10.1016/j.dci.2020.103676 . hal-02516411

HAL Id: hal-02516411

<https://hal.inrae.fr/hal-02516411>

Submitted on 20 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 Partner-specific induction of *Spodoptera frugiperda* immune genes in
2 response to the entomopathogenic nematobacterial complex *Steinernema*
3 *carpocapsae*-*Xenorhabdus nematophila*

4

5 Louise Huot¹, Audrey Bigourdan¹, Sylvie Pagès¹, Jean-Claude Ogier¹, Pierre-Alain
6 Girard¹, Nicolas Nègre^{1,*} and Bernard Duvic^{1,*}

7 ¹ DGIMI, Univ Montpellier, INRAE, Montpellier, France

8

9 * Co-corresponding authors

10 E-mail: nicolas.negre@umontpellier.fr (NN) ; bernard.duvic@umontpellier.fr (BD)

11 Abstract

12 The *Steinernema carpocapsae*-*Xenorhabdus nematophila* association is a nematobacterial complex used
13 in biological control of insect crop pests. The infection success of this dual pathogen strongly depends on
14 its interactions with the host's immune system. Here, we used the lepidopteran pest *Spodoptera*
15 *frugiperda* to analyze the respective impact of each partner in the induction of its immune responses.
16 First, we used previously obtained RNAseq data to construct the immunome of *S. frugiperda* and analyze
17 its induction. We then selected representative genes to study by RT-qPCR their induction kinetics and
18 specificity after independent injections of each partner. We showed that both *X. nematophila* and *S.*
19 *carpocapsae* participate in the induction of stable immune responses to the complex. While *X.*
20 *nematophila* mainly induces genes classically involved in antibacterial responses, *S. carpocapsae* induces
21 lectins and genes involved in melanization and encapsulation. We discuss putative relationships between
22 these differential inductions and the pathogen immunosuppressive strategies.

23 Keywords

24 *Steinernema carpocapsae*, *Xenorhabdus nematophila*, *Spodoptera frugiperda*, nematobacterial complex,
25 immunome, transcriptional response

26 Abbreviations¹

27 1. Introduction

28 The *Steinernema*-*Xenorhabdus* nematobacterial complexes (NBCs) are natural symbiotic associations
29 between nematodes and enterobacteria that are pathogenic for insects. The soil-living nematodes infest
30 insects through the respiratory and/or the intestinal tract (Dowds and Peters, 2002) and reach the
31 hemocoel, the internal body cavity, where they release their intestinal symbionts. The bacteria then grow
32 extracellularly in the hemolymph, the insect equivalent of blood, and improve the nematodes'
33 pathogenicity as well as their ability to reproduce in the host dead body (Forst and Clarke, 2002). Until
34 now, about 90 species of *Steinernema* have been identified, among which several are usable as biological
35 control agents against diverse insect crop pests (Labaude and Griffin, 2018; Lacey et al., 2015). In
36 consequence, their interactions with insects have been extensively studied for about 50 years (Poinar and
37 Grewal, 2012). These studies have shown that in addition to ecological and morphological parameters
38 (Labaude and Griffin, 2018), the NBCs' interactions with the host's immune system is one of the most

¹ NBC: nematobacterial complex; AMP: antimicrobial peptides; NBTA: Nutrient bromothymol blue agar;
PBS: phosphate buffered saline; CFU: Colony-forming unit; GNBP: Gram-negative binding protein;
PGRP: Peptidoglycan recognition protein; LLP: Lysozyme-like protein; PPAE: Prophenoloxidase-
activating enzyme; IMPI: Insect metalloproteinase inhibitor protein; TEP: Thiol-ester protein

crucial factors influencing their ability to infest and kill a given insect (Li et al., 2007; Thurston et al., 1994; Wang et al., 1994).

Insects possess an elaborate immune system, which is able to respond appropriately to diverse types of pathogens and of infections. Firstly, this system involves protective external barriers such as the cuticle, or the peritrophic matrix in the midgut (Kristensen and Chauvin, 2012; Lehane, 1997). It then relies on local defenses of the surface epithelia, which repair efficiently (Ferrandon, 2013; Galko and Krasnow, 2004; Rowley and Ratcliffe, 1978) and produce toxic factors such as antimicrobial peptides (AMPs) (Brey et al., 1993; Tingvall et al., 2001; Tzou et al., 2000; Wu et al., 2010) and reactive oxygen species (Ha et al., 2009). The third line of defense of insects is provided by the hemocytes, which are the circulating immune cells. They can produce diverse types of immune responses, including AMP synthesis, phagocytosis, nodulation, encapsulation, coagulation and melanization (Strand, 2008). Nodulation and encapsulation are cellular immune responses respectively consisting in the engulfment of bacterial aggregates and of large invaders via hemocytes aggregation (Strand, 2008). Together with coagulation, these responses are coupled with a melanization process consisting of series of phenolic compounds oxidations resulting in synthesis of reactive molecules and melanin that participate in pathogens trapping and killing (Jiravanichpaisal et al., 2006; Nappi and Christensen, 2005). Finally, the fat body, a functional equivalent of the mammalian liver, produces potent systemic humoral immune responses involving a massive secretion of AMP cocktails in the hemolymph. These responses can be induced by two major signaling pathways of insect immunity; the Imd pathway, which is mainly activated by Gram negative bacteria, and/or the Toll pathway, which is mainly activated by Gram positive bacteria, fungal organisms and by proteases released by pathogens (Ferrandon et al., 2007; Issa et al., 2018).

The *Steinernema-Xenorhabdus* NBC whose interactions with the immune system have been the most extensively studied is the *S. carpocapsae-X. nematophila* association. These interactions have firstly been studied from the NBC point of view, which allowed the identification of a multitude of immunoevasive and immunosuppressive strategies. For instance, studies in *Rhynchophorus ferrugineus* and *Galleria mellonella* have respectively shown that the cuticle of *S. carpocapsae* is not recognized by the host's immune system (Binda-Rossetti et al., 2016; Mastore et al., 2015) and that the nematode secretes protease inhibitors impairing the coagulation responses (Toubarro et al., 2013a; Toubarro et al., 2013b). Studies in diverse insect models have also shown that both partners produce factors impairing melanization (Balasubramanian et al., 2009; Balasubramanian et al., 2010; Crawford et al., 2012; Eom et al., 2014), hemocyte's viability (Brivio et al., 2018; Kim et al., 2005; Ribeiro et al., 1999; Ribeiro et al., 2003; Vigneux et al., 2007) and the production of cellular immune responses by several ways (Balasubramanian et al., 2009; Balasubramanian et al., 2010; Eom et al., 2014; Park and Kim, 2000; Park and Stanley, 2006; Toubarro et al., 2013b). Finally, both *X. nematophila* and *S. carpocapsae* secrete proteolytic factors

74 degrading cecropin AMPs (Caldas et al., 2002; Gotz et al., 1981) and the bacterium has also been shown
 75 to reduce more globally the hemolymph antimicrobial activity, as well as AMP transcription in
 76 lepidopteran models (Binda-Rossetti et al., 2016; Caldas et al., 2002; Duvic et al., 2012; Ji and Kim,
 77 2004).
 78 On the other hand, the description of these interactions from the hosts' points of view is at its beginning.
 79 This aspect has mainly been studied in the *Drosophila melanogaster* model, with a first transcriptomic
 80 analysis of the whole larva responses to infestations by entire NBCs and by axenic nematodes (Yadav et
 81 al., 2017). This analysis has shown that several immune processes are induced by both pathogens at the
 82 transcriptional level. For instance, the authors found in each case an overexpression of genes related to
 83 the Imd and Toll pathways that was accompanied by the induction of a few AMP genes. They also found
 84 an upregulation of genes related to melanization, coagulation, or involved in the regulation of cellular
 85 immune responses (Yadav et al., 2017). Complementary gene knockout experiments in this model
 86 demonstrated an involvement of the Imd pathway in the response against *X. nematophila* (Aymeric et al.,
 87 2010) and revealed a possible involvement of the Imaginal Disc Growth Factor-2, the intestinal serine
 88 protease Jonah 66Ci (Yadav and Eleftherianos, 2019) as well as TGF- β and JNK pathways members in
 89 the regulation of anti-nematode immunity (Yadav and Eleftherianos, 2018; Yadav et al., 2018).
 90 In order to improve our understanding of the dialogue that takes place between this NBC and its host, we
 91 recently published a topologic transcriptomic analysis of the response of the lepidopteran model
 92 *Spodoptera frugiperda* to the infestation (Huot et al., 2019). This analysis was focused on the three main
 93 immunocompetent tissues that are confronted to the NBC, which are the midgut (the main entry site in
 94 the hemocoel), the hemocytes and the fat body. The RNAseq experiment showed that there was no potent
 95 or well-defined transcriptional response in the midgut. However, we observed dramatic transcriptional
 96 responses in the fat body and the hemocytes at 15 h post-infestation, which is a middle time point of the
 97 infection. In agreement with the results obtained in *D. melanogaster* whole larvae (Yadav et al., 2017),
 98 global analysis of these responses showed they are dominated by immune processes. The objective of the
 99 present study is to go further in the analysis of these induced immune responses. In order to describe them
 100 with high accuracy, we first examine the expression variations of all the immune genes that have been
 101 identified in the insect's genome. We then use tissue RT-qPCR experiments to analyze the temporal
 102 dynamics and the relative contribution of each NBC partner in the identified immune responses. Our
 103 results show that a large number of immune genes are responsive in either one or the two tissues during
 104 the infestation, with activation of antimicrobial and cellular immunities, of melanization, coagulation and
 105 of metalloprotease inhibition. These responses were found to be stable over the time post-infestation and
 106 to consist in combinations of *X. nematophila*-induced and *S. carpocapsae*-induced responses in each
 107 tissue. The *X. nematophila*-induced responses mainly correspond to genes that are classically involved in
 108 antibacterial immunity, whereas the *S. carpocapsae*-induced ones mainly include lectins and genes

109 potentially involved in melanization and encapsulation. In addition, our RT-qPCR experiments show that
110 two previously identified candidate clusters of uncharacterized genes (Huot et al., 2019) also present
111 partner-specific induction profiles. Our hypothesis is that they may correspond to new types of anti-
112 nematode and antibacterial immune factors found in *Spodoptera* genus and lepidopteran species,
113 respectively.

114 **2. Materials and Methods**

115 **2.1. Insect rearing**

116 Corn-strain *Spodoptera frugiperda* (Lepidoptera : Noctuidae) were fed on corn-based artificial diet
117 (Poitout and Buès, 1970). They were reared at 23°C +/- 1°C with a photoperiod of 16 h/8 h (light/dark)
118 and a relative humidity of 40 % +/- 5 %. *Galleria mellonella* (Lepidoptera : Pyralidae) were reared on
119 honey and pollen at 28°C in dark.

120 **2.2. Production and storage of nematobacterial complexes**

121 *Steinernema carpocapsae*-*Xenorhabdus nematophila* complexes (strain SK27 isolated from Plougastel,
122 France) were renewed by infestation of one month-old *Galleria mellonella* larvae. They were collected on
123 White traps (White, 1927) and stored at 8°C in aerated Ringer sterile solution with 0.1 % formaldehyde.
124 The maximal time of storage was limited to 4 weeks to avoid pathogenicity losses.

125 **2.3. Production of axenic nematodes**

126 Gravid *S. carpocapsae* females were extracted from *G. mellonella* dead bodies at day 4 to 6 after
127 infestation by nematobacterial complexes (NBCs). After 5 washing steps in Ringer sterile solution, the
128 females were surface-sterilized by 20 min incubation in 0.48 % (wt/vol) sodium hypochlorite and 3 h
129 incubation in Ringer sterile solution supplemented with antibiotics (150 µg/mL polymyxin, 50 µg/mL
130 colistin, 50 µg/mL nalidixic acid). The eggs were extracted by female crushing with sterile glass pestles
131 and then washed by centrifugation (2 min, 16000 g) in Ringer sterile solution, disinfected by incubation
132 in 0.48 % sodium hypochlorite for 5 min, and washed again twice. After microscopic observation, the
133 intact eggs were placed on liver-agar (40 g/L Tryptycase Soja Agar [BioMérieux], 5 g/L Yeast Extract
134 [Difco], 100 g/L porc liver) plates supplemented with antibiotics (150 µg/mL polymyxin, 50 µg/mL
135 colistin and 50 µg/mL nalidixic acid). The plates were maintained inside a dark humid chamber for 1
136 month to allow nematodes development. The nematodes were then suspended in Ringer sterile solution
137 and infective juvenile stages (IJs) were sorted by pipetting under a microscope (Leica). The IJs were
138 rinsed twice by centrifugation (2 min, 3000 g) in 1 mL Ringer sterile solution and used within minutes for
139 experimental infection.

Nematodes' axenicity was verified by DNA extraction and PCR amplification. Nematodes were suspended in 200 μ L milliQ water supplemented with 200 μ L glass beads ($\varnothing \leq 106 \mu\text{m}$) (Sigma). They were ground for 2 x 40 sec at 4.5 ms speed with a FastPrep homogenizer (MP Biomedicals). The debris were discarded by centrifugation (2 min, 16000 g) and 150 μ L supernatant were mixed with 200 μ L lysis buffer (Quick extract kit, Epi-centre) for a second grinding. To ensure bacterial cell lysis, the samples were incubated at room temperature for 48 h with 2 μ L Ready-Lyse Lysozyme solution at 30000 U/ μ L (Epi-centre). Protein denaturation was then performed by 10 min incubation at 90°C, and RNA was removed by 10 min incubation at 37°C with 20 μ L RNase A (20 mg/mL) (Invitrogen). DNA was extracted by successive addition of 500 μ L phenol-chloroform-isoamyl alcohol and 500 μ L chloroform, followed by centrifugations (10 min, 16000 g) and aqueous phase collections. DNA was precipitated with 500 μ L 100% ethanol supplemented with 20 μ L sodium acetate and by freezing at -80°C for 2 h. After defrosting, DNA was concentrated by centrifugation (30 min, 16000 g) and the precipitates were washed twice by centrifugation (15 min, 16000 g) in 500 μ L 70% ethanol. DNA was finally suspended in 50 μ L sterile milliQ water and left at room temperature for a few hours to ensure precipitate dissolution. After DNA quantification with a Qubit fluorometer (Invitrogen), *X. nematophila* presence was assessed by PCR amplification with *Xenorhabdus*-specific primers (Xeno_F: 5'-ATG GCG CCA ATA ACC GCA ACT A-3'; Xeno_R: 5'-TGG TTT CCA CTT TGG TAT TGA TGC C-3'), which target a region of the XNC1_0073 gene encoding a putative TonB-dependent heme-receptor (Cambon et al., 2020). The presence of other bacteria was assessed by 16S rRNA gene amplification with universal primers (Tailliez et al., 2006). Thirty cycles of PCR were performed using Taq polymerase (Invitrogen) in a Biorad thermocycler, with hybridization temperatures of 55°C and 50°C respectively. PCR products were then analyzed by agarose gel electrophoresis with DNAs from *X. nematophila*, *Pseudomonas protegens* and *S. carpocapsae*-*X. nematophila* complexes as controls (Supplementary Fig. 1).

2.4. Experimental infections

Experimental infestations with NBCs were carried out on individual 2nd day 6th instar *S. frugiperda* larvae according to Huot et al. (2019). Briefly, 12-well plates were coated with pieces of filter paper (Whatman) and larvae were individually introduced in each well with a cube of artificial diet (Poitout and Buès, 1970). For infestations, 150 +/- 20 NBCs in 150 μ L Ringer sterile solution were poured in each well and the plates were then maintained at 23°C. 150 μ L Ringer sterile solution were used for control experiments.

For intra-hemocoelic injection experiments, pathogens were injected in larvae's abdomens after local application of 70% ethanol with a paintbrush. Injections were performed using a syringe pump (Delta labo) with 1 mL syringes (Terumo) and 25G needles (Terumo). The injected larvae were then kept in 12-well plates at 23 °C with cubes of artificial diet (Poitout and Buès, 1970).

X. nematophila suspensions were prepared as described in Sicard et al. (2004). Bacterial culture was diluted in PBS and 20 µL containing 200 +/- 50 bacterial cells were injected in the hemocoel at a rate of 1.67 mL/min. 20 µL sterile PBS were used for control larvae. The purity and number of injected *X. nematophila* were verified by plating 20 µL of the bacterial suspension on NBTA (Boemare et al., 1997).

For NBC and axenic nematode injections, 10 +/- 3 nematodes in 20 µL 70% Ringer and 30% glycerol solution were injected at a rate of 2.23 mL/min. Syringes were frequently renewed in order to limit nematodes concentration and sedimentation and the number of injected nematodes was verified by 10 simulations of injection in Petri dishes followed by nematode counting under a stereomicroscope (Zeiss). Sterile 70% Ringer and 30% glycerol solutions were used for control larvae. To avoid accidental *per os* infections, the injected larvae were then briefly washed in sterile PBS and dried on paper towel before being placed in 12-well plates.

The pathogens efficacies were checked by monitoring the survival of 12 control and 12 infected larvae for 72 h after infestation or after injection.

2.5. Production and storage of bacterial symbionts

X. nematophila strain F1 isolated from NBCs strain SK27 was conserved at -80°C. Within 3 weeks before each experiment, they were grown for 48 h at 28°C on NBTA supplemented with erythromycin (15 µg/mL) to which the strain is resistant (Sicard et al., 2004). The colonies were then conserved at 15°C and used for overnight culture at 28°C in 5 mL Luria-Bertani broth (LB) before experiments.

2.6. Caterpillar RNA extraction

RNAs were prepared as described in Huot et al. (2019). Briefly, nine larvae per technical replicate were bled in anti-coagulant buffer (van Sambeek and Wiesner, 1999). Hemocytes were recovered by centrifugation (1 min, 800 g) at 4°C and the pellet was immediately flash-frozen with liquid nitrogen. The larvae were then dissected for fat body and midgut sampling and the tissues were flash-frozen in eppendorf tubes with liquid nitrogen. After storage at -80°C for at least 24 h, 1 mL Trizol (Life technologies) was added to the pooled tissues. The tissues were then ground by using a TissueLyzer 85210 Rotator (Qiagen) with one stainless steel bead (Ø : 3 mm) at 30 Hz for 3 min. For optimal cell lyses, ground tissues were left at room temperature for 5 min. To extract nucleic acids, 200 µL chloroform (Interchim) were added and the preparations were left at room temperature for 2 min with frequent vortex homogenization. After centrifugation (15 min, 15,000 g) at 4°C, the aqueous phases were transferred in new tubes and 400 µL 70% ethanol were added. RNA purifications were immediately performed with the RNeasy mini kit (Qiagen) and contaminant DNA was removed with the Turbo DNA-freeTM kit (Life Technologies).

RNA yield and preparation purity were analyzed by measuring the ratios A_{260}/A_{280} and A_{260}/A_{230} with a Nanodrop 2000 spectrophotometer (Thermo Scientific). RNA integrity was verified by agarose gel electrophoresis and RNA preparations were conserved at - 80 °C.

2.7. RNAseq experiments

RNAseq raw data originate from Huot et al. (2019). In brief, during 3 independent experiments, 9 larvae infested with 150 *S. carpocapsae-X. nematophila* complexes and 9 uninfected control larvae were bled and dissected at time 15 hours post-contact with NBCs or Ringer sterile solution (see Materials and Methods 2.4.). Total RNA from their hemocytes and fat bodies were extracted (see Materials and Methods 2.6.) and the corresponding libraries were prepared by MGX GenomiX (IGF, Montpellier, France) with the TruSeq Stranded mRNA sample preparation kit (Illumina). The libraries were then validated on Fragment Analyzer with a Standard Sensitivity NGS kit (Advanced Analytical Technologies, Inc) and quantified by qPCR with a Light Cycler 480 thermal cycler (Roche Molecular diagnostics). cDNAs were then multiplexed by 6 and sequenced on 50 base pairs in a HiSeq 2500 system (Illumina) with a single-end protocol. Image analysis and base calling were performed with the HiSeq Control and the RTA software (Illumina). After demultiplexing, the sequences quality and the absence of contaminant were checked with the FastQC and the FastQ Screen softwares. Data were then submitted to a Purity Filter (Illumina) to remove overlapping clusters.

For each sample, the reads were pseudoaligned on the *S. frugiperda* reference transcriptome version OGS2.2 (Gouin et al., 2017) using the Kallisto software (Bray et al., 2016). Differential expression between infested and control conditions were then assessed for each time point and tissue with the Sleuth software (Pimentel et al., 2017). Wald tests were used with a q-value (equivalent of the adjusted p-value) threshold of 0.01 and a beta value (biased equivalent of the log2 fold change) threshold of 1. Only transcripts with normalized counts over 5 in all three replicates of the infested and/or of the control condition were considered as reliably differentially expressed.

Previously annotated immune transcripts (Gouin et al., 2017) were then checked for significant expression changes and not annotated differentially expressed ones were researched with the Blast2GO software by blastx on the NCBI nr and drosophila databases (Conesa et al., 2005). To avoid mistakes related to genome fragmentation, the immune transcripts were gathered by unique gene after careful examination of their sequences and of the available genomic data (Gouin et al., 2017). The induction levels of the transcripts were then averaged by unique gene before graphical representation of the results.

2.8. RT-qPCR experiments

RT-qPCR experiments were performed with extracted total RNA (see Materials and Methods 2.6.) from hemocytes and fat bodies of 9 larvae infested or infected by injection and of 9 uninfected control larvae during 3 independent experiments.

For each sample, cDNAs were synthesized from 1 µg of RNA with the SuperScript II Reverse Transcriptase (Invitrogen), according to the manufacturer's protocol. The primers (Supplementary Table 1) were designed with the Primer3Web tool (Untergasser et al., 2012). Their efficiency was estimated by using serial dilutions of pooled cDNA samples and their specificity was verified with melting curves analyses. Amplification and melting curves were analyzed with the LightCycler 480 software (Roche Molecular diagnostics). RT-qPCRs were carried out in technical triplicate, with the LightCycler 480 SYBR Green I Master kit (Roche). For each replicate and primer pair, 1.25 µL of sample containing 50 ng/µL of cDNA and 1.75 µL of Master mix containing 0.85 µM of primers were distributed in multiwell plates by an Echo 525 liquid handler (Labcyte). The amplification reactions were then performed in a LightCycler 480 thermal cycler (Roche) with an enzyme activation step of 15 min at 95°C, and 45 cycles of denaturation at 95°C for 5 sec, hybridization at 60°C for 10 sec and elongation at 72°C for 15 sec. Crossing points were determined using the Second Derivative Maximum method with the LightCycler 480 software (Roche) and relative expression ratios between control and infected conditions were manually calculated according to the method of Ganger et al. (2017). The ratios were normalized to the RpL32 housekeeping gene relative levels and the EF1 gene was used as an internal control. Statistical analyses of the data were all performed with the R software (R Core Team, 2017). Differential expression significance between the control and infected conditions was assessed by paired one-tailed t-tests on ΔCq values. Multiple comparisons of fold changes were assessed by one-way ANOVA on $\Delta\Delta Cq$ values followed by post hoc Tukey tests. P-values under 0.05 were considered as significant for all the above tests. The gplots package was used to draw the heatmaps and the clusters were built from a dissimilarity matrix based on Pearson correlation coefficients.

2.9. Quantification of nematodes in the midgut lumen

NBCs in the midgut lumen were quantified at several times after infestation by nematode counting in the alimentary bolus. For 3 independent experiments, 3 infested larvae were dissected and the midguts' alimentary bolus were extracted. Each alimentary bolus was then dissolved in 3 mL sterile PBS in a Petri dish (Ø : 35 mm) and motile nematodes were counted with a stereomicroscope (Leica).

2.10. Quantification of *X. nematophila* in the hemolymph

The concentration of *X. nematophila* strain F1 in the hemolymph was estimated by CFU counting on selective medium. For 3 independent infestation or infection by injection experiments and 3 technical replicates, hemolymph was collected by bleeding of 3 caterpillars in 200 µL PBS supplemented with phenylthiourea (Sigma). The volumes of hemolymph were then estimated by pipetting and serial dilutions of the samples were plated on NBTA with 15 µg/mL erythromycin. CFU were counted after 48 h

incubation at 28°C and the counts were reported to the estimated hemolymph volumes in order to calculate the bacterial concentrations. Hemolymph of naive larvae was also plated for control.

2.11. Insect survival kinetics

Survival kinetics were measured by 3 independent experiments on pools of 20 larvae infested with NBCs or injected with NBCs, with axenic *S. carpocapsae* nematodes or with *X. nematophila* symbionts. Survival was monitored from 0 to 72 hours after contact or injection.

2.12. Parasitic success measurement

Parasitic success was measured during 3 independent experiments on pools of 20 NBCs or axenic nematodes-injected larvae. Dead larvae were individually placed on white traps (White, 1927) approximately 2 days after their deaths. The emergence of nematodes was assessed at day 40 after injection by observation of the collection liquid with a stereomicroscope (Leica). Parasitic success was then calculated as the percentage of larvae with nematode emergence among the infected larvae.

3. Results & Discussion

3.1. Hemocytes' and fat body's immune responses

In order to get an accurate picture of the *S. frugiperda* transcriptional immune responses to the NBC infestation, we first used a previously published list of immune genes identified by sequence homology in the *S. frugiperda* genome (Gouin et al., 2017). We then used previously obtained RNAseq data (Huot et al., 2019) to look at their expression variations in the fat body and in the hemocytes (Supplementary Table 2A) and we completed the repertoire with additional putative immune genes that we directly identified from lists of differentially expressed genes upon NBC infestation (Supplementary Table 2B). In total, we present the annotation of 226 immune or putative immune genes of which 132 were significantly modulated at 15 h post-infestation (hpi) (Sleuth, p-value < 0.01; |Beta| > 1; all count values > 5 in at least one condition) in one or both tissues (Fig. 1). Among them, 62 were involved in antimicrobial responses (Fig. 1A), 18 were related to melanization (Fig. 1B), 23 were involved in cellular responses (Fig. 1C) and the 29 remaining genes were grouped in a category called “diverse” due to pleiotropic or poorly characterized functions (Fig. 1D).

3.1.1. Antimicrobial responses

In the antimicrobial response category, 58 genes were found to be upregulated in at least one of the two tissues (Fig. 1A). The signaling genes encoded 3 and 8 members of the Imd and Toll pathways, respectively, as well as 5 short catalytic peptidoglycan recognition proteins (PGRP-S), which are probably involved in the regulation of these pathways by peptidoglycan degradation (Myllymaki et al., 2014; Valanne et al., 2011) (Fig. 1A). Four other genes were considered as involved in recognition. They

305 encoded Gram negative binding proteins (GNBPs), which have been reported to recognize
 306 peptidoglycans or β -glucans and participate in the further activation of the Toll pathway (Ferrandon et al.,
 307 2007) and/or of the melanization response (Nakhleh et al., 2017) (Fig. 1A). Finally, the effector genes
 308 encoded 33 antimicrobial peptides (AMPs) belonging to all the *S. frugiperda*'s AMP families (Gouin et
 309 al., 2017) plus 4 lysozymes and lysozyme-like proteins (LLPs) (Fig. 1A). Depending on their families and
 310 on the insect species, AMPs can present varied activity spectra, ranging from antiviral or antibacterial
 311 activities to anti-fungal and anti-parasitic ones (Yi et al., 2014). Varied activity spectra have also been
 312 found for several insects' lysozymes and LLPs (Chen et al., 2018; Gandhe et al., 2007; Satyavathi et al.,
 313 2018; Sowa-Jasilek et al., 2014; Yu et al., 2002). Interestingly, all of the categories and subcategories
 314 cited above were represented in the two tissues, indicating that their antimicrobial responses are
 315 diversified and that the factors responsible for their disappearance in the hemolymph (Binda-Rossetti et
 316 al., 2016; Duvic et al., 2012) probably act at a post-transcriptional level. About a half of the genes
 317 presented similar and significant induction profiles in the hemocytes and in the fat body. This is for
 318 instance the case of the usually anti-Gram negative bacteria attacin, cecropin and gloverin AMPs (Yi et
 319 al., 2014), which were all highly induced in the two tissues (Fig. 1A), suggesting they both respond to the
 320 bacterial partner *X. nematophila*. On the other hand, all the induced GGBP, lysozyme and LLP genes
 321 were found to be either significantly induced in the hemocytes or in the fat body, and in the AMP
 322 category, tissue-specificities were observed for diapausin, defensin-like and most moricin genes (Fig.
 323 1A).
 324 Only 8 antimicrobial response genes were found to be significantly downregulated (Fig. 1A).
 325 Interestingly, 4 were involved in the Imd pathway whereas the 4 remaining ones were dispersed between
 326 the AMP, GGBP and lysozyme categories (Fig. 1A). The Imd pathway downregulated factors included
 327 *sickie* and the *akirin* in the hemocytes and *SMARCC2* and *BAP60* in the fat body (Fig. 1A). In *D.*
 328 *melanogaster*, Sickie participates in the activation of Relish, the transcription factor of the Imd pathway
 329 (Foley and O'Farrell, 2004) and the akirin acts together with the Brahma chromatin-remodeling complex,
 330 containing BAP60 and SMARCC2, as cofactor of Relish to induce the expression of AMP genes (Bonnay
 331 et al., 2014). Only few data can be found in current literature on the transcriptional regulation of these
 332 factors (Bonnay et al., 2014; Cao et al. 2015; Chen et al., 2018). It is thus difficult to determine whether
 333 the downregulations we observed correspond to a normal negative feedback of the humoral immune
 334 response or to an immunosuppressive effect of the NBC. In a previous study on the *Spodoptera exigua*
 335 model, Hwang et al. (2013) have shown injection of live *X. nematophila* induces lower transcriptional
 336 inductions of several AMP genes than injection of dead bacteria (Hwang et al., 2013). It would be
 337 interesting to determine, either in the *S. exigua* or in the *S. frugiperda* model, whether reduction of AMP
 338 induction in response to live bacteria co-occurs with a downregulation of the Relish cofactors and
 339 activators we identified in the present analysis.

340 To summarize, the antimicrobial responses are potent and diversified in the two tissues, with a common
341 induction of genes that probably respond to *X. nematophila*. Yet unexplained tissue-specific responses
342 were observed and the results show a down-regulation of Imd pathway members that could be related to a
343 previously described transcriptional immunosuppressive effect of the NBC. However, since at 15hpi, the
344 full repertoire of *S. frugiperda* AMP is still expressed in response to naturally infesting NBC (Fig. 1A),
345 this effect might not be potent enough to suppress the humoral response at this time point, suggesting that
346 the NBC probably uses additional resistance and/or immunosuppressive strategies to survive the insect's
347 humoral immune response.

348 **3.1.2. Melanization**

349 In the melanization category, 16 genes were found to be upregulated in at least one of the two tissues
350 (Fig. 1B). These genes firstly encoded 6 serine proteases (Fig. 1B) that were considered as members of
351 the prophenoloxidase (proPO) system. The proPO system is an extracellular proteolytic cascade ending in
352 the maturation of the proPO zymogen into PO, which initiates the melanization process (Nakhleh et al.,
353 2017). Among the upregulated serine proteases, PPAE2 is the only one that is known to take part in
354 proPO processing whereas the other proteases were included in this category because of their
355 characteristic CLIP domains and of their low homology with the serine proteases acting upstream of the
356 Toll pathway in *D. melanogaster* (Veillard et al., 2016). The other upregulated genes in this category
357 included 3 serpins, which are known to regulate the proPO system in several model insects (Nakhleh et
358 al., 2017), 3 melanization enzymes, DDC, Yellow-like 1 and Punch-like (De Gregorio et al., 2001; Tang,
359 2009) as well as 4 genes, Reeler-1 and 3 Hdd23 homologs, that are involved in melanization and nodule
360 formation in other models (Bao et al., 2011; Qiao et al., 2014) (Fig. 1B). Despite of tissue-specific
361 induction patterns, serine proteases and serpins were found in the two tissues (Fig. 1B), suggesting that
362 both participate in the stimulation of the proPO system, which is consistent with results obtained in other
363 interaction models (Yuan et al., 2017; Zou et al., 2015; Zou et al., 2010). However, with the exception of
364 the DDC, all the melanization enzymes as well as the nodulation-related genes were specifically induced
365 in the hemocytes (Fig. 1B), which is consistent with the very localized nature of this immune response
366 (Tang, 2009) that is mainly mediated by hemocyte subtypes.

367 Finally, only 2 genes, PPAE1 and Yellow-like 2, were found to be significantly down-regulated in this
368 category (Fig. 1B). Both were specifically repressed in the hemocytes, which could be due to functional
369 interferences with their upregulated homologs (PPAE2 and Yellow-like 1) and/or to an expression in
370 specific hemocyte subtype(s) whose proportion in the total hemocyte population would influence the
371 observed log2 fold changes. There is also the possibility of an immunosuppressive effect of the NBC on
372 the expression of these genes.

373 In summary, our results suggest that both the hemocytes and the fat body participate in induction and
374 regulation of melanization in response to the NBC and no clear sign of immunosuppression at the

transcriptional level is detected for this response. These results are in agreement with the previous identification of diverse inhibitors of PO activity in both *S. carpocapsae* (Balasubramanian et al., 2009; Balasubramanian et al., 2010) and *X. nematophila* (Crawford et al., 2012; Eom et al., 2014) that act at the post-transcriptional level.

3.1.3. Cellular responses

In the hemocytes, 19 upregulated genes were placed in the cellular responses category (Fig. 1C). The signaling ones encoded 3 homologs of the transcription factor Krüppel (Kr) (Fig. 1C). In *D. melanogaster*, Kr and Kr homologs are involved in several developmental processes such as embryo patterning (Schmucker et al., 1992), organogenesis (Fichelson et al., 2012; Harbecke and Janning, 1989; Hoch and Jackle, 1998), and cell differentiation (Ivy et al., 2015). More specifically in the hemocytes, Kr has been shown to take part in hemocytes' differentiation and/or activation (Stofanko et al., 2008), a crucial step for the induction of cellular immune responses. The recognition genes encoded 3 cellular receptors of the Scavenger (SR) and Integrin families plus the hemolin, a secreted immunoglobulin-containing protein (Fig. 1C). Both Scavenger receptors and integrins are known to act as membrane receptors in phagocytosis of bacteria and apoptotic cells (Nazario-Toole and Wu, 2017). In addition, integrins are involved in diverse processes, including cell motility and adhesion, and encapsulation (Levin et al., 2005; Melcarne et al., 2019). The hemolin is known to act as an opsonin by increasing phagocytosis and nodulation of bacteria in *Manduca sexta* (Eleftherianos et al., 2007). Among the effector genes, we first identified 5 upregulated genes corresponding to conserved intracellular phagocytosis-related proteins. They included Ced-6, the Rabenosyn-5 (Rbsn-5-like), a V-ATPase subunit (ATP6V0A2-like) and 2 small GTPase Activating Proteins (Rabex-5-like, CdGAPr-like) (Nazario-Toole and Wu, 2017) (Fig. 1C). We also found genes encoding membrane proteins, such as the immunoglobulin-containing hemicentin (HMCN-like) (Barat-Houari et al., 2006) and 4 tetraspanin-like (Tsp-like) proteins (Hemler, 2008) (Fig. 1C), that could participate in cell-cell adhesion and cellular immune responses. Interestingly, one of the upregulated tetraspanins (Tsp-like 3) presented 79.5% identity with the *Manduca sexta* (Lepidoptera : Noctuidae) tetraspanin D76, which takes part in hemocytes aggregation during capsule formation by trans-interacting with a specific integrin (Zhuang et al., 2007). Finally, 2 genes encoding proteins similar to the *D. melanogaster* clotting factors GP150 (Korayem et al., 2004) and a transglutaminase (Tg-like) (Lindgren et al., 2008) were also found upregulated (Fig. 1C). Only 2 genes (Ced-6-like, Rbsn-5-like) of the cellular responses category were found to be upregulated in the fat body (Fig. 1C) and both encoded intracellular proteins that are probably not related to immunity in this tissue. All the 4 down-regulated putative cellular immunity-related genes were specifically modulated in the hemocytes (Fig. 1C). They encoded 2 Rho GTPase Activating Proteins (RhoGAP-like), a scavenger receptor similar to the *D. melanogaster* Croquemort receptor (SR-B3) and a homolog of the *D. melanogaster* integrin α -PS1. In *D. melanogaster*, Croquemort has been shown to take part in

phagocytosis of apoptotic cells and of the Gram positive bacterium *Staphylococcus aureus* but not of the Gram negative bacterium *Escherichia coli* (Melcarne et al., 2019; Stuart et al., 2005). Integrin α -PS1 is a ligand of the extracellular matrix protein laminin (Gotwals et al., 1994). It is involved in migration and differentiation of several cell types during development (Delon and Brown, 2009; Roote and Zusman, 1996; Urbano et al., 2011) but does not seem to be required for any immune process. Their down-regulations are thus probably due to their uselessness in the context of the response to the NBC. Overall, the results suggest that all types of cellular responses are transcriptionally induced at 15 hpi, including phagocytosis and nodulation, as well as encapsulation that would be adapted to the bacterial partner or the nematode, respectively. In addition, the induction of coagulation responses is particularly interesting, since many clotting factors participate in *D. melanogaster* resistance to infestation by another type of NBC, the *Heterorhabditis bacteriophora*-*Photorhabdus luminescens* association (Arefin et al., 2014; Hyrsl et al., 2011; Kucerova et al., 2016; Wang et al., 2010). Moreover, despite *S. carpocapsae* does not pierce the insects' cuticles as *H. bacteriophora* (Dowds and Peters, 2002), it has been shown to express at least two secreted proteases with inhibitory activities towards the formation of clot fibers and coagulation-associated pathogen trapping (Toubarro et al., 2013a; Toubarro et al., 2013b). Once again, the induction of such immune responses is consistent with the previous identification of several virulence factors of the NBC targeting cellular immunity (Balasubramanian et al., 2009; Balasubramanian et al., 2010; Brivio et al., 2018; Eom et al., 2014; Kim et al., 2005; Park and Kim, 2000; Park and Stanley, 2006; Ribeiro et al., 1999; Ribeiro et al., 2003; Toubarro et al., 2013a; Vigneux et al., 2007).

3.1.4. Diverse immunity-related genes

A total of 29 modulated genes were involved in other diverse immune processes. They included 10 up- or down-regulated signaling genes, 7 upregulated recognition genes, 8 upregulated effector genes and 5 upregulated genes of unknown functions that are known to be modulated after immune challenge (Fig. 1D).

The signaling genes firstly encoded 2 insulin-like growth factor (IGF-II-like) and 2 insulin receptor substrate homologs (IRS1-like) (Fig. 1D). Insulin signaling is known to have a deleterious impact on the induction of systemic immune responses in the fat body of *D. melanogaster* (Lee and Lee, 2018) whereas insulin increases hemocyte proliferation in the hemolymph of mosquitoes (Castillo et al., 2011) as well as in the hematopoietic organs of the lepidopteran model *Bombyx mori* (Nakahara et al., 2006). In agreement with these assertions, we found that 2 of these genes were down-regulated in the fat body, but all 4 genes were upregulated in the hemocytes (Fig. 1D). Two other signaling genes were found to be specifically overexpressed in the hemocytes. The first one is a homolog of the *Litopenaeus vannamei* (Decapoda: Penaeidae) leucine-rich repeat flightless-I-interacting protein 2 (LRRFIP2-like) (Fig. 1D), which has been shown to upregulate AMP expression in *L. vannamei* as well as in *D. melanogaster* (Zhang et al., 2013). On the other hand, 3 signaling genes were found to be strictly down-regulated (Fig. 1D). Interestingly,

these genes included a member of the TGF- β pathway (BAMBI-like) in the hemocytes and a member of the JNK pathway in the fat body (Basket), two pleiotropic pathways that are currently suspected to take a part in the *D. melanogaster* immune response to nematodes after NBC infestation (Eleftherianos et al., 2016; Ozakman and Eleftherianos, 2019; Patrnogic et al., 2018; Yadav et al., 2018). The third down-regulated gene was found in the fat body and encoded MASK, an inducer of the Jak/Stat pathway (Fisher et al., 2018). In the fat body, the Jak-Stat pathway has mainly been shown to induce the expression of cytokines (Pastor-Pareja et al., 2008) and of a putative opsonin belonging to the TEP family (Lagueux et al., 2000). Remarkably, several TEP genes have been shown to participate in antibacterial immunity after NBC infestation in *D. melanogaster* (Arefin et al., 2014; Shokal and Eleftherianos, 2017; Shokal et al., 2017; Shokal et al., 2018). All of these down-regulations could thus impair the insect's immune response to the NBC. However, more detailed analyses of their functions and modulations would be required to hypothesize immunosuppressive effects of the NBCs.

All 7 upregulated recognition genes encoded lectins (Fig. 1D). Five of them encoded C-type lectins (CLECT), which are known to be involved in binding of diverse pathogens (Xia et al., 2018), including bacteria and nematodes (Yu and Kanost, 2004). This binding can then stimulate several immune responses, such as bacterial aggregation, melanization, phagocytosis, nodulation and encapsulation (Xia et al., 2018). The 2 others encoded galectins, which are involved in diverse aspects of mammalian immunity, including pathogens binding (Baum et al., 2014), and are considered as relevant candidate immune proteins in insects (Pace and Baum, 2002). Despite a larger set of upregulated lectins was identified in the fat body, members of these protein families were found upregulated in the two tissues.

In the hemocytes, the upregulated effector genes firstly encoded a homolog of the superoxide dismutase (SOD-like), a conserved detoxifying enzyme involved in responses to reactive oxygen species (Wang et al., 2018) (Fig. 1D). The 7 remaining genes encoded proteins with similarity to insect metalloproteinase inhibitors (IMPI-like) (Fig. 1D), whose functions have only been studied in the lepidopteran model *Galleria mellonella*. The only characterized IMPI encodes two proteins of which one is probably involved in the regulation of extracellular matrix remodeling and the second specifically targets metalloproteinases from pathogens (Wedde et al., 1998; Wedde et al., 2007). *S. carpocaspae* and *X. nematophila* both express several secreted serine proteases as well as metalloproteinases during the infectious process (Caldas et al., 2002; Chang et al., 2019; Dillman et al., 2015; Hao et al., 2010; Jing et al., 2010; Lu et al., 2017; Massaoud et al., 2011). The induction of such immune responses could interfere with some of these proteinases to impair the NBC's virulence and/or survival. Interestingly, all but one of these IMPI homologs were found to be specifically upregulated in the hemocytes, a tissue-specificity that had not been highlighted in previous reports (Griesch et al., 2000; Vertyporokh and Wojda, 2017).

Finally, the remaining genes of unknown function encoded Spod-x-tox, a protein without antimicrobial activity which contains tandem repeats of defensin-like motifs (Destoumieux-Garzon et al., 2009), 3

480 REPAT genes, which are known to be induced in the midgut after exposure to toxins, viruses and
481 intestinal microbiota perturbations in the close species *S. exigua* (Herrero et al., 2007; Navarro-Cerrillo et
482 al., 2012; Navarro-Cerrillo et al., 2013), and Hdd1, which is induced in response to bacteria and
483 peptidoglycan in the lepidopteran models *Hyphantria cunea* and *Bombyx mori* (Shin et al., 1998; Zhang
484 et al., 2017) (Fig. 1D).

485 In summary, we found an important additional mobilization of several relevant candidate immune genes,
486 including mainly insulin signaling factors and IMPIs in the hemocytes and lectins in the fat body. In
487 addition, these results suggest that the candidate immune pathways TGF- β , JNK and Jak/Stat could be
488 down-regulated. Such down-regulations are in disagreement with the results of Yadav and colleagues
489 (Yadav et al., 2017) in *D. melanogaster* and thus would require further investigation.

490 **3.2. Temporal analysis of the induced immune responses**

491 In order to put the *S. frugiperda* immune responses in relation with the infectious process, we then
492 described their temporal dynamics in each analyzed immunocompetent tissue. To this aim, we monitored
493 with RT-qPCR experiments the induction levels of selected representative immune genes from 5 hpi, the
494 mean time at which nematodes release *X. nematophila* in the hemocoel, to 20 hpi, which is about 9 hours
495 before the first insect deaths (Supplementary Fig. 2).

496 In the hemocytes, the selected genes included 15 genes of the antimicrobial response, 2 genes involved in
497 melanization, 5 cellular response genes, 2 lectins and one IMPI-like gene. At 5 hpi, only 2 genes,
498 encoding a leucocin antibacterial (Yi et al., 2014) AMP (Lebocin 2) and the negative regulator Pirk of the
499 Imd pathway (Kleino et al., 2008), were found to be significantly upregulated. However, most of the
500 selected genes that are strongly induced at later time points also presented positive log2 fold changes at
501 this time point (Fig. 2A). From 10 to 20 hpi, all selected genes but few exceptions (*cecropin D*, *Tg-like*
502 and *Integrin β -like*) due to biological variability were significantly upregulated at each time point (Fig.
503 2A). Clustering analyses based on Pearson coefficients however revealed 3 distinct clusters of
504 covariations. The first one contained 13 genes belonging to all the categories cited above and
505 corresponded to very stable induction patterns (Fig. 2A). The second one, which contained 8 genes
506 involved antimicrobial and cellular responses plus the selected C-type lectin (*CLECT (ccBV)*),
507 corresponded to slightly increasing patterns (Fig. 2A). Finally, the third one, which contained the Relish
508 and Pelle members of the Imd and Toll pathways (Ferrandon et al., 2007), an integrin and the DDC
509 melanization enzyme (Huang et al., 2005) genes, corresponded to slightly decreasing patterns (Fig. 2A).

510 In the fat body, the selected genes included 15 genes of the antimicrobial response, 2 genes involved in
511 melanization, one galectin gene (*Galectin 1*) and an IMPI-like gene (*IMPI-like 3*). At 5 hpi, all 7 selected
512 AMPs, *PGRP-S1* and *Galectin 1* were found to be upregulated (Fig. 2B). All these genes were among the
513 most strongly overexpressed at later time points. Such as in the hemocytes, most of the selected genes
514 were then significantly upregulated from 10 to 20 hpi (Fig. 2B). In this tissue, the genes only subdivided

515 into two main covariation clusters: a cluster of genes with stable induction patterns and a cluster of genes
516 with increasing induction patterns. The first cluster contained 10 genes of which 8 were involved in
517 antimicrobial responses, one encoded a melanization-related serine protease (Snake-like 2) and one
518 encoded the Galectin 1 (Fig. 2B). The second cluster contained 9 genes, of which 7 were involved in
519 antimicrobial responses, one encoded the DDC melanization enzyme (Huang et al., 2005) and the last one
520 encoded the IMPI-like 3 (Fig. 2B).

521 Altogether, the results obtained for the two tissues show that most of the transcriptional immune
522 responses induced at 15 hpi take place between 0 and 10 hpi, which is comparable to timings observed in
523 other interaction models (Boutros et al., 2002; Erler et al., 2011; Lemaitre et al., 1997). The results also
524 indicate that these responses are globally stable across the time post-infestation despite some distinct gene
525 induction patterns in each category of response. Interestingly, while we were hoping to discriminate
526 between an early response, probably activated by the nematode presence, and a later response, probably
527 reacting to bacterial growth, we did not find any clear link between the gene inductions' dynamics and
528 the different immune processes and pathways that were represented in our selection.

529 **3.3. Evaluation of each NBC partner's part in the induced immune responses**

530 In order to identify each NBC partner's relative participation in the fat body's and hemocytes' immune
531 responses, we used RT-qPCR to compare the induction levels of the selected immune genes after
532 independent infections by the whole NBC, the axenic nematode or the bacterial symbiont. To this aim, we
533 decided to use a more standardized protocol of direct injection of the pathogens into the hemocoel,
534 thereby limiting putative side effects such as early hemocoel colonization by intestinal microorganisms.
535 Importantly, we previously compared the kinetics of *X. nematophila* growth and of *S. frugiperda* survival
536 after injection of the entire NBC and of 200 *X. nematophila* (Supplementary Fig. 3A,B). This comparison
537 showed that both kinetics are very similar and thus that any difference of induction level between the 2
538 conditions would not reflect differences in bacterial load or physiological state. However, the putative
539 impact of axenization on the nematode's physiology could not be assessed by the same way due to
540 technical limitations and to its avirulence in absence of its bacterial symbiont (Supplementary Fig. 3B,C).
541 In the hemocytes, 14 genes presented higher induction levels in response to *X. nematophila* than in
542 response to the axenic nematode (Fig. 3). In the antimicrobial category, they included the negative
543 regulator *Pirk* of the Imd pathway (Kleino et al., 2008), all the selected attacin, cecropin, gloverin,
544 lebocin and gallerimycin AMP genes, the 2 selected PGRP-S genes, and also probably the Imd pathway
545 transcription factor *Relish* despite non-significant statistics (Ferrandon et al., 2007) (Fig. 3A). As
546 indicated above, the Imd pathway, as well as the attacin, cecropin and gloverin AMP families, are known
547 to take part in anti-Gram negative bacteria immune responses (Ferrandon, 2013; Yi et al., 2014). Their
548 induction patterns thus indicate that the antimicrobial *X. nematophila*-induced responses are well adapted
549 to the nature of the pathogen. Moreover, these results are in agreement with the study of Aymeric et al.

(2010) showing that the Imd pathway functions in the *D. melanogaster* immune response to *X. nematophila*. In the other categories, the *X. nematophila*-induced genes encoded the DDC melanization enzyme (Huang et al., 2005), the hemolin antibacterial opsonin (Eleftherianos et al., 2007), the IMPI-like 3, and also probably the selected integrin (Integrin β -like) (Fig. 3B-D). Once again, all of these genes are susceptible to play a part in an immune response to a pathogenic bacterium even though most of them could act on diverse types of invaders. Surprisingly, we found that *X. nematophila* strongly over-induces the transglutaminase (Tg-like) putative clotting factor gene (Lindgren et al., 2008) (Fig. 3C). This result could suggest that the bacterium is actually the main responsible for tissue damages at this time point and/or that *Tg-like* expression is induced in response to bacteria. Importantly, this result is in agreement with the study of Yadav and colleagues (Yadav et al., 2017), who showed that the *D. melanogaster* Fondue clotting factor was induced after infestation by the NBC but not after infestation by axenic nematodes. Remarkably, most of the genes that were mostly induced by *X. nematophila* presented higher induction values in response to the bacterium alone than in response to the whole NBC. However, this observation cannot be directly interpreted as an antagonistic effect of the nematode partner since it could be due to changes in the relative proportions of each hemocyte subtype, which would not necessarily reflect absolute variations in their numbers. In addition, the nematode partner specifically induced the overexpression of the selected C-type lectin (*CLECT (ccBV)*) and was probably the main inducer of the *Galectin 1*, the tetraspanin D76 homolog (*Tsp-like 3*) and the selected diapausin AMP (*Diapausin 5*) (Fig. 3A,C,D). As mentioned before, the *M. sexta* tetraspanin D76 is known to take part in encapsulation (Zhuang et al., 2007) and some lectins can bind nematodes and participate in melanization (Yu and Kanost, 2004) as well as in all types of cellular immune responses. Once again, their induction patterns are consistent with the nature of the pathogen, since both types of molecules could be involved in classical anti-nematode immune responses, such as cellular or melanotic encapsulation (Eleftherianos et al., 2017). Finally, 5 genes, encoding the Toll pathway members Pelle and Cactus (Ferrandon et al., 2007), the selected moricin AMP (Moricin 2), the melanization-related PPAE2 and the Krüppel-like transcription factor (Kr-like factor 1), were similarly induced by each of the three pathogens (Fig. 3A-C), suggesting that these responses are induced by the 2 partners without any additive effect.

In the fat body, statistical analysis of the results firstly revealed that the induction levels of *Pirk* as well as of the selected cecropin and gloverin AMP genes were significantly lower in response to the axenic nematode than in response to the NBC and to *X. nematophila* (Fig. 4A), suggesting the bacterial partner is the main responsible for their inductions. In addition, despite non-significant statistics, the results for the selected attacin AMPs, PGRP-S6 and GNB3 genes showed similar induction patterns (Fig. 4A). As for the hemocytes, the induction patterns of *Pirk* and of the attacin, cecropin and gloverin AMP genes suggest that the fat body's antimicrobial response to *X. nematophila* is well adapted to the type of pathogen that is met. On the contrary, the induction levels of the melanization-related serine protease

(*Snake-like 2*) was significantly lower in response to *X. nematophila* than in response to the NBC and to the axenic nematode (Fig. 4B), suggesting that the nematode partner is the main responsible for its induction. Similar induction patterns were obtained for the Toll pathway members *Toll* and *Cactus* (Ferrandon et al., 2007) as well as for *Galectin 1* (Fig. 4A,C). As mentioned for the hemocytes, the induction of lectins and melanization-related genes in response to the nematode is consistent with the nature of the pathogen since both could participate in classical anti-nematode immune responses (Eleftherianos et al., 2017). The induction of Toll pathway members is more difficult to relate with known anti-nematode immune responses and Yadav et al. (2018) found that the inactivation of this pathway does not impact the *D. melanogaster* survival to infestation by the whole NBC or by axenic *S. carpocapsae*. However, the involvement of this immune pathway in anti-nematode immune responses may depend on the downstream effectors and thus be variable between insect species. Finally, the other genes did not show any clear difference of induction level after injection of the 3 pathogens, except for the gallerimycin AMP, PGRP-S1 and the DDC melanization enzyme genes, which presented a lower induction when each NBC partner was injected alone (Fig. 4A,B). These results suggest synergistic effects of the nematode and of the bacterium on the induction of these genes.

In summary, we found in the 2 tissues that most of the selected genes presented partner-specific induction patterns, suggesting that the immune response to the NBC corresponds to combinations of responses induced by each partner. The detailed analysis of these genes indicates that *X. nematophila* is the main inducer of most of the selected genes, and especially of the well-known antibacterial ones. On the other hand, *S. carpocapsae* is the main inducer of some melanization and encapsulation-related genes and of the selected lectins, which could all take part in classical anti-nematode immune responses. The results thus globally suggest that the hemocytes and the fat body both respond appropriately to each NBC partner despite some yet unexplained results, such as an induction of Toll pathway members in the fat body by the nematode partner.

3.4. Expression patterns of two new clusters of candidate immune genes

During our first analysis of the RNAseq data, we identified 2 new clusters of candidate immune genes (Huot et al., 2019). The first one, named the Unknown (Unk) cluster, was localized close to *Tamozhennic*, a gene encoding a nuclear porin involved in the nucleation of Dorsal, the transcription factor of the Toll pathway (Minakhina et al., 2003). It contained 5 genes predicted to encode secreted peptides and short proteins that were all highly overexpressed in the midgut, fat body and hemocytes at 15 hpi and of which 4 were the unique mobilized genes at 8 hpi in the fat body. The second cluster, named the Genes with Bacterial Homology (GBH) cluster, contained 3 genes located inside a defensin-like AMP genes cluster in the *S. frugiperda* genome. The 3 genes were predicted to encode secreted proteins similar to each other and one of them was also found highly induced at 15 hpi in the 3 tissues. The particularity of these genes is that homologs are found only in lepidopteran species as well as, intriguingly, in Gram positive bacteria.

Here, we reexamined the expression patterns of the *Unk* and *GBH* genes and found that the 5 *Unk* genes were mainly expressed in the fat body whereas 2 of the 3 *GBH* genes were mainly expressed and induced in the hemocytes (Supplementary Table 3).

In order to learn more about their putative functions, we decided to analyze, as we did for the known immune genes, their induction patterns across the time post-infestation and in response to each NBC partner in the corresponding tissues. In both cases, we found that the induction dynamics of the genes were very similar to those of immune genes, with an upregulation that becomes significant at 5 or 10 hpi and with globally stable induction patterns from 10 to 20 hpi (Fig. 5A,B).

In the case of the *GBH* cluster, the results that we got for the 2 NBC-responsive genes (*GBH1* and *GBH3*) in the hemocytes indicate that they are significantly less induced after axenic nematode injection than after NBC and *X. nematophila* injections, suggesting that the bacterium is the main responsible for their up-regulation (Fig. 5C). We could hypothesize an acquisition by horizontal gene transfer from bacteria of the *GBH* genes. In this case, their putative involvement in the antibacterial immune response would be particularly interesting, since bacterial genes hijacking for immune purpose has only been reported once in metazoans, in the tick *Ixodes scapularis* (Chou et al., 2015). Such a hypothesis however requires functional confirmation.

In the case of the *Unk* cluster, we found that the 4 most induced genes in the fat body (*Unk2* to 5) are all strongly and similarly induced by the NBC and by the axenic nematode whereas they are not induced by *X. nematophila* (Fig. 5D). The results are very similar for the least expressed *Unk* gene (*Unk1*), for which we only found a significant induction for the injection of axenic nematodes (Fig. 5D). This partner-specific induction pattern suggests that the *Unk* genes are involved in specific aspects of the insect responses to the infestation. In addition, the putative involvement of the *Unk* genes in the response to the nematode partner seems to be in agreement with their early mobilization during the infectious process and with their overexpression in the midgut (Huot et al., 2019), which is the entry site of the nematode. In our previous study, we had hypothesized the *Unk* genes may encode new types of immune effectors (Huot et al., 2019). However, given their low levels of conservation in species as close as *S. litura* or *S. littoralis* (Supplementary Fig. 4) another hypothesis would be that they correspond to regulatory long non-coding RNAs (Johnsson et al., 2014; Qu and Adelson, 2012). In both cases, the further functional characterization of these genes could be very promising given our current lack of knowledge of the immune pathways and molecular effectors of insect anti-nematode immunity.

4. Conclusion

Here, we provide a very deep and contextualized analysis of the *S. frugiperda*'s hemocytes' and fat body's transcriptional immune responses to infestation by the *S. carpocapsae*-*X. nematophila* NBC. Our topologic analysis of these responses at 15 hpi firstly confirmed the induction of very potent and

diversified immune responses towards the pathogen, such as suggested by our previous analysis of the transcriptomic data (Huot et al., 2019) as well as by the study of Yadav et al. (2017) in the *D. melanogaster* model. The present work now shows that these responses are very stable across the post-infestation time and that they correspond to combinations of *X. nematophila*- and *S. carpocapsae*-induced responses that seem to be well adapted to the nature of each partner (Figure 6).

Together with results obtained in other insect models, the pieces of information collected during these analyses are of great interest for the study of the dialogue that takes place between each NBC partner and *S. frugiperda*'s immune systems. First, our results strongly suggest that the NBC immunosuppressive strategies globally have a low impact on the induction of immune responses at the transcriptional level in *S. frugiperda*. They also indicate that the nematode and/or its effects on the host are detected by the insect's immune system that in return seems to induce appropriate immune responses towards the pathogen. Such observations could help to determine the limits and/or universality of previously described immunosuppressive and immuno-evasive strategies of the NBC that have been described in other models. For example, they suggest that the suppressive effect of *X. nematophila* on the expression of AMP genes (Hwang et al., 2013; Ji and Kim, 2004; Park et al., 2007) as well as the camouflage strategy of *S. carpocapsae* (Binda-Rossetti et al., 2016; Mastore et al., 2015) are probably far from sufficient to explain their success towards the *S. frugiperda*'s immune system, which could also be true for other insect species. On the other hand, we found several unexplained down-regulations of signaling genes, such as of members of the Imd, JNK, TGF- β and Jak-Stat pathways, that had never been reported before and which could open new working trails for the study of the molecular basis of the NBC's immunosuppressive strategies. Finally, this study allowed the identification of very large panels of candidate immune genes involved in all the main components of insect immunity as well as of some yet uncharacterized genes that could encode new immune factors involved in the response to NBCs, such as the *GBH* and the *Unk* genes, which are respectively specific to lepidopterans and to some noctuid species (Huot et al., 2019).

Continuing this work with more functional and mechanistic approaches is now required to get an accurate picture of the molecular dialogue between the NBC and the immune system. In the longer term, using such detailed and contextualized approaches on diverse insect species could help to identify the precise causes of immune systems' failure or success towards this NBC and thus the conditions that are required for an adequate use of this pathogen against insect pests.

Acknowledgments

We thank the quarantine insect platform (PIQ), member of the Vectopole Sud network, for providing the infrastructure needed for pest insect experimentations. We are also grateful to Clotilde Gibard, Raphaël Bousquet and Gaëtan Clabots for maintaining the insect collections of the DGIMI laboratory in

688 Montpellier. This work was supported by grants from the French Institut National de la Recherche
689 Agronomique.

690 **Authors' contribution**

691 L.H., N.N. and B.D. conceived this study. N.N. and B.D. directed this study. L.H. and P.-A.G. performed
692 the infestation experiments. L.H., P.-A.G. performed dissections. L.H. and A.B. extracted and purified the
693 RNA. J.-C.O. designed the *X. nematophila* specific primers. S.P. produced the axenic nematodes and
694 checked their axenization. L.H. and A.B. performed the qPCRs. L.H., N.N. and B.D. analysed the data.
695 L.H. wrote the manuscript. L.H., N.N. and B.D. revised the manuscript. All authors have read and
696 approved the manuscript.

697 **Additional Information**

698 **Competing Interests:** The authors declare no competing interests.

- Arefin, B., Kucerova, L., Dobes, P., Markus, R., Strnad, H., Wang, Z., Hyrs, P., Zurovec, M., Theopold, U., 2014. Genome-wide transcriptional analysis of *Drosophila* larvae infected by entomopathogenic nematodes shows involvement of complement, recognition and extracellular matrix proteins. *J Innate Immun* 6, 192-204, 10.1159/000353734.
- Aymeric, J.L., Givaudan, A., Duvic, B., 2010. Imd pathway is involved in the interaction of *Drosophila melanogaster* with the entomopathogenic bacteria, *Xenorhabdus nematophila* and *Photorhabdus luminescens*. *Mol Immunol* 47, 2342-2348, 10.1016/j.molimm.2010.05.012.
- Balasubramanian, N., Hao, Y.J., Toubarro, D., Nascimento, G., Simoes, N., 2009. Purification, biochemical and molecular analysis of a chymotrypsin protease with prophenoloxidase suppression activity from the entomopathogenic nematode *Steinernema carpocapsae*. *Int J Parasitol* 39, 975-984, 10.1016/j.ijpara.2009.01.012.
- Balasubramanian, N., Toubarro, D., Simoes, N., 2010. Biochemical study and *in vitro* insect immune suppression by a trypsin-like secreted protease from the nematode *Steinernema carpocapsae*. *Parasite Immunol* 32, 165-175, 10.1111/j.1365-3024.2009.01172.x.
- Bao, Y.Y., Xue, J., Wu, W.J., Wang, Y., Lv, Z.Y., Zhang, C.X., 2011. An immune-induced Reeler protein is involved in the *Bombyx mori* melanization cascade. *Insect Biochem Mol Biol* 41, 696-706, 10.1016/j.ibmb.2011.05.001.
- Barat-Houari, M., Hilliou, F., Jousset, F.X., Sofer, L., Deleury, E., Rocher, J., Ravallec, M., Galibert, L., Delobel, P., Feyereisen, R., Fournier, P., Volkoff, A.N., 2006. Gene expression profiling of *Spodoptera frugiperda* hemocytes and fat body using cDNA microarray reveals polydnavirus-associated variations in lepidopteran host genes transcript levels. *BMC Genomics* 7, 10.1186/1471-2164-7-160.
- Baum, L.G., Garner, O.B., Schaefer, K., Lee, B., 2014. Microbe-host interactions are positively and negatively regulated by galectin glycan interactions. *Front Immunol* 5, 10.3389/Fimmu.2014.00284.
- Binda-Rossetti, S., Mastore, M., Protasoni, M., Brivio, M.F., 2016. Effects of an entomopathogen nematode on the immune response of the insect pest red palm weevil: Focus on the host antimicrobial response. *J Invertebr Pathol* 133, 110-119, 10.1016/j.jip.2015.11.001.
- Boemare, N., Thaler, J.O., Lanois, A., 1997. Simple bacteriological tests for phenotypic characterization of *Xenorhabdus* and *Photorhabdus* phase variants. *Symbiosis* 22, 167-175.
- Bonnay, F., Nguyen, X.H., Cohen-Berros, E., Troxler, L., Batsche, E., Camonis, J., Takeuchi, O., Reichhart, J.M., Matt, N., 2014. Akirin specifies NF- κ B selectivity of *Drosophila* innate immune response via chromatin remodeling. *EMBO J* 33, 2349-2362, 10.15252/embj.201488456.
- Boutros, M., Agaisse, H., Perrimon, N., 2002. Sequential activation of signaling pathways during innate immune responses in *Drosophila*. *Dev Cell* 3, 711-722, 10.1016/S1534-5807(02)00325-8.
- Bray, N.L., Pimentel, H., Melsted, P., Pachter, L., 2016. Near-optimal probabilistic RNA-seq quantification. *Nat Biotechnol* 34, 525-527, 10.1038/nbt.3519.
- Brey, P.T., Lee, W.J., Yamakawa, M., Koizumi, Y., Perrot, S., Francois, M., Ashida, M., 1993. Role of the integument in insect immunity - Epicuticular abrasion and induction of cecropin synthesis in cuticular epithelial-cells. *Proc Natl Acad Sci USA* 90, 6275-6279, 10.1073/pnas.90.13.6275.
- Brivio, M.F., Toscano, A., De Pasquale, S.M., Barbaro, A.D., Giovannardi, S., Finzi, G., Mastore, M., 2018. Surface protein components from entomopathogenic nematodes and their symbiotic bacteria: effects on immune responses of the greater wax moth, *Galleria mellonella* (Lepidoptera: Pyralidae). *Pest Manag Sci* 74, 2089-2099, 10.1002/ps.4905.
- Caldas, C., Cherqui, A., Pereira, A., Simoes, N., 2002. Purification and characterization of an extracellular protease from *Xenorhabdus nematophila* involved in insect immunosuppression. *Appl Environ Microbiol* 68, 1297-1304, 10.1128/Aem.68.3.1297-1304.2002.
- Cambon, M., Lafont, P., Frayssinet, M., Lanois, A., Ogier, J.C., Pagès, S., Parthuisot, N., Ferdy, J.B., Gaudriault, S., 2020. Bacterial community profile after the lethal infection of *Steinernema-Xenorhabdus* pairs into soil-reared *Tenebrio molitor* larvae. *FEMS Microbiol Ecol*, 10.1093/femsec/fiaa009.
- Cao, X., He, Y., Hu, Y., Wang, Y., Chen, Y.R., Bryant, B., Clem, R.J., Schwartz, L.M., Blissard, G., Jiang, H., 2015. The immune signaling pathways of *Manduca sexta*. *Insect Biochem Mol Biol* 62, 64-74, 10.1016/j.ibmb.2015.03.006.
- Castillo, J., Brown, M.R., Strand, M.R., 2011. Blood feeding and insulin-like peptide 3 stimulate proliferation of hemocytes in the mosquito *Aedes aegypti*. *PLoS Pathog* 7, e1002274, 10.1371/journal.ppat.1002274.

- Chang, D.Z., Serra, L., Lu, D., Mortazavi, A., Dillman, A.R., 2019. A core set of venom proteins is released by entomopathogenic nematodes in the genus *Steinernema*. *PLoS Pathog* 15, e1007626, 10.1371/journal.ppat.1007626.
- Chen, T.T., Tan, L.R., Hu, N., Dong, Z.Q., Hu, Z.G., Jiang, Y.M., Chen, P., Pan, M.H., Lu, C., 2018. C-lysozyme contributes to antiviral immunity in *Bombyx mori* against nucleopolyhedrovirus infection. *J Insect Physiol* 108, 54-60, 10.1016/j.jinsphys.2018.05.005.
- Chou, S., Daugherty, M.D., Peterson, S.B., Biboy, J., Yang, Y.Y., Jutras, B.L., Fritz-Laylin, L.K., Ferrin, M.A., Harding, B.N., Jacobs-Wagner, C., Yang, X.F., Vollmer, W., Malik, H.S., Mougous, J.D., 2015. Transferred interbacterial antagonism genes augment eukaryotic innate immune function. *Nature* 518, 98-101, 10.1038/nature13965.
- Conesa, A., Gotz, S., Garcia-Gomez, J.M., Terol, J., Talon, M., Robles, M., 2005. Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics* 21, 3674-3676, 10.1093/bioinformatics/bti610.
- Crawford, J.M., Portmann, C., Zhang, X., Roeffaers, M.B.J., Clardy, J., 2012. Small molecule perimeter defense in entomopathogenic bacteria. *Proc Natl Acad Sci USA* 109, 10821-10826, 10.1073/pnas.1201160109.
- De Gregorio, E., Spellman, P.T., Rubin, G.M., Lemaitre, B., 2001. Genome-wide analysis of the *Drosophila* immune response by using oligonucleotide microarrays. *Proc Natl Acad Sci USA* 98, 12590-12595, 10.1073/pnas.221458698.
- Delon, I., Brown, N.H., 2009. The integrin adhesion complex changes its composition and function during morphogenesis of an epithelium. *J Cell Sci* 122, 4363-4374, 10.1242/jcs.055996.
- Destoumieux-Garzon, D., Brehelin, M., Bulet, P., Boublik, Y., Girard, P.A., Baghdiguian, S., Zumbihl, R., Escoubas, J.M., 2009. *Spodoptera frugiperda* X-tox protein, an immune related defensin rosary, has lost the function of ancestral defensins. *PLoS One* 4, e6795, 10.1371/journal.pone.0006795.
- Dillman, A.R., Macchietto, M., Porter, C.F., Rogers, A., Williams, B., Antoshechkin, I., Lee, M.M., Goodwin, Z., Lu, X.J., Lewis, E.E., Goodrich-Blair, H., Stock, S.P., Adams, B.J., Sternberg, P.W., Mortazavi, A., 2015. Comparative genomics of *Steinernema* reveals deeply conserved gene regulatory networks. *Genome Biol* 16, 10.1186/s13059-015-0746-6.
- Dowds, B.C.A., Peters, A., 2002. Virulence Mechanisms, in *Entomopathogenic Nematology*, Gaugler, R. (Ed.). CABI Publishing, pp. 79-93.
- Duvic, B., Jouan, V., Essa, N., Girard, P.A., Pages, S., Khattar, Z.A., Volkoff, N.A., Givaudan, A., Destoumieux-Garzon, D., Escoubas, J.M., 2012. Cecropins as a marker of *Spodoptera frugiperda* immunosuppression during entomopathogenic bacterial challenge. *J Insect Physiol* 58, 881-888, 10.1016/j.jinsphys.2012.04.001.
- Eleftherianos, I., Castillo, J.C., Patrino, J., 2016. TGF- β signaling regulates resistance to parasitic nematode infection in *Drosophila melanogaster*. *Immunobiology* 221, 1362-1368, 10.1016/j.imbio.2016.07.011.
- Eleftherianos, I., Gokcen, F., Felfoldi, G., Millichap, P.J., Trenczek, T.E., French-Constant, R.H., Reynolds, S.E., 2007. The immunoglobulin family protein Hemolin mediates cellular immune responses to bacteria in the insect *Manduca sexta*. *Cell Microbiol* 9, 1137-1147, 10.1111/j.1462-5822.2006.00855.x.
- Eleftherianos, I., Shokal, U., Yadav, S., Kenney, E., Maldonado, T., 2017. Insect immunity to entomopathogenic nematodes and their mutualistic bacteria. *Curr Top Microbiol Immunol* 402, 123-156, 10.1007/82_2016_52.
- Eom, S., Park, Y., Kim, Y., 2014. Sequential immunosuppressive activities of bacterial secondary metabolites from the entomopathogenic bacterium *Xenorhabdus nematophila*. *J Microbiol* 52, 161-168, 10.1007/s12275-014-3251-9.
- Erler, S., Popp, M., Lattorff, H.M.G., 2011. Dynamics of immune system gene expression upon bacterial challenge and wounding in a social insect (*Bombus terrestris*). *PLoS One* 6, e18126, 10.1371/journal.pone.0018126.
- Ferrandon, D., 2013. The complementary facets of epithelial host defenses in the genetic model organism *Drosophila melanogaster*: from resistance to resilience. *Curr Opin Immunol* 25, 59-70, 10.1016/j.coi.2012.11.008.
- Ferrandon, D., Imler, J.L., Hetru, C., Hoffmann, J.A., 2007. The *Drosophila* systemic immune response: sensing and signalling during bacterial and fungal infections. *Nat Rev Immunol* 7, 862-874, 10.1038/nri2194.
- Fichelson, P., Brigui, A., Pichaud, F., 2012. Orthodenticle and Krüppel homolog 1 regulate *Drosophila* photoreceptor maturation. *Proc Natl Acad Sci USA* 109, 7893-7898, 10.1073/pnas.1120276109.

804 Fisher, K.H., Fragiadaki, M., Pugazhendhi, D., Bausek, N., Arredondo, M.A., Thomas, S.J., Brown, S., Zeidler, M.P.,
805 2018. A genome-wide RNAi screen identifies MASK as a positive regulator of cytokine receptor stability. *J*
806 *Cell Sci* 131, 10.1242/jcs.209551.

807 Foley, E., O'Farrell, P.H., 2004. Functional dissection of an innate immune response by a genome-wide RNAi
808 screen. *PLoS Biol* 2, 1091-1106, 10.1371/journal.pbio.0020203.

809 Forst, S., Clarke, D., 2002. Bacteria–nematode symbiosis, in *Entomopathogenic Nematology*, Gaugler, R. (Ed.).
810 CABI Publishing, pp. 57-78.

811 Galko, M.J., Krasnow, M.A., 2004. Cellular and genetic analysis of wound healing in *Drosophila* larvae. *PLoS Biol* 2,
812 1114-1126, 10.1371/journal.pbio.0020239.

813 Gandhe, A.S., Janardhan, G., Nagaraju, J., 2007. Immune upregulation of novel antibacterial proteins from
814 silkmoths (Lepidoptera) that resemble lysozymes but lack muramidase activity. *Insect Biochem Mol Biol* 37,
815 655-666, 10.1016/j.ibmb.2007.03.013.

816 Ganger, M.T., Dietz, G.D., Ewing, S.J., 2017. A common base method for analysis of qPCR data and the application
817 of simple blocking in qPCR experiments. *BMC Bioinformatics* 18, 534, 10.1186/s12859-017-1949-5.

818 Gotwals, P.J., Fessler, L.I., Wehrli, M., Hynes, R.O., 1994. *Drosophila* PS1 integrin is a laminin receptor and differs
819 in ligand specificity from PS2. *Proc Natl Acad Sci USA* 91, 11447-11451, 10.1073/pnas.91.24.11447.

820 Gotz, P., Boman, A., Boman, H.G., 1981. Interactions between insect immunity and an insect-pathogenic
821 nematode with symbiotic bacteria. *Proc R Soc B* 212, 333-350, 10.1098/rspb.1981.0043.

822 Gouin, A., Bretaudeau, A., Nam, K., Gimenez, S., Aury, J.M., Duvic, B., Hilliou, F., Durand, N., Montagne, N.,
823 Darboux, I., Kuwar, S., Chertemps, T., Siaussat, D., Bretschneider, A., Mone, Y., Ahn, S.J., Hanniger, S.,
824 Grenet, A.S.G., Neunemann, D., Maumus, F., Luyten, I., Labadie, K., Xu, W., Koutroumpa, F., Escoubas, J.M.,
825 Llopis, A., Maibeche-Coisne, M., Salasc, F., Tomar, A., Anderson, A.R., Khan, S.A., Dumas, P., Orsucci, M.,
826 Guy, J., Belser, C., Alberti, A., Noel, B., Couloux, A., Mercier, J., Nidelet, S., Dubois, E., Liu, N.Y., Boulogne, I.,
827 Mirabeau, O., Le Goff, G., Gordon, K., Oakeshott, J., Consoli, F.L., Volkoff, A.N., Fescemyer, H.W., Marden,
828 J.H., Luthe, D.S., Herrero, S., Heckel, D.G., Wincker, P., Kergoat, G.J., Amselem, J., Quesneville, H., Groot,
829 A.T., Jacquin-Joly, E., Negre, N., Lemaitre, C., Legeai, F., d'Alençon, E., Fournier, P., 2017. Two genomes of
830 highly polyphagous lepidopteran pests (*Spodoptera frugiperda*, Noctuidae) with different host-plant
831 ranges. *Sci Rep* 7, 11816, 10.1038/S41598-017-10461-4.

832 Griesch, J., Wedde, M., Vilcinskas, A., 2000. Recognition and regulation of metalloproteinase activity in the
833 haemolymph of *Galleria mellonella*: a new pathway mediating induction of humoral immune responses.
834 *Insect Biochem Mol Biol* 30, 461-472, 10.1016/S0965-1748(00)00010-2.

835 Ha, E.M., Lee, K.A., Park, S.H., Kim, S.H., Nam, H.J., Lee, H.Y., Kang, D., Lee, W.J., 2009. Regulation of DUOX by the
836 Gαq-phospholipase Cβ-Ca²⁺ pathway in *Drosophila* gut immunity. *Dev Cell* 16, 386-397,
837 10.1016/j.devcel.2008.12.015.

838 Hao, Y.J., Montiel, R., Abubucker, S., Mitreva, M., Simoes, N., 2010. Transcripts analysis of the entomopathogenic
839 nematode *Steinernema carpocapsae* induced *in vitro* with insect haemolymph. *Mol Biochem Parasit* 169,
840 79-86, 10.1016/j.molbiopara.2009.10.002.

841 Harbecke, R., Janning, W., 1989. The segmentation gene Krüppel of *Drosophila melanogaster* has homeotic
842 properties. *Gene Dev* 3, 114-122, 10.1101/Gad.3.1.114.

843 Hemler, M.E., 2008. Targeting of tetraspanin proteins - potential benefits and strategies. *Nat Rev Drug Discov* 7,
844 747-758, 10.1038/nrd2659.

845 Herrero, S., Ansems, M., Van Oers, M.M., Vlak, J.M., Bakker, P.L., de Maagd, R.A., 2007. REPAT, a new family of
846 proteins induced by bacterial toxins and baculovirus infection in *Spodoptera exigua*. *Insect Biochem Mol*
847 *Biol* 37, 1109-1118, 10.1016/j.ibmb.2007.06.007.

848 Hoch, M., Jackle, H., 1998. Krüppel acts as a developmental switch gene that mediates Notch signalling-
849 dependent tip cell differentiation in the excretory organs of *Drosophila*. *EMBO J* 17, 5766-5775,
850 10.1093/emboj/17.19.5766.

851 Huang, C.Y., Chou, S.Y., Bartholomay, L.C., Christensen, B.M., Chen, C.C., 2005. The use of gene silencing to study
852 the role of dopa decarboxylase in mosquito melanization reactions. *Insect Mol Biol* 14, 237-244,
853 10.1111/j.1365-2583.2004.00552.x.

854 Huot, L., George, S., Girard, P.A., Severac, D., Negre, N., Duvic, B., 2019. *Spodoptera frugiperda* transcriptional
855 response to infestation by *Steinernema carpocapsae*. *Sci Rep* 9, 12879, 10.1038/s41598-019-49410-8.

856 Hwang, J., Park, Y., Kim, Y., Hwang, J., Lee, D., 2013. An entomopathogenic bacterium, *Xenorhabdus nematophila*,
857 suppresses expression of antimicrobial peptides controlled by Toll and Imd pathways by blocking
858 eicosanoid biosynthesis. Arch Insect Biochem Physiol 83, 151-169, 10.1002/arch.21103.

859 Hyrsi, P., Dobes, P., Wang, Z., Hauling, T., Wilhelmsson, C., Theopold, U., 2011. Clotting factors and eicosanoids
860 protect against nematode infections. J Innate Immun 3, 65-70, 10.1159/000320634.

861 Issa, N., Guillaumot, N., Lauret, E., Matt, N., Schaeffer-Reiss, C., Van Dorsselaer, A., Reichhart, J.M., Veillard, F.,
862 2018. The circulating protease Persephone is an immune sensor for microbial proteolytic activities
863 upstream of the *Drosophila* Toll pathway. Mol Cell 69, 539-550, 10.1016/j.molcel.2018.01.029.

864 Ivy, J.R., Drechsler, M., Catterson, J.H., Bodmer, R., Ocorr, K., Paululat, A., Hartley, P.S., 2015. Klf15 is critical for
865 the development and differentiation of *Drosophila* nephrocytes. PLoS One 10, e0134620,
866 10.1371/journal.pone.0134620.

867 Ji, D.J., Kim, Y., 2004. An entomopathogenic bacterium, *Xenorhabdus nematophila*, inhibits the expression of an
868 antibacterial peptide, cecropin, of the beet armyworm, *Spodoptera exigua*. J Insect Physiol 50, 489-496,
869 10.1016/j.jinsphys.2004.03.005.

870 Jing, Y.J., Toubarro, D., Hao, Y.J., Simoes, N., 2010. Cloning, characterisation and heterologous expression of an
871 astacin metalloprotease, Sc-AST, from the entomoparasitic nematode *Steinernema carpocapsae*. Mol
872 Biochem Parasit 174, 101-108, 10.1016/j.molbiopara.2010.07.004.

873 Jiravanichpaisal, P., Lee, B.L., Soderhall, K., 2006. Cell-mediated immunity in arthropods: Hematopoiesis,
874 coagulation, melanization and opsonization. Immunobiology 211, 213-236, 10.1016/j.imbio.2005.10.015.

875 Johnsson, P., Lipovich, L., Grander, D., Morris, K.V., 2014. Evolutionary conservation of long non-coding RNAs:
876 Sequence, structure, function. Biochim Biophys Acta Gen Subj 1840, 1063-1071,
877 10.1016/j.bbagen.2013.10.035.

878 Kim, Y., Ji, D., Cho, S., Park, Y., 2005. Two groups of entomopathogenic bacteria, *Photorhabdus* and *Xenorhabdus*,
879 share an inhibitory action against phospholipase A₂ to induce host immunodepression. J Invertebr Pathol
880 89, 258-264, 10.1016/j.jip.2005.05.001.

881 Kleino, A., Myllymaki, H., Kallio, J., Vanha-aho, L.M., Oksanen, K., Ulvila, J., Hultmark, D., Valanne, S., Ramet, M.,
882 2008. Pirk is a negative regulator of the *Drosophila* Imd pathway. J Immunol 180, 5413-5422,
883 10.4049/jimmunol.180.8.5413.

884 Korayem, A.M., Fabbri, M., Takahashi, K., Scherfer, C., Lindgren, M., Schmidt, O., Ueda, R., Dushay, M.S.,
885 Theopold, U., 2004. A *Drosophila* salivary gland mucin is also expressed in immune tissues: evidence for a
886 function in coagulation and the entrapment of bacteria. Insect Biochem Mol Biol 34, 1297-1304,
887 10.1016/j.ibmb.2004.09.001.

888 Kristensen, N., Chauvin, G., 2012. Lepidoptera, moths and butterflies: Morphology, physiology, and development.
889 Integument, in Handbook of Zoology, De Gruyter, W. (Ed.). Berlin: De Gruyter, pp. 1-8.

890 Kucerova, L., Broz, V., Arefin, B., Maaroufi, H.O., Hurychova, J., Strnad, H., Zurovec, M., Theopold, U., 2016. The
891 *Drosophila* chitinase-like protein IDGF3 is involved in protection against nematodes and in wound healing. J
892 Innate Immun 8, 199-210, 10.1159/000442351.

893 Labaude, S., Griffin, C.T., 2018. Transmission success of entomopathogenic nematodes used in pest control.
894 Insects 9, E72, 10.3390/insects9020072.

895 Lacey, L.A., Grzywacz, D., Shapiro-Ilan, D.I., Frutos, R., Brownbridge, M., Goettel, M.S., 2015. Insect pathogens as
896 biological control agents: Back to the future. J Invertebr Pathol 132, 1-41, 10.1016/j.jip.2015.07.009.

897 Lagueux, M., Perrodou, E., Levashina, E.A., Capovilla, M., Hoffmann, J.A., 2000. Constitutive expression of a
898 complement-like protein in Toll and JAK gain-of-function mutants of *Drosophila*. Proc Natl Acad Sci USA 97,
899 11427-11432, 10.1073/pnas.97.21.11427.

900 Lee, K.A., Lee, W.J., 2018. Immune-metabolic interactions during systemic and enteric infection in *Drosophila*.
901 Curr Opin Insect Sci 29, 21-26, 10.1016/j.cois.2018.05.014.

902 Lehane, M.J., 1997. Peritrophic matrix structure and function. Annu Rev Entomol 42, 525-550,
903 10.1146/annurev.ento.42.1.525.

904 Lemaitre, B., Reichhart, J.M., Hoffmann, J.A., 1997. *Drosophila* host defense: Differential induction of
905 antimicrobial peptide genes after infection by various classes of microorganisms. Proc Natl Acad Sci USA 94,
906 14614-14619, 10.1073/pnas.94.26.14614.

907 Levin, D.M., Breuer, L.N., Zhuang, S.F., Anderson, S.A., Nardi, J.B., Kanost, M.R., 2005. A hemocyte-specific
908 integrin required for hemocytic encapsulation in the tobacco hornworm, *Manduca sexta*. Insect Biochem
909 Mol Biol 35, 369-380, 10.1016/j.ibmb.2005.01.003.

Li, X.Y., Cowles, R.S., Cowles, E.A., Gaugler, R., Cox-Foster, D.L., 2007. Relationship between the successful infection by entomopathogenic nematodes and the host immune response. *Int J Parasitol* 37, 365-374, 10.1016/j.ijpara.2006.08.009.

Lindgren, M., Riaz, R., Lesch, C., Willielinsson, C., Theopold, U., Dushay, M.S., 2008. Fondue and transglutaminase in the *Drosophila* larval clot. *J Insect Physiol* 54, 586-592, 10.1016/j.jinsphys.2007.12.008.

Lu, D., Macchietto, M., Chang, D., Barros, M.M., Baldwin, J., Mortazavi, A., Dillman, A.R., 2017. Activated entomopathogenic nematode infective juveniles release lethal venom proteins. *PLoS Pathog* 13, e1006302, 10.1371/journal.ppat.1006302.

Massaoud, M.K., Marokhazi, J., Venekei, I., 2011. Enzymatic characterization of a serralsin-like metalloprotease from the entomopathogen bacterium, *Xenorhabdus*. *Biochim Biophys Acta-Proteins Proteom* 1814, 1333-1339, 10.1016/j.bbapap.2011.05.008.

Mastore, M., Arizza, V., Manachini, B., Brivio, M.F., 2015. Modulation of immune responses of *Rhynchophorus ferrugineus* (Insecta: Coleoptera) induced by the entomopathogenic nematode *Steinernema carpocapsae* (Nematoda: Rhabditida). *Insect Sci* 22, 748-760, 10.1111/1744-7917.12141.

Melcarne, C., Lemaitre, B., Kuran, E., 2019. Phagocytosis in *Drosophila*: From molecules and cellular machinery to physiology. *Insect Biochem Mol Biol* 109, 1-12, 10.1016/j.ibmb.2019.04.002.

Minakhina, S., Yang, J., Steward, R., 2003. Tamo selectively modulates nuclear import in *Drosophila*. *Genes Cells* 8, 299-310, 10.1046/j.1365-2443.2002.00634.x.

Myllymaki, H., Valanne, S., Ramet, M., 2014. The *Drosophila* Imd signaling pathway. *J Immunol* 192, 3455-3462, 10.4049/jimmunol.1303309.

Nakahara, Y., Matsumoto, H., Kanamori, Y., Kataoka, H., Mizoguchi, A., Kiuchi, M., Kamimura, M., 2006. Insulin signaling is involved in hematopoietic regulation in an insect hematopoietic organ. *J Insect Physiol* 52, 105-111, 10.1016/j.jinsphys.2005.09.009.

Nakhleh, J., El Moussawi, L., Osta, M.A., 2017. Chapter 3 - The melanization response in insect immunity, in *Advances in Insect Physiology*, Ligoxygakis, P. (Ed.). Academic Press, pp. 83-109.

Nappi, A.J., Christensen, B.M., 2005. Melanogenesis and associated cytotoxic reactions: Applications to insect innate immunity. *Insect Biochem Mol Biol* 35, 443-459, 10.1016/j.ibmb.2005.01.014.

Navarro-Cerrillo, G., Ferre, J., de Maagd, R.A., Herrero, S., 2012. Functional interactions between members of the REPAT family of insect pathogen-induced proteins. *Insect Mol Biol* 21, 335-342, 10.1111/j.1365-2583.2012.01139.x.

Navarro-Cerrillo, G., Hernandez-Martinez, P., Vogel, H., Ferre, J., Herrero, S., 2013. A new gene superfamily of pathogen-response (repat) genes in *Lepidoptera*: classification and expression analysis. *Comp Biochem Physiol B-Biochem Physiol* 164, 10-17, 10.1016/j.cbpb.2012.09.004.

Nazario-Toole, A.E., Wu, L.P., 2017. Phagocytosis in insect immunity, in *Advances in Insect Physiology*, Volume 52, Ligoxygakis, P. (Ed.). Kruze, K., pp. 35-73.

Ozakman, Y., Eleftherianos, I., 2019. TGF- β signaling interferes with the *Drosophila* innate immune and metabolic response to parasitic nematode infection. *Front Physiol* 10, 10.3389/fphys.2019.00716.

Pace, K.E., Baum, L.G., 2002. Insect galectins: Roles in immunity and development. *Glycoconjugate J* 19, 607-614, 10.1023/B:Glyc.0000014092.86763.2f.

Park, Y., Herbert, E.E., Cowles, C.E., Cowles, K.N., Menard, M.L., Orchard, S.S., Goodrich-Blair, H., 2007. Clonal variation in *Xenorhabdus nematophila* virulence and suppression of *Manduca sexta* immunity. *Cell Microbiol* 9, 645-656, 10.1111/j.1462-5822.2006.00815.x.

Park, Y., Kim, Y., 2000. Eicosanoids rescue *Spodoptera exigua* infected with *Xenorhabdus nematophilus*, the symbiotic bacteria to the entomopathogenic nematode *Steinernema carpocapsae*. *J Insect Physiol* 46, 1469-1476, 10.1016/S0022-1910(00)00071-8.

Park, Y., Stanley, D., 2006. The entomopathogenic bacterium, *Xenorhabdus nematophila*, impairs hemocytic immunity by inhibition of eicosanoid biosynthesis in adult crickets, *Gryllus firmus*. *Biol Control* 38, 247-253, 10.1016/j.biocontrol.2005.11.002.

Pastor-Pareja, J.C., Wu, M., Xu, T., 2008. An innate immune response of blood cells to tumors and tissue damage in *Drosophila*. *Dis Model Mech* 1, 144-154, 10.1242/dmm.000950.

Patnogi, J., Heryanto, C., Eleftherianos, I., 2018. Wounding-induced upregulation of the Bone Morphogenic Protein signaling pathway in *Drosophila* promotes survival against parasitic nematode infection. *Gene* 673, 112-118, 10.1016/j.gene.2018.06.052.

963 Pimentel, H., Bray, N.L., Puente, S., Melsted, P., Pachter, L., 2017. Differential analysis of RNA-seq incorporating
 964 quantification uncertainty. *Nat Methods* 14, 687-690, 10.1038/nmeth.4324.
 965 Poinar, G.O., Grewal, P.S., 2012. History of entomopathogenic nematology. *J Nematol* 44, 153-161.
 966 Poitout, S., Buès, R., 1970. Elevage de plusieurs espèces de Lépidoptères Noctuidae sur milieu artificiel riche et
 967 sur milieu artificiel simplifié. *Ann Zool Ecol Anim* 2, 79-91.
 968 Qiao, C., Li, J., Wei, X.H., Wang, J.L., Wang, Y.F., Liu, X.S., 2014. SRP gene is required for *Helicoverpa armigera*
 969 prophenoloxidase activation and nodulation response. *Dev Comp Immunol* 44, 94-99,
 970 10.1016/j.dci.2013.11.016.
 971 Qu, Z.P., Adelson, D.L., 2012. Identification and comparative analysis of ncRNAs in human, mouse and zebrafish
 972 indicate a conserved role in regulation of genes expressed in brain. *PLoS One* 7, e52275,
 973 10.1371/journal.pone.0052275.
 974 R Core Team, 2017. R: A language and environment for statistical computing. R Foundation for Statistical
 975 Computing, Vienna, Austria, URL <https://www.R-project.org/>.
 976 Ribeiro, C., Duvic, B., Oliveira, P., Givaudan, A., Palha, F., Simoes, N., Brehelin, M., 1999. Insect immunity - effects
 977 of factors produced by a nematobacterial complex on immunocompetent cells. *J Insect Physiol* 45, 677-685,
 978 10.1016/S0022-1910(99)00043-8.
 979 Ribeiro, C., Vignes, M., Brehelin, M., 2003. *Xenorhabdus nematophila* (Enterobacteriaceae) secretes a cation-
 980 selective calcium-independent porin which causes vacuolation of the rough endoplasmic reticulum and cell
 981 lysis. *J Biol Chem* 278, 3030-3039, 10.1074/jbc.M210353200.
 982 Roote, C.E., Zusman, S., 1996. Alternatively spliced forms of the *Drosophila* α PS2 subunit of integrin are sufficient
 983 for viability and can replace the function of the α PS1 subunit of integrin in the retina. *Development* 122,
 984 1985-1994.
 985 Rowley, A.F., Ratcliffe, N.A., 1978. Histological study of wound-healing and hemocyte function in wax moth
 986 *Galleria mellonella*. *J Morphol* 157, 181-199, 10.1002/jmor.1051570206.
 987 Satyavathi, V.V., Mohamed, A.A., Kumari, S., Mamatha, D.M., Duvic, B., 2018. The IMD pathway regulates
 988 lysozyme-like proteins (LLPs) in the silkworm *Antheraea mylitta*. *J Invertebr Pathol* 154, 102-108,
 989 10.1016/j.jip.2018.04.006.
 990 Schmucker, D., Taubert, H., Jackle, H., 1992. Formation of the *Drosophila* larval photoreceptor organ and its
 991 neuronal differentiation require continuous Krüppel gene activity. *Neuron* 9, 1025-1039, 10.1016/0896-
 992 6273(92)90063-J.
 993 Shin, S.W., Park, S.S., Park, D.S., Kim, M.G., Kim, S.C., Brey, P.T., Park, H.Y., 1998. Isolation and characterization of
 994 immune-related genes from the fall webworm, *Hyphantria cunea*, using PCR-based differential display and
 995 subtractive cloning. *Insect Biochem Mol Biol* 28, 827-837.
 996 Shokal, U., Eleftherianos, I., 2017. The *Drosophila* Thioester containing Protein-4 participates in the induction of
 997 the cellular immune response to the pathogen *Photobacterium*. *Dev Comp Immunol* 76, 200-208,
 998 10.1016/j.dci.2017.06.008.
 999 Shokal, U., Kopydlowski, H., Eleftherianos, I., 2017. The distinct function of Tep2 and Tep6 in the immune defense
 1000 of *Drosophila melanogaster* against the pathogen *Photobacterium*. *Virulence* 8, 1668-1682,
 1001 10.1080/21505594.2017.1330240.
 1002 Shokal, U., Kopydlowski, H., Harsh, S., Eleftherianos, I., 2018. Thioester-containing proteins 2 and 4 affect the
 1003 metabolic activity and inflammation response in *Drosophila*. *Infect Immun* 86, 10.1128/IAI.00810-17.
 1004 Sicard, M., Brugirard-Ricaud, K., Pages, S., Lanois, A., Boemare, N.E., Brehelin, M., Givaudan, A., 2004. Stages of
 1005 infection during the tripartite interaction between *Xenorhabdus nematophila*, its nematode vector, and
 1006 insect hosts. *Appl Environ Microbiol* 70, 6473-6480, 10.1128/AEM.70.11.6473-6480.2004.
 1007 Sowa-Jasilek, A., Zdybicka-Barabas, A., Staczek, S., Wydrych, J., Mak, P., Jakubowicz, T., Cytrynska, M., 2014.
 1008 Studies on the role of insect hemolymph polypeptides: *Galleria mellonella* anionic peptide 2 and lysozyme.
 1009 Peptides 53, 194-201, 10.1016/j.peptides.2014.01.012.
 1010 Stofanko, M., Kwon, S.Y., Badenhorst, P., 2008. A misexpression screen to identify regulators of *Drosophila* larval
 1011 hemocyte development. *Genetics* 180, 253-267, 10.1534/genetics.108.089094.
 1012 Strand, M.R., 2008. The insect cellular immune response. *Insect Sci* 15, 1-14, 10.1111/j.1744-7917.2008.00183.x.
 1013 Stuart, L.M., Deng, J.S., Silver, J.M., Takahashi, K., Tseng, A.A., Hennessy, E.J., Ezekowitz, R.A.B., Moore, K.J., 2005.
 1014 Response to *Staphylococcus aureus* requires CD36-mediated phagocytosis triggered by the COOH-terminal
 1015 cytoplasmic domain. *J Cell Biol* 170, 477-485, 10.1083/jcb.200501113.

1016 Tailliez, P., Pages, S., Ginibre, N., Boemare, N., 2006. New insight into diversity in the genus *Xenorhabdus*,
1017 including the description of ten novel species. *Int J Syst Evol Microbiol* 56, 2805-2818, 10.1099/ij.s.0.64287-
1018 0.

1019 Tang, H.P., 2009. Regulation and function of the melanization reaction in *Drosophila*. *Fly* 3, 105-111,
1020 10.4161/Fly.3.1.7747.

1021 Thurston, G.S., Yule, W.N., Dunphy, G.B., 1994. Explanations for the low susceptibility of *Leptinotarsa*
1022 *decemlineata* to *Steinernema carpocapsae*. *Biol Control* 4, 53-58, 10.1006/bcon.1994.1010.

1023 Tingvall, T.O., Roos, E., Engstrom, Y., 2001. The Imd gene is required for local Cecropin expression in *Drosophila*
1024 barrier epithelia. *EMBO Rep* 2, 239-243.

1025 Toubarro, D., Avila, M.M., Hao, Y.J., Balasubramanian, N., Jing, Y.J., Montiel, R., Faria, T.Q., Brito, R.M., Simoes, N.,
1026 2013a. A serpin released by an entomopathogen impairs clot formation in insect defense system. *PLoS One*
1027 8, e69161, 10.1371/journal.pone.0069161.

1028 Toubarro, D., Avila, M.M., Montiel, R., Simoes, N., 2013b. A pathogenic nematode targets recognition proteins to
1029 avoid insect defenses. *PLoS One* 8, e75691, 10.1371/journal.pone.0075691.

1030 Tzou, P., Ohresser, S., Ferrandon, D., Capovilla, M., Reichhart, J.M., Lemaitre, B., Hoffmann, J.A., Imler, J.L., 2000.
1031 Tissue-specific inducible expression of antimicrobial peptide genes in *Drosophila* surface epithelia.
1032 *Immunity* 13, 737-748, 10.1016/S1074-7613(00)00072-8.

1033 Untergasser, A., Cutcutache, I., Koressaar, T., Ye, J., Faircloth, B.C., Remm, M., Rozen, S.G., 2012. Primer3--new
1034 capabilities and interfaces. *Nucleic Acids Res* 40, e115, 10.1093/nar/gks596.

1035 Urbano, J.M., Dominguez-Gimenez, P., Estrada, B., Martin-Bermudo, M.D., 2011. PS integrins and laminins: Key
1036 regulators of cell migration during *Drosophila* embryogenesis. *PLoS One* 6, e23893,
1037 10.1371/journal.pone.0023893.

1038 Valanne, S., Wang, J.H., Ramet, M., 2011. The *Drosophila* Toll Signaling Pathway. *J Immunol* 186, 649-656,
1039 10.4049/jimmunol.1002302.

1040 van Sambeek, J., Wiesner, A., 1999. Successful parasitism of locusts by entomopathogenic nematodes is
1041 correlated with inhibition of insect phagocytes. *J Invertebr Pathol* 73, 154-161, 10.1006/jipa.1998.4823.

1042 Veillard, F., Troxler, L., Reichhart, J.M., 2016. *Drosophila melanogaster* clip-domain serine proteases: Structure,
1043 function and regulation. *Biochimie* 122, 255-269, 10.1016/j.biochi.2015.10.007.

1044 Vertyporokh, L., Wojda, I., 2017. Expression of the insect metalloproteinase inhibitor IMPI in the fat body of
1045 *Galleria mellonella* exposed to infection with *Beauveria bassiana*. *Acta Biochim Pol* 64, 273-278,
1046 10.18388/abp.2016_1376.

1047 Vigneux, F., Zumbihl, R., Jubelin, G., Ribeiro, C., Poncet, J., Baghdiguian, S., Givaudan, A., Brehelin, M., 2007. The
1048 xaxAB genes encoding a new apoptotic toxin from the insect pathogen *Xenorhabdus nematophila* are
1049 present in plant and human pathogens. *J Biol Chem* 282, 9571-9580, 10.1074/jbc.M604301200.

1050 Wang, Y., Branicky, R., Noe, A., Hekimi, S., 2018. Superoxide dismutases: Dual roles in controlling ROS damage
1051 and regulating ROS signaling. *J Cell Biol* 217, 1915-1928, 10.1083/jcb.201708007.

1052 Wang, Y., Gaugler, R., Cui, L.W., 1994. Variations in immune response of *Popillia japonica* and *Acheta domesticus*
1053 to *Heterorhabditis bacteriophora* and *Steinernema* species. *J Nematol* 26, 11-18.

1054 Wang, Z., Wilhelmsson, C., Hyrsil, P., Loof, T.G., Dobes, P., Klupp, M., Loseva, O., Morgelin, M., Ikle, J., Cripps, R.M.,
1055 Herwald, H., Theopold, U., 2010. Pathogen entrapment by transglutaminase -- a conserved early innate
1056 immune mechanism. *PLoS Pathog* 6, e1000763, 10.1371/journal.ppat.1000763.

1057 Wedde, M., Weise, C., Kopacek, P., Franke, P., Vilcinskis, A., 1998. Purification and characterization of an
1058 inducible metalloproteinase inhibitor from the hemolymph of greater wax moth larvae, *Galleria mellonella*.
1059 *Eur J Biochem* 255, 535-543, 10.1046/j.1432-1327.1998.2550535.x.

1060 Wedde, M., Weise, C., Nuck, R., Altincicek, B., Vilcinskis, A., 2007. The insect metalloproteinase inhibitor gene of
1061 the lepidopteran *Galleria mellonella* encodes two distinct inhibitors. *Biol Chem* 388, 119-127,
1062 10.1515/Bc.2007.013.

1063 White, G.F., 1927. A method for obtaining infective nematode larvae from cultures. *Science* 66, 302-303,
1064 10.1126/science.66.1709.302-a.

1065 Wu, S., Zhang, X.F., He, Y.Q., Shuai, J.B., Chen, X.M., Ling, E.J., 2010. Expression of antimicrobial peptide genes in
1066 *Bombyx mori* gut modulated by oral bacterial infection and development. *Dev Comp Immunol* 34, 1191-
1067 1198, 10.1016/j.dci.2010.06.013.

1068 Xia, X., You, M., Rao, X.J., Yu, X.Q., 2018. Insect C-type lectins in innate immunity. *Dev Comp Immunol* 83, 70-79,
1069 10.1016/j.dci.2017.11.020.

1070 Yadav, S., Daugherty, S., Shetty, A.C., Eleftherianos, I., 2017. RNAseq analysis of the *Drosophila* response to the
1071 entomopathogenic nematode *Steinernema*. *G3 (Bethesda)* 7, 1955-1967, 10.1534/g3.117.041004.

1072 Yadav, S., Eleftherianos, I., 2018. The Imaginal Disc Growth Factors 2 and 3 participate in the *Drosophila* response
1073 to nematode infection. *Parasite Immunol* 40, 10.1111/pim.12581.

1074 Yadav, S., Eleftherianos, I., 2019. Participation of the serine protease Jonah66Ci in the *Drosophila* antinematode
1075 immune response. *Infect Immun* 87, 10.1128/IAI.00094-19.

1076 Yadav, S., Gupta, S., Eleftherianos, I., 2018. Differential regulation of immune signaling and survival response in
1077 *Drosophila melanogaster* larvae upon *Steinernema carpocapsae* nematode infection. *Insects* 9,
1078 10.3390/Insects9010017.

1079 Yi, H.Y., Chowdhury, M., Huang, Y.D., Yu, X.Q., 2014. Insect antimicrobial peptides and their applications. *Appl*
1080 *Microbiol Biotechnol* 98, 5807-5822, 10.1007/s00253-014-5792-6.

1081 Yu, K.H., Kim, K.N., Lee, J.H., Lee, H.S., Kim, S.H., Cho, K.Y., Nam, M.H., Lee, I.H., 2002. Comparative study on
1082 characteristics of lysozymes from the hemolymph of three lepidopteran larvae, *Galleria mellonella*, *Bombyx*
1083 *mori*, *Agrius convolvuli*. *Dev Comp Immunol* 26, 707-713, 10.1016/S0145-305x(02)00027-7.

1084 Yu, X.Q., Kanost, M.R., 2004. Immulectin-2, a pattern recognition receptor that stimulates hemocyte
1085 encapsulation and melanization in the tobacco hornworm, *Manduca sexta*. *Dev Comp Immunol* 28, 891-
1086 900, 10.1016/j.dci.2004.02.005.

1087 Yuan, C.F., Xing, L.S., Wang, M.L., Wang, X., Yin, M.Y., Wang, Q.R., Hu, Z.H., Zou, Z., 2017. Inhibition of
1088 melanization by serpin-5 and serpin-9 promotes baculovirus infection in cotton bollworm *Helicoverpa*
1089 *armigera*. *PLoS Pathog* 13, e1006645, 10.1371/journal.ppat.1006645.

1090 Zhang, K., Pan, G.Z., Zhao, Y.Z., Hao, X.W., Li, C.Y., Shen, L., Zhang, R., Su, J.J., Cui, H.J., 2017. A novel immune-
1091 related gene HDD1 of silkworm *Bombyx mori* is involved in bacterial response. *Mol Immunol* 88, 106-115,
1092 10.1016/j.molimm.2017.06.023.

1093 Zhang, S., Yan, H., Li, C.Z., Chen, Y.H., Yuan, F.H., Chen, Y.G., Weng, S.P., He, J.G., 2013. Identification and function
1094 of leucine-rich repeat flightless-I-interacting protein 2 (LRRFIP2) in *Litopenaeus vannamei*. *PLoS One* 8,
1095 e57456, 10.1371/journal.pone.0057456.

1096 Zhuang, S.F., Kelo, L.S., Nardi, J.B., Kanost, M.R., 2007. An integrin-tetraspanin interaction required for cellular
1097 innate immune responses of an insect, *Manduca sexta*. *J Biol Chem* 282, 10.1074/Jbc.M700341200.

1098 Zou, F.M., Lee, K.S., Kim, B.Y., Kim, H.J., Gui, Z.Z., Zhang, G.Z., Guo, X.J., Jin, B.R., 2015. Differential and spatial
1099 regulation of the prophenoloxidase (proPO) and proPO-activating enzyme in cuticular melanization and
1100 innate immunity in *Bombyx mori* pupae. *J Asia-Pac Entomol* 18, 757-764, 10.1016/j.aspen.2015.09.007.

1101 Zou, Z., Shin, S.W., Alvarez, K.S., Kokoza, V., Raikhell, A.S., 2010. Distinct melanization pathways in the mosquito
1102 *Aedes aegypti*. *Immunity* 32, 41-53, 10.1016/j.immuni.2009.11.011.

1103

1104 Figure Legends

1105 **Fig 1.** Expression variations of the differentially expressed immune genes after infestation by the
1106 nematobacterial complex. Heatmaps showing the expression variations of the differentially expressed
1107 immune genes in the hemocytes and in the fat body at a middle time point of 15 h post-infestation.
1108 RNAseq raw data were retrieved from the study of Huot et al. (2019) in which three independent
1109 experiments were performed with N=9 larvae exposed to 150 NBCs or to a Ringer sterile solution in each
1110 sample. Differential expression between the infested and the control conditions was analyzed with the
1111 Sleuth software (following pseudoalignment with the Kallisto software) using statistical thresholds of
1112 0.01 for q-value (equivalent of adjusted p-value), -1 and +1 for Beta value (biased equivalent of log2 fold
1113 change) and 5 for pseudocount means. The immune genes were identified by homology and classified as
1114 (A) antimicrobial immunity-related, (B) melanization-related, (C) cellular immunity-related and (D)
1115 diverse immune responses. Black dots indicate genes with statistically non-significant variations to the
1116 controls in the corresponding tissue; HC : Hemocytes, FB : Fat body.

1117 **Fig 2.** Temporal dynamics of the identified immune responses after infestation by the nematobacterial
1118 complex. Heatmaps showing the temporal evolution of the induction levels of representative immune
1119 genes in the hemocytes (A) and in the fat body (B) after infestation by the NBC. Three independent
1120 experiments were performed with N=9 larvae exposed to 150 NBCs and N=9 larvae exposed to Ringer
1121 sterile solution. RT-qPCR experiments were performed in triplicate and the RpL32 housekeeping gene
1122 was used as reference for relative quantifications. Differential expression between the infested and the
1123 control conditions was assessed according to the method of Ganger et al. (2017) and with Student t tests
1124 on ΔCq . Black dots indicate genes with statistically non-significant variations to the controls in the
1125 corresponding tissue (p-value > 0.05). The dendrograms represent clustering analyses based on Pearson
1126 correlation coefficients.

1127 **Fig 3.** Relative participations of *S. carpocapsae* and *X. nematophila* in the hemocytes' immune responses.
1128 Histograms showing the induction levels (+/- SEM) of representative immune genes in the hemocytes at
1129 13 h after independent injections of either whole NBCs, axenic nematodes or bacterial symbionts. Three
1130 independent experiments were performed with N=9 control larvae and N=9 larvae injected with 10 NBCs
1131 (NBC), 10 axenic *S. carpocapsae* (*S.c.*) or 200 *X. nematophila* (*X.n.*). For NBC and axenic nematode
1132 injections control larvae were injected with 70 % Ringer - 30 % glycerol sterile solutions whereas they
1133 were injected with sterile PBS for bacterial injections. RT-qPCR experiments were performed in triplicate
1134 and the RpL32 housekeeping gene was used as reference for relative quantifications. Differential
1135 expression values between the infected and the corresponding control conditions were calculated
1136 according to the method of Ganger et al. (2017) and statistical differences between the three types of
1137 infection were assessed by one-way ANOVA and Tukey tests on $\Delta\Delta Cq$. Letters indicate the statistical
1138 groups resulting from the Tukey tests (p-value > 0.05). The genes were then gathered by type of immune

1139 response with (A) antimicrobial immunity-related, (B) melanization-related, (C) cellular immunity-
1140 related and (D) diverse immune responses.

1141 **Fig 4.** Relative participations of *S. carpocapsae* and *X. nematophila* in the fat body's immune responses.
1142 Histograms showing the induction levels (+/- SEM) of representative immune genes in the fat body at 13
1143 h after independent injections of either whole NBCs, axenic nematodes or bacterial symbionts. Three
1144 independent experiments were performed with N=9 control larvae and N=9 larvae injected with 10 NBCs
1145 (NBC), 10 axenic *S. carpocapsae* (*S.c.*) or 200 *X. nematophila* (*X.n.*). For NBC and axenic nematode
1146 injections control larvae were injected with 70 % Ringer - 30 % glycerol sterile solutions whereas they
1147 were injected with sterile PBS for bacterial injections. RT-qPCR experiments were performed in triplicate
1148 and the RpL32 housekeeping gene was used as reference for relative quantifications. Differential
1149 expression values between the infected and the corresponding control conditions were calculated
1150 according to the method of Ganger et al. (2017) and statistical differences between the three types of
1151 infection were assessed by one-way ANOVA and Tukey tests on $\Delta\Delta Cq$. Letters indicate the statistical
1152 groups resulting from the Tukey tests (p-value > 0.05). The genes were then gathered by type of immune
1153 response with (A) antimicrobial immunity-related, (B) melanization-related and (C) diverse immune
1154 responses.

1155 **Fig 5.** Transcriptional induction patterns of putative new immune genes. (A, C) Histograms showing the
1156 induction levels (+/- SEM) of 2 GBH genes in the hemocytes (A) and of the 5 Unk genes in the fat body
1157 (C) at several times after infestation by the NBC. Three independent experiments were performed with
1158 N=9 larvae exposed to 150 NBCs and N=9 larvae exposed to Ringer sterile solution. RT-qPCR
1159 experiments were performed in triplicate and the RpL32 housekeeping gene was used as reference for
1160 relative quantifications. Differential expression between the infested and the control conditions was
1161 assessed according to the method of Ganger et al. (2017) and with Student t tests on ΔCq . Black dots
1162 indicate genes with statistically non-significant variations to the controls (p-value > 0.05). (B, D)
1163 Histograms showing the induction levels (+/-SEM) of 2 GBH genes in the hemocytes (B) and of the 5
1164 Unk genes in the fat body (D) at 13 h after independent injections of either whole NBCs, axenic
1165 nematodes or bacterial symbionts. Three independent experiments were performed with N=9 control
1166 larvae and N=9 larvae injected with 10 NBCs (NBC), 10 axenic *S. carpocapsae* (*S.c.*) or 200 *X.*
1167 *nematophila* (*X.n.*). For NBC and axenic nematode injections control larvae were injected with 70 %
1168 Ringer - 30 % glycerol sterile solutions whereas they were injected with sterile PBS for bacterial
1169 injections. RT-qPCR experiments were performed in triplicate and the RpL32 housekeeping gene was
1170 used as reference for relative quantifications. Differential expression values between the infected and the
1171 corresponding control conditions were calculated according to the method of Ganger et al. (2017) and
1172 statistical differences between the three types of infection were assessed by one-way ANOVA and Tukey
1173 tests on $\Delta\Delta Cq$. Letters indicate the statistical groups resulting from the Tukey tests (p-value > 0.05).

1174 **Fig 6.** Hypothetical structure of the *S. frugiperda* larva's immune response to the NBC. Graphical
1175 abstract illustrating the main hypotheses we can emit from the present RNAseq and RT-qPCR data and
1176 from our current knowledge of *S. frugiperda* immunity. Dark green letters, lines and arrows indicate
1177 responses that seem to be mainly induced by the nematode partner *S. carpocapsae* whereas orange ones
1178 indicate responses that seem to be mainly induced by the bacterial symbiont *X. nematophila*. The arrows'
1179 thicknesses and the letter sizes refer to the relative strengths of the induced transcriptional responses.
1180

1181 **Supporting Information Legends**

1182 **Supplementary Table 1. Primers sequences and genes used in this study.**

1183 **Supplementary Table 2. Hemocytes and fat body RNAseq results for *S. frugiperda*'s immune genes.**

1184 RNAseq raw data were retrieved from the study of Huot et al. (2019) in which three independent
1185 experiments were performed with N=9 larvae exposed to 150 NBCs or to a Ringer sterile solution for 15
1186 hours. Following pseudoalignment with the Kallisto software, differential expression between the infested
1187 and the control conditions was analyzed with the Sleuth software using statistical thresholds of 0.01 for q-
1188 value (equivalent of adjusted p-value), -1 and +1 for Beta value (biased equivalent of log2 fold change)
1189 and 5 for pseudocount means. The statistics corresponding to the transcripts that were considered as
1190 significantly up- or down-regulated are highlighted in red and blue, respectively. The normalized
1191 pseudocounts are also indicated for each individual sample, with HCn15 and FBn15 corresponding to
1192 control larvae and HCi15 and FBi15 corresponding to infested larvae. Blast hits on the *Drosophila* and nr
1193 NCBI databases were obtained by blastx with the Blast2GO software. (A) Immune genes previously
1194 annotated in the *S. frugiperda*'s genome. (B) Newly identified *S. frugiperda*'s immune genes.

1195 **Supplementary Table 3. Hemocytes and fat body RNAseq results for the Unk and GBH putative**

1196 **new immune genes.** RNAseq raw data were retrieved from the study of Huot et al. (2019) in which three
1197 independent experiments were performed with N=9 larvae exposed to 150 NBCs or to a Ringer sterile
1198 solution for 15 hours. Following pseudoalignment with the Kallisto software, differential expression
1199 between the infested and the control conditions was analyzed with the Sleuth software using statistical
1200 thresholds of 0.01 for q-value (equivalent of adjusted p-value), -1 and +1 for Beta value (biased
1201 equivalent of log2 fold change) and 5 for pseudocount means. The statistics corresponding to the
1202 transcripts that were considered as significantly upregulated are highlighted in red. The normalized
1203 pseudocounts are also indicated for each individual sample, with HCn15 and FBn15 corresponding to
1204 control larvae and HCi15 and FBi15 corresponding to infested larvae.

1205 **Supplementary Fig 1. Verification of *S. carpocapsae* axenicity.** Electrophoresis gel showing the
1206 absence of bacterial contaminants in the axenized nematodes used for experimental infections. Total
1207 DNA from ground infective stage nematodes was extracted a few hours after each experimental infection
1208 (fresh axenic *S.c.*) and after several days of storage without antibiotics (stored axenic *S.c.*). The absence
1209 of bacterial contaminations was assessed by agarose gel electrophoresis after PCR amplification of the
1210 16S rRNA gene with universal primers and of the *Xenorhabdus*-specific XNC1_0073 gene (encoding a
1211 putative TonB-dependent heme-receptor). Whole NBCs (NBC) and a pure suspension of *X. nematophila*
1212 (*X.n.*) were used as positive controls. A pure suspension of *Pseudomonas protegens* (*P.p.*) was used as
1213 negative control of putative TonB-dependent heme-receptor amplification. The results indicate no
1214 bacterial contamination is detected in the sterile axenic nematodes used for experimental infections and

1215 that even after several days of storage without antibiotics, the *X. nematophila* population did not recover
1216 within these nematodes.

1217 **Supplementary Fig 2. Temporal monitoring of nematobacterial infestation parameters.** (A) Dotplot
1218 showing the number of *S. carpocapsae* nematodes detected in the midgut alimentary bolus at several
1219 times after contact with 150 NBCs. Dot colors correspond to 3 independent experiments on N=3 larvae
1220 per time point. (B) Curve showing the temporal evolution of *X. nematophila* concentration (+/-SEM) in
1221 the hemolymph across the time post-infestation. Three independent infestation experiments were
1222 performed with 3 pools of 3 larvae per time point. Hemolymph was collected by bleeding and *X.*
1223 *nematophila* was quantified by CFU counting on selective culture medium containing erythromycin. (C)
1224 Curve showing the temporal evolution of *S. frugiperda* larvae survival percentage (+/- SEM) across the
1225 time post-infestation. Three independent infestation experiments were performed on N=20 larvae per
1226 experiment.

1227 **Supplementary Fig 3. Comparison of the main infection parameters after independent injections of**
1228 **the nematobacterial complex, of axenic *S. carpocapsae* and of *X. nematophila*.** (A) Curves showing
1229 the temporal evolution of *X. nematophila* concentration (+/-SEM) after independent injections of either
1230 10 NBCs or 200 *X. nematophila* (*X.n.*). Three independent injection experiments were performed with 3
1231 pools of 3 larvae per time point. Hemolymph was collected by bleeding and *X. nematophila* was
1232 quantified by CFU counting on selective culture medium containing erythromycin. (B) Curves showing
1233 the temporal evolution of *S. frugiperda* larvae survival percentage (+/- SEM) after independent injections
1234 of either 10 NBCs , 10 axenic *S. carpocapsae* (*S.c.*) or 200 *X. nematophila* (*X.n.*). Three independent
1235 injection experiments were performed on N=20 larvae per experiment. No insect death was reported for
1236 control buffer-injected larvae. (C) Histogram showing the parasitic success (+/- SEM) (i.e.: number of
1237 larvae with NBC emergence on total number of infested larvae) after independent injections of either 10
1238 NBCs or 10 axenic *S. carpocapsae* (*S.c.*). Three independent injection experiments were performed on
1239 N=20 larvae per infection type and per experiment.

1240 **Supplementary Fig 4. Alignment of deduced amino acid sequences of Unks from *S. frugiperda* with**
1241 **those of *S. litura* and *S. littoralis*.** Nucleotide sequences were retrieved by blastn on *S. litura* and *S.*
1242 *littoralis* genomes.

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6

