



**HAL**  
open science

## Human societies began to play a significant role in global sediment transfer 4,000 years ago

Jean-Philippe Jenny, Sujan Koirala, Irene Gregory-Eaves, Pierre Francus,  
Bernhard Ahrens, Victor Brovkin, Antti Ojala, Bernd Zolitschka, Juergen  
Bader, Nuno Carvalhais

► **To cite this version:**

Jean-Philippe Jenny, Sujan Koirala, Irene Gregory-Eaves, Pierre Francus, Bernhard Ahrens, et al.. Human societies began to play a significant role in global sediment transfer 4,000 years ago. *Proceedings of the National Academy of Sciences of the United States of America*, 2020, 117 (11), pp.5571-5572. 10.1073/pnas.1922723117 . hal-02527731

**HAL Id: hal-02527731**

**<https://hal.inrae.fr/hal-02527731v1>**

Submitted on 2 Jul 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0  
International License

## Reply to Li et al.: Human societies began to play a significant role in global sediment transfer 4,000 years ago

Jenny J-P., S. Koirala, I. Gregory-Eaves, P. Francus, C. Niemann, B. Ahrens, V. Brovkin, A. Baud, A. E. K. Ojala, A. Normandeau, B. Zolitschka, J. Bader, N. Carvalhais

In a recent study (1), we conducted a global synthesis of long term records of reconstructed sediment accumulation rates (SAR) in lakes to diagnose that a dominant anthropogenic imprint on soil erosion rates started ~4,000 years ago. Our approach was based on the assumption that lake SARs are watershed integrators of terrestrial biogeochemical responses to atmospheric (climate) and land surface dynamics spanning several millennia (2–4). Our statistical analysis was based on SAR reconstructions for the last 12,000 years (1), global pollen records spanning the same period to infer past vegetation changes (5), and model-based climate reconstructions from transient Holocene simulation with MPI-ESM1.2-LR developed at the Max Planck Institute for Meteorology (6), which has been previously used for analysis of coupled climate-carbon-vegetation dynamics (7).

Li et al. (8) suggest that complex 'human-vegetation-climate' interactions in the Late Holocene may have biased pollen inferred climate reconstructions. Hence, these pollen records may not be appropriate in our study to reconstruct climate variations in addition to land cover changes. Climate can be reconstructed from pollen records (see e.g., 9), but we have chosen not applying those data because they were already used in our study for land cover reconstruction. Yet, to avoid circularity and standardize our approach, climate time series for the past 8,000 years (i.e., average, minimum and maximum of annual precipitation, surface air temperature, and wind) were instead collected for all sites from model simulation with the MPI-ESM1.2-LR, which does not include or assimilate the pollen records.

Li et al. (8) are questioning the climate model selection in our study Forcing (greenhouse gas vs orbital) is minor, consequently climate changes are also minor in all models. Other model reconstructions can be investigated, but regardless of which climate reconstructions are used, the climate changes over the last 4,000 years are minor. Thus, the anthropogenic factor (i.e., significant tree cover reduction synchronous to the increases in SAR trends) emerges as the dominant driver of SARs.

In our study we show that pollen records from European and North American lakes have been affected by human activities during the Holocene, hence recording a combination of a climate and a human signal. Li et al. suggest that pollen records from European and North American sites are less appropriate for climate reconstructions compared with records from remote areas. This discussion goes beyond the conclusions of our paper, but our subset of sites is including lake systems only, hence representing only a small fraction of all pollen archives stored in the pollen database, e.g. not including bogs, ponds, swamps, soils, estuaries or alluvial systems. Hence, our subset of pollen records is not representative of the total pollen records available and seems not appropriate to discuss the climate reconstruction methodologies. Other papers discuss climate reconstruction methodologies from pollen records (9), but this was not the scope of our paper.

### References

1. J.-P. Jenny, *et al.*, Human and climate global-scale imprint on sediment transfer during the Holocene. *PNAS* (2019) <https://doi.org/www.pnas.org/cgi/doi/10.1073/pnas.1908179116>.
2. K. J. Edwards, G. Whittington, Lake sediments, erosion and landscape change during the Holocene in Britain and Ireland. *CATENA* **42**, 143–173 (2001).
3. J. A. Dearing, Lake sediment records of erosional processes. *Hydrobiologia* **214**, 99–106 (1991).
4. C. E. Williamson, J. E. Saros, W. F. Vincent, J. P. Smol, Lakes and reservoirs as sentinels, integrators, and regulators of climate change. *Limnol. Oceanogr.* **54**, 2273–2282 (2009).

5. A. C. Martin, W. J. Harvey, The Global Pollen Project: a new tool for pollen identification and the dissemination of physical reference collections. *Methods Ecol. Evol.* **8**, 892–897 (2017).
6. T. Mauritsen, *et al.*, Developments in the MPI-M Earth System Model version 1.2 (MPI-ESM1.2) and its response to increasing CO<sub>2</sub>. *J. Adv. Model. Earth Syst.* (2018) (November 22, 2018).
7. V. Brovkin, *et al.*, What was the source of the atmospheric CO<sub>2</sub> increase during the Holocene? *Biogeosciences* **16**, 2543–2555 (2019).
8. Y. Li, *et al.*, Complex ‘human-vegetation-climate’ interactions in the late Holocene and their significance for paleotemperature reconstructions. *Proc. Natl. Acad. Sci.* (2020).
9. B. A. S. Davis, S. Brewer, A. C. Stevenson, J. Guiot, The temperature of Europe during the Holocene reconstructed from pollen data. *Quat. Sci. Rev.* **22**, 1701–1716 (2003).