



HAL
open science

SCHENGEN receptor module drives localized ROS production and lignification in plant roots.

Satoshi Fujita, Damien de Bellis, Kai Edel, Philipp Köster, Tonni Grube Andersen, Emanuel Schmid-Siegert, Valérie Dénervaud Tendon, Alexandre Pfister, Peter Marhavý, Robertas Ursache, et al.

► To cite this version:

Satoshi Fujita, Damien de Bellis, Kai Edel, Philipp Köster, Tonni Grube Andersen, et al.. SCHENGEN receptor module drives localized ROS production and lignification in plant roots.. EMBO Journal, 2020, 39 (9), pp.e103894. 10.15252/emj.2019103894 . hal-02529862

HAL Id: hal-02529862

<https://hal.inrae.fr/hal-02529862v1>

Submitted on 21 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

ROS production by localized SCHENGEN receptor module drives lignification at subcellular precision

Authors: Satoshi Fujita^{1, †, *}, Damien De Bellis^{1,2}, Kai H. Edel³, Philipp Köster^{3, †}, Tonni Grube Andersen^{1, ‡}, Emanuel Schmid-Siegert⁴, Valérie Dénervaud Tendon¹, Alexandre Pfister¹, Peter Marhavý^{1, ††}, Robertas Ursache¹, Verónica G. Doblas^{1, §}, Marie Barberon^{1, §§}, Jean Daraspe^{1,2}, Audrey Creff⁵, Gwyneth Ingram⁵, Jörg Kudla³ and Niko Geldner^{1,*}

Affiliations:

1 Department of Plant Molecular Biology, Biophore, Campus UNIL-Sorge, University of Lausanne, CH-1015 Lausanne, Switzerland

2 Electron Microscopy Facility, University of Lausanne, 1015 Lausanne, Switzerland

3 Institut für Biologie und Biotechnologie der Pflanzen, Westfälische Wilhelms-Universität Münster, Schlossplatz 7, Münster 48149, Germany

4 Vital-IT Competence Center, Swiss Institute of Bioinformatics, Lausanne, Switzerland

5 Laboratoire Reproduction et Développement des Plantes, Univ Lyon, ENS de Lyon, CNRS, INRAE, Lyon, France

‡ Present address: National Institute of Genetics, 1111 Yata, Mishima, Shizuoka 411-8540, Japan

† Present address: Department of Plant and Microbial Biology, University of Zurich, Zollikerstrasse 107, CH-8008 Zurich, Switzerland

†† Present address: Umeå Plant Science Centre (UPSC), Department of Forest Genetics and Plant Physiology, Swedish University of Agricultural Sciences (SLU), 90183, Umeå, Sweden.

‡ Present address: Max Planck Institute for Plant Breeding Research, 50829 Cologne, Germany

§ Present address: Institut Jean-Pierre Bourgin, INRAE, AgroParisTech, Université Paris-Saclay, 78000, Versailles, France

§§ Present address: Department of Botany and Plant Biology Quai Ernest-Ansermet 30 CH-1211 Geneva Switzerland

*Correspondence to: satoshi.fujita@nig.ac.jp (S.F.) and niko.geldner@unil.ch (N.G.)

Abstract:

Production of reactive-oxygen species (ROS) by NADPH oxidases (NOXs) impacts many processes in animals and plants and many plant receptor pathways involve rapid, NOX-dependent increases of ROS. Yet, their general reactivity has made it challenging to pinpoint the precise role and immediate molecular action of ROS. A well-understood ROS action in plants is to provide the co-substrate for lignin peroxidases in the cell wall. Lignin can be deposited with exquisite spatial control, but the underlying mechanisms have remained elusive. Here we establish a kinase signaling relay that exerts direct, spatial control over ROS production and lignification within the cell wall. We show that polar localization of a single kinase component is crucial for pathway function. Our data indicates that an intersection of more broadly localized components allows for micrometer-scale precision of lignification and that this system is triggered through initiation of ROS production as a critical peroxidase co-substrate.

As in animals, NADPH oxidase-produced ROS in plants is important for a multitude of processes and the number of NADPH oxidase genes (10 in Arabidopsis, called *RESPIRATORY BURST OXIDASE HOMOLOGs*, *RBOHs*, *A-J*) suggests a high complexity of regulation of ROS production in plants. Among its many roles, ROS-dependent regulation of plant cell wall structure and function is considered to be among its most important (Kärkönen & Kuchitsu, 2015). The cell wall is the nano-structured, sugar-based, pressure-resisting extracellular matrix of plants and NOXs are thought to be the predominant ROS source in this compartment (also termed apoplast) (Kärkönen & Kuchitsu, 2015).

48 A staggering number of kinases have been shown to regulate plant NOXs and the activation mechanism
49 of NOX-dependent ROS production is well established, especially in response to microbial pattern-recognition
50 by immune receptors (Zipfel, 2014). However, the specific role and direct molecular targets of ROS during
51 microbial pattern-recognition have remained elusive (Qi *et al*, 2017). The same applies to the central role of ROS
52 in tip growing cells, such as root hairs or pollen tubes, where ROS is thought to be part of an intricate oscillation
53 of cell wall stiffening and loosening, aimed at allowing cell wall expansion without catastrophic collapse
54 (Boisson-Dernier *et al*, 2013; Monshausen *et al*, 2007). In this case, ROS is proposed to be important for
55 counteracting cell wall loosening pH decreases, but it is again unclear what direct targets of ROS would mediate
56 cell wall stiffening. Cell wall lignification by apoplastic peroxidases, can therefore be considered as the most
57 well-established role of ROS, where the peroxidases themselves are the direct “ROS targets”, using it as a co-
58 substrate for the oxidation of mono-lignols (Barbosa *et al*, 2019; Liu, 2012). In the case of lignification, however,
59 a molecularly-defined signaling pathway that induces ROS production during lignification has not been defined.
60 A few years ago, our group identified a specific NADPH oxidase, RBOHF, to be required for the localized
61 formation of lignin in the root endodermis (Lee *et al*, 2013). Lignin is a poly-phenolic polymer that is generated
62 by the radical-coupling of mono-lignols, oxidized through the action of ROS-dependent peroxidases, as well as
63 laccases (Liu, 2012). The hydrophobic lignin polymer impregnates the cellulosic cell wall of plants, rendering it
64 unextendible and highly resistant to degradation. Lignin in the root endodermis is deposited in a central,
65 longitudinal band around every endodermal cell. Named Casparian strips (CS), these ring-like lignin structures
66 fuse into a supracellular network, establishing a tissue-wide, extracellular diffusion barrier (Fig. 1A), analogous
67 to epithelial tight junctions in animals (Geldner, 2013; Barberon & Geldner, 2014). Functionality of this barrier
68 can be easily visualized by a block of penetration of a fluorescent cell wall dye, propidium iodide, into the
69 vasculature (Alassimone *et al*, 2010; Naseer *et al*, 2012). CS localization occurs through the action of
70 CASPARIAN STRIP DOMAIN PROTEINS (CASPs), 4-TM proteins, which form a highly scaffolded
71 transmembrane protein platform, assembling RBOHF and cell wall peroxidases and other proteins at the
72 Casparian strip domain (CSD) (Lee *et al*, 2013; Hosmani *et al*, 2013a; Roppolo *et al*, 2011). CSD formation and
73 lignin deposition are coordinated such that the aligned rings of endodermal neighbors’ fuse, leading to a
74 supracellular network that seals the extracellular space between endodermal cells, generating a tissue-wide
75 diffusion barrier.

76 More recently, we identified a pair of peptide ligands, a leucine-rich repeat receptor-like kinase (LRR-
77 RLK) and a cytoplasmic kinase, whose phenotypes, genetic interaction and specific subcellular localizations led
78 us to propose that they combine into a barrier surveillance pathway. Previous reports had shown that the CIF1/2
79 (CASPARIAN STRIP INTEGRITY FACTORS 1/2) peptides are SCHENGEN3 (SGN3) (also called GASSHO1
80 (GSO1)) ligands and that the SGN1 and SGN3 kinases govern Casparian strip integrity (Alassimone *et al*, 2016;
81 Doblbas *et al*, 2017; Nakayama *et al*, 2017; Pfister *et al*, 2014). *cif1 cif2* and *sgn1, sgn3* mutants have similar,
82 discontinuous CS, caused by a discontinuous CSD, as well as a conspicuous absence, or strong attenuation of,
83 compensatory lignification and suberization observed in other CS mutants (Doblbas *et al*, 2017; Hosmani *et al*,
84 2013a; Kalmbach *et al*, 2017; Kamiya *et al*, 2015; Li *et al*, 2017; Pfister *et al*, 2014). Their phenotypic similarities
85 suggested that these factors act in one pathway. CIF1 and 2 peptides do not express in the endodermis, where
86 the CS is formed, but in the stele. In contrast, both SGN1 and SGN3 present specific localization on the
87 endodermal plasma membrane; SGN3 receptor-like kinase resides along both sides of the CS, while
88 palmitoylated SGN1 polarly localizes on cortex facing (outer) plasma membranes (Doblbas *et al*, 2017).
89 Remarkably, their localization overlaps only in a small region next to the cortex-facing side of the CS
90 (Alassimone *et al*, 2016). This would require peptides from the stele to diffuse across the CS in order to access
91 the signaling complex. This is only possible while the CS is still permeable (Doblbas *et al*, 2017) (Fig. 1E).

92 This pathway would therefore provide a mechanism that allows to probe the tissue-wide integrity of
93 an extracellular diffusion barrier and to respond to barrier defects through compensatory over-lignification
94 (Doblbas *et al*, 2017). Here we demonstrate that the receptor, cytoplasmic kinase and NADPH oxidase
95 molecularly connect into one pathway, with great resemblance to plant immune signaling pathways, but whose
96 direct action is to locally produce ROS for spatially restricted lignification. We establish the crucial importance
97 of the restricted subcellular localization of its components and demonstrate that stimulation of this signaling
98 pathway additionally leads to strong transcriptional activation of target genes that further drive and sustain

99 endodermal lignification, as well as suberization and endodermal sub-domain formation and differentiation. We
100 thus provide a molecular circuitry in which an endogenous peptide from the stele stimulates localized signaling
101 kinases and NADPH oxidases in the endodermis, causing extracellular ROS production at micrometer-scale
102 precision and a precisely localized lignification of the plant cell wall.

103 RESULTS

104 We generated a new *cif1-2 cif2-2* double mutant allele by CRISPR-Cas9 in a pure Col background,
105 because the previous *cif1-1 cif2-1* double mutant allele (Nakayama *et al*, 2017) was a mixture of Ws and Col
106 alleles. The new CRISPR allele was complemented by CIF1 or CIF2 application, visualized by PI (propidium
107 iodide) uptake assays or reconstitution of CASP1 membrane domain connectivity (Fig. EV1 D,E and Appendix
108 Fig. S1A).

109 Apolar SGN1 kinase leads to constitutive barrier defect signaling

111 Central to the barrier surveillance model is the polar localization of SGN1, which is thought to limit
112 the potential for signal activation to the cortex side of the endodermis, requiring passage of CIF peptides across
113 the CS region (Fig. 1A,E). Consistently, application of peptide ligand to the media, leading to stimulation from
114 the outside, causes overlignification at the cortex-facing endodermal edges (Fig. 1B). In an attempt to falsify the
115 model we had proposed and to interrogate the importance of SGN1 polar localization, we generated a SGN1
116 variant that localized in an apolar fashion, by adding a myristoylation (*myr*) and palmitoylation (*palm*) motifs
117 on the N terminus (Vermeer *et al*, 2004). This myrpalm SGN1-mCitrinerine (Citrine) variant was expressed
118 under the control of the endodermis-specific CASP1 promoter, which is strongly active during Casparian strip
119 formation and complemented the *sgn1* barrier phenotype (Fig. EV 1A). *In planta*, the wild-type SGN1-Citrine
120 variant resides polarly on the cortex-facing side of endodermal cells, while myrpalm SGN1-Citrine was observed
121 at both sides of the endodermal plasma membranes, even though preferentially accumulation at the cortex side
122 could still be observed (Fig. 1C, 1D). Both variants were excluded from the central position where the Casparian
123 strip domain is formed (Fig. EV 1B) and the localization patterns of the two variants did not change when
124 introgressed into *sgn1*, *sgn3* or *cif1 cif2* mutants (Fig. 1C, 1D). An apolar SGN1 localization would allow SGN3
125 to encounter SGN1 also on the stele-facing side, not only on the cortex side (Fig. 1E). This should lead to
126 constitutive signal activation in the absence of barrier defects, because the CIF peptides would now be able to
127 access a SGN3/SGN1 signaling module on the stele-facing side without crossing the barrier. Indeed, we found
128 that the apolar SGN1 variant caused both, ectopic lignin deposition and precocious suberization in endodermal
129 cells (Figs. 1D, EV 1F,G), as previously described for endodermal barrier mutants. Yet, no barrier defect was
130 observed in the lines complemented with apolar SGN1 (Fig. EV 1A) and, consistently, we found CASP1-
131 mCherry distribution to be normal in these lines, forming a continuous band in the central position of the
132 endodermal cells, indistinguishable from wild-type (Fig. EV 1B). This indicates that presence of SGN1 at the
133 plasma membrane to the inside of the CS leads to signaling in the absence of barrier defects. CASP1 promoter-
134 driven wild type SGN1 lines as a control, complemented the mutant (Fig. EV 1A) and did not cause any changes
135 in lignin accumulation pattern (Fig. 1C). Interestingly, the apolar SGN1 lines accumulated lignin mainly on the
136 stele-facing edges of the endodermal cell walls, as opposed to the cortical lignin deposition observed by ectopic
137 ligand treatment (compare Fig. 1B with Fig. 1D, arrows). The ectopic lignin deposition in apolar SGN1 lines
138 was fully dependent on the presence of receptor and ligand, as both in *sgn3* and *cif1 cif2* mutants, no excess
139 lignification at the stele-facing side could be observed in apolar SGN1 lines (Fig. 1D). This strongly suggests
140 that mislocalized SGN1 does not become constitutively active, but leads to continuous, ectopic transduction of
141 CIF1/2 signals through SGN3. An apolar, but kinase-dead variant of SGN1 was also unable to induce ectopic
142 lignification (Fig. EV 1C), suggesting that a phosphorylation relay downstream of SGN1 is necessary for
143 lignification.

144 SGN1 is a downstream component of the CIF/SGN3 pathway

145
146 Previous data showed that *sgn1* is less sensitive to high doses of externally applied CIF peptide,
147 consistent with a role of SGN1 downstream of the SGN3 receptor (Doblas *et al*, 2017), at least with respect to

148 CIF-induced excess lignification. In order to address whether SGN1 is indeed a generally required downstream
149 component of the CIF/SGN3 pathway during CS formation, we evaluated whether the *sgn1* mutant is resistant
150 to complementation by CIF2 peptide treatment. In contrast to the full complementation of the discontinuous CS
151 domain of *cif1 cif2* double mutants, the *sgn1* mutant did not show full restoration of domain integrity, neither on
152 10 nM nor 100 nM CIF2 medium (Fig. 2A, Appendix Fig. S1A). While some degree of rescue occurred, only
153 about 50% of discontinuities were rescued. This was corroborated by testing CS functionality using the
154 propidium iodide (PI) assay. Only a weak complementation of barrier formation was observed even when grown
155 on 100 nM CIF2 medium (Fig. 2B). Our results indicate that SGN1 functions downstream of CIFs/SGN3, but
156 suggest that additional factors can partially compensate for its absence, most probably homologs of the extended
157 RLCKVII family to which SGN1 belongs.

158
159 We then tested for direct connectivity between SGN3 receptor kinase and SGN1 by carrying out an *in*
160 *vitro* kinase assay. Some RLCKVII members are reported to be phosphorylated and activated by LRR receptor
161 kinases (Kim *et al.*, 2011, 1; Lu *et al.*, 2010, 1). A Glutathione S-transferase (GST)-fused SGN3 kinase domain
162 was incubated with a kinase-dead form of a trigger factor (TF)-SGN1 fusion protein - which does not have
163 autophosphorylation activity - in the presence of radioactive ATP. Kinase-dead TF-SGN1 was efficiently
164 phosphorylated by SGN3 kinase domain, but not by a kinase-dead form of SGN3 *in vitro* (Fig. 2C, Appendix
165 Fig. S1B). This further supports our model that SGN1 is a direct downstream component of the SGN3/CIF
166 pathway.

167 168 **Two NADPH oxidases are absolutely required for CIF-induced lignification in the endodermis**

169
170 Our generation of an apolar SGN1 thus appears to have reconstituted a functional CIF/SGN3 pathway
171 at the stele-facing (inner) endodermal surface, causing ectopic lignification in the absence of barrier defects. Yet,
172 the second intriguing aspect of this manipulation is the observation that lignification occurs almost exclusively
173 at the inner endodermal edges (Fig. 1D), the side where endogenous CIF peptide must be assumed to be present.
174 External treatment, by contrast, leads to predominant lignification at the outer endodermal edges (Fig. 1B). This
175 surprising spatial correlation between the site of signal perception and localized lignification suggests a very
176 direct molecular connection between the two events that would allow to maintain spatial information. In a cell
177 primed for lignification, i.e. with mono-lignol substrates available and polymerizing enzymes expressed,
178 lignification could be simply “switched on” by activating ROS production through NADPH oxidases. Previously,
179 we had found one of the NOXs, RBOHF, to be crucial for lignification at the CS. Intriguingly, RBOHF is the
180 only transmembrane protein known to accumulate at the Casparian strip domain, save the CASPs themselves
181 (Lee *et al.*, 2013) (Fig. 3A). Moreover, homologous NADPH oxidases, such as RBOHB or RBOHD, also present
182 in the endodermal plasma membrane, are excluded from this domain (Lee *et al.*, 2013) (Fig. 3A). We therefore
183 asked whether CIF peptides would induce lignification through activation of RBOHF as a downstream
184 component. To our surprise, CIF treatment still led to induced lignification in *rbohF*, despite the fact that RBOHF,
185 is strictly required for CS lignification in untreated conditions. Yet, when RBOHD was knocked-out in addition
186 to RBOHF, a complete absence of lignification upon CIF treatment was observed (Fig. 3B). The fact that
187 RBOHD single mutant neither showed defects in developmental CS lignification, nor in CIF-induced
188 lignification, indicates that RBOHF is required for both processes, but that upon strong stimulation with
189 exogenous CIF peptide application, RBOHD can be additionally used. Indeed, using the general NADPH
190 oxidase inhibitor diphenyleneiodonium (DPI) in short-term co-treatment with CIF was also able to fully block
191 CIF-induced ectopic lignification in wild-type with intact Casparian strips, further supporting that NADPH
192 oxidases act downstream in the CIF/SGN3 pathway (Fig. EV 2A).

193 194 **CIF2 induces highly localized ROS production through RBOHF and RBOHD**

195
196 The direct activity of NOX enzymes is not lignification, but production of superoxide (O_2^-) that becomes
197 dismutated to hydrogen peroxide (H_2O_2) in the apoplast (Kärkönen & Kuchitsu, 2015). We had previously
198 established that the different subcellular distribution of the SGN3 receptor and the SGN1 kinase only intersect
199 at a very restricted domain at the outer (cortex-facing) edge of the CS (Alassimone *et al.*, 2016)(Fig. 4A). We
200 therefore attempted to visualize whether ROS might be produced locally in response to CIF treatment. Many

ways exist to visualize and quantify ROS, but only few allow for high spatial resolution and for discrimination between extracellular and intracellular ROS. An older method, based on ROS-induced cerium precipitation that can be detected using transmission electron microscopy (TEM), has been used by us previously to demonstrate that a highly localized ROS production indeed occurs at the CS and is dependent on NOX activity (Bestwick *et al*, 1997a; Lee *et al*, 2013)(Fig. 4B). Using this method, we could detect a strong ROS production in response to CIF2, exclusively at the regions of endodermal-endodermal cell walls outside of the CS, but nowhere along the endodermis-cortex cell walls, which are equally reached by the cerium chloride and where NOX enzymes are also present in the plasma membrane (Fig. 4B,C). Based on this striking spatial coincidence between the SGN3/SGN1 overlap region (Fig. 4A) (Allassimone *et al*, 2016) and CIF2-induced ROS production, we developed a procedure to quantitatively assess cerium precipitates and checked whether this localized ROS production is indeed dependent on the SGN3 pathway (Figs. 4D,E, EV 3A, see also Experimental Procedures section). For SGN3, we found that already steady-state ROS production was undetectable in the mutant and that there was no increase upon CIF2-treatment (Fig. 4D,E). The *sgn1* mutant showed lower, but still detectable steady state ROS levels, but no significant increase upon CIF2-treatment (Fig. 4D,E). Thus, the highly localized ROS accumulation induced by CIF2 is entirely dependent on the localized SGN3/SGN1 receptor module, suggesting that the localization of the module determines the spatial extent of ROS production. As in the case of lignification, we observed that CIF2-induced ROS could be produced by either RBOHF or RBOHD, as only the double mutant caused a complete absence of ROS after CIF-stimulation. As expected, but not previously demonstrated, the steady-state ROS production at the CS observed before stimulation was exclusively dependent on RBOHF, but not RBOHD (Fig. 4D,E).

In contrast to lignin polymerization, suberin might not require NADPH oxidase activity. It was shown previously that CIF2 triggers excess suberization in WT (Fig. EV 2B,C)(Doblas *et al*, 2017). We found excess suberin deposition in *rbohD* and *rbohF* upon peptide treatment, although a slight enhancement is already observed in *rbohF*. In *rbohDF*, excess suberin deposition is very strong, even without treatment, likely due to a strong activation of the surveillance system due to complete absence of a CS (Fig. EV 2B,C). These data indicate that CIF2-triggered suberin accumulation is not affected by NADPH oxidases and suggests existence of a ROS-independent branch of the SCHENGEN pathway that regulates this process (see below).

SGN1 can directly activate RBOHF and RBOHD via phosphorylation

The above data strongly suggest a direct connection between the SGN3/SGN1 kinase module and the two NOX enzymes. We therefore conducted an *in vitro* kinase assay in order to ask whether SGN1 can directly phosphorylate the N-terminal cytoplasmic region of RBOHF and RBOHD. We found that recombinantly expressed TF-SGN1 could phosphorylate both the recombinant N-terminal part of RBOHF and RBOHD (Fig. 5A). Plant NOX regulation has been intensively studied and shown to be highly complex, requiring possibly interdependent activities of kinases, as well as small GTPases (Kadota *et al*, 2015). We therefore tested whether SGN1 might be sufficient for activation of RBOHF activity in a cellular context. To do so, we made use of a heterologous reconstitution system in human HEK293T cells, which show very low endogenous ROS production and for which it had been previously demonstrated that plant NADPH oxidases can be expressed and their activation mechanism be studied (Han *et al*, 2019). As a positive control the previously described calcium-dependent kinase complex of calcineurin B-like (CBL) interacting protein kinases 26 (CIPK26) and CBL1 was used and shown to be active (Fig. 5B). When expressing wild-type SGN1 in this cell line, we noticed that it did not activate RBOHF or RBOHD, but that it also did not localize to the plasma membrane as in plant cells (Fig. EV 4A). However, when we used the functional, constitutively plasma membrane-localized myrpal-SGN1 version, a significant induction of ROS production was observed for RBOHF and to a lesser extent for RBOHD (Figs. 5B, EV 4B).

CIF peptide-induced CASP domain growth requires new protein synthesis, but not ROS production

The direct phospho-relay from SGN3 receptor, to SGN1 kinase to RBOHF and RBOHD outlined above draws a direct molecular connection from perception of a peptide hormone stimulus to cell wall lignification. Moreover, it accounts for both, the highly localized ROS production that we observe upon CIF

254 stimulation, as well as the observation that localization of lignification is correlated with the site of active SGN3
255 signaling. Yet, the massive enhancement of lignification observed upon CIF stimulation (Fig. 3B), the increase
256 of CASP accumulation and ectopic patch formation (Appendix Fig. S1A), as well as the non-localized formation
257 of precocious and enhanced suberin (Fig. EV 2B,C), are additional outcomes of SGN3 pathway stimulation and
258 should be driven by transcriptional changes. Indeed, some degree of transcriptional upregulation of CASP genes
259 has been reported previously upon CIF1 treatment (Nakayama *et al*, 2017). CIF stimulation does not only lead
260 to enhanced accumulation and ectopic patches of CASP1-GFP, absence of SGN3 signaling also leads to
261 discontinuous CASP1-GFP signals that could be explained by insufficient amount of CASPs and other factors
262 being produced during endodermal differentiation (Fig. 6A) (Doblas *et al*, 2017; Pfister *et al*, 2014). Neither the
263 single, nor the double NADPH oxidase mutants display discontinuous CASP1-GFP signals, indicating that ROS
264 production is not required for this aspect of the CIF/SGN3 pathway (Fig. 6A). In order to directly demonstrate
265 that formation of a continuous CASP domain requires newly formed gene products, we treated our *cif1 cif2*
266 double mutant with CIF2 peptide in the presence or absence of protein synthesis inhibitor cycloheximide (CHX).
267 CASP1-GFP signal strongly increased during 8 hours of CIF treatment, during which the discontinuous CASP1-
268 GFP domains became continuous. This effect was abrogated by CHX treatment (Fig. 6 B,C, Movie EV1).

269 **The NADPH oxidase-independent branch of the CIF/SGN3 pathway is associated with MAP kinase** 270 **stimulation and causes strong activation of gene expression.** 271

272
273 In pattern-triggered immune receptor signaling, gene activation is thought to depend in large parts on
274 activation of Mitogen-activated protein (MAP) kinases (Dodds & Rathjen, 2010). We therefore tested whether
275 CIF-treatment leads to MAP kinase phosphorylation and indeed found that CIFs can induce MAP kinase
276 phosphorylation in a SGN3-dependent manner, further extending the molecular parallels between immune
277 receptor signaling and the CIF/SGN3 pathway and suggesting that gene induction in the CIF pathway might
278 equally depend on MAP kinase signaling (Fig. EV 5B).

279
280 We then undertook an RNA profiling of seedling roots at different time points (30, 120 and 480 min)
281 after CIF stimulation, using wild-type, *cif1 cif2* and *sgn3* mutants as genotypes. 930 genes were found to be
282 differentially expressed across any of the genotypes and time points combined, using a stringent cut-off (adj.-
283 pval. ≤ 0.05 ; $\log_{2}FC \geq 1$ or $\log_{2}FC \leq -1$) (Table EV1). After normalization of batch effects, the three replicates
284 clustered closely with a large degree of variance explained by the peptide treatment in wild-type and *cif1 cif2*
285 (Fig. EV 5A). Wild-type and *cif1 cif2* samples showed nearly identical responses (co-relation efficiency values
286 were 0.90, 0.89, 0.94 at 30, 120, 480 min respectively), with *cif1 cif2* displaying a slightly stronger overall
287 amplitude (Fig. 6D). Importantly, *sgn3* had virtually no differentially expressed genes across treatments,
288 indicating the specificity of the CIF responses and further corroborating that SGN3 is the single, relevant receptor
289 for CIF responses in roots (Figs. 6D, EV 5A, Table EV1). Our data extends on the previous data of Nakayama
290 *et al.* (2017), by showing that all 5 *CASP* genes are differentially regulated upon CIF-treatment (Fig. EV 5C).
291 Moreover, we observed upregulation of *MYB36*, a central transcription factor for Casparian strip formation and
292 *CASP* expression, thus potentially accounting for the increases in *CASP1-4* expression (Kamiya *et al*, 2015;
293 Liberman *et al*, 2015)(Fig. EV 5C). The fact that CIF stimulates *MYB36* expression also is consistent with the
294 recent report that CIF treatment can enhance ectopic endodermal differentiation, driven by overexpression of the
295 SHR transcription factor (Drapek *et al*, 2018).

296
297 Of the four response clusters defined by k-mean clustering, the two largest clusters (1 and 3) group
298 early response genes (cluster 1) and later response genes (cluster 3). Cluster 2 groups a smaller fraction of genes
299 showing strong and largely sustained responses over the time scale of the experiment, while cluster 4 groups a
300 minor fraction of genes downregulated at later time points. GO term analysis indicates that many of the most
301 significant, overrepresented terms in the “early response” (1) and “strong and sustained” (2) gene clusters are
302 related to immune and defense responses (response to chitin, bacterium, callose, etc.), as well as responses to
303 oxidative stress (Table EV2). The top GO categories of the late response cluster (3) were suberin/cutin
304 biosynthesis, supporting our observation that suberin accumulation is a late effect of CIF stimulation (Figs. EV
305 5D, EV 2B,C)(Doblas *et al*, 2017). Additional, enriched terms were related to oxidative stress and cell wall
306 remodeling, nicely fitting the CIF2-induced ROS production we observed. Intriguingly, the cluster showing late

307 downregulation of gene expression contains many overrepresented GO terms related to a wide variety of
308 transport processes, reaching from water, nitrate and ammonium transport, to primary and secondary metabolite
309 or hormone transport (Fig. 6D, Table EV2). We interpret this as a consequence of CIF-induced lignification and
310 suberization, which must profoundly impact endodermal ability for shuttling organic and inorganic compounds,
311 as well as signaling molecules between stele and cortical root cell layers (Barberon & Geldner, 2014).
312

313 While GO terms related to lignification were also overrepresented, they were not as highly ranked as
314 expected from the strong induction of lignification upon CIF treatment. A number of reasons might account for
315 this: Firstly, GO terms for pathogen, defense and oxidative stress responses, are all bound to overlap with and
316 contain genes that mediate lignification. Moreover, already elevated levels of developmental, lignin-related gene
317 expression in the endodermis might lead to less-pronounced fold-changes. Finally, the multiple roles for lignin
318 biosynthetic genes in secondary metabolism further obscure clear categorization. Nevertheless, upregulation of
319 lignin biosynthesis by CIF-treatment became evident when clustering a set of laccases and peroxidases, the two
320 main families of enzymes implicated in lignin polymerization in the apoplast (Liu, 2012). A significant number
321 of both are strongly upregulated (Fig. EV 5E,F). Moreover, *MYB15*, a transcription factor shown to be involved
322 in stress and MAMP(microbe-associated molecular pattern)-induced lignification (Chezem *et al*, 2017, 15) is
323 about 6-fold induced after 30 min and about 17-fold after 120 min in both treated wild-type and *cif1 cif2*, nicely
324 correlating with the upregulation of peroxidases and laccases (Fig. EV 5E,F, Table EV1). We then made use of
325 two of the most highly differentially expressed genes upon CIF2-treatment (*PER15* and *PER49*) in order to
326 establish whether SGN1 and the NADPH oxidases are also required for gene regulation downstream of SGN3
327 activation. In qPCR analysis both peroxidase genes still showed a slight, but significant upregulation upon CIF-
328 treatment in the *sgn1* mutant (compared to the complete absence of response in *sgn3*) (Fig. 6E-H), clearly
329 indicating that, as for ROS production, SGN1 is an important, yet not absolutely required, downstream
330 component in CIF-regulated gene expression. When directly plotting peroxidase fold-inductions in *rbohF*,
331 *rbohD* and double mutant, a seemingly strong attenuation was observed (Fig. 6F,H). However, when
332 independently plotting gene expression in treated and untreated conditions (Fig. 6E,G), this lower fold-induction
333 is largely explained by an already significant basal increase in marker gene expression in *rbohF* and *rbohD*
334 *rbohF* double mutants. Indeed, both mutants display an absence of Casparian strips, which should lead to an
335 endogenous stimulation of the SGN3 pathway, nicely supporting the barrier surveillance model. Absolute
336 expression of both marker genes in the NADPH oxidase mutants is in effect higher than in wild-type after CIF
337 treatment, leading us to conclude that ROS production and gene activation in response to CIF are two
338 independent branches of this signaling pathway, with the branching occurring downstream of the SGN1 kinase.
339 We finally wanted to know whether the mislocalization of the SGN1 kinase reported initially is only affecting
340 CIF/SGN3 signaling at the plasma membrane, or whether it also affects gene activation. We found that both
341 *PER15* (Fig. 6I) and *PER49* (Fig. 6J) are dramatically upregulated in the non-polar SGN1 lines, independent of
342 exogenous peptide application, strongly corroborating our model whereby the SCHENGEN pathway function
343 crucially depends on the correct subcellular localization of its downstream kinase.
344

345 DISCUSSION

346
347 The data presented here sketch out an entire signaling pathway. Previously, SGN3 had been established
348 as the receptor for CIF1 and 2, but its connection to SGN1 had exclusively been based on genetic evidence. Here
349 we show that SGN3, SGN1 and RBOHD/F are, biochemically and functionally, part of a signal transduction
350 chain, leading directly from localized peptide perception to localized ROS production and lignification (Fig. 7A).
351 Moreover, the pathway branches downstream of SGN1, leading to MAP kinase activation and stimulation of
352 gene expression. Some of the most strongly induced genes being peroxidases and laccases that would further
353 enhance and sustain lignification. The SCHENGEN pathway therefore elegantly integrates fast, plasma
354 membrane-based responses that maintain positional information (ROS and lignin are produced close to where
355 the ligand is perceived) and slower, gene expression-based responses, which have lost positional information,
356 but would allow to enhance and maintain the ROS-burst-controlled activation of peroxidases (Fig. 7A).
357

358 The SCHENGEN pathway bears a striking overall resemblance to well-established signaling pathways
359 for perception of MAMPs (Fig. 7B). In MAMP perception, structurally similar receptor kinases, such as FLS2

360 or EFR bind to microbial patterns and interact with the SERK family of co-receptors (Chinchilla *et al*, 2007), as
361 is the case for SGN3 (Okuda *et al*, 2020). The signal is then transduced through kinases of the RLCKVII family,
362 such as BIK1 or PBLs, which are homologs of SGN1 (Liang & Zhou, 2018). BIK1 in turn was shown to
363 phosphorylate RBOHD, driving the well-described MAMP-induced ROS burst (Kadota *et al*, 2014, 1; Li *et al*,
364 2014, 1). Moreover, it has recently been shown that kinases of the RLCKVII family directly phosphorylate
365 MAPKKs, which now mechanistically explains how MAMP perception induces MAP kinase phosphorylation
366 (Bi *et al*, 2018). It is thus tempting to speculate that the SCHENGEN pathway represents an ancient neo-
367 functionalization of an immune receptor pathway. Indeed, this has been proposed recently and it was pointed out
368 that the closest receptor homologs to SGN3 and GSO2 are PEPR1 and PEPR2 (Creff *et al*, 2019). The latter are
369 receptors to an endogenous plant peptides (AtPEPs), whose activities resemble that of MAMPs and are best
370 thought of as “phyto cytokines”, i.e. agents able to induce an immune-like response in cells that have not yet
371 encountered MAMPs (Gust *et al*, 2017). Such an original phyto cytokine might then have been neo-
372 functionalized to induce pre-formed defensive cell wall barriers in a developmental context, in the absence of
373 any actual biotic or abiotic stress. MAMP perception and MAMP-induced ROS production are found already in
374 mosses and clearly precede the wide-spread adoption of lignin as a major, cell-wall-reinforcing polymer
375 (Bressendorff *et al*, 2016; Weng & Chapple, 2010). Moreover, lignin was speculated to originate from
376 phenylpropanoid-derived defense compounds, making it plausible that immune signaling pathways have been at
377 the origin of developmental regulation of lignification. The intriguing innovation of the SCHENGEN pathway
378 would then reside in the subcellular arrangement of its signaling components, especially that of SGN1. Indeed,
379 we demonstrate that polar localization of SGN1 to the outside of the Casparian strip domain is crucial for barrier
380 function - since its simple mislocalization to the inside leads to ligand- and receptor-dependent overlignification
381 at inner cell corners. Thus, the only feature that arrests signaling in wild-type is the formation of a lignified
382 diffusion barrier in the cell wall, preventing access of the stele-produced peptides to SGN3 receptors at the outer
383 domain - as the only population able to stimulate the polar SGN1 kinase. Since we have shown here that a main
384 read-out of the SCHENGEN pathway is cell wall lignification itself, this designs a fascinating, spatial negative-
385 feedback loop, in which SCHENGEN pathway-stimulated lignification feedback-regulates itself once enough
386 lignin has been produced to form a tight diffusion barrier and to fully prevent further CIF peptide penetration to
387 the outside.

388
389 It will be important to further describe the extent to which the SCHENGEN pathway uses components
390 of MAMP signaling (as for example the requirement for SERK-family co-receptors, which has not been
391 demonstrated yet for SGN3), but we propose that our findings on the SCHENGEN pathway function can already
392 be of considerable interest for MAMP receptor signaling. The particular, restricted spatial overlap of receptor
393 and downstream kinase has allowed us to visualize that, even if NADPH oxidases are non-localized (as in the
394 case of RBOHD), ROS can be locally produced in a micrometer-scale region at the plasma membrane, through
395 localized receptor stimulation. In MAMP receptor signaling, the non-localized nature of both MAMP receptor
396 and NADPH oxidase would not have allowed to visualize this unexpected degree of spatial control. Yet, during
397 actual microbial infections, highly-localized receptor stimulation and ROS production might indeed occur and
398 be relevant for the outcome of the immune response. In addition, while it is established that MAMP-stimulated
399 ROS production is an important part of the immune response, its direct molecular downstream action is not well
400 understood. Lignin production has long been associated with immune responses, as well as responses to cell wall
401 damage, yet a direct molecular connection from MAMP perception to lignification has rarely been drawn
402 (Chezem *et al*, 2017, 15). It will be intriguing to investigate whether and how much MAMP-induced ROS
403 production is actually used for lignification during defense and to which degree this explains the importance of
404 ROS in plant defense responses.

405
406 The SCHENGEN pathway might not be limited to the regulation of lignified diffusion barriers, as it
407 has been shown that it is also important in the formation of the embryonic cuticle. The peptide ligand used in
408 this context has not been identified and it remains unclear whether an equally precise barrier surveillance
409 mechanism is also acting to ensure separation of endosperm and embryo during embryonic cuticle formation
410 (Creff *et al*, 2019). In this context, the SCHENGEN pathway might be used to drive production and deposition
411 of cutin instead of lignin and suberin as in the case of the endodermis. In the future, it will be fascinating to
412 investigate whether the SCHENGEN pathway is of even broader developmental significance and to understand

413 the molecular basis that enables its distinct, organ and cell-type specific activities.

414 MATERIAL AND METHODS

416 **Plant material and growth conditions.**

417 For all experiments, *Arabidopsis thaliana* (ecotype Columbia) was used. The T-DNA tagged lines, *sgn1-2*
418 (SALK_055095C) and *sgn3-3* (SALK_043282) were obtained from NASC (Alassimone *et al*, 2016; Pfister *et al*,
419 2014). *rbohD dSpm*, *rbohF dSpm* and *rbohD dSpm rbohF dSpm* are described in (Tissier *et al*, 1999; Torres *et al*,
420 2002). *esb1-1* was kindly shared by Prof. David Salt's group (Baxter *et al*, 2009; Hosmani *et al*, 2013b). The *cif1-2*
421 *cif2-2* double mutant was generated by CRISPR-Cas9 method (Fauser *et al*, 2014) using 5'- gctttggttagactggag -3'
422 as a protospacer sequence to target both CIF1 and CIF2 loci.

423
424
425 *pCASP1::CASP1-GFP* marker line was crossed into *sgn1-2*, *sgn3-3* or *cif1-2 cif2-2*. The *pCASP1::myrpalm SGN1*
426 construct was independently transformed into each mutant background using the floral dip method(Clough & Bent,
427 1998). For observations and histological analysis, seeds were kept for 2 days at 4°C in the dark for stratification, then
428 grown for 5 days at 22°C under 16 h light/8 h dark vertically on solid half-strength Murashige–Skoog (MS) medium.
429 Cycloheximide (Sigma) and Diphenyleiodonium chloride (Sigma) were prepared as 50 mM water stock solution
430 and as 20 mM DMSO stock solution, respectively.

431 **Plasmid construction**

432 To generate myrpalm SGN1, GGCFSSK (5'- GGAGGATGCTTCTCTAAGAAG -3') (Vermeer *et al*, 2004) was
433 added to SGN1 cDNA right after the start codon by inverse PCR. This SGN1 variant was combined with pCASP1
434 and mCitrine coding sequence by LR clonase (Thermo Fisher Scientific). Coding sequences of RBOHF 2-383 a.a.
435 and RBOHD 1-376 a.a. were inserted between BamHI and XhoI sites of a modified pET24(+) vector, in which GST
436 (glutathione S-transferase) was added to the N-terminus. For constructing a GST-SGN3 kinase domain vector, a
437 cDNA fragment coding for 899 a.a. to 1249 a.a. of SGN3 was fused into pDEST15 by Gateway LR reaction. TF-
438 SGN1 was as described previously (Alassimone *et al*, 2016). Kinase-dead mutations in SGN1 (K134E) and SGN3
439 (K979E) were introduced by site-directed mutagenesis.

441 **Protein expression in bacteria and purification**

442 GST or Trigger factor (TF) proteins were expressed from pGex6p-1 or pCold-TF vectors in BL21 (DE3) CodonPlus
443 RIL and purified in a glutathione sepharose column (Thermo Fischer Scientific) or a Co-Sepharose column
444 (Clontech) according to the manufacturer's protocols. GST-RBOHD N-ter and GST-RBOHF N-ter were produced
445 and purified as described in (Drerup *et al*, 2013). Bacterially expressed GST-SGN3 kinase domain (WT or kinase-
446 dead) in BL21 (DE3) CodonPlus RIL was purified with a glutathione sepharose column following the manufacture's
447 protocol. For TF-SGN1 (WT or kinase-dead), we followed the method in the previous report (Alassimone *et al*,
448 2016). The buffer of all proteins was changed to 50 mM HEPES-KOH pH7.5 and 1 mM DTT with PD-10 or NAP-
449 5. Purified kinase protein solutions were mixed with glycerol to 30% [v/v] final concentration and kept at -20°C until
450 use.

451 **In vitro kinase assay**

452
453 *In vitro* kinase assay was done as described previously (Alassimone *et al*, 2016) with small modifications. Purified
454 250 ng GST-SGN3 kinase domain was incubated with 250 ng TF-SGN1 or TF proteins in reaction buffer (50 mM
455 HEPES-KOH pH7.5, 1 mM MnCl₂, 1 mM DTT, 1 mM ATP, 185,000 Bq [γ -³²P] ATP) at 30°C for 30 min. 250 ng
456 TF-SGN1 and 1 μ g GST or 500 ng GST-N-terminal cytoplasmic domains of RBOHD or F were treated in the same
457 way as above. The reaction was stopped by adding 4 x LDS sample buffer (Invitrogen) and heating at 75°C for 10
458 min. The samples were separated on 4-12 % gradient Nu-PAGE gels or 10 % Nu-PAGE gels (Invitrogen). After
459 drying the gels, signal was detected using Typhoon FLA7000 (GE Healthcare).

461 **Western blotting**

462 Plants were grown in liquid MS medium supplemented with 0.5% sucrose with cell strainer for 5 days under long
463 day (18 h light /6 h dark) condition after 2-day vernalization at 4°C. Hydroponically grown seedlings were
464 transferred into fresh liquid MS medium containing 0.5% sucrose with or without 1 μ M CIF2 peptide and incubated
465 for 15 min. The seedlings were immediately frozen in liquid nitrogen and ground by TissueLyser II (Qiagen).
466 Extraction buffer (20 mM Na phosphate pH 7.4, , 150 mM NaCl, 1 mM EDTA,, 0.1% Tween, 50 mM β

468 glycerophosphate, 0.5 mM PMSF, 100 μ M sodium orthovanadate, 5 mM Na fluoride, Complete cocktail) was added
469 to the frozen samples. The samples were briefly mixed by vortex and centrifuged at 15,000 rpm for 15 min at 4°C.
470 The supernatant was transferred in new tubes and measured the concentration by Bradford method (Thermo Fischer
471 Scientific). Equal amount of protein was loaded (20 μ g / lane) and separated in 10% acrylamide gel (Eurogentec).
472 After the electrophoresis, the separated proteins were transferred onto nitrocellulose membrane (Amersham, GE
473 Healthcare) by XCell SureLock (Invitrogen) and stained with Ponceau S to show the loading control. The blot was
474 incubated in blocking buffer (3% skim milk in TBS) for 1 hour and probed with anti-phospho-p42, p44 antibody
475 (1:1000 dilution, Cell signaling technologies #9101) in 0.5% skim milk in TBS-0.1% Tween for overnight at 4°C.
476 After washing the blot three times with TBS-0.1% Tween, the blot was treated with anti-rabbit secondary antibody
477 (1:30,000 dilution, Agrisera) in 0.5% skim milk in TBS-0.1% Tween for 1 hour and washed three times with TBS-
478 0.1% Tween. Signals were detected on X-ray film with SuperSignal West Femto Kit (Thermoscientific). For the
479 MPK6 detection, the membrane was stripped (Pierce), incubated in blocking buffer for 1 hour and probed with anti-
480 MPK6 antibody (1:10000 dilution, Sigma) in blocking buffer for 1 hour at room temperature. After washing the blot
481 three times with TBS-0.1% Tween, the membrane was treated with anti-rabbit secondary antibody (1:30000,
482 Agrisera) in blocking buffer and washed three times. Signal were detected with SuperSignal West Femto kit
483 (Thermo Scientific).

484 485 **Confocal microscopes**

486 Confocal pictures were obtained using Leica SP8 or Zeiss LSM 880 confocal microscopes. The excitation and
487 detection window settings to obtain signal as follows: When using Leica SP8 (excitation, detection window), GFP
488 (488 nm, 500–550 nm), mCitrine, (514 nm, 518–560 nm), mCitrine/mCherry, (514 nm and 594 nm, 518–560 nm,
489 600–650 nm, sequential scan), when using Zeiss LSM880 (excitation, detection window), GFP (488 nm, 500–550
490 nm), Calcofluor White (405 nm, 425-475 nm), Basic Fuchsin (561 nm, 570-650 nm) and Fluorol Yellow (488 nm,
491 500-550 nm).

492 493 **PI assay**

494 PI assay was done as described previously (Lee *et al*, 2013) with small changes. Seedlings were incubated in water
495 containing 10 μ g /mL PI for 10 min and transferred into fresh water. The number of endodermal cells were scored
496 using a Zeiss LSM 700 confocal microscope (excitation 488 nm, SP640, split at 570 nm) from the onset of cell
497 elongation (defined as endodermal cell length being more than two times than width in the median, longitudinal
498 section) until PI could not penetrate into the stele.

500 501 **Scoring discontinuities in CS domain**

502 CASP1-GFP signal was obtained by Leica SP8 as 1 μ m step z-stack images from 5-day old seedlings. After the
503 images were projected as maximum projections, total lengths of CS domains were measured in maximum
504 projections and the number of discontinuities in the domain was counted manually.

505 506 **Lignin and cell wall staining**

507 ClearSee adapted cell wall staining was performed as described in recent publications (Kurihara *et al*, 2015; Ursache
508 *et al*, 2018). Briefly, 7-8 five-day-old seedlings were fixed in 3 mL 1 x PBS containing 4% para-formaldehyde for 1
509 hour at room temperature in 12-well plates and washed twice with 3 mL 1 x PBS. Following fixation, the seedlings
510 were cleared in 3 mL ClearSee solution under gentle shaking. After overnight clearing, the solution was exchanged
511 to new ClearSee solution containing 0.2% Fuchsin and 0.1% Calcofluor White for lignin and cell wall staining
512 respectively. The dye solution was removed after overnight staining and rinsed once with fresh ClearSee solution.
513 The samples were washed in new ClearSee solution for 30 min with gentle shaking and washed again in another
514 fresh ClearSee solution for at least one overnight before observation.

515 516 **Methanol-based Fluorol Yellow staining of Arabidopsis root suberin**

517 Vertically grown 5-day old seedlings were incubated in methanol for three days at room temperature. The cleared
518 seedlings were transferred to a freshly prepared solution of Fluorol Yellow 088 (0.01%, in methanol) and incubated
519 for 1 hour. The stained seedlings were rinsed shortly in methanol and transferred to a freshly prepared solution of
520 aniline blue (0.5%, in methanol) for counterstaining. Finally, the seedlings were washed for 2-3min in water and
521 transferred to a chambered coverglass (Thermo Scientific), covered with a piece of 1% half-strength MS agar and
522 imaged using a Zeiss LSM 880 confocal microscope as described above.

523 **Detection of H₂O₂ production *in situ* using transmission electron microscopy**

524 Visualization of H₂O₂ production around Casparian strip was done by cerium chloride method as described
525 previously (Bestwick *et al*, 1997b; Lee *et al*, 2013) with some modifications. 4-day-old Arabidopsis seedlings were
526 transferred onto fresh 1/2 MS solid medium with or without 1 μM CIF2 and incubated for 24 hours. After the
527 peptide treatment, the seedlings were incubated in 50 mM MOPS pH7.2 containing 10 mM CeCl₃ for 30 min.
528 After incubation with CeCl₃, seedlings were washed twice in MOPS buffer for 5 min and fixed in glutaraldehyde
529 solution (EMS, Hatfield, PA) 2.5% in 100 mM phosphate buffer (pH 7.4) for 1 hour at room temperature. Then, they
530 were post-fixed in osmium tetroxide 1% (EMS) with 1.5% of potassium ferrocyanide (Sigma, St. Louis, MO) in
531 phosphate buffer for 1 hour at room temperature. Following that, the plants were rinsed twice in distilled water and
532 dehydrated in ethanol solution (Sigma) at gradient concentrations (30% 40 min; 50% 40 min; 70% 40 min; two
533 times (100% 1 hour). This was followed by infiltration in Spurr resin (EMS) at gradient concentrations (Spurr 33%
534 in ethanol, 4 hours; Spurr 66% in ethanol, 4 hours; Spurr two times (100% 8 hours) and finally polymerized for 48
535 hours at 60°C in an oven. Ultrathin sections 50 nm thick were cut transversally at 1.3 ± 0.1 mm from the root tip, on
536 a Leica Ultracut (Leica Mikrosysteme GmbH, Vienna, Austria) and picked up on a copper slot grid 2x1 mm (EMS)
537 coated with a polystyrene film (Sigma). Micrographs were taken with a transmission electron microscope FEI
538 CM100 (FEI, Eindhoven, The Netherlands) at an acceleration voltage of 80kV and 11000x magnification (pixel size
539 of 1.851nm, panoramic of 17x17 pictures), exposure time of 800ms, with a TVIPS TemCamF416 digital camera
540 (TVIPS GmbH, Gauting, Germany) using the software EM-MENU 4.0 (TVIPS GmbH, Gauting, Germany). All the
541 pictures were taken using the same beam intensity, and panoramic aligned with the software IMOD (Kremer *et al*,
542 1996). For quantification, thresholding of cerium precipitate on pictures was scored using IMOD software (Kremer
543 *et al*, 1996). Briefly, all pictures subjected to the quantification were normalized to one picture, a section from non-
544 treated wild-type seedlings, by comparing the gray value of plastids in pericycle cells. After normalization, cell wall
545 spaces from the beginning of the Casparian strip at the pericycle side until the cortex corner between endodermal
546 cells were selected and one identical threshold setting was applied to all pictures in order to highlight signals around
547 Casparian strips. The values were shown as a percentage of thresholded pixels to selected area.
548

549 **ROS production assay in HEK cells**

550 Measurements of RBOHD and RBOHF activity were performed as described in (Han *et al*, 2019) with some
551 modifications. RBOHD or RBOHF were transiently expressed in HEK293T cells with or without coexpression of
552 SGN1 or myrpalm SGN1. HEK293T cells were cultivated at 5 % CO₂ and 37 °C in Dulbecco's modified Eagle's
553 medium (DMEM) mixture F-12HAM (Sigma) enriched with 10% fetal bovine serum. Before transfection, lysine
554 coated white 96-well plates were inoculated with HEK293T cells and incubated for 24 h. For the transient
555 transfection of the cells, GeneJuice transfection reagent (Novagen) was used according to the manufacturer's
556 guidelines. The transfected expression cassettes were cloned into modified pEF1 vectors (Drerup *et al*. 2013). Each
557 well was transfected with 110 ng of plasmid mixture (50 ng pEF1-RBOH; 30 ng of each effector; empty pEF1
558 vector to ensure equal loading). Transfected cells were incubated for 48 h and subsequently measured in a buffer
559 consisting of Hank's Balanced Salt Solution (Gibco) with 62 μM L-012 and 60 μg / mL HRP. To stimulate SGN1,
560 calyculin A, a phosphatase inhibitor, was added at the final concentration of 0.1 μM directly before the start of the
561 measurements. ROS production was measured in a LB 943 Mithras² (Berthold) microplate reader and is presented as
562 relative luminescence units s⁻¹ (RLU s⁻¹) in graphs. For all experiments, the values were gained from 6 wells
563 measured in parallel and average values were plotted with S.D. Experiments have been repeated three times.
564

565 **Sample preparation for RNA-seq experiments**

566 Wild type, *sgn3-3* and *cif1-2 cif2-2* were grown on solid half-MS medium with mesh for 5 days and transferred onto
567 fresh half MS medium with or without 100 nM CIF2. After 30, 120, or 480 min incubation, aerial parts were cut off
568 and whole roots were collected. Samples were immediately frozen in liquid nitrogen and RNA was extracted using a
569 Trizol-adapted Reliaprep RNA extraction Kit (Promega).
570

571 **RNA-seq library preparation and sequencing**

572
573 RNA-seq libraries were prepared as described in (Jan *et al*, 2019) in using 1000 ng of total RNA. Libraries were
574 sequenced using the Illumina TruSeq SR Cluster Kit v4 reagents on the Illumina HiSeq 2500 and processed using
575 the Illumina Pipeline Software version 2.20.

576 **RNA-seq data processing and analysis**

577
578 Data processing was performed by the Lausanne Genomic Technologies Facility using their in-house RNA-seq
579 pipeline. It includes purity-filtered read trimming for adapters and low quality sequences with Cutadapt (v. 1.8)

580 (Martin, 2011). Removal of reads matching ribosomal RNA sequences with fastq_screen (v. 0.11.1), followed by
581 low complexity read filtering with reaper (v. 15-065) (Davis et al, 2013). Cleaned reads were aligned against
582 Arabidopsis thaliana TAIR10 genome using STAR (v. 2.5.3a) (Dobin et al, 2013). Read counts were obtained per
583 gene locus with htseq-count (v. 0.9.1) (Anders et al, 2015) using A. thaliana TAIR10 Ensembl 39 gene annotation.
584 The quality of the data alignment was evaluated with RSeQC (v. 2.3.7) (Wang et al, 2012).
585

586 Gene-level statistical analysis was performed in R (R version 3.4.3). Filtering of genes with low counts was
587 achieved based on the rule of 1 count per million (cpm) in at least 1 sample. TMM normalization was applied for
588 library sizes scaling followed by log-transformation into counts per million or CPM (EdgeR package version 3.20.8)
589 (Robinson et al, 2010). PCA was computed using normalized values corrected for batch effect using limma function
590 removeBatchEffect.

591 Differential expression was computed with limma-trend approach (Ritchie *et al*, 2015) by fitting all samples into one
592 linear model. The batch factor was added to model matrix.

593 - Pairwise comparisons treated vs untreated per time point were assessed using moderated t-tests. The adjusted p-
594 value is computed by the Benjamini-Hochberg method, controlling for false discovery rate (FDR or adj.P.Val).

595 - Differential expression of untreated mutant vs wt per time point was assessed using moderated F-test and Post-Hoc
596 classification. The adjusted p-value is computed by the Benjamini-Hochberg method, controlling for false discovery
597 rate (FDR or adj.P.Val).

598 - Differential expression of treated vs untreated over time paired by genotype was assessed using moderated F-test
599 and Post-Hoc classification. The adjusted p-value is computed by the Benjamini-Hochberg method, controlling for
600 false discovery rate (FDR or adj.P.Val).

601 - Interaction between time and treatment (paired by genotype, excluding SGN3 of the model) was assessed using
602 moderated F-test. The adjusted p-value is computed by the Benjamini-Hochberg method, controlling for false
603 discovery rate (FDR or adj.P.Val).

604 - Time effect in untreated conditions (paired by genotype, excluding SGN3 of the model) was assessed using
605 moderated F-test. The adjusted p-value is computed by the Benjamini-Hochberg method, controlling for false
606 discovery rate (FDR or adj.P.Val).
607

608 Genes were considered significant in further analysis if the adjusted p-value was equal or below 0.05 and the log2-
609 fold change was ≥ 1 . Heatmaps were constructed using the package ComplexHeatmap (v1.99.4, pearson distance
610 for row clustering) (Gu *et al*, 2016). Genes were clustered using kmeans (factoextra v1.0.5,
611 <https://github.com/kassambara/factoextra>) using a non-supervised approach resulting in 3 suggested clusters. We
612 found that 4 inferred clusters split one gene cluster in a more sensible way and adjusted the number of clusters
613 accordingly to four. GO analysis were conducted using the package topGO (v. 2.34.0, weight01 algorithm) (Alexa &
614 Rahnenfuhrer, 2019). GO annotations were obtained through Biomart (version 2.40.0, Ensembl Plants release 43 -
615 April 2019) (Durinck *et al*, 2009). Genes for pathway specific heatmaps were obtained from the corresponding GO
616 term through the Ensembl Plant database.
617

618 qPCR analysis

619
620 For the CIF2 peptide treatment, 5-days-old seedlings grown on half MS were moved to fresh 1/2 MS medium in the
621 absence or presence of 100 nM CIF2 and incubated for 30 or 120 min. Otherwise, the plants were grown with mesh.
622 Only root parts (around 100 mg) were collected at each time point and total RNA was extracted using a Trizol-
623 adapted ReliaPrep RNA Tissue Miniprep Kit (Promega). Reverse transcription was carried out with PrimeScript RT
624 Master Mix (Takara). All steps were done as indicated in the manufacturer's protocols. The qPCR reaction was
625 performed on an Applied Biosystems QuantStudio3 thermocycler using a MESA BLUE SYBR Green kit
626 (Eurogentech). All transcripts are normalized to *Clathrin adaptor complexes medium subunit family protein*
627 (*AT4G24550*) expression. All primer sets are indicated in star methods.
628

629 Statistical analysis for experiments

630 All statistic-related analysis was done with R software (R Core Team, 2013) (<https://www.r-project.org/>). For multi-
631 comparison analysis, one-way ANOVA was carried out and Tukey's test was subsequently performed.

632 Data and materials availability

633 All data to support the conclusions of this manuscript are included in the main text and supplementary materials.
634 The full RNA seq dataset was deposited in GEO (accession GSE144182,
635 <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE144182>).

636
637
638
639
640

641
642
643
644
645
646

647
648
649
650
651

652

653
654
655
656
657
658
659

660
661
662
663
664

Acknowledgments: We thank the Central Imaging Facility (CIF), Genome Technology Facility (GTF), particularly Sandra Calderon, and Electron Microscopy facility (EMF) of the University of Lausanne for expert technical support. We also thank Hiroko Uchida for expert graphical support.

Funding: This work was supported by funds to N.G. from an ERC Consolidator Grant (GA-Nº: 616228 – ENDOFUN), and two SNSF grants (CRSII3_136278 and 31003A_156261), a Federation of European Biochemical Sciences Postdoctoral Long-Term Fellowship to P.M. and an EMBO Long-term postdoctoral fellowship to R.U., an overseas research fellowship from JSPS to S.F., a Marie Curie postdoctoral fellowship to T.G.A., a fellowship of the Fundación Alfonso Martín Escudero to V.G.D and a DFG grant (Ku931/14-1) to J.K.

Author contributions: S.F. and N.G. conceived the project. S.F., T.G.A and N.G. designed the experiments. S.F., D.D.B., K.H.E, P.K, T.G.A, V.D.T, A.P., P.M., R.U., V.G.D., M.B. and A.C. performed the experimental work. D.D.B., T.G.A., E.S.S. and J.D. performed image quantification and RNA-seq analysis. S.F., and N.G. wrote the manuscript. G.I., J.K. and all other authors revised the manuscript and were involved in the discussion of the work.

Conflict of interest: The authors declare no competing interests.

REFERENCES

Automatic citation updates are disabled. To see the bibliography, click Refresh in the Zotero tab.

Supporting information

Appendix file
Tables EV1-2
Movie EV1




Figure1:

666 **Fig. 1 Apolar SGN1 leads to ectopic lignin accumulation in endodermal cells**

667

668 (A) Schematic of Casparian strip development (magenta). Casparian strips start to appear as centrally aligned
669 discontinuous dots in the endodermal cell layer, progressing into a network of fused rings functioning as a root
670 apoplastic barrier.

671

672 (B) Lignin accumulation patterns at endodermal surface or median positions with or without the 100 nM CIF2
673 (Casparian strip integrity factor 2) ligand treatment. Lignin and cellulosic (unmodified) cell walls are stained
674 with Basic Fuchsin and Calcofluor White, shown in magenta and white respectively. Schematics are indicating
675 the position of optical sections in a 3D illustration. For each condition, at least 10 roots were tested and showed
676 similar results in two independent experiments. White arrows indicate sites of excess lignification on the cortex
677 facing (outer) side. Scale Bar = 5 μm .

678

679 (C,D) Localization of SGN1-Citrine and lignin deposition patterns in *pCASPI::SGN1-Citrine* lines in wild-type
680 (Col) and different mutant backgrounds (*sgn1*, *sgn3*, *cif1 cif2*) (C). *myrpalm*-SGN1-Citrine localization and
681 lignin deposition patterns in *pCASPI::myrpalm-SGN1-Citrine* lines (D). Lignin (Basic Fuchsin) and cell walls
682 (Calcofluor White) are shown in magenta and white respectively. For this experiment, two or three independent
683 lines were tested. From each transgenic line, 2 positions from 12 roots were observed and representative pictures
684 are shown in the figure. Schematics are indicating the position of optical sections in a 3D illustration. White
685 arrows in (D) highlight excess lignification on the pericycle-facing (inner) side. Scale bars = 5 μm .

686

687 (E) Schematic illustrating how signal activation can be governed by SGN1 localization and peptide ligand
688 diffusion from the stele.

689

690 Data information: “inner” designates the stele-facing endodermal surface, “outer”, the cortex-facing surface.

691




Figure2

Fig. 2 SGN1 acts as a transducer of CIF2 signaling and is phosphorylated by the SGN3 receptor

(A) Quantification of defects in CSD formation as number of holes per 100 μm in the CASP1-GFP domain at around 10 cells after onset of CASP1-GFP expression in 5-day-old seedlings. In the box plot, boxes indicate ranges from first to third quartiles, bold central lines display median. Upper and lower whiskers extend to maximum or minimum values no further than 1.5 times IQR (interquartile range, the distance between the first and third quartiles). One-way ANOVA was performed followed by Tukey's test. Different letters show significant statistical differences. ($p < 0.05$, One-way ANOVA and Tukey's test, 12 roots in total were observed for each condition in two independent assays).

(B) Propidium iodide (PI) penetration assay in the presence or the absence of CIF2. CS barrier function was scored as exclusion of PI signal from the inner side of endodermal cells. In the box plot, boxes indicate ranges from first to third quartiles, bold central lines display median. Upper and lower whiskers extend to the maximum or minimum values no further than 1.5 times IQR. Different letters show significant statistical differences. ($p < 0.05$, One-way ANOVA and Tukey's test. During two independent experiments, 10 roots in total were tested for each condition).

(C) [γ - ^{32}P]ATP radioactive *in vitro* kinase assay of SGN3 kinase domain against SGN1. Autoradiograph is shown on top. Coomassie stained-gel below illustrates presence and equal loading of recombinant proteins. Note that a kinase-dead SGN1 variant was used to avoid auto-phosphorylation activity of SGN1. Also note that trigger factor represents a very big tag protein, accounting for the high migration of TF-SGN1. Representative result of three independent experiments is shown.





Figure3

Fig. 3 Both RBOHD and F are required for CIF2-induced excess lignin accumulation

(A) Localization of Citrine-RBOHD (left) or Citrine-RBOHF (right) in endodermal (en) cells. Both proteins were expressed under the control of pCASP1, an endodermis-specific promoter. Representative pictures are shown, 2 positions from 10 roots for each transgenic line were inspected. Scale bar = 5 μ m.

(B) Lignin accumulation in WT and *rbohD* and *rbohF* single mutants and a double mutant with or without 2-hour 100 nM CIF2 peptide treatment. Arrowheads indicate excess lignification. Pictures are shown as overviews (maximum projection) or median sections. Lignin and cell walls are shown with magenta (stained with Basic Fuchsin) and gray (stained with Calcofluor White) respectively. Representative pictures are shown, 12 roots (overview) and 2 positions in 12 roots (median section) were inspected. Scale bars = 20 μ m (lignin overviews), 5 μ m (median sections). “inner” designates the stele-facing endodermal surface, “outer”, the cortex-facing surface.



731 **Fig. 4 ROS production is enhanced by SGN3/CIFs and requires RBOHD and F**

732

733 (A) Co-visualization of *pCASPI::SGN1-mCherry* and *pCASPI::SGN3-GFP*. Note that their respective
734 localization at the PM has only restricted overlap at the cortical side of the CS domain (Arrow). Representative
735 picture is shown, 3-6 roots in 4 different transgenic lines were inspected. Scale bar = 10 μm . “inner” designates
736 the stele-facing endodermal surface, “outer”, the cortex-facing surface.

737

738

739 (B,C) Overview of endodermal cells after with or without 1 μM CIF2 treatment for 24 hours. Red arrowheads
740 are indicating ROS production sites and boxes in dotted lines are corresponding to the regions in (sky blue and
741 creme boxes in C and black boxes in D). Similar patterns were obtained in 39 or 41 cells from 5 roots with or
742 without the peptide treatment. Scale bars = 500 nm.

743

744 (D) *In situ* H_2O_2 detection at Casparian strips in WT, *sgn3*, *sgn1*, *rbohF*, *rbohD* and *rbohDF* with or without 24-
745 hour treatment of 1 μM CIF2. Brackets and arrowheads indicate Casparian strips (seen as uniformly whitish cell
746 wall areas) and H_2O_2 production sites (black area) respectively. Scale bar = 1000 nm.

747

748 (E) Quantification of ROS production as a number of dark pixels area (n = 23 – 44 sites from 5 roots of each
749 condition in one experiment.). For quantification methods, see fig. EV 3. In the box plot, boxes indicate ranges
750 from first to third quartiles, bold central lines display median. Upper and lower whiskers extend to the maximum
751 or minimum values no further than 1.5 times IQR. Different letters mean significant statistical differences.
752 ($p < 0.01$, One-way ANOVA and Tukey’s test.)

753

754





Figure5

755
756

757 **Fig. 5 SGN1 directly activates NADPH oxidases in a cellular context**

758

759 (A) [γ - 32 P]ATP radioactive *in vitro* kinase assay of TF-SGN1 against GST-N-terminal cytoplasmic domains of
760 RBOHD or F. Autoradiograph is shown on top. Coomassie stained-gel below illustrates presence and equal
761 loading of recombinant proteins. Experiments were done independently three times with similar results.

762

763 (B) HEK293Tcell based NOX activation assay. Cells were transfected with the indicated plasmid combinations.
764 The phosphatase inhibitor Calyculin A was added directly before the start of the measurements. Each data point
765 represents the mean of six wells analyzed in parallel, bars indicate S.D.. Experiments were repeated three times
766 (another set of results is shown in Fig. EV 4B).

767

768




Figure 6

771 **Fig. 6 CIF2 induces large-scale transcriptional changes for cell wall remodeling**

772

773 (A) CASP1-GFP and lignin deposition in WT, *sgn3*, *rbohD*, *rbohF* and *rbohDF*. CASP1-GFP and lignin
774 (fuchsin) are presented in green and magenta respectively. Pictures were obtained from more than 10 roots from
775 each background with similar results. Scale bar = 10 μ m.

776

777 (B) Time lapse imaging of single- or cotreatment of 10 nM CIF2 with 25 μ M cycloheximide (CHX) on CASP1-
778 GFP in *cif1 cif2*. Seedlings were pretreated with or without CHX for 30 min and transferred onto each medium.
779 Scale bar = 10 μ m (See also movie EV1).

780

781 (C) Quantification of (B). Relative numbers of holes in CASP1-GFP domain after single- or co-treatment with
782 CIF2 or CHX from the pictures in (B). Bars are S.D. * indicates statistical significance from all other conditions
783 ($p < 0.01$) after one-way ANOVA and Tukey test. 6 roots in total for each condition were observed during two
784 independent tests.

785

786 (D) Fold-change of 930 genes ($p < 0.05$ and $\log_2(\text{Fold change}) \geq 1$ or ≤ -1 at least one time point in one genotype)
787 after CIF2 treatment at indicated time points in WT, *cif1,2* and *sgn3*. Degree of the fold changes are shown in
788 color code as indicated.

789

790 (E) Relative expression levels of *PER15* to *CLATHRIN* control in each genotype with or without 2-hour CIF2
791 treatment. Bars are S.D. (n=3). Different characters indicate statistically significant differences ($p < 0.01$,
792 ANOVA and Tukey test).

793

794 (F) Fold changes of *PER15* in each genotype with or without 2-hour CIF2 treatment. Bars are S.D. (n=3).
795 Different characters indicate statistical significance differences ($p < 0.01$, ANOVA and Tukey test).

796

797 (G) Relative expression levels of *PER49* to *CLATHRIN* control in each genotype with or without 2-hour CIF2
798 treatment. Bars are S.D. (n=3). Different characters indicate statistically significant differences ($p < 0.01$,
799 ANOVA and Tukey test).

800

801 (H) Fold changes of *PER49* in each genotype with or without 2-hour CIF2 treatment. Bars are S.D. (n=3).
802 Different characters indicate statistically significant differences ($p < 0.01$, ANOVA and Tukey test).

803

804 (I) Relative fold changes of *PER15* in *pCASP1::SGN1-Citrine* and *pCASP1::myrpalm-SGN1-Citrine* compared
805 to the expression level in WT. Bars are S.D. (n=3). Different characters indicate statistically significant
806 differences ($p < 0.01$, ANOVA and Tukey test).

807

808 (J) Relative fold changes of *PER49* in *pCASP1::SGN1-Citrine* and *pCASP1::myrpalm-SGN1-Citrine* compared
809 to the expression level in WT. Bars are S.D. (n=3). Different characters indicate statistically significant
810 differences ($p < 0.01$, ANOVA and Tukey test).

811




Figure 7 Overview of the plasma membrane-based and nuclear branches of the SGN3 pathway

813 **Fig. 7 Overview of the plasma membrane-based and nuclear branches of the SGN3 pathway**

814 A, B (A) Schematic of the spatially-restricted activation of ROS production at the plasma membrane after CIF
815 stimulation (B) Comparison of the components of the cytoplasmic/nuclear signaling cascades induced by the
816 flg22 bacterial pattern peptide (left) and the CIF peptides (right). Note that all of the signaling components in
817 the two pathways belong to the same gene families, with the exception of the transcription factors. Components
818 for which there is currently no direct experimental evidence are marked in gray, as are arrows indicating
819 activation events that have not yet been experimentally established.

820

821

822 **Table EV1.**

823 Differentially expressed genes from heatmap presented in Fig.6D

824 **Table EV2.**

825 A list of enriched GO terms of clusters presented in Fig.6D

826

827