



HAL
open science

Effects of a bactericide on the structure and survival of benthic diatom communities

Soizic Morin, L. Proia, M. Ricart, C. Bonnineau, A. Geiszinger, F. Ricciardi, H. Guasch, A.M. Romani, Sergi Sabater

► **To cite this version:**

Soizic Morin, L. Proia, M. Ricart, C. Bonnineau, A. Geiszinger, et al.. Effects of a bactericide on the structure and survival of benthic diatom communities. *Vie et Milieu / Life & Environment*, 2010, 60 (2), pp.109-116. hal-02593829

HAL Id: hal-02593829

<https://hal.inrae.fr/hal-02593829>

Submitted on 19 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 EFFECTS OF A BACTERICIDE ON THE STRUCTURE AND SURVIVAL OF BENTHIC
2 DIATOM COMMUNITIES

3 Soizic Morin^{1,2}, Lorenzo Proia², Marta Ricart², Chloé Bonnineau², Anita Geiszinger²,
4 Francesco Ricciardi², Helena Guasch², Anna Romani², Sergi Sabater^{2,3}

5

6 ¹ Cemagref, UR REBX, 50 avenue de Verdun, 33612 Cestas cedex, France

7 ² Institut d'Ecologia Aquàtica, Campus Montilivi, 17073 Girona, Spain

8 ³ Catalan Institute for Water Research (ICRA), Scientific and Technologic Park of the
9 University of Girona, 17003, Girona, Spain

10

11 Running title: Effects of a bactericide on diatom communities

12

13 ABSTRACT

14 We studied the adverse effects of triclosan, a widely used biocide commonly reported in
15 surface waters, on the structure and function of benthic diatom communities. Laboratory-
16 grown biofilms were exposed (i) to chronic contamination by increasing concentrations of
17 triclosan and (ii) to a short-pulse of sublethal triclosan concentrations followed by a 2-week
18 recuperation period.

19 The first experiment was performed using 6 nominal concentrations ranging from 0.05 to 500
20 µg/L triclosan to obtain the concentration – effect relationships for benthic diatom
21 communities. Here effects at the highest triclosan concentration in the diatom community
22 consisted of a 63% increase in diatom mortality, with respect to control conditions. The
23 second experiment aimed at determining the long-term effects of the toxicant and biofilm
24 recuperation after addition of 60 µg/L triclosan for 48h exposure. After two weeks the
25 sublethal pulse had caused a decrease in diatom growth rates and a significant delay in the

26 exponential phase of growth. The triclosan pulse provoked a decrease in diatom species
27 richness and diversity. The diatom communities were dominated by 6 species, with
28 *Achnantheidium minutissimum* being highly preponderant, and variations were not large
29 enough to provide information about sensitivities / tolerance to triclosan in different species.

30

31 KEYWORDS

32 Bacillariophyceae; benthic communities; triclosan; dose-response; bioassays; live and dead
33 cells; recovery

34

35 INTRODUCTION

36 Residues of pharmaceutical and personal care products (PPCPs) have been detected in
37 freshwater bodies for decades, but assessing their impact on the ecosystem and on the
38 biological communities inhabiting these ecosystems (especially non-target organisms) is
39 relatively new (Fent *et al.* 2006). The use of triclosan (TCS), a broad-spectrum bactericide,
40 disinfectant and fungicide, in PPCPs is increasing and the compound is, together with its
41 transformation products, frequently found in surface waters after biological degradation. The
42 degradation products include methyltriclosan (Balmer *et al.* 2004; Singer *et al.* 2002; Wind *et*
43 *al.* 2004) and others, originating from photolytic and abiotic degradation, such as 2,8-
44 dichlorodibenzo-p-dioxin (Aranami & Readman 2007; Buth *et al.* 2009) and chlorophenols
45 (Canosa *et al.* 2005). Further degradation compounds, still not completely identified, are
46 produced by direct photolysis (Wong-Wah-Chung *et al.* 2007) and their effects are still
47 completely unknown.

48 Triclosan has several modes of action against bacteria such as the inhibition of lipid synthesis
49 (Sivaraman *et al.* 2004) and modifications of the cell membrane permeability (Villalaín *et al.*
50 2001). Though triclosan is intended for use as a bactericide, it has also been shown to have

51 deleterious effects on various other aquatic organisms. Recent reviews (Capdevielle *et al.*
52 2008; Sanderson & Thomsen 2009) reported a high sensitivity of algae to TCS, with growth
53 inhibition EC₅₀ values less than 5 µg/L. However, significant differences in sensitivity were
54 observed among algal groups (Orvos *et al.* 2002) and algal life forms (Franz *et al.* 2008).
55 Quick recovery of algal cultures following triclosan contamination events has also been
56 reported (Orvos *et al.* 2002).
57 Though evidence exists that triclosan may affect algal growth, the potential sensitivity of
58 fluvial (benthic) diatom assemblages is unknown. The arrival of this pharmaceutical into
59 stream systems through sewage waters may cause relevant effects in the diversity, sensitivity
60 and recovery capacity of diatom communities (Ricciardi *et al.* 2009). Here, we propose to use
61 a dose-response test to determine the sensitivity of fluvial diatom assemblages to triclosan.
62 Further, using a sublethal concentration, the recuperation potential of benthic diatom
63 communities to this toxicant was assessed through functional and structural descriptors, i.e.
64 chlorophyll-a *in vivo* fluorescence, live / dead diatom cell ratio and taxonomic composition.

65

66 MATERIALS AND METHODS

67 1. Experimental design

68 *Experiment 1: Dose – response relationships*

69 Benthic algal communities were collected from rocks in a small stream and inoculated in
70 aerated incubators with flow-like water circulation to allow settlement of biofilms on glass
71 slides (1 cm² each). After 3 weeks, the colonised slides were deposited in eight artificial
72 channels supplied with micro-filtered water to which nutrients (N and P) were added
73 continuously. The channels received a constant flow of about 1.2 L/min and a light/dark cycle
74 of 12/12h (as described in Ricart *et al.* submitted). The biofilms were acclimated for three
75 days to the new conditions before being exposed to an increasing range of triclosan

76 concentrations for 48 hours. The stock triclosan solution was prepared by dissolving triclosan
77 powder (Irgasan DP-300) in 100% methanol and was stored in the dark to avoid
78 photodegradation. From this, working standard solutions were prepared in the range of 0.05-
79 500 µgTCS /L, in a solution containing 25% methanol. One of the eight artificial channels
80 was used as a “control” *sensu stricto* (no additions to the initial culture medium), and a
81 second one was used as the “control solvent” (culture medium + 0.05% methanol). The other
82 six channels received triclosan at one of the nominal concentrations: 0.05, 5, 25, 125, 250 and
83 500 µgTCS/L. At the end of the 48-hour exposure period, 4 glass slides per channel were
84 collected at random for chlorophyll-a *in vivo* fluorescence measurements and diatom
85 enumerations.

86

87

88 *Experiment 2: Long-term effects of a 48h-pulse of triclosan*

89 Eight independent incubators equipped with bubbling systems were used to grow biofilms in
90 as described in Experiment 1. Colonisation lasted for 4 weeks, with a 12/12h light/dark cycle.
91 The incubators were filled with liquid medium (1.5L each, detailed composition in Table 1).
92 The liquid was replaced in each incubator twice per week, in order to avoid nutrient depletion.
93 Then, in 4 of the 8 incubators triclosan was added, to reach the nominal concentration of 60
94 µgTCS/L. In order to maintain the contamination level, water was renewed every 3 hours
95 during illuminated periods. After the 48-hour exposure with triclosan, all the incubators were
96 re-filled with regularly replaced unpolluted liquid medium (as described above), and the
97 biofilms incubated for 2 weeks.

98 Biofilm samples were taken just before triclosan addition (called T0d), at the end of the 48-
99 hour exposure (T2d), and then after 7 days (T9d) and 13 days (T15d) in uncontaminated
100 medium. For each sampling, 4 glass slides were randomly taken from each incubator and

101 analysed for *in vivo* fluorescence, the diatoms were enumerated and live and dead cells
102 distinguished. Then, 4 replicate slides for each incubator were pooled for taxonomic purposes,
103 to give one integrated sample per incubator.

104

105 2. Determination of the physicochemical characteristics

106 Temperature, pH, conductivity and dissolved oxygen in the channels and incubators were
107 measured using a multiparametric probe (WTW, Weilheim, Germany). Water samples were
108 collected and filtered (nylon membrane filters 0.2 µm, Whatman International Ltd,
109 Maidstone, England) prior to analysis. Nutrient concentrations were determined using ion
110 chromatography (761 Compact IC, Metrohm Ltd., Herisau, Switzerland). Soluble reactive
111 phosphate was measured following Murphy and Riley (1962).

112 Triclosan and methyl-triclosan standards were purchased from Sigma–Aldrich (St. Louis,
113 MO, USA). Standard solutions were prepared in pure methanol. Water samples (500 mL)
114 were filtered through Millipore 0.45 µm membrane filters and pre-concentrated immediately
115 at a flow rate of 5 mL/min on Waters C18 6-mL solid phase extraction cartridges (Oasis HLB,
116 Waters, Milford, MA), previously activated and conditioned with 5 mL of acetone, methanol
117 water (1 mL/min). Finally, the SPEs were eluted with 4 mL of methanol (1 mL/min) and the
118 sample was directly injected.

119 The HPLC system consisted of a Waters 717 autosampler and a Waters 1525 binary pump.

120 HPLC separation was achieved on a 5-µm, 150 × 4 mm i.d. C18 reversed-phase column

121 (SunFire). The injection volume was set at 50 µL, and the flow rate was 1 mL/min of 90%

122 methanol - 10% water with isocratic flow. Detection was carried out using a UV–vis detector

123 (Waters 2489) at 280 nm.

124

125 3. Diatom analyses

126 *In vivo fluorescence measurements*

127 An estimation of diatom-specific chlorophyll concentration was obtained by measuring the
128 specific F₀ (minimum fluorescence yield of dark-adapted cells) by means of a PhytoPAM
129 fluorometer (Heinz Walz, Effeltrich, Germany) at 520 nm (Schmitt-Jansen & Altenburger
130 2007). The F₀ was determined after applying a weak measuring light to the periphyton
131 samples (directly on the colonised slides), after a 20-minute incubation in the dark to ensure
132 that all the reaction centres were open. The photosynthetic capacity of the diatom community
133 (Y_{max}) was also determined on dark-adapted samples. After being kept in the light for 20
134 minutes, the samples were exposed to actinic light for the determination of their
135 photosynthetic efficiency (Y_{eff}). PhytoPAM measurements for each parameter were repeated
136 5 times per slide and averaged.

137 *Quantitative estimates of diatom community structure and live / dead ratio*

138 After fluorescence measurements, the glass slides were scraped using polyethylene cell lifters
139 (Corning Inc., NY, USA), and cells preserved with a drop of formalin solution and diluted to
140 a final volume of 5 mL. Samples were ultrasonicated for 7 minutes to separate the aggregated
141 cells without destroying the frustules. 125 µL of each sample were then pipetted onto a
142 Nageotte counting chamber to count the total number of diatom cells in 10 microscope fields
143 (1.25 µL each, 0.5 mm depth) selected at random, using light microscopy at a 10×
144 magnification (photomicroscope Nikon Eclipse 80i, Nikon Co., Tokyo, Japan). Data were
145 recorded as cells per unit area of sampled substrate (number of cells/cm²). Countings were
146 separated into 2 types: empty cells that were considered as ‘dead’, and cells occupied by
147 chloroplasts that, whatever their color (from pale yellow to green or brown), shape and
148 number, were considered as ‘alive’ (Cox 1996). From live diatom counts, growth rates (as

149 expressed in cell divisions/day) of the diatom community were calculated according to
150 Guillard (1973).

151 The optimal use of the live / dead cell indicator (L/D ratio) required:

152 1) The observations / countings to be carried out within 1 month after sampling. Chlorophyll
153 content has been shown to decrease very quickly in formalin-preserved samples (Dell'Anno *et*
154 *al.* 1999), and a decrease in pigment content would make it harder to distinguish between live
155 and dead cells. Storing the formalin-fixed samples in the dark will however delay chlorophyll
156 degradation (M.-J. Chrétiennot-Dinet, pers. comm.). Additionally, we used the same dilution
157 per unit substratum surface area in a single study, in order to maximize comparability
158 between data. Moreover, as the difference between live and dead cells may be subjective in
159 some cases (e.g. very slight coloration of the cell content), all observations were performed by
160 a single operator;

161 2) The proportion of live vs. dead diatoms was based on counting entire cells, *i.e.* those that
162 exhibit an entire frustule. This required the ultrasonication to be limited to a maximum of 7
163 minutes, in order to minimise frustule damage.

164 *Relative abundances of the diatom species*

165 Diatom identifications were performed only in experiment 2, after having prepared permanent
166 slides following European standard NF EN 13946. About 500 frustules were counted per slide
167 at 1,000× magnification and diatoms were identified to the lowest taxonomic level possible
168 using standard references and recent nomenclature updates. References consisted of:

169 Krammer & Lange-Bertalot (1986 - 1991), Sala et al. (1993), Round & Bukhtiyarova (1996),
170 Krammer (1997), Compère (2001), Bukhtiyarova (2006) and Potapova (2006).

171 Diatom life forms were separated following the guilds as defined by Passy (2007). Species
172 with solitary life forms were divided into 3 groups: motile, erected and prostrate, according to

173 the growth forms described by Cox (1996), Hudon & Legendre (1987), Katoh (1992), and
174 Kelly et al. (2005).

175

176 4. Processing the data

177 The values of chlorophyll-a *in vivo* fluorescence and L/D ratios with increasing triclosan
178 concentrations (Experiment 1) were analysed for significant differences using 1-way ANOVA
179 performed with Statistica v5.1 (StatSoft Inc. Tulsa, Oklahoma). The EC₅₀ of the L/D ratio was
180 estimated from the dose / response curve, using the 3 replicates of the 2 controls and 6
181 measured triclosan concentrations.

182 The variation of the same parameters with treatment and duration of the experiment
183 (Experiment 2) were explored by means of a linear mixed-effect model (LME) for repeated
184 measurements. For this purpose, the treatments and sampling dates were considered as fixed
185 effects and the replicate samples (*i.e.* the samples taken from replicate incubators) as random
186 effects. These statistical analyses were computed with the package nlme in the statistical
187 modelling environment R (Ihaka & Gentleman 1996).

188 Calculating the ratio between live and dead cells was quite independent of the number of cells
189 counted. Indeed, relative counts allowed minimization of the influence of microhabitat factors
190 (as shown by the reduced error in the intra-treatment L/D values compared to absolute cell
191 density data), especially in the case of small-sized substrates as used in these experiments,
192 where algal development is highly variable.

193

194 RESULTS

195 *Experiment 1: Dose – response relationships*

196 Apart from the TCS additions, all channels were fed with the same water; its characteristics
197 are given in Table 1. TCS concentrations measured in the channels (Table 2) differed slightly

198 from the nominal concentrations. The highest concentration was 461.29 $\mu\text{g/L}$. Concentrations
199 of methyltriclosan, a degradation product of triclosan, always remained below the detection
200 limit.

201 After a 48-hour exposure, chlorophyll-a *in vivo* fluorescence parameters did not display any
202 significant trend related to triclosan addition, with mean values of $F_0 = 100.2 \pm 3.0$ ($n = 36$),
203 $Y_{\text{max}} = 0.43 \pm 0.01$ ($n = 36$) and $Y_{\text{eff}} = 0.30 \pm 0.01$ ($n = 36$).

204 Live diatom densities tended to decrease with increasing TCS concentrations but did not
205 express statistically significant differences ($p = 0.09$). Up to $130,000 \pm 24,000$ cells/cm²
206 occurred in the controls, while $91,000 \pm 9,000$ cells/cm² were counted in the maximum TCS
207 concentration, representing a decrease of about 30% (Figure 1). The highest diatom densities
208 were observed in the channel exposed to 5.41 $\mu\text{gTCS/L}$, that reached ca. $170,000 \pm 37,000$
209 cells/cm².

210 The ratio between live and dead diatoms ranged from 6.2 ± 0.2 in the controls down to $3.5 \pm$
211 0.3 in the most contaminated channel. Diatom L/D ratio expressed as a percentage of the
212 mean control values fitted the sigmoidal curve reported in Figure 2. The dose-response curve
213 was used to evaluate the EC_{50} of TCS for the diatom community. An EC_{50} value of 560 $\mu\text{g/L}$
214 was derived from the regression equation of the curve (confidence interval: 534-584 $\mu\text{g/L}$).
215 This value was taken as a criterion to use a 10-fold lower concentration (60 $\mu\text{g/L}$) to perform
216 Experiment 2.

217

218 *Experiment 2: Long term response after a 48h-pulse of triclosan*

219 Among the *in vivo* parameters, the linear mixed-effect model did not show any statistical
220 differences in F_0 and Y_{eff} values between treatments, duration of exposure, nor combined
221 effects. Indeed, T0d values were $F_0 = 138.4 \pm 26.0$ ($n = 40$) and $Y_{\text{eff}} = 0.37 \pm 0.03$ ($n = 40$),
222 with maximum variations of 29% of the control (T15d, where a slight date x treatment effect

223 was observed, $p \sim 0.05$) and 4 % (T2d) respectively. Conversely, the photosynthetic capacity
224 (Y_{max}) expressed differences both due to sampling date, treatment ($p < 0.0001$) and date x
225 treatment ($p < 0.01$). Y_{max} increased with time (from 0.47 ± 0.04 at T0d), with values
226 slightly higher in the TCS-treated samples (0.52 ± 0.02) than in the controls (0.48 ± 0.02).
227 Diatom communities in the controls showed exponential growth during the 15 days of the
228 experiment ($R^2 = 0.93$; $p < 0.001$, see Figure 3), from $117,000 \pm 12,000$ cells/cm² (T0d) to
229 $690,000 \pm 88,000$ cells/cm² (T15d). In contrast, diatom densities in the incubators receiving
230 triclosan were quite stable throughout the 9 first days following the beginning of exposure
231 ($106,000 \pm 22,000$ cells/cm²) and then reached $447,000 \pm 55,000$ cells/cm² at T15d. Whatever
232 the treatment, the increase in cell densities from T0d to T15d was highly significant ($p <$
233 0.0001), and a date x treatment effect was also observed at the last dates of the experiment (p
234 < 0.05).

235 Before TCS exposure, the ratio between live and dead diatoms averaged 10.9 ± 0.4 in the
236 incubators. This ratio decreased down to 7.4 ± 0.7 (more than 30% decrease) in the TCS-
237 exposed systems in comparison to the control values (treatment x date effect: $p < 0.0001$ at
238 T9d) and then increased at T15d but remained significantly lower than the controls (-16.6%, p
239 < 0.05).

240 A total of 32 diatom species were identified during this study, but only 6 of them accounted
241 for more than 1% in at least one of the samples (Figure 4). In all treatments, diatom
242 communities were dominated by *Achnantheidium minutissimum* Kützing (87.3 ± 1.4 %),
243 followed by *Gomphonema angustatum* (Kützing) Rabenhorst (5.5 ± 0.9 %), *Achnantheidium*
244 *pyrenaicum* (Hustedt) Kobayasi (1.9 ± 0.5 %), *Gomphonema minutum* (Agardh) Agardh (1.0
245 ± 0.2 %), *Ulnaria ulna* (Nitzsch) Compère (0.8 ± 0.2 %) and *Fragilaria capucina*
246 Desmazières var. *capucina* (0.6 ± 0.3). Relative abundances of these main species remained
247 quite stable in the control incubators during the course of the experiment, and slight changes

248 occurred in the TCS incubators after the 2-day exposure. Indeed, relative abundances of *A.*
249 *minutissimum* decreased by about 6%, whereas increased percentages of *G. angustatum*
250 (almost reaching 10 %) and *F. capucina* (from 0.2 to 2.7 %) were observed. Twisted valves of
251 *F. capucina*, *G. angustatum* and *U. ulna* were also recorded in the TCS-exposed communities
252 but cumulative abundances of abnormal cells did not exceed 2 % (at T2d). At the end of the
253 experiment, diatom community structure was comparable between treatments in terms of
254 dominant species ($p = 0.095$).

255 Diatom species richness and Shannon diversity displayed differences between controls and
256 TCS-incubators. At T0d, Shannon index was of 0.70 ± 0.10 ; it slightly decreased in the
257 controls down to about 0.61 at T15d and was much lower (0.33) in the TCS-treated samples.
258 Species richness also suffered a significant decrease in the treated incubators (13.7 ± 0.8
259 taxa), whereas it was constant through time in the controls (16.7 ± 0.5 taxa).

260

261 Discussion

262 *TCS dose – response relationships*

263 As a bactericide, triclosan inhibits the enzyme enoyl-acyl carrier protein reductase (ENR),
264 which is involved in the bacterial lipid biosynthesis (Levy *et al.* 1999; McMurry *et al.* 1998).
265 However, it is also known as a broad-spectrum antimicrobial agent and as such it could
266 generate effects on non-target components of the fluvial community (Wilson *et al.* 2003). The
267 results obtained in this study reinforced this hypothesis and showed effects on non-target
268 organisms within the biofilm community. Chlorophyll-a *in vivo* fluorescence parameters were
269 expected to show significant variations among the concentrations tested (Franz *et al.* 2008),
270 that were not observed in our study. However, diatom countings provided information about
271 some possible unexpected effects of TCS towards algae. The total number of live cells
272 enumerated per unit surface area seemed to indicate that, at quite low concentrations, diatoms

273 could be favoured (higher cell densities), probably reflecting a decrease in competition
274 between bacteria and diatoms (Grover 2000) due to the bactericidal effects of TCS (Escalada
275 *et al.* 2005). Indeed, bacterial mortality was significantly higher in the TCS-treated channels
276 compared to the controls ($p < 0.001$, data not shown). Toxicant effects on non-target
277 organisms are common in biofilms, where algae and bacteria interact in a small space (Ricart
278 *et al.* 2009). The L/D ratio (Figure 2) confirmed the adverse effects of TCS on diatom
279 communities, inhibiting diatom community growth and / or increasing mortality at the highest
280 concentrations tested.

281 In the first experiment, performed in artificial channels, we observed 50% diatom mortality at
282 concentrations far higher than those obtained by Sanderson and Thomsen (2009), who found
283 values for 24h-LC₅₀ of 4.5 µgTCS/L for algal growth in laboratory cultures. However, our
284 results were consistent with the concentration effects determined by Franz *et al.* (2008). This
285 variation of results between experiments underlines the relevance of abiotic factors
286 (experimental conditions, and in particular water flow regime modifying boundary layer
287 effects, and thus TCS bioavailability) in the assessment of the hazards of toxicants. Besides,
288 Franz *et al.* (2008) compared the tolerance of monospecific diatom suspensions and semi-
289 naturally grown biofilms and reported different responses depending on biotic factors (e.g.
290 single species tests *vs.* life forms and community structure). Increased resistance of a natural
291 algal community (the object of our study) by comparison to laboratory culture could be linked
292 to the intrinsic characteristics of the biofilms, in which the cells are embedded in a
293 polysaccharide matrix acting as a protective barrier. The thickness of the matrix also leads to
294 diffusion gradients within the biofilms, limiting the penetration of triclosan in the inner layers.
295

296 *Long term effects of a 48h-pulse of triclosan*

297 The moderate increase of the *in vivo* fluorescence parameter Y_{max} in the exposed samples
298 could be attributed to a hormetic response of the community (Calabrese 2005), *i.e.* to the
299 capacity of the system to initiate damage-repair processes. This adaptative response could
300 either be linked to enhanced photosynthetic capacity of the diatoms, or to structural changes
301 in the community favouring species with higher efficiency (Sabater *et al.* 2007). Previous
302 studies (Bérard & Pelte 1999; Ricart *et al.* 2009; Serra *et al.* 2009) have already described
303 increases in chlorophyll concentration and photosynthetic capacity under sublethal pulses of
304 toxicants. Here the persistent hormetic response suggested that the compensation process was
305 not totally achieved, or that exposure durably selected for species with higher photosynthetic
306 potential, able to modify community tolerance to further exposure (Blanck *et al.* 1988).

307 The two quantitative diatom estimates (cell densities, L/D ratios) indicated deleterious effects
308 of a 60 µg/L triclosan concentration on diatoms, lasting in time. The exponential growth of
309 diatoms (as seen in the controls) was significantly delayed by the 48-hour exposure to
310 triclosan. Cell densities after 2 weeks were not as high in the TCS-treated incubators as they
311 were in the controls, which could also indicate a persistence of toxicant exposure linked to
312 TCS or methylTCS bioaccumulation within biofilms (Coogan *et al.* 2007). The L/D ratio also
313 expressed an increased mortality of the communities exposed, with a 6.1 ± 0.5 % decrease in
314 the ratio after the 48-h exposure. This value fitted the curve of Experiment 1 ($R^2=0.90$; $p <$
315 0.001), suggesting that the response of this endpoint was not significantly modified by the
316 differences in flow conditions between Experiments 1 and 2. After 15 days, the communities
317 recovered slightly, although they did not attain the L/D ratio of the controls. This was also
318 seen in the diatom community composition, which was quite comparable between controls
319 and treated incubators at the end of the experiment. However, when looking at the life forms
320 (Figure 5), some differences were observed, the treated samples having a lower proportion of

321 species forming clumps (4%, like *Ulnaria ulna* or *Gomphonema* sp.) or filaments (0.6%, e.g.
322 *Melosira varians* and *Fragilaria* sp.), than in the controls (7.5 and 1.5% resp.). Lawrence *et*
323 *al.* (2009) observed that the biofilm architecture was affected by relatively low concentrations
324 of TCS (10 µg/L), that caused a less tightly packed structure and the disappearance of
325 filamentous species. In the present study, TCS exposure was also followed by a decrease in
326 colony-forming diatom species. Moreover, the control glass slides were covered by other
327 organisms with filamentous physiognomy, which were almost absent in the TCS-treated
328 incubators, even after the 2-week recovery period (L. Proia, unpubl. data).

329

330 Data about TCS effects on non-target organisms such as algal communities are scarce, and
331 apparently contradictory. However, the results of this study indicate that high concentrations
332 of pharmaceuticals occurring in sewage water may have unexpected effects on the receiving
333 aquatic ecosystems. Given the current lack of knowledge about sensitivities to PPCPs and the
334 multiplicity of toxic modes of action, it cannot be expected at this stage that any index of
335 toxicity based on diatom community structure can be immediately developed but these results
336 confirm the high sensitivity of diatoms to various environmental stressors, including
337 pharmaceutical inputs. Diatom mortality as inferred from the L/D ratio proved to be sensitive
338 to TCS contamination, and could be a useful tool for future biomonitoring purposes.

339

340

341 Acknowledgements

342 This study was partially funded by the European projects Modelkey (511237, GOCE) and
343 Keybioeffects (MRTN-CT-2006-035695). The authors are grateful to M. Laviale and S.
344 Boutry for their help in analysing the data.

345

346

347 References

348 Aranami K & Readman JW 2007. Photolytic degradation of triclosan in freshwater and

349 seawater. *Chemosphere* 66(6):1052-1056

350 Balmer ME, Poiger T, Droz C, Romanin K, Bergqvist P-A, Muller MD & Buser H-R 2004.

351 Occurrence of methyl triclosan, a transformation product of the bactericide triclosan, in fish

352 from various lakes in Switzerland. *Environ. Sci. Technol.* 38(2):390-395

353 Bérard A & Pelte T 1999. Les herbicides inhibiteurs du photosystème II, effets sur les

354 communautés algales et leur dynamique. *Rev. Sci. Eau* 12(2):333-361

355 Berney M, Hammes F, Bosshard F, Weilenmann H-U & Egli T 2007. Assessment and

356 interpretation of bacterial viability by using the LIVE/DEAD BacLight Kit in combination

357 with flow cytometry. *Appl. Environ. Microbiol.* 73(10):3283-3290

358 Blanck H, Wängberg SA & Molander S 1988. Pollution-Induced Community Tolerance - A

359 new ecotoxicological tool. *In* Functional Testing of Aquatic Biota for estimating Hazards of

360 Chemicals, Cairns J. Jr, Pratt J.R. (Eds), Philadelphia, 219-230 p.

361 Bukhtiyarova LN 2006. Additional data on the diatom genus *Karayevia* and a proposal to

362 reject the genus *Kolbesia*. *Nova Hedw. Beih.* 130:85-96

363 Buth JM, Grandbois M, Vikesland PJ, McNeill K & Arnold WA 2009. Aquatic

364 photochemistry of chlorinated triclosan derivatives: Potential source of polychlorodibenzo-*p*-

365 dioxins. *Environ. Toxicol. Chem.* 28(12):2555-2563

366 Calabrese EJ 2005. Paradigm lost, paradigm found: The re-emergence of hormesis as a

367 fundamental dose response model in the toxicological sciences. *Environ. Pollut.* 138(3):378-

368 411

369 Canosa P, Morales S, Rodríguez I, Rubí E, Cela R & Gómez M 2005. Aquatic degradation of
370 triclosan and formation of toxic chlorophenols in presence of low concentrations of free
371 chlorine. *Anal. Bioanal. Chem.* 383(7):1119-1126

372 Capdevielle M, Van Egmond R, Whelan M, Versteeg D, Hofmann-Kamensky M, Inauen J,
373 Cunningham V & Woltering D 2008. Consideration of exposure and species sensitivity of
374 triclosan in the freshwater environment. *Integr. Environ. Assess. Manag.* 4(1):15-23

375 Compère P 2001. *Ulnaria* (Kützinger) Compère, a new genus name for *Fragilaria* subgen.
376 *Alterasynedra* Lange-Bertalot with comments on the typification of *Synedra* Ehrenberg. In
377 Lange-Bertalot-Festschrift: Studies on Diatoms, Jahn R, Kociolek JP, Witkowski A, Compère
378 P. ARG Gantner Verlag, Ruggell, 97-102 p.

379 Coogan MA, Edziyie RE, La Point TW & Venables BJ 2007. Algal bioaccumulation of
380 triclocarban, triclosan, and methyl-triclosan in a North Texas wastewater treatment plant
381 receiving stream. *Chemosphere* 67(10):1911-1918

382 Cox EJ 1996. Identification of freshwater diatoms from live material. Chapman & Hall
383 London.

384 Dell'Anno A, Fabiano M, Bompadre S, Armeni M, Leone L & Danovaro R 1999.
385 Phytopigment and DNA determinations in long-time formalin-preserved trap samples. *Mar.*
386 *Ecol. Prog. Ser.* 191:71-77

387 Escalada MG, Russell AD, Maillard JY & Ochs D 2005. Triclosan-bacteria interactions:
388 single or multiple target sites? *Lett. Appl. Microbiol.* 41(6):476-481

389 Fent K, Weston AA & Caminada D 2006. Ecotoxicology of human pharmaceuticals. *Aquat.*
390 *Toxicol.* 76(2):122-159

391 Franz S, Altenburger R, Heilmeyer H & Schmitt-Jansen M 2008. What contributes to the
392 sensitivity of microalgae to triclosan? *Aquat. Toxicol.* 90(2):102-108

393 Grover JP 2000. Resource competition and community structure in aquatic micro-organisms:
394 experimental studies of algae and bacteria along a gradient of organic carbon to inorganic
395 phosphorus supply. *J. Plankton Res.* 22(8):1591-1610

396 Guillard RRL 1973. Division rates. *In* Culture methods and growth measurements. Handbook
397 of phycological methods, Stein JR ed, Cambridge University Press, Cambridge, 289-311 p.

398 Hudon C & Legendre P 1987. The ecological implications of growth forms in epibenthic
399 diatoms. *J. Phycol.* 23(3):434-441

400 Ihaka R & Gentleman R 1996. R: A language for data analysis and graphics. *J. Comput.*
401 *Graph. Statist.* 5299-314

402 Katoh K 1992. Correlation between cell density and dominant growth form of epilithic diatom
403 assemblages. *Diat. Res.* 777-86

404 Kelly MG, Bennion H, Cox EJ, Goldsmith B, Jamieson J, Juggins S, Mann DG & Telford RJ.
405 2005. Common freshwater diatoms of Britain and Ireland: an interactive key. *In* Environment
406 Agency, Bristol

407 Krammer K 1997. Die cymbelloiden Diatomeen. Teil 2: Encyonema part., Encyonopsis und
408 Cymbellopsis. Bibliotheca Diatomologica Band 37. J. Cramer, Stuttgart

409 Krammer K & Lange-Bertalot H 1986 - 1991. Bacillariophyceae 1. Teil: Naviculaceae; 2.
410 Teil: Bacillariaceae, Epithemiaceae, Surirellaceae; 3. Teil: Centrales, Fragilariaceae,
411 Eunotiaceae; 4. Teil: Achnanthaceae. Kritische Ergänzungen zu *Navicula* (Lineolatae) und
412 *Gomphonema*. G. Fischer Verlag., Stuttgart

413 Lawrence JR, Zhu B, Swerhone GDW, Roy J, Wassenaar LI, Topp E & Korber DR 2009.
414 Comparative microscale analysis of the effects of triclosan and triclocarban on the structure
415 and function of river biofilm communities. *Sci. Total Environ.* 407(10):3307-3316

416 Levy CW, Roujeinikova A, Sedelnikova S, Baker PJ, Stuitje AR, Slabas AR, Rice DW &
417 Rafferty JB 1999. Molecular basis of triclosan activity. *Nature* 398(6726):383-384

418 McMurry LM, Oethinger M & Levy SB 1998. Overexpression of *marA*, *soxS*, or *acrAB*
419 produces resistance to triclosan in laboratory and clinical strains of *Escherichia coli*. *FEMS*
420 *Microbiol. Lett.* 166(2):305-309

421 Murphy J & Riley JP 1962. A modified single solution method for the determination of
422 phosphate in natural waters. *Anal. Chim. Acta* 27:31-36

423 Orvos DR, Versteeg DJ, Inauen J, Capdevielle M, Rothenstein A & Cunningham V 2002.
424 Aquatic toxicity of triclosan. *Environ. Toxicol. Chem.* 21(7):1338-1349

425 Potapova M 2006. *Achnantheidium zhakovschikovii* sp. nov. (Bacillariophyta) and related
426 species from rivers of Northwestern Russia. *Nova Hedw.* 82(3-4):399-408

427 Ricart M, Barceló D, Geiszinger A, Guasch H, López de Alda M, Romaní AM, Vidal G,
428 Villagrasa M, Sabater S 2009. Effects of low concentrations of the phenylurea herbicide
429 diuron on biofilm algae and bacteria. *Chemosphere* 76(10):1392-1401

430 Ricart M, Guasch H, Alberch M, Bonnineau C, Farré M, Ferrer J, Geiszinger A, Morin S,
431 Proia L, Ricciardi F, Romaní AM, Sala LI, Sureda D & Sabater S. Effects of short term
432 exposure of triclosan on biofilm communities (submitted)

433 Ricciardi F, Bonnineau C, Faggiano L, Geiszinger A, Guasch H, Lopez J, Muñoz I, Proia L,
434 Ricart M, Romaní AM, Sabater S 2009. Is the chemical contamination linked to the diversity
435 of biological communities in rivers? *Trends Analyt. Chem.* 28(5):592-602

436 Round FE & Bukhtiyarova L 1996. Four new genera based on *Achnanthes* (*Achnantheidium*)
437 together with a re-definition of *Achnantheidium*. *Diat. Res.* 11:345-361

438 Sabater S, Guasch H, Ricart M, Romaní AM, Vidal G, Klünder C, Schmitt-Jansen M 2007.
439 Monitoring the effects of chemicals on biological communities. The biofilm as an interface.
440 *Anal. Bioanal. Chem.* 387(4):1425-1434

441 Sala SE, Guerrero JM, Ferrario ME 1993. Redefinition of *Reimeria sinuata* (Gregory)
442 Kociolek & Stoermer and recognition of *Reimeria uniseriata* nov. spec. *Diat. Res.* 8:439-446

443 Sanderson H & Thomsen M 2009. Comparative analysis of pharmaceuticals versus industrial
444 chemicals acute aquatic toxicity classification according to the United Nations classification
445 system for chemicals. Assessment of the (Q)SAR predictability of pharmaceuticals acute
446 aquatic toxicity and their predominant acute toxic mode-of-action. *Toxicol. Lett.* 187(2):84-93

447 Schmitt-Jansen M & Altenburger R 2007. The use of pulse-amplitude modulated (PAM)
448 fluorescence-based methods to evaluate effects of herbicides in microalgal systems of
449 different complexity. *Toxicol. Environ. Chem.* 89(4):665-681

450 Serra A, Corcoll N & Guasch H 2009. Copper accumulation and toxicity in fluvial
451 periphyton: The influence of exposure history. *Chemosphere* 74(5):633-641

452 Singer H, Muller S, Tixier C & Pillonel L 2002. Triclosan: Occurrence and fate of a widely
453 used biocide in the aquatic environment: Field measurements in wastewater treatment plants,
454 surface waters, and lake sediments. *Environ. Sci. Technol.* 36(23):4998-5004

455 Sivaraman S, Sullivan TJ, Johnson F, Novichenok P, Cui G, Simmerling C & Tonge PJ 2004.
456 Inhibition of the bacterial enoyl reductase FabI by triclosan: A structure-reactivity analysis of
457 FabI inhibition by triclosan analogues. *J. Med. Chem.* 47(3):509-518

458 Villalaín J, Mateo CR, Aranda FJ, Shapiro S & Micol V 2001. Membranotropic effects of the
459 antibacterial agent triclosan. *Arch. Biochem. Biophys.* 390(1):128-136

460 Wilson BA, Smith VH, deNoyelles F & Larive CK 2003. Effects of three Pharmaceutical and
461 Personal Care Products on natural freshwater algal assemblages. *Environ. Sci. Technol.*
462 37(9):1713-1719

463 Wind T, Werner U, Jacob M & Hauk A 2004. Environmental concentrations of boron, LAS,
464 EDTA, NTA and Triclosan simulated with GREAT-ER in the river Itter. *Chemosphere*
465 54(8):1145-1154

466 Wong-Wah-Chung P, Rafqah S, Vyard G & Sarakha M 2007. Photochemical behaviour of
467 triclosan in aqueous solutions: Kinetic and analytical studies. *J. Photochem. Photobiol. A*
468 *Chem.* 191(2-3):201-208
469

470 Table 1. Mean values \pm standard error (number of samples analysed) of physicochemical
471 parameters during the course of both experiments. n.c. nominal concentration.

472

	Experiment 1	Experiment 2
pH	7.7 ± 0.0 ($n=8$)	7.9 ± 0.5 ($n=11$)
Temperature ($^{\circ}\text{C}$)	15.8 ± 0.0 ($n=8$)	17.5 ± 1.1 ($n=11$)
Conductivity ($\mu\text{S}/\text{cm}$)	289.9 ± 0.2 ($n=8$)	143.4 ± 34.1 ($n=11$)
Dissolved oxygen (mg/L)	9.0 ± 0.1 ($n=8$)	9.2 ± 0.2 ($n=11$)
Oxygen saturation (%)	94.3 ± 0.4 ($n=8$)	7.9 ± 0.5 ($n=11$)
NO_3 (mg/L)	1.85 ± 0.17 ($n=16$)	0.75 (<i>n.c.</i>)
PO_4 ($\mu\text{g}/\text{L}$)	77.3 ± 8.6 ($n=16$)	16.9 ± 2.1 ($n=12$)

473

474 Table 2. Nominal and effective concentrations of triclosan ($\mu\text{g/L}$) in the channels (Experiment
475 1). d.l.: detection limit.

476

Nominal	0 (Control)	0 (Control solvent)	0.05	5	25	125	250	500
Measured	< d.l.	< d.l.	4.8	5.4	13.9	120.7	300.0	461.3


477

478 Figure captions

479 Figure 1

480 Densities of live diatom cells (number of cells/cm²) after a 48h-exposure, for the different
481 concentrations of triclosan measured in the channels.

482 Note the logarithmic scale on the X-axis



483 Figure 2


484 Triclosan dose-response curve for diatom live / dead ratio (in percentage of average control)

485 after a 48h-exposure ($R^2 = 0.90$; $p < 0.001$).

486 Note the logarithmic scale on the X-axis

487

488



489 Figure 3

490 Diatom cell densities (in ln cells/cm²) in the control (○) and treated (▲) incubators during the

491 course of Experiment 2.

492

493

494

495

496

497


498

499

500

501

502




503 Figure 4

504 Changes in species relative abundances during the long-term response experiment. Species

505 abbreviations: ADMI: *Achnantheidium minutissimum*, ADPY: *A. pyrenaicum*, FCAP:

506 *Fragilaria capucina* var. *capucina*, GANG: *Gomphonema angustatum*, GMIN: *G. minutum*,

507 UULN: *Ulnaria ulna*.




508

509 Figure 5

510 Distribution of diatom life-forms with respect to taxon abundances and specific cell

511 biovolume during the course of Experiment 2.



512