



HAL
open science

Development of q-PCR approaches to assess water quality: Effects of direct cadmium exposure on gene expression of the diatom *Eolimna minima*

S. Kim Tiam, A. Feurtet Mazel, François Delmas, Nicolas Mazzella, Soizic Morin, G. Daffe, P. Gonzalez

► To cite this version:

S. Kim Tiam, A. Feurtet Mazel, François Delmas, Nicolas Mazzella, Soizic Morin, et al.. Development of q-PCR approaches to assess water quality: Effects of direct cadmium exposure on gene expression of the diatom *Eolimna minima*. 21st SETAC Europe Annual Meeting, May 2011, Milan, Italy. pp.1, 2011. hal-02595066

HAL Id: hal-02595066

<https://hal.inrae.fr/hal-02595066>

Submitted on 15 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Development of q-PCR approaches to assess water quality: Effects of direct cadmium exposure on gene expression of the diatom *Eolimna minima*.



Sandra Kim Tiam^{1,2}, Agnès Feurtet-Mazel², François Delmas¹, Nicolas Mazzella¹, Soizic Morin¹, Guillemine Daffe², Patrice Gonzalez²

¹ Cemagref, UR REBX, 50 avenue de Verdun, F-33612 Cestas cedex, France; Tél : +33 (0)557-89-09-81; E-mail : sandra.kim-tiam@cemagref.fr

² EPOC, UMR CNRS 5805, Place du Dr Peyneau, 33120 Arcachon, France

Introduction and objectives

In regard to the degradation of water quality since the last decades, appropriate diagnostic tools have been implemented. In France, water agencies in collaboration with Cemagref have developed a diatom index to estimate global water quality. The Biological Diatom Index (BDI, Coste *et al.* 2009) is routinely used for monitoring applications in European territories and has been standardized (NF T 90-354).

Nevertheless, indices currently used for water quality assessment don't take in consideration metal contamination in their conception, despite the high bioaccumulation and impact shown on periphytic diatom communities (da Silva *et al.*, 2009, Cunningham *et al.*, 2005, Feurtet-Mazel *et al.*, 2003). Moreover these methods are time consuming and require important taxonomic knowledge. In this context development of q-PCR approaches are of particular interest.

In our study we have developed a new RNA extraction method for diatoms, 8 genes of interest for the diatom *Eolimna minima* have been selected, sequenced and deposited in the GenBank. The responses of the q-PCR tools developed have been tested after Cd exposure on *Eolimna minima*. Bioaccumulation and population kinetics were also followed.

Materials and methods

Molecular methodological developments:

- Use of a vortex and glass beads for diatoms RNA extraction.
- Sequencing of 8 genes of interest for the diatom *Eolimna minima* and design of qPCR primers.

Assessment of cadmium toxicity on *Eolimna minima*:

- Eolimna minima*: a metal-tolerant freshwater diatom commonly collected within periphytic biofilm samples in running water and especially in metal-contaminated areas (Morin *et al.*, in press).
- Exposure by direct route to 0, 10 and 100 µgCd/L during 14 days (noted C₀, C₁ and C₂).




Results and discussion



Molecular methodological developments:

- RNA extraction method: efficient, rapid, simple and low cost.
- Sequencing, deposit on the Genbank and design of specific primers.

Gene name	Accession number	Primer (5'-3')
<i>sodMn</i>	HM 449706	GGTAGTAGGGGTGCTCCC ⁺ CCAGGACAACCCGCTC ⁺
<i>cox1</i>	HM 449704	CAGTAATTCTCAC ⁺ TGCCAGC ⁺ CCGTGTACCCACGGTGT ⁺
<i>nad5</i>	HM 449708	TCAACTGGTTTGCATCATGGC ⁺ TTGAACATACTGTTGTGGAAGC ⁺
<i>d1</i>	HM 449711	ACCACCAATACACCGCAAC ⁺ GGTCTTGGATTCTGTAGC ⁺
<i>psaA</i>	HM 449705	CATAAAGCGGCACCCAAAC ⁺ CTTGGATATACTGACTCATTAACTCAGG ⁺
<i>act</i>	HM 449707	GGCTCCACAAAACCCCAAG ⁺ GCGGTACCCCTCGTAGAT ⁺
<i>12s</i>	HM 449710	GCGGTAATACGGAGGATG ⁺ AGTGCCTTCGCCATCGG ⁺
<i>18s</i>	HM 449712	CATTGTCAAGGTGAAATCTTGGAA ⁺ CCCCGAACCCAAAAGT ⁺



Assessment of cadmium toxicity on *Eolimna minima*:



Functions	Genes	Cadmium contaminated experimental units								
		C1 (10.0 ± 3.2 µg/l)				C2 (96.0 ± 34.2 µg/l)				
		1	2	7	14	1	2	7	14	
Mitochondrial metabolism	<i>cox1</i>	/	/	/	/	/	/	/	9.5	/
	<i>nad5</i>	/	/	/	2.5	/	/	/	/	9.5
	<i>12s</i>	/	/	/	/	/	/	/	15	/
Oxidative stress	<i>sodMn</i>	/	/	/	/	/	/	/	/	/
Photosynthesis	<i>d1</i>	/	/	/	2	/	/	/	5.5	24
	<i>psaA</i>	/	/	/	2.5	/	/	/	7.5	48

1/ Growth

- Not significantly different in C₀ and C₁ over the whole duration of the experiment.
- Diatom cell density 3.2 times higher in C₀ than in C₂.
- Null diatom growth with C₂.

2/ Bioaccumulation

- Significant increase between days 7 and 14 for the 2 contaminated treatments.
- High levels of total accumulated Cd (C₁: 430.1 ± 86.4 µgCd/g dw and C₂: 734.1 ± 70 µgCd/g dw)

3/ Genetic response

- Amplification of 5 genes observed (*cox1*, *nad5*, *d1*, *psaA* and *12S*).
- Different genetic responses expressed as a function of time and concentration of exposure.
- Effect on mitochondrial and photosynthetic metabolism, observed after 7 days of exposure only at the highest contamination pressure, and for both contamination pressures on day 14.

Conclusions and perspectives

In this study, a new glass-bead RNA extraction technique for diatoms was successfully developed and optimized. Eight genes of interest were sequenced for *Eolimna minima* allowing the application of q-PCR tools to this species.

Our results underlined the toxicity of Cd towards *E. minima* population kinetics only for the highest concentration, while q-PCR analyses revealed an impact on mitochondrial metabolism and the chloroplast photosystem for both Cd exposure concentrations. Genetic expression of *nad5*, *cox1*, *12S*, *d1* and *psaA* by q-PCR could thus constitute an early warning biomarker of metal pollution. Future studies should investigate sequences of genes coding for catalase or glutathione peroxidase in order to study the response to oxidative stress.

The present study is the first reported use of q-PCR on river benthic diatoms and the results obtained are extremely promising. The techniques developed were successfully tested using simplified mixtures of diatom species. Further interesting steps for the early and sensitive assessment of metal pollution would firstly involve validating the results obtained by examining sensitive vs tolerant diatom species response levels, then finding or confirming genetic biomarkers for use on natural multispecific biofilms for impact assessment of toxic pollution.

Reference:

Kim-Tiam, S., Feurtet-Mazel A., Delmas F., Mazzella N., Morin S., Daffe G., Gonzalez P, (submitted to Water Research). Development of q-PCR approaches to assess water quality: Effects of direct cadmium exposure on gene expression of the diatom *Eolimna minima*.