



HAL
open science

Sediment bed destabilization induced by oscillating horizontal pressure gradients

Céline Berni, H. Michallet, Elizabeth Barthélemy

► **To cite this version:**

Céline Berni, H. Michallet, Elizabeth Barthélemy. Sediment bed destabilization induced by oscillating horizontal pressure gradients. EGU General Assembly, Apr 2014, Vienne, Austria. pp.1, 2014. hal-02600969

HAL Id: hal-02600969

<https://hal.inrae.fr/hal-02600969>

Submitted on 16 May 2020


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Céline Berni^{1,2}, Hervé Michallet¹, Eric Barthélemy¹


1. LEGI - UMR 5519, University of Grenoble, France,
2. Now at Irstea, UR HHLY, Centre de Lyon-Villeurbanne, France.

1. The LEGI wave flume : physical model of a beach with Shields and Rouse scaling



Measure of
- velocity (ADVP)
- pore pressure
- bed destabilization (acoustic & optical)
- free-surface elevation

2. Sleath model of plug flow



p pore pressure, T friction on the still bed,
 h_L height of the plug, τ shear stress

Horizontal equilibrium : $-h_L \frac{\partial p}{\partial x} dx + \tau dx - T dx = 0$

Friction law : $|T| = K_f \times C^* (\rho_s - \rho) g h_L$


Motion occurs if [2]

$$\left| \frac{d_{50}}{h_L} \theta - S \right| \geq K_f C^* \quad \text{condition for plug flow initiation}$$

in our experiments $K_f C^* \sim 0.3$


where $S = \frac{\partial p}{\partial x} / ((\rho_s - \rho)g)$; $\theta = \frac{\tau}{(\rho_s - \rho)g d_{50}}$

3. Bichromatic waves conditions



sequence repeated (>50 times) \Rightarrow ensemble averaged [3]:
turbulence and orbital flows


4. Turbulence and shear stress




Decomposition of shear stress in 3 terms [4].
Predominance of the orbital term; viscous and turbulence terms can be neglected.

$$\tau \approx -\rho u w$$

5. Time series



6. When is erosion triggered ?



Erosion is triggered by high S (in agreement with the field study [5]), mostly in the wave troughs, even for low shear.

[1] Grasso, F., Michallet, H., Barthélemy, E. & Certain, R. (2009) Physical modeling of intermediate cross-shore beach morphology: Transients and equilibrium states. *J. Geophys. Res.*, 114, C09001 114.
[2] Sleath, J. F. A. (1999) Conditions for plug formation in oscillatory flow. *Continental shelf Res.* 19 (13), 1643-1664.
[3] Berni, C., Barthélemy, E. & Michallet, H. (2013) Surf zone cross-shore boundary layer velocity asymmetry and skewness: an experimental study on a mobile bed. *J. Geophys. Res.*: Oceans 118, 2188-2200.
[4] Nielsen, P. (1992) *Coastal Bottom Boundary Layers and Sediment Transport*. Singapore: World Scientific.
[5] Foster, D. L., Bowen, A. J., Holman, R. A. & Natto, P. (2006) Field evidence of pressure gradient induced incipient motion. *J. Geophys. Res.*, 111, C05004.
This study was partly funded by the project MODLIT (DGA-SHOM/INSU-RELIEFS) and the project BARCAN (INSU-LEFE).