



HAL
open science

Contrasting abilities of metal bioaccumulation in Gammarus populations with different exposure histories

Nastassia Urien, Jérémie D. Lebrun, Antoine Farfarana, Arnaud Chaumot,
Lise C. Fechner, Olivier Geffard

► **To cite this version:**

Nastassia Urien, Jérémie D. Lebrun, Antoine Farfarana, Arnaud Chaumot, Lise C. Fechner, et al.. Contrasting abilities of metal bioaccumulation in Gammarus populations with different exposure histories. SETAC Europe 25th Annual Meeting, May 2015, Barcelone, Spain. Chair: Karel A.C. and De Schamphelaere, pp.1, 2015. hal-02601293

HAL Id: hal-02601293

<https://hal.inrae.fr/hal-02601293v1>

Submitted on 16 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Contrasting abilities of metal bioaccumulation in *Gammarus* populations with different exposure histories

N. Urien¹, J.D. Lebrun¹, A. Farfarana¹, A. Chaumot², L. Fechner¹, O. Geffard²

¹ Irstea, UR HBAN - Ecotoxicology, 1 rue Pierre-Gilles de Gennes, 92160 Antony, France

² Irstea, UR MALY, Ecotoxicology, 5 rue de la Doua, 69100 Villeurbanne, France

email: nastassia.urien@irstea.fr

SequAdapt
PROJECT



ED 341-E2M2
Evolution Ecosystèmes
Microbiologie
Modélisation
Ecole Doctorale

INTRODUCTION

- Kinetic models are used to link metal exposure to metal contamination in aquatic organisms and enable also the prediction of metal bioaccumulation.
- Kinetic models are characterised by kinetic parameters (uptake and elimination rate constants) which can be determined in controlled conditions.
- “Global” kinetic parameters for Cd, Pb and Ni have been established in five naïve populations of gammarids.
- BUT gammarids can live in rivers exhibiting metal contamination to which organisms may adapt physiologically.
- Adaptation can result in the modulation of bioaccumulation abilities i.e. kinetic parameters.
- This may limit the environmental relevance of kinetic models and the reliability of bioaccumulation predictions.

OBJECTIVES

Assess the bioaccumulation abilities of metals in gammarids chronically exposed to metals *in situ*.

1. Determine the kinetic parameters describing Cd, Pb and Ni bioaccumulation in populations of gammarids chronically exposed to metals *in situ*.

2. Compare the kinetic parameters with “global” kinetic parameters already determined in five naïve populations of gammarids.




MATERIALS & METHODS

1 Collection of 3 gammarids populations from metal-contaminated sites

2 Exposure of gammarids to metals in controlled conditions

3 Fitting of the kinetic model to the bioaccumulation data

4 Determination of kinetic parameters for each contaminated population



C_a : Metal concentration in gammarids ($\mu\text{g g}^{-1}$)
 k_{in} : Uptake rate constant ($\text{L g}^{-1} \text{d}^{-1}$)
 k_{out} : Elimination rate constant (d^{-1}) / t: time (d)
 C_{water} : Dissolved metal concentration ($\mu\text{g L}^{-1}$)

→ k_{in} and k_{out}

5 Comparison with the “global” kinetic parameters already established in five naïve populations living in pristine sites but exhibiting contrasted geochemical parameters

(See Plateform: «Between-population variability of waterborne metal bioaccumulation in *Gammarus* sp. from uncontaminated freshwaters», at 2:30 PM, ID 149, room 131/132)

RESULTS & DISCUSSION

1. Bioaccumulation kinetics




Fig. 1. Accumulation and elimination kinetics of Cd in gammarids from contaminated rivers after exposure to Cd in the laboratory. A point represents a pool of 5 gammarids.

For GAL and BIE:

- k_{in} was successfully determined for Cd.
- k_{out} was not determined because of the high variability between replicates and the inadequate elimination phase length.

For ARDI:

- High basal Cd content
- No new accumulation
- No kinetic parameters




Fig. 2. Accumulation and elimination kinetics of Pb in gammarids from contaminated rivers after exposure to Pb in the laboratory. A point represents a pool of 5 gammarids.

For GAL, BIE and ARDI:

- k_{in} and k_{out} were successfully determined for Pb and Ni.





Fig. 3. Accumulation and elimination kinetics of Ni in gammarids from contaminated rivers after exposure to Ni in the laboratory. A point represents a pool of 5 gammarids.

2. Comparison with the “global” kinetic parameters

k_{in} - Uptake rate constant



Cd:

- $k_{in} < k_{in}$ in naïve populations

GAL and BIE: Populations chronically exposed to metals *in situ* decrease their Cd uptake compared to naïve populations.

→ Physiological adaptation

ARDI: High basal Cd content in organisms chronically exposure to Cd *in situ*.

→ Storage under detoxified forms

Pb:

- $k_{in} > k_{in}$ in naïve populations
- $k_{out} < k_{out}$ in naïve populations

GAL, BIE, ARDI: Population chronically exposed to metals *in situ* tend to accumulate more Pb and to eliminate less Pb compared to naïve populations.

→ Storage abilities


Ni:

- $k_{in} < k_{in}$ in naïve populations
- $k_{out} < k_{out}$ in naïve populations

GAL, BIE, ARDI: Populations chronically exposed to metals *in situ* accumulate and eliminate less Ni compared to naïve populations.

→ Regulation abilities

k_{out} - Elimination rate constant



Legend: GAL (green), BIE (orange), ARDI (pink), Global kinetic parameters (dashed line), [95% IC-inf - 95% IC-sup] (shaded area)

Fig. 4. Kinetic parameters, k_{in} and k_{out} , determined for each population of gammarids from contaminated sites (with 95% confidence interval) and comparison with the “global” kinetic parameters determined in 5 naïve populations and represented by the upper and lower values of 95% confidence interval.

These results suggest that tolerance mechanisms occur in chronically contaminated populations with metals.

CONCLUSION

- Bioaccumulation strategies are specific to the metal considered.
- Populations chronically exposed to metals *in situ* modulate their metal bioaccumulation abilities compared to naïve populations.
- The modulation of metal bioaccumulation does not seem to be specific to the type of metal encountered *in situ* but to a global metallic pressure.

- Predictive models require to well defined the population used to determined k_{in} and k_{out} and, thus, to calibrate models.
- Further studies are needed to assess the physiological mechanisms involved in the modulation of metal bioaccumulation.