



HAL
open science

Towards understanding the organisation of metacommunities in highly dynamic ecological systems

T. Datry, N. Bonada, J. Heino

► **To cite this version:**

T. Datry, N. Bonada, J. Heino. Towards understanding the organisation of metacommunities in highly dynamic ecological systems. *Oikos*, 2016, 125 (2), pp.149-159. 10.1111/OIK.02922 . hal-02601969

HAL Id: hal-02601969

<https://hal.inrae.fr/hal-02601969v1>

Submitted on 16 Aug 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Towards understanding the organisation of metacommunities in highly**
2 **dynamic ecological systems**

3 Thibault Datry^{1,2}, Núria Bonada³ & Jani Heino⁴

4 ¹ IRSTEA, UR-MALY, centre de Lyon-Villeurbanne, 5 rue de la Doua CS70077, 69626
5 VILLEURBANNE Cedex, France.

6 ² UMR “BOREA” CNRS 7208/IRD 207/MNHN/UPMC, DMPA, Museum National
7 d’Histoire Naturelle, 43 rue Cuvier, 75231 Paris Cedex, France

8 ³ Grup de Recerca Freshwater Ecology and Management (FEM), Departament
9 d’Ecologia. Facultat de Biologia. Universitat de Barcelona (UB), Diagonal 643, 08028-
10 Barcelona, Catalonia, Spain.

11 ⁴ Finnish Environment Institute, Natural Environment Centre, Biodiversity, Paavo
12 Havaksen Tie 3, FI-90570 Oulu, Finland.

13

14 **Running head:** Metacommunities in dynamic ecological systems

15

16 **Key words:** Coexistence, dispersal, species sorting, niche, instabilities, intermittent
17 rivers, environmental heterogeneity

18

19 **Type of article:** Forum

20

21 **Corresponding author:** T. Datry, IRSTEA, UR-MALY, 5 rue de la Doua CS70077
22 69626 VILLEURBANNE Cedex, France. Tel (33) 4.72.20.87.55 Fax (33)
23 4.78.47.78.75

24

25

26 **Abstract: 189 words**
27 **Text (including references and illustrations): 8142 words**
28 **Main text: 5072 words**
29 **4 figures and 2 Tables**
30 **85 references**
31 **Authorship: TD, NB and JH conceived and co-wrote the manuscript.**

32

33 **Abstract**

34 Community ecology recognises today that local biological communities are not only
35 affected by local biotic interactions and abiotic environmental conditions, but also by
36 regional processes (e.g. dispersal). While much is known about how metacommunities
37 are organised in space in terrestrial, marine and freshwater ecological systems, their
38 temporal variations remain poorly studied. Here, we address the question of the
39 dynamics of metacommunities in highly variable systems, using intermittent rivers
40 (IRs), those rivers which temporarily stop flowing or dry up, as a model system. We
41 first review how habitat heterogeneity in space and time influences metacommunity
42 organisation. Second, we compare the metacommunities in IRs to those in perennial
43 rivers (PRs) and develop the idea that IRs could undergo highly dynamic shifts due to
44 the temporal variability in local and regional community processes. Third, we develop
45 the idea that in IRs, metacommunities of the wet and dry phases of IRs are closely
46 intertwined, thereby increasing even more their respective temporal dynamics. Last, we
47 provide a roadmap to stimulate further conceptual and empirical developments of
48 metacommunity research and identify possible applications for improving the
49 management of IRs and other highly dynamic ecological systems.

50

51 **1. Introduction**

52 Community ecology has progressed rapidly in recent years owing to the recognition
53 that local communities are not spatially closed and temporally stable (Leibold et al.
54 2004; Ricklefs 2008). Current views thus emphasise that local communities are not only
55 affected by local abiotic environmental conditions and biotic interactions, but also by
56 processes external to local ecological systems and operating at a regional scale, such as
57 speciation, extinction, immigration and emigration (Hubbell 2001). Although the
58 foundation of this idea dates back several decades (MacArthur & Wilson 1967), a shift
59 from purely local views of community organisation to those that also acknowledge the
60 importance of regional processes has been increasingly evident in the past years
61 (Hubbell 2001; Leibold et al. 2004). Sets of local communities linked by dispersal, or
62 metacommunities (**Table 1**), have been studied intensively in various ecological
63 systems (Logue et al. 2011), including terrestrial (e.g. Meynard et al. 2013), marine (e.g.
64 Moritz et al. 2013), and freshwater (e.g. Heino 2013) systems, and the number of
65 studies on the topic continues to increase rapidly (Heino et al. 2015). The
66 metacommunity framework has thus become a conceptually sound and empirically
67 well-explored framework to study the spatio-temporal organisation of communities.

68

69 As an outcome of developments in spatial and dispersal ecology, different
70 metacommunity models have been proposed to explain how both environmental
71 filtering (i.e. local biotic interactions and abiotic environmental conditions, **Table 1**),
72 and dispersal processes (i.e. the movement of individuals from one site to another
73 within a region, **Table 1**), interact to shape local community structure (Vellend 2010,
74 Logue et al. 2011; Winegardner et al. 2012). For example, “mass effects” models
75 predict that high rates of dispersal can obscure the effect of environmental filtering,

76 while the “patch dynamic” models assume that the best dispersers arrive first and
77 occupy the patches (sensu Logue et al. 2011) as long as no more competitive species
78 have arrived (Logue et al. 2011). Although the potential differences between these
79 various models have often been emphasised (Leibold et al. 2004; Logue et al. 2011;
80 Winegardner et al. 2012), very few observational studies have succeeded in offering
81 unambiguous explanations for their relative importance in nature (Cottenie 2005;
82 Beisner et al. 2006; Bonada et al. 2012). One explanation stems from the fact that it is
83 highly difficult to measure dispersal directly and that separating the effects of
84 environmental filtering vs. dispersal processes is complex since most environmental
85 factors are spatially autocorrelated (Legendre et al. 2005; Tuomisto & Ruokolainen
86 2006; Jacobson & Peres-Neto 2010, Gilbert & Bennett 2010; Bonada et al. 2012).
87 Another explanation is that a number of metacommunity studies considered spatial
88 patterns of local communities as static within a given landscape (Leibold et al. 2004;
89 Presley et al. 2010), while communities can be very dynamic with abrupt and constant
90 change in richness and composition over very short scales (Chesson & Huntly 1989;
91 Azeria & Kolasa 2008; Erös et al. 2012; Aiken & Navarette 2014; Fernandes et al.
92 2014).

93

94 Logue et al. (2011) suggested that broadening the types of focal habitats and
95 organisms in metacommunity studies would provide a better understanding of the
96 variability of metacommunity organisation (**Table 1**). They specifically pointed out that
97 metacommunity studies should go beyond those focusing on insular habitats with
98 discrete boundaries and those using large organisms as models. Different types of
99 aquatic systems (e.g. marine, coastal, temporary ponds, estuaries, and running waters)
100 provide excellent opportunities for such additional studies because they harbour highly

101 disparate organismal groups, vary widely in their degree of connectivity, and exhibit
102 wide spatial and temporal variability in local habitat conditions (Heino et al. 2015).
103 However, previous focus on purely aquatic communities has provided limited
104 perspectives on highly dynamic ecological systems, such as intermittent rivers (IRs), the
105 rivers which cease to flow or dry up in time and space. IRs provide especially suitable
106 arenas for examining metacommunity organisation in highly dynamic ecological
107 systems because they are mosaics of aquatic and terrestrial habitats shifting constantly
108 in time and space (Larned et al. 2010; Datry et al. 2014a). This dynamism, combined
109 with the fact that nearly 50% of lengths of rivers across the globe are characterized by
110 intermittent flow (Acuña et al. 2014; Datry et al. 2014a), calls for addressing
111 metacommunity organisation in IRs and other highly dynamic systems.

112

113 Here, we address the question of how communities are organized in time and space
114 in highly dynamic systems. We use IRs as model ecological systems because of their
115 high dynamism and apply the metacommunity concept because of the possibility to
116 distinguish between local and regional processes in river systems (e.g. Brown & Swan
117 2010; Logue et al. 2011; Heino 2013). We first review the effects of habitat variability
118 on metacommunities in various dynamic systems. Second, we compare
119 metacommunities in IRs to those in perennial rivers (PRs) and develop the idea that IRs
120 could undergo constant and severe shifts due to the variability in environmental filtering
121 and dispersal processes. Third, we develop the idea that in IRs, metacommunities of the
122 wet and dry phases of IRs are closely intertwined, which enhances their respective
123 temporal dynamics. Last, we provide a roadmap to stimulate further conceptual and
124 empirical developments of metacommunity research and identify possible applications
125 for improving the management of IRs and other highly dynamic ecological systems.

126

127 **2. The importance of spatial and temporal dynamics for metacommunity**
128 **organisation**

129 In the metacommunity framework, landscapes are considered as networks of
130 interconnected focal habitats in a matrix of unsuitable habitats, i.e. they are spatially
131 heterogeneous (Leibold et al. 2004; Holyoak et al. 2005). However, many conceptual
132 and empirical developments have considered metacommunities as relatively stable
133 entities (e.g. Leibold et al. 2004; Presley et al. 2010; Carrara et al. 2012; Altermatt
134 2013), although local communities and their environments can be temporally variable
135 (Azeria & Kolasa 2008; Erös et al. 2012; Aiken & Navarette 2014; Fernandes et al.
136 2014). Factors that control metacommunity organisation, including habitat availability,
137 local environmental conditions, and spatial connectivity, vary in space and time (Aiken
138 & Navarette 2014). The most extreme aquatic systems are arguably those that alternate
139 between aquatic and terrestrial conditions on a short time scale, for which high
140 variability may be the predominant rule. This category includes rocky marine
141 shorelines, tidal zones, small freshwater rock pools, temporary wetlands, vernal pools,
142 floodplains, and IRs. In such systems, environmental conditions vary both spatially and
143 temporally on a short-term basis. For example, the surface areas and connectivity of
144 floodplain aquatic habitats vary considerably during periods of weeks to months
145 between alternating wet and dry phases (Fernandes et al. 2014). Tide cycles generate
146 short pulses of aquatic habitat expansion and contraction on a daily basis (Kirwan &
147 Murray 2007). In such systems, and more generally in other systems exhibiting high
148 environmental variability, a static view of metacommunity organisation is likely to be
149 inaccurate.

150

151 In dynamic systems (i.e., systems experiencing constant and severe changes) the
152 various mechanisms shaping local communities, as well as their respective importance,
153 are constantly varying. For example, the contribution of dispersal will increase heavily
154 after a disturbance to allow for recolonization of a given patch (defined as a discrete
155 area with favourable environmental conditions, typically a flowing section in a river for
156 a fish or aquatic invertebrates, but see Cavanaugh et al. 2014). Yet, after most colonists
157 have reached the patch, environmental filtering will become more important in
158 explaining the local community organisation in the absence of mass effects. In systems
159 experiencing variable environmental conditions, such shifts between environmental
160 filtering vs. dispersal processes may be common and generate dynamic patterns in
161 community structure including short-term instabilities (Drake 1990; Chesson & Huntly
162 1989; Aiken & Navarrete 2014).

163

164 Understanding and predicting the dynamics of metacommunities will be essential to
165 managing, conserving, and restoring biodiversity in all ecological systems, including in
166 freshwater ecological systems which have been drastically altered by global changes
167 (Dudgeon et al. 2006). Because extreme climatic events and disturbances to ecological
168 systems are occurring more frequently (Easterling et al. 2000; Parmesan, 2006), current
169 static views of community organisation are becoming less and less useful and, given
170 these limitations, we may fail to predict biodiversity loss accurately in disturbed
171 systems. Moreover, most current management, conservation and restoration applications
172 do not fully recognize the fact that maintaining the spatial and temporal dynamics of
173 entire ecological systems is essential (Heino 2013; Tonkin et al. 2014). This recognition
174 is certainly very important in the context of highly dynamic systems, where
175 communities are restructured again and again within a short period of time.

176

177 **3. Intermittent rivers as model systems to explore the dynamics of**
178 **metacommunities**

179 IRs occur under all types of climates on all continents, including Antarctica, and
180 make up the majority of river networks in terms of length in many regions (Acuña et al.
181 2014; Datry et al. 2014a). Globally, IRs represent 69% of the low-order streams south
182 of 60°N latitude and from 30 to 40% of the larger river basins (Raymond et al. 2013). In
183 the coming decades, the number and length of IRs will increase in many regions that
184 experience drying trends due to climate and land-cover change, and increasing water
185 abstraction for irrigation and other economic uses (Palmer et al. 2008; Larned et al.
186 2010).

187

188 Most conceptual and empirical developments in freshwater ecology, including the
189 application of metacommunity models (e.g. Brown & Swan 2010; Altermatt et al. 2011;
190 Heino et al. 2015), have emerged from and been produced for perennial river systems
191 (Datry et al. 2014a). Consequently, metacommunity ecology is still in its infancy in IRs,
192 and this contrasts with temporary lentic systems which have been widely used to test
193 predictions from metapopulation and metacommunity ecology (e.g. Kolasa & Romanuk
194 2005; Vanschoenwinkel et al. 2007) or have provided new insights into these fields (e.g.
195 Smol & Douglas 2007; Vanschoenwinkel et al. 2010). One important difference with
196 temporary lentic systems is that rivers occur as dendritic networks, which consist of
197 linearly-arranged, hierarchical and branching habitat elements (Fagan 2002; Grant et al.
198 2007; Altermatt 2013). Headwaters are more isolated than mainstem reaches and this
199 possibly results in spatial differences in the relative importance of community assembly
200 processes (Brown & Swan 2010). Moreover, unidirectional flow and directionally-

201 biased dispersal can exacerbate the effects of disturbance (drying) on species
202 coexistence (Altermatt et al. 2011).

203

204 IRs are dynamic shifting habitat mosaics of flowing, non-flowing and dry patches
205 (**Figure 1**), the extent and connectivity of which constantly vary across drainage basins
206 in response to river discharge and groundwater levels (Stanley et al. 1997; Jaeger et al.
207 2014; Datry et al. 2015, **Figure 2**). This spatially complex and temporally dynamic
208 habitat template is inhabited by both aquatic (e.g. fish, invertebrates, microbes, fungi,
209 algae) and terrestrial (e.g. birds, mammals, arthropods, microbes, fungi) organisms, the
210 communities of which alternate, coexist, interact, and experience extreme disturbances
211 in terms of drying and rewetting phases (Stanley et al. 1997; Datry et al. 2014a).
212 Aquatic and terrestrial communities can show intense biotic interactions. Predation by
213 terrestrial organisms on aquatic organisms trapped in drying pools can be an important
214 mortality factor in addition to interactions among fully aquatic organisms (Larned et al.
215 2010). Some large, lentic specialist predators, such as giant water bugs or diving
216 beetles, can also colonize the pools to prey on the stranded organisms, thereby
217 enhancing aquatic interactions (Gasith & Resh 1999; Boulton 2003). Altogether, local
218 environmental conditions, biotic interactions and accessibility to patches vary
219 continuously in IRs, challenging the current static views of metapopulation and
220 metacommunity approaches. Here, we make the case that IRs provide suitable arenas to
221 explore the temporal dynamics of metacommunities and notably the idea that
222 communities can experience highly dynamic shifts in structure and composition due to
223 the variability in environmental filtering and dispersal processes.

224

225 **4. Uniqueness of intermittent rivers and implications for metacommunity**
226 **dynamics**

227 IRs have some unique features that challenge current views of metacommunity
228 organisation in river systems, which are mostly derived from research in PRs. While
229 **Table 2** synthesises these features and their possible implications for metacommunity
230 organisation, the different hydrological phases through which IR communities have to
231 persist are presented below.

232 **Flow cessation: shifts from lotic to lentic conditions**

233 The most striking difference between PRs and IRs is that the flow ceases
234 periodically in IRs, and surface water can disappear completely from IR channels
235 (**Figure 1, Table 2**). Flow cessation gradually converts flowing river channels into
236 chains of disconnected pools of standing waters, which face increased water
237 temperatures and solute concentrations, and decreased pH and dissolved oxygen levels
238 (Boulton 2003). As these pools decrease in size, the densities of organisms can increase
239 dramatically, leading to strong intra- and interspecific interactions, such as competition
240 or predation. Some large pools can persist throughout dry phases and represent essential
241 refugia for aquatic organisms (Sheldon et al. 2010) and can be important “stepping
242 stones” facilitating the continued existence of populations and communities across an
243 IR network by connecting refuges and allowing recolonization of re-wetted sites by
244 actively flying or passively dispersing aquatic organisms (Bogan & Boersma 2012;
245 Cañedo-Argüelles et al. 2015). These shifts from lotic to lentic conditions occurring in
246 many IRs challenge the view that riverine systems, in general, are dominated by
247 physical constraints related to unidirectional flow (e.g. water velocity, shear stress)
248 (Fagan 2002; Altermatt et al. 2011). During the lentic phase, biotic interactions may be

249 the dominant processes structuring aquatic communities (Gasith & Resh 1999; Boulton
250 2003).

251 **Streambed drying**

252 In many IR systems, riffles dry first, after which remnant pools progressively dry
253 up, thus converting river channel into terrestrial habitat and leading to the disappearance
254 of aquatic communities (**Table 2**). Physiological tolerance to loss of water has been
255 well documented in organisms of temporary ponds and pools (Wiggins et al. 1980) and
256 a variety of species have also evolved physiological resistance strategies to cope with
257 desiccation in IRs (**Table 2**). This “invertebrate seedbank” contributes to community
258 recovery upon flow resumption, although its efficiency in such contribution is variable
259 and not completely understood (Warner & Chesson 1985; Snyder 2006; Stubbington &
260 Datry 2013). Described as the storage effect (**Table 1**), this mechanism contributes
261 strongly to species coexistence when resources are limiting and recruitment fluctuates
262 (Warner & Chesson 1985; Snyder 2006), thus promoting the resistance (defined as the
263 capacity to persist unchanged through a disturbance) of IR communities to drying.
264 During these dry phases, dry riverbeds are also being colonised by rich and abundant
265 terrestrial biotas, including microbes, plants, arthropods, birds, and mammals (Steward
266 et al. 2012). In spite of a recent increase in research on dry riverbed communities, they
267 still represent *terra incognita* from an ecological perspective (Steward et al. 2012).

268 **Rewetting**

269 Rewetting of previously dry patches is often sudden and unpredictable and can take
270 the form of impressive flood bores (a video can be found here: www.irstea.fr/en/datry).
271 Myriads of terrestrial invertebrates have been reported on these advancing fronts, with
272 densities typically reaching > 5000 individuals/m³ (Corti & Datry 2012; Rosado et al.
273 2014). Many of these organisms have the ability to float or resist submersion for short

274 periods of time and are thus deposited alive further downstream. Such events could be
275 mass dispersal events for terrestrial organisms.

276

277 **5. A dynamic view of metacommunities**

278 The above features (**Table 2**) and resulting effects on environmental filtering and
279 dispersal processes can be used to build a new conceptual model of metacommunity
280 organisation in highly dynamic systems. We propose that (1) metacommunities are very
281 dynamic owing to temporal variations of habitat availability, environmental
282 heterogeneity, and connectivity between patches; (2) these dynamics vary spatially
283 within systems, depending on where environmental variability is the highest, where
284 potential sources of colonists are located, and how much their dispersal is limited by
285 distances between patches; and (3) metacommunities from aquatic and terrestrial phases
286 interact and may have intertwined dynamics.

287

288 **5.1. Metacommunity organisation can be very variable**

289 Alternating wet and dry cycles create contrasting terrestrial and aquatic habitat
290 phases in IRs (**Figure 3a**). As presented above, each of these phases is associated with
291 the predominance of community processes operating locally or regionally. For example,
292 shifts from lotic to lentic conditions are followed by an immediate increase in the
293 importance of environmental filtering processes, including adaptations to lentic
294 conditions, enhanced biotic interactions within contracting pools and very strong
295 predation pressure by terrestrial organisms (**Table 2**). Later, the relative importance of
296 dispersal processes to explain community structure and composition increases with the
297 arrival of large specialist predators such as dragonflies (Odonata), diving beetles
298 (Coleoptera) and some true bugs (Heteroptera), which are generally strong fliers and

299 colonize suitable habitats rapidly (Bogan & Boersma 2012; Bonada et al. 2012).
300 Conversely, upon rewetting, dispersal processes may first predominate to allow
301 colonisation of rewetted habitats by dispersal from refuges (Datry et al. 2014b). Soon
302 after, the communities may be influenced by storage effects from the emergence of
303 desiccation-resistant forms from the rewetted sediments (Stubbington & Datry 2013).
304 After the initial colonisation events, environmental filtering should become gradually
305 more important (**Figure 3b**). High dynamism may also characterize terrestrial
306 communities in IRs, although community-structuring processes involved during each
307 phase shift have been inadequately explored.

308

309 Over time, the structure of metacommunities should show imprints of the high
310 temporal variability in the relative roles of community assembly processes. The most
311 obvious example is probably the respective portions, at a given location, of lotic, lentic,
312 and terrestrial species (**Figure 3c**). Community structure may thus vary sharply during
313 the different phases, with notable dominance by lotic species during flowing phases,
314 dominance by lentic species during non-flowing phases, and dominance by terrestrial
315 species during dry phases (**Figure. 3c**). Although this is speculative, there are some
316 datasets supporting these ideas in IRs (e.g. Bonada et al. 2007; Anna et al. 2008; Corti
317 & Datry 2015). For example, Corti & Datry (2015) described how aquatic and terrestrial
318 successions alternate following hydrological phases in one French IR. Other metrics of
319 community structure, such as taxonomic richness, species abundance, or the proportion
320 of predatory species should also change abruptly and include “a step-change” following
321 phase shift.

322

323 The strong temporal variability in the relative roles of community assembly
324 processes should also generate predictable spatial patterns of metacommunities. During
325 phases dominated by dispersal processes from patches to other patches, communities
326 should be predominantly nested, particularly for weak to moderate dispersers, while
327 species turnover may dominate in phases dominated by environmental filtering
328 operating locally to determine species coexistence (**Figure 3d**). However, these patterns
329 are likely to alternate on short time scales, jeopardizing attempts to infer on
330 communities processes from snap-shot spatial views of metacommunities. Lines of
331 evidence from various systems support these predictions. For example, frequent
332 hurricanes temporarily reduce the degree of nestedness of gastropod assemblages in
333 tropical wet forests (Bloch et al. 2007). Fish in seasonal floodplains show differences in
334 the metacommunity structure between the initial and late phases of the flooding period
335 in response to a shift in the importance of connectivity versus local environmental
336 conditions in structuring local communities (Fernandes et al. 2014). In PRs, different
337 metacommunity models apply along a gradient of disturbance level (Campbell et al.
338 2015). In temporary ponds, shifts from terrestrial to aquatic phases modulate
339 community processes, and the importance of environmental filtering decreases with
340 inundation time (Vanschoenwinkel et al. 2010).

341

342 **5.2. The temporal dynamics of metacommunities in dendritic structures**

343 In dendritic structures experiencing drying events, the spatial scale of
344 metacommunities can be defined through the competitive process and/or the spatial
345 scale of the disturbance, if wet and dry phases occur more rapidly than competitive
346 exclusion (Massol et al. 2011). In this context, the dynamics of metacommunities likely
347 varies spatially within ecosystems with hierarchical structure, notably in case of

348 directionally-biased dispersal (Brown & Swan 2010; Altermatt et al. 2011). For
349 example, in the case of IR networks, the temporal variability of communities may differ
350 according to where drying events (i.e. disturbance) prevail. There are five types of
351 spatial drying configurations in river systems (**Figure 4**), each being potentially
352 associated to a different spatial structuration of the temporal community dynamics.
353 Contrary to the paradigm that headwater communities are being driven purely by
354 environmental filtering and those of downstream, lowland reaches by mass effects due
355 to convergence of all branches and downstream water flow (Brown & Swan 2010),
356 more complex patterns may emerge for each drying configuration. For example, rivers
357 drying completely or partly in their headwaters should have headwater communities
358 driven essentially by dispersal (and perhaps by storage effects), as the source of
359 colonists may be located downstream or in the saturated or dry underlying sediments
360 (see section 3). For the different drying configurations, the degree of connectivity and
361 resulting dispersal rates between patches is probably a key factor, as very high dispersal
362 rates may decouple communities from purely local environmental control (Mouquet &
363 Loreau 2003; Ng et al. 2009; for riverine systems, see also Heino & Peckarsky 2014).
364 At the river network scale, the communities may show contrasting spatial patterns, with
365 community nestedness being more predominant in mid-reach or downstream drying
366 systems (e.g. Datry et al. 2014b), and environmental filtering dominating in headwaters
367 or complete drying systems (e.g. Grant et al. 2007; Clarke et al. 2008; Brown & Swan
368 2010).

369

370 **5.3. Terrestrial and aquatic metacommunities have intertwined dynamics**

371 Biotic interactions between species forming aquatic and terrestrial communities in
372 IRs are localised in space (i.e. drying and rewetting sections) and time (i.e. drying and

373 rewetting phases) (see section 4). These discrete and punctuated interactions may
374 enhance the temporal variability in the structure of both aquatic and terrestrial
375 metacommunities. For example, aquatic species in drying reaches may be heavily
376 preyed upon by terrestrial predators (e.g. beetles, spiders), further reducing the number
377 of species able to survive flow cessation events (McHugh et al. 2014). Conversely, the
378 myriad of terrestrial invertebrates colonising dry riverbeds are entrained by advancing
379 rewetting fronts to downstream river sections, where they may provide subsidies to
380 aquatic food webs and influence aquatic community dynamics (Corti & Datry 2012;
381 Rosado et al. 2014). From a terrestrial perspective, these mass dispersal events could
382 also homogenize terrestrial metacommunities, which could thus be dominated by
383 dispersal processes in IRs rather than by local processes. Yet, aquatic and terrestrial
384 metacommunities show contrasting spatial organisations in IRs, with aquatic
385 metacommunities being longitudinally organised along river networks and terrestrial
386 metacommunities being laterally organised by riparian and upland processes away from
387 river channels (Corti & Datry 2015). Understanding the complex interaction between
388 the respective temporal dynamics and spatial organisation of aquatic and terrestrial
389 metacommunities offers a unique opportunity for integrating aquatic and terrestrial
390 ecology (Datry et al. 2014a).

391

392 **6. A roadmap for future research on metacommunities in intermittent rivers and** 393 **other highly dynamic ecological systems**

394 As a first step, the questions associated with our conceptual models shown above
395 should be tested, particularly because datasets may be now or will be soon available
396 following the growing interest in IRs (e.g. Larned et al. 2010; Datry et al. 2014a, Acuña
397 et al. 2014). Below, we list specific research questions and indicate their relevance to

398 furthering our understanding of metacommunities in IRs. Most of these research
399 questions could be also adapted to other highly dynamic ecological systems.

400

401 *How temporally variable are metacommunities of dynamic systems?*

402 Fine-scale descriptions of the temporal dynamics of metacommunities during
403 typical flowing/non-flowing/dry/flowing phase cycles (**Figure 3a**) are necessary to
404 identify if high dynamism of local communities and metacommunities is a general rule.
405 This would be useful for both aquatic and terrestrial communities and address current
406 limitations in the static view of metacommunities (Erös et al. 2012, 2014). Such
407 temporal descriptions of communities across multiple systems would help to determine
408 if thresholds in phase duration, frequency or timing leading to alternative states of
409 community organisation exist (Bogan & Lytle 2011). Describing and understanding
410 such thresholds is crucial to predict biodiversity change in the context of climate change
411 (Parmesan, 2006; Palmer et al. 2008; Jaeger et al. 2014). Empirical developments
412 should in turn foster conceptual development of metacommunity models suitable for
413 highly dynamic systems.

414

415 *How the dendritic nature of river systems influences the dynamics of metacommunities?*

416 Previous attempts to explore the spatial dynamics of communities in ecosystems,
417 including rivers (e.g., Brown & Swan 2010) have been limited due to the low power of
418 distance matrix-based approaches (Legendre et al. 2005). The development of spatial
419 modelling in the context of constrained ordination (Cottenie 2005; Legendre et al. 2005;
420 Muneeppeerakul et al. 2007; Heino et al. 2015) and in particular those accounting for
421 directionally-biased dispersal (Blanchet et al. 2008) provide better opportunities to
422 understand the spatial organisation of communities in dynamic ecosystems. Yet,

423 constrained ordination and associated spatial models assume a sort of equilibrium in
424 communities, and further developments, such as cost distance-based methods used to
425 model the dispersal of large terrestrial species (Larkin et al. 2004; LaRue & Nieslen
426 2008; Erős & Campbell Grant 2015), represent a more relevant alternative approach.

427

428 *How significant is the storage effect in promoting community dynamics in IRs?*

429 The contribution of the invertebrate seedbank to community recovery has been
430 shown to be significant, although highly variable across IR systems (Stubington &
431 Datry 2013). Yet, its role in driving metacommunity dynamics in IRs is still unknown
432 and notably, whether or not it can compensate for dispersal is an open question (Snyder
433 2006). Manipulative approaches, such as common-garden experiments manipulating the
434 invertebrate seedbank and/or the dispersal of organisms in water and the air could be
435 helpful to address this question. Alternatively, cross-system comparisons of
436 metacommunity dynamics in systems with contrasting drying patterns (i.e. with and
437 without perennial refuges) may help disentangle the respective role of storage effects
438 and dispersal in promoting community dynamics. From a modelling perspective,
439 mechanistic approaches, for example using multi-occupancy models (e.g. Lamy et al.
440 2013) applied to more than one species, would allow accounting for storage effects in
441 estimating colonization and persistence rates, thus assessing the extent of cryptic
442 dormant stages in metacommunities.

443

444 *Do interactions between aquatic and terrestrial metacommunities alter their respective*
445 *dynamics?*

446 Understanding how biotic interactions between aquatic and terrestrial
447 metacommunities influence their respective temporal dynamics requires further

448 quantification in IRs. Yet, studies describing synoptically aquatic and terrestrial
449 metacommunities in IRs or other aquatic-terrestrial systems (e.g. wetlands, tidal
450 marshes) remain scarce (but see Corti & Datry 2015). Considering simultaneously
451 aquatic and terrestrial metacommunities in IRs would provide a way forward to account
452 for the linkages between these two components, as well as to bring terrestrial and
453 aquatic ecologists together in metacommunity research.

454

455 *Is there mass dispersal of terrestrial organisms during rewetting events?*

456 While recent studies have reported how dramatic rewetting events in IRs can be and
457 how much large quantities of terrestrial organisms are being entrained downstream of a
458 river network, it is still unknown how strong roles these events play in terrestrial or
459 aquatic community dynamics (Corti & Datry 2012; Rosado et al. 2014). Typically, such
460 events could produce a mass effect for downstream riparian communities, thereby
461 obscuring local environmental filtering processes. To address this question, field
462 experiments could monitor the fate of the entrained organisms on downstream
463 communities using stable isotopes or molecular approaches. Also, the amount and type
464 of terrestrial inputs to these rewetting fronts could be manipulated to examine their
465 effects on aquatic and terrestrial communities.

466

467 *How can the metacommunity perspective help managers to conserve the biodiversity of*
468 *IRs?*

469 The metacommunity perspective is a mechanistic route to relating biodiversity
470 patterns to landscape features and exploring biodiversity conservation plans (Economo
471 2011). Identifying the relative roles of environmental filtering and dispersal processes in
472 metacommunities can help managers to better conserve these dynamic ecological

473 systems. If local communities are mainly governed by environmental filtering
474 processes, management initiatives to maintain local habitats should be prioritized. If
475 dispersal processes are predominant instead, management strategies should maintain
476 landscape connectivity and natural disturbance regimes to promote “source” patches of
477 biodiversity (Bengtsson 2010). In the specific case of IRs, this would require preserving
478 the natural mosaic of shifting habitat types (Datry et al. 2015) and the local conditions
479 of perennial headwater patches, which are considered as important sources of
480 biodiversity (Cañedo-Argüelles et al. 2015). Additionally, the metacommunity
481 framework applied to IRs will help to pinpoint keystone species or habitat-patches in
482 the mosaic, on which managers should focus conservation and restoration efforts
483 (Mouquet et al. 2013). Empirical studies analysing the metacommunity processes in IRs
484 are thus urged to implement conservation actions specifically tailored to these dynamic
485 systems.

486

487 **6. Conclusions**

488 While community ecology has progressed rapidly in recent years owing to the
489 recognition that local communities are not spatially closed but form metacommunities
490 (Logue et al. 2011; Winegardner et al. 2012), empirical studies have often reported
491 ambiguous findings about the relative importance of underlying processes (Cottenie
492 2005; Beisner et al. 2006; Logue et al. 2011). This could be partly due to the common
493 consideration in metacommunity research that both biological communities and their
494 habitats as relatively stable, whereas many ecological systems are actually highly
495 dynamic in nature. Because IRs are common ecological systems across the globe and
496 are dynamic shifting mosaics of lotic, lentic, and terrestrial habitats, they are ideal
497 arenas for addressing the spatio-temporal variability of metacommunities in highly

498 dynamic settings. As developed here for IRs, the respective importance of
499 environmental filtering and dispersal processes may shift abruptly or gradually over
500 time in highly dynamic ecological systems, leading local community structure to vary
501 constantly. This temporal variability is certainly enhanced by discrete and punctuated
502 biotic interactions between aquatic and terrestrial communities in IRs and other coupled
503 aquatic-terrestrial systems. In the case of dendritic systems with directionally-biased
504 dispersal (Fagan 2002; Grant et al. 2007; Altermatt et al. 2011), the location and spatial
505 extent of disturbances, such as drying, may interact with the temporal variations in
506 community assembly processes to produce complex spatio-temporal variability in local
507 community structure. We contend that addressing these questions in IRs may 1)
508 substantially advance the metacommunity theory (Erös et al. 2012, 2014); 2) offer a
509 unique opportunity for bridging the gap between aquatic and terrestrial community
510 ecology (Datry et al. 2014a; Soininen et al. 2015); and 3) improve the management of
511 ecological systems (Heino 2013; Tonkin et al. 2014). In the context of increasing
512 extreme climatic events and ecosystem disturbances (Easterling et al. 2000; Parmesan
513 2006), understanding how metacommunities are organised in highly dynamic systems is
514 becoming a key research topic. Also, translating such research into efficient
515 management guidelines is urgently needed.

516

517 **Acknowledgments**

518 Earlier versions of this manuscript benefited from comments by Mike T. Bogan,
519 Tibor Erös, Bernard Hugueny, Adriano S. Melo, the Oikos Subject Editor François
520 Massol and three anonymous reviewers. TD and NB were supported by the IRBAS
521 (Intermittent River Biodiversity Analysis and Synthesis) working group supported by
522 CESAB (Center for Synthesis and Analysis of Biodiversity), which is funded jointly by

523 FRB (French Foundation for Research & Biodiversity) and ONEMA (French National
524 Agency for Water & Aquatic Environments). JH was supported by a grant from the
525 Academy of Finland for the project entitled “Scaling biodiversity in tropical and boreal
526 streams: Implications for diversity mapping and environmental assessment”.

527

528 **Reference list**

529 Acuña, V., et al. 2014. Why should we care about temporary waterways? - *Science* 343:
530 1080-1082.

531 Altermatt, F. 2013. Diversity in riverine metacommunities: a network perspective. -
532 *Aquatic Ecology* 47: 365-377.

533 Altermatt, F., et al. 2011. Interactive effects of disturbance and dispersal directionality
534 on species richness and composition in metacommunities. -*Ecology* 92(4): 859-
535 870.

536 Anna, A., et al. 2009. Do intermittent and ephemeral Mediterranean rivers belong to the
537 same river type? -*Aquatic Ecology* 43(2): 465-476.

538 Azeria, E. T., and Kolasa, J. 2008. Nestedness, niche metrics and temporal dynamics of
539 a metacommunity in a dynamic natural model system. *Oikos* 117(7): 1006-1019.

540 Baselga, A. 2010. Partitioning the turnover and nestedness components of beta
541 diversity. -*Global Ecology and Biogeography* 19(1): 134-143.

542 Beisner, B. E., et al. 2006. The role of dispersal in structuring lake communities from
543 bacteria to fish. -*Ecology* 87: 2895-2991.

544 Bengtsson, J. 2010. Applied (meta)community ecology: diversity and ecosystem
545 services at the intersection of local and regional processes. In *Community*
546 *Ecology: Processes, Models, and Applications* (eds H. A. Verhoef and P. L.
547 Morin), pp. 115-130. Oxford University Press, New York Oxford.

548 Blanchet, F. G., et al. 2008. Modelling directional spatial processes in ecological data. -
549 Ecological Modelling 215(4): 325-336.

550 Bloch, C. P., et al. 2007. Effects of large-scale disturbance on metacommunity structure
551 of terrestrial gastropods: temporal trends in nestedness. -Oikos 116(3): 395-406.

552 Bogan, M. T. and Lytle, D. A. 2011. Severe drought drives novel community
553 trajectories in desert stream pools. -Freshwater Biology 56(10): 2070-2081.

554 Bogan, M. T. and Boersma, K. S. 2012. Aerial dispersal of aquatic invertebrates along
555 and away from arid-land streams. -Freshwater Science 31: 1131-1144.

556 Bonada, N., et al. 2007. Taxonomic and biological trait differences of stream
557 macroinvertebrate communities between mediterranean and temperate regions:
558 implications for future climatic scenarios. -Global Change Biology 13: 1658-
559 1671.

560 Bonada, N., et al. 2012. Spatial autocorrelation patterns of stream invertebrates:
561 exogenous and endogenous factors. -Journal of Biogeography 39: 56-68.

562 Boulton, A. J. 2003. Parallels and contrasts in the effects of drought on stream
563 macroinvertebrate assemblages. -Freshwater Biology 48(7): 1173-1185.

564 Brown, B. L. and Swan, C. M. 2010. Dendritic network structure constrains
565 metacommunity properties in riverine ecosystems. -Journal of Animal Ecology
566 79: 571–580.

567 Cañedo-Arguelles, M., et al. 2015. Dispersal strength determines meta-community
568 structure in a dendritic riverine network. -Journal of Biogeography
569 doi: 10.1111/jbi.12457.

570 Campbell, R. E., et al. 2015. Flow-related disturbance creates a gradient of
571 metacommunity types within stream networks. Landscape Ecology 30(4): 667-
572 680.

573 Carrara, F., et al. 2012. Dendritic connectivity controls biodiversity patterns in
574 experimental metacommunities. -Proceedings of the National Academy of
575 Sciences 109: 5761-5766.

576 Clarke, A., et al. 2008. Macroinvertebrate diversity in headwater streams: a review. -
577 Freshwater biology 53(9): 1707-1721.

578 Corti, R. and Datry, T. 2012. Invertebrates and sestonic matter in an advancing wetted
579 front travelling down a dry river bed (Albarine, France). -Freshwater Science
580 31(4): 1187-1201.

581 Corti, R. and Datry, T. 2015. Ecological ambivalence: aquatic and terrestrial
582 invertebrate communities in an intermittent river. -Freshwater Biology (In
583 press).

584 Cottenie, K. 2005. Integrating environmental and spatial processes in ecological
585 community dynamics. -Ecology letters 8(11): 1175-1182.

586 Datry, T., et al. 2014a. Intermittent rivers: a challenge for freshwater ecology. -
587 BioScience bit027.

588 Datry, T., et al. 2014b. Broad-scale patterns of invertebrate richness and community
589 composition in temporary rivers: effects of flow intermittence. -Ecography 37:
590 94-104.

591 Datry T., et al. 2015. A landscape, mosaic perspective of intermittent rivers: ecological
592 implications. -Freshwater Biology (in press).

593 Drake, J. A. 1990. The mechanics of community assembly and succession. -Journal of
594 Theoretical Biology 147(2): 213-233.

595 Dudgeon, D., et al. 2006. Freshwater biodiversity: importance, threats, status and
596 conservation challenges. -Biological Reviews 81(2): 163-182.

597 Easterling, D. R., et al. 2000. Climate extremes: observations, modeling, and impacts. -
598 Science 289: 2068-2074.

599 Economo, E. P. 2011. Biodiversity conservation in metacommunity networks: linking
600 pattern and persistence. -American Naturalist 177: 167-180.

601 Erős, T., et al. 2012. Temporal variability in the spatial and environmental determinants
602 of functional metacommunity organisation—stream fish in a human-modified
603 landscape. -Freshwater Biology 57: 1914-1928.

604 Erős, T., et al. 2014. Quantifying temporal variability in the metacommunity structure
605 of stream fishes: the influence of non-native species and environmental drivers. -
606 Hydrobiologia 722(1): 31-43.

607 Erős, T., and Campbell Grant, E.H., 2015. Unifying research on the fragmentation of
608 terrestrial and aquatic habitats: patches, connectivity and the matrix in
609 riverscapes. -Freshwat Biol, in press.

610 Fagan, W. F. 2002. Connectivity, fragmentation, and extinction risk in dendritic
611 metapopulations. -Ecology 83(12): 3243-3249.

612 Fernandes, I. M., et al., 2014. Spatiotemporal dynamics in a seasonal metacommunity
613 structure is predictable: the case of floodplain-fish communities. -Ecography
614 37(5): 464-475.

615 Fukui, D. A. I., et al. 2006. Effect of emergent aquatic insects on bat foraging in a
616 riparian forest. -Journal of Animal Ecology 75(6): 1252-1258.

617 Gasith, A. and Resh, V. H. 1999. Streams in Mediterranean climate regions: abiotic
618 influences and biotic responses to predictable seasonal events. -Annual Review
619 of Ecology and Systematics 51-81.

620 Gilbert, B., Bennett, J. R. 2010. Partitioning variation in ecological communities: do the
621 numbers add up? -Journal of Applied Ecology 47(5): 1071-1082.

622 Grant, E. H. C., et al. 2007. Living in the branches: population dynamics and ecological
623 processes in dendritic networks. -Ecology Letters 10: 165-175.

624 Heino, J. 2013. The importance of metacommunity ecology for environmental
625 assessment research in the freshwater realm. -Biological Reviews 88: 166-178.

626 Heino, J. and Peckarsky, B. L. 2014. Integrating behavioral, population and large-scale
627 approaches for understanding stream insect communities. -Current Opinion in
628 Insect Science 2: 7-13.

629 Heino, J., et al. 2015. Metacommunity organisation, spatial extent and dispersal in
630 aquatic systems: patterns, processes and prospects. -Freshwater Biology 60: 845-
631 869.

632 Holyoak, M., et al. 2005. Metacommunities: Spatial dynamics and ecological
633 communities. University of Chicago Press, Chicago.

634 Hubbell, S. J. 2001. The unified neutral theory of biodiversity and biogeography.
635 Princeton University Press, Princeton.

636 Jacobson, B., and Peres-Neto, P. R. 2010. Quantifying and disentangling dispersal in
637 metacommunities: how close have we come? How far is there to go? -Landscape
638 Ecology 25: 495-507.

639 Jaeger, K. L., et al. 2014. Climate change poised to threaten hydrologic connectivity and
640 endemic fishes in dryland streams. -Proceedings of the National Academy of
641 Sciences: 10.1073/pnas.1320890111.

642 Kolasa, J., Romanuk, T. N. 2005. Assembly of unequals in the unequal world of a rock
643 pool metacommunity. *Metacommunities: spatial dynamics and ecological*
644 *communities. University of Chicago Press, Chicago, Illinois, USA, 212-232.*

645 Kirwan, M. L. and Murray, A. B. 2007. A coupled geomorphic and ecological model of
646 tidal marsh evolution. *Proceedings of the National Academy of Sciences* 104:
647 6118-6122.

648 Lamy, T., et al. 2013. Metapopulation Dynamics of Species with Cryptic Life Stages. -
649 *The American Naturalist* 181: 479-491.

650 Larned, S. T., et al. 2010. Emerging concepts in temporary river ecology. -*Freshwater*
651 *Biology* 55: 717-738.

652 Larkin, J.L., et al. 2004. Landscape linkages and conservation planning for the black
653 bear in west-central Florida. -*Animal Conservation* 7: 23-34.

654 LaRue, M.A. and Nielsen, C.K., 2008. Modelling potential dispersal corridors for
655 cougars in midwestern North America using least-cost path methods. -
656 *Ecological Modelling* 212: 371–381.

657 Legendre, P. et al. 2005. Analyzing beta diversity: partitioning the spatial variation of
658 community composition data. -*Ecological Monographs*, 75(4): 435-450.

659 Leibold, M. A., et al. 2004. The metacommunity concept: a framework for multi-scale
660 community ecology. -*Ecology Letters* 7: 601-613.

661 Logue, J. B., et al. 2011. Empirical approaches to metacommunities: a review and
662 comparison with theory. -*Trends in Ecology and Evolution* 26(9): 482-491.

663 MacArthur, R. H. and Wilson, E. O. 1967. *The Theory of Island Biogeography*.
664 Princeton, Princeton University Press.

665 Massol, F., et al. 2011. Linking ecosystem and community dynamics through spatial
666 ecology. -*Ecology Letters* 14: 313-323.

667 McHugh, P. A., et al. 2014. Habitat size influences food web structure in drying
668 streams. -*Ecography*, doi: 10.1111/ecog.01193.

- 669 Meynard, C. N., 2013. Disentangling the drivers of metacommunity structure across
670 spatial scales. -*Journal of Biogeography* 40(8): 1560-1571.
- 671 Moritz, C., et al. 2013. Disentangling the role of connectivity, environmental filtering,
672 and spatial structure on metacommunity dynamics. -*Oikos* 122: 1401-1410.
- 673 Mouquet, N. and Loreau, M. 2003. Community patterns in source-sink
674 metacommunities. -*American Naturalist* 162: 544–557.
- 675 Mouquet, N. et al. 2013. Extending the concept of keystone species to communities and
676 ecosystems. -*Ecology Letters* 16: 1-8.
- 677 Munepeeraikul, R., S., et al. 2007. On biodiversity in river networks: A trade-off
678 metapopulation model and comparative analysis. -*Water Resources Research* 43,
679 W07426.
- 680 Ng, I.S.Y., et al. 2009. Hierarchical zooplankton metacommunities: distinguishing
681 between high and limiting dispersal mechanisms. -*Hydrobiologia* 619: 133–143.
- 682 Palmer, M. A., et al. 2008. Climate change and the world's river basins: anticipating
683 management options. -*Frontiers in Ecology and the Environment* 6(2): 81-89.
- 684 Parmesan, C. 2006. Ecological and evolutionary responses to recent climate change. -
685 *Annual Review of Ecology, Evolution, and Systematics* 37: 637-669.
- 686 Presley, S. J., et al. 2010. A comprehensive framework for the evaluation of
687 metacommunity structure. -*Oikos* 119(6): 908-917.
- 688 Raymond, P. A., et al. 2013. Global carbon dioxide emissions from inland waters. -
689 *Nature* 503(7476): 355-359.
- 690 Ricklefs, R. E. 2008. Disintegration of the ecological community. -*American Naturalist*
691 172(6): 741-750.
- 692 Rosado, J., et al. 2014. Mass dispersal of terrestrial organisms during first flush in a
693 temporary stream. -*River Research and Applications*: doi: 10.1002/rra.2791.

694 Sheldon, F., et al. 2010. Ecological roles and threats to aquatic refugia in arid
695 landscapes: dryland river waterholes. -*Marine and Freshwater Research* 61: 885-
696 895.

697 Smol, J. P., and Douglas, M. S. 2007. Crossing the final ecological threshold in high
698 Arctic ponds. -*Proceedings of the national Academy of Sciences* 104(30):
699 12395-12397.

700 Snyder, R. E. 2006. Multiple risk reduction mechanisms: can dormancy substitute for
701 dispersal? -*Ecology Letters* 9(10): 1106-1114.

702 Soininen, J., et al. 2015. Towards more integrated ecosystem research in aquatic and
703 terrestrial environments. -*BioScience* 65: 174-182.

704 Stanley, E. H., et al. 1997. Ecosystem expansion and contraction in streams. -
705 *BioScience*: 427-435.

706 Steward, A. L., et al. 2012. When the river runs dry: human and ecological values of dry
707 riverbeds. -*Frontiers in Ecology and the Environment* 10(4): 202-209.

708 Stubbington, R. and Datry, T. 2013. The macroinvertebrate seedbank promotes
709 community persistence in temporary rivers across climate zones. -*Freshwater*
710 *Biology* 58(6): 1202-1220.

711 Tonkin, J. D., et al. 2014. Dispersal distance and the pool of taxa, but not barriers,
712 determine the colonisation of restored river reaches by benthic invertebrates. -
713 *Freshwater Biology* 59: 1843–1855.

714 Tuomisto, H., Ruokolainen, K. 2006. Analyzing or explaining beta diversity?
715 Understanding the targets of different methods of analysis. -*Ecology* 87(11):
716 2697-2708.

717 Vanschoenwinkel, B., et al. 2007. The role of metacommunity processes in shaping
718 invertebrate rock pool communities along a dispersal gradient. - *Oikos* 116(8):
719 1255-1266.

720 Vanschoenwinkel, B., et al. 2010. Species sorting in space and time-the impact of
721 disturbance regime on community assembly in a temporary pool
722 metacommunity. -*Journal of the North American Benthological Society* 29:
723 1267-1278.

724 Vellend, M. 2010. Conceptual synthesis in community ecology. -*Quarterly Review of*
725 *Biology* 85: 183-206.

726 Warner, R. R., Chesson, P. L. 1985. Coexistence mediated by recruitment fluctuations:
727 a field guide to the storage effect. -*American Naturalist*: 769-787.

728 Wiggins, G. B., et al. 1980. Evolutionary and ecological strategies of animals in annual
729 temporary pools. -*Archiv fur Hydrobiologie Supplement* 58: 97-206.

730 Winegardner, A. K., et al. 2012. The terminology of metacommunity ecology. -*Trends*
731 *in Ecology and Evolution* 27: 253–254.

732

733

734

735

736

737

738

739

740

741

742 **Table 1. Definitions of terms used throughout the manuscript.**

Term	Definition
Community (or local community)	The individuals of all species that potentially interact within a single patch or local area of habitat (Leibold et al. 2004).
Dispersal	A regional process influencing metacommunity dynamics involving the movement of individuals from one site (i.e., emigration) to another (i.e., immigration) within a region (see also Leibold et al. 2004).
Storage effect	A local process involving the recruitment of many individuals from a single generation, allowing species coexistence in systems prone to disturbance (see also Warner & Chesson 1985; Snyder 2006).
Flow intermittence	Proportion of a given period, generally a year, during which a site in a river network is either under lentic or terrestrial phases (Datry et al. 2014).
Local community structure	A general term used to describe community characteristics resulting from the processes involved in its formation (e.g. species richness, dominance, etc.).
Metacommunity	A set of local communities that are linked by dispersal of multiple potentially interacting species (Leibold et al. 2004).
Metacommunity dynamics	The dynamics that arise within metacommunities; these consist of spatial dynamics, temporal dynamics, and community dynamics (multispecies interactions or the emergent properties arising from them within communities), and the interaction of these three dynamics (see also Leibold et al. 2004).
Metacommunity organisation	A term that refers to the processes that explain metacommunities, i.e., environmental filtering or dispersal (Heino et al. 2015).
Environmental filtering	A local, niche-based process influencing metacommunity dynamics and encompassing (i) the effects of local abiotic factors on species survival and (ii) local species interactions (see also species sorting in Leibold et al. 2004).
Nestedness	The species of communities with smaller numbers of species are subsets of species-richer communities, reflecting a non-random process of species loss or gain as a consequence of any factor that promotes the orderly disaggregation (or aggregation) of community (Baselga 2010).
Turnover	The replacement of some species by others between communities is a consequence of environmental filtering or spatial and historical constraints. Different from nestedness, the gain of one species is necessarily accompanied by a loss of a second species (Baselga 2010).

743

744

745

746

747

