



**HAL**  
open science

## Numerical modelling contribution to sedimentary sedynamisation projects in a lowland gravel-bed river: the Ain River, France

G. Naudet, B. Camenen, Jérôme Le Coz, André Paquier, H. Piegay

### ► To cite this version:

G. Naudet, B. Camenen, Jérôme Le Coz, André Paquier, H. Piegay. Numerical modelling contribution to sedimentary sedynamisation projects in a lowland gravel-bed river: the Ain River, France. *I.S. Rivers*, Jun 2015, Lyon, France. 2015. <hal-02602356>

**HAL Id: hal-02602356**

**<https://hal.inrae.fr/hal-02602356v1>**

Submitted on 16 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

# Numerical modelling contribution to sedimentary redynamisation projects

## in a lowland gravel-bed river: the Ain River, France

Grégoire Naudet, Benoît Camenen, Jérôme Le Coz, André Paquier, Hervé Piégay

ZABR



### Introduction


Ain river characteristics :  
 mean annual discharge : 120 m<sup>3</sup>/s  
 Q2 : 860 m<sup>3</sup>/s    Q10 : 1400 m<sup>3</sup>/s

**Sedimentary deficit ~ 15,000 m<sup>3</sup>/year**

Excavation and reconnection of old channels  
 Injection of the mined sediments

Objectives :  
**Estimate the efficiency of these redynamisation scenarii as well as their associated risks thanks to numerical modelling.**


Numerical software developed by Irstea:  
 - 1D hydraulic: MAGE  
 - 1D hydraulic and sedimentary: RubarBE  
 - 2D hydraulic and sedimentary: Rubar20TS




Characterise the reference status

Determine the connection level of the channel



Determine the new hydraulic conditions



Estimate the critical discharge for incipient motion and describe the erosion processes




Critical discharge for incipient motion :  
 - sensitive to the grain size ( $d_{50}$ )  
 - quite frequent (350 to 525 m<sup>3</sup>/s)




Determine the evolution of the river morphology


The grid that represents the injection area does not seem fine enough to describe correctly the gravel bar evolution



Evaluate the transport distances and solid discharge



Difficulty in discriminating the sediments of the bar from the others  
 Modelled transport distances (~ 4 to 5 km for 1 year) > Observed travel distances of PIT tags (800m) (Arnaud, 2014)



### References

Arnaud, F., Piégay H., Michel K., Bultingaire L., Collery P., Drissi N., Tissot C. (2014). Caractérisation de la dynamique sédimentaire et des habitats aquatiques d'une rivière régulée: l'exemple de l'Ain dans sa haute et basse vallée, IS Rivers 2015.

Béraud, C. (2012). Modélisation numérique des impacts de recharges sédimentaires en rivière aménagée. Cas du Vieux-Rhin entre Kembs et Breisach. Thèse de doctorat. Université Claude Bernard – Lyon 1

Rollet A.J. (2007). Étude et gestion de la dynamique sédimentaire d'un tronçon fluvial à l'aval d'un barrage : le cas de la Basse Vallée de l'Ain. Thèse de doctorat Université Jean Moulin Lyon 3

### Perspectives

- Refine the grid in order to improve the representation of the sedimentary processes
- Analyse the evolutions of the river morphology and evaluate the influence of the injections on the grain size distribution
- Apply these results in a biological model (to evaluate the evolutions of the habitat quality for fish using water depth, current speed, and substratum composition)