



**HAL**  
open science

## Decoding Metal Associations in an Arid Urban Environment with Active and Legacy Mining: the Case of Copiapó, Chile

M. Moya, P. Pasten, Marina Coquery, C.A. Bonilla, A. Vega, A. Carkovic, M. Calcagni

► **To cite this version:**

M. Moya, P. Pasten, Marina Coquery, C.A. Bonilla, A. Vega, et al.. Decoding Metal Associations in an Arid Urban Environment with Active and Legacy Mining: the Case of Copiapó, Chile. American Geophysical Union, Dec 2015, San Francisco, United States. pp.1, 2015. hal-02604503

**HAL Id: hal-02604503**

**<https://hal.inrae.fr/hal-02604503>**

Submitted on 16 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Decoding Metal Associations in an Arid Urban Environment with Active and Legacy Mining: the Case of Copiapó, Chile

Pablo Moya<sup>1,2</sup>, Pablo Pasten<sup>1,2</sup>, Marina Coquery<sup>3</sup>, Carlos A Bonilla<sup>1,2</sup>, Alejandra Vega<sup>1,2</sup>, Athena Carkovic<sup>1,2</sup> and Magdalena Calcagni<sup>1,2</sup>

Abstract number: GC51G-1177

## Introduction:

Copiapó city in the arid Atacama desert has **more than 30 abandoned mine tailings** (Fig. 1) and one **active copper smelter** (Fig. 2).

**Soils and street dust** could be enriched in **heavy metals** due to different dispersion mechanisms, such as **wind, rain and landslides**, leaving the population exposed to high **health risk**.

In March of this year **many landslides happened** in Copiapó city affecting tailings and cover with muds part of the city. It is vital to learn **what sources are affecting the city, which ones are the most dangerous and what is their range of dispersion**.

Study Site: Copiapó, Atacama Region, Chile

Population: 166.751 inhabitants

Area: 47,77 km<sup>2</sup>

Rainfall: 28 mm annual mean

Mean temperature: 15,2 °C annual mean

Main economic activity: Mining and agriculture

Number of Tailings in the region: 164




Fig. 1: Location of some of the mine tailings in Copiapó

## Methodology:

- **71 street dust, 42 soils, 41 mud (transect method after landslides), 83 tailings** in two samples campaigns.
- Samples were dried (40 °C) and sieved (< 2 mm). Elemental composition was determined by **pXRF**. Measures was validated with ICP-MS comparison (replicated and reference material).
- Values of **As, Co, Cu, Fe, Mn, Pb** and **Zn** were considered valid for the pXRF method.
- **Positive Matrix Factorization (PMF)** is a model that uses the concentration and uncertainty of each sample to determine the **source's fingerprint** and the **apportionment to these samples** [1,2]: Was used with the **pXRF measurements of the street dust** samples.
- The **source contribution** was represented spatially and graphically using the **Kriging** method.




Fig. 2: Paipote Copper Smelter nearby population

## Results:

PMF model identified four main factors or sources (Fig. 3) related to:

- **Industries emission:** 87% of Cu variation (copper smelter and industries).
- **Traffic emission:** 85% of Pb variation.
- **Crustal:** 55% of Co variation and 45% of Fe variation, but Co concentration were very low so its considered not enriched by anthropogenic activities.
- **Tailings:** 67% of Mn, 59% of As and 52% of Zn variations.




Fig. 3: Source's fingerprint related to street dust pXRF measurements

Two zones had been **contributed by the industrial factor** (Fig. 4), one is **nearly the copper smelter** and the second one is the **industrial zone of Copiapó** in the southwest of the city.




Fig. 4: Spatial distribution of contributions from industrial factor.

## Landslides (March, 2015):

Heavy rains were the cause of many landslides in Atacama region. In the city the landslide went through the riverbed of Copiapó river and through the Paipote ravine, covering **40% of the urban area** with muds (Fig. 5).

The main concern was that some **tailings could be dragged by the landslide** and could be deposited in the streets of Copiapó, exposing the population to a high health risk.



Fig. 5: Copiapó's street after the landslides

- Some muds samples were above the Brazilian and Canada intervention limit for Cu, Zn, Hg, As and Pb in soils. Muds were taken off as soon as possible.
- **Two tailings were affected by the landslide.**
- Fig. 6 shows that **Cu and Zn** concentration rises while the muds go trough the city and the river bed, suggesting a process of **enrichment**. Also the **fine fraction** shows a tendency to decrease.




Fig. 6: As, Cu, Pb and Zn concentration obtained by pXRF on muds of the Copiapó river axis

## Conclusions:

- Two contaminating sources were clearly identified, **traffic emissions and the industrial emissions**.
- The muds that pass trough Copiapó City and Copiapó riverbed **were enriched by Zn and Cu**.
- The dragged tailings material was diluted **without changing the concentration of elements on muds**.
- The **pXRF technique has limitations** on the quantity of elements correctly measured and in limits of quantification.

## Further work and perspectives:

- **Other techniques** such as X-ray diffraction, Cluster and Principal Component Analysis, ratios of elements and increase the elements measured could **reinforce the identification of sources**.
- Analyze the effect of the **change on percentage of finest fraction in the element concentrations** from the landslide muds.
- We hope that this work can help **planners and regulators** to make better decisions considering the zones that are affected by anthropogenic contamination.

## References:

- [1] Paatero, P. and Tapper, U. (1994) Positive matrix factorization: a non-negative factor model with optimal utilization of error estimates of data values. *Environmetrics*, 5, pp. 111–126.
- [2] Norris et al., 2014G. Norris, R. Duvall, S. Brown, S. BaiEPA Positive Matrix Factorization (PMF) 5.0 Fundamentals and User Guide.

1. Pontificia Universidad Católica de Chile, Av. Vicuña Mackenna 4860, Macul, Santiago, Chile.

2. Centro de Desarrollo Urbano Sustentable CEDEUS, El Comendador 1916, Providencia, Santiago, Chile

3. Irstea, U.R. MALY, 5 rue de la Doua, CS70077, 69626 Villeurbanne Cedex, France.

