



HAL
open science

Natural attenuation of metals from acid drainage in surface waters: effects of organic matter in the association of arsenic to hydrous Al and Fe oxyhydroxides and their particle size distribution

G.J. Arce, P. Pasten, Marina Coquery, M. I. Abarca, M. Montecinos

► To cite this version:

G.J. Arce, P. Pasten, Marina Coquery, M. I. Abarca, M. Montecinos. Natural attenuation of metals from acid drainage in surface waters: effects of organic matter in the association of arsenic to hydrous Al and Fe oxyhydroxides and their particle size distribution. American Geophysical Union (AGU), Dec 2015, San Francisco, United States. pp.1, 2015. hal-02604504

HAL Id: hal-02604504

<https://hal.inrae.fr/hal-02604504>

Submitted on 16 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

 **AGU FALL MEETING**

San Francisco | 14 – 18 December 2015

H13L-1758**Natural Attenuation of Metals from Acid Drainage in Surface Waters: Effects of Organic Matter in the Association of Arsenic to Hydrous Al and Fe Oxyhydroxides and Their Particle Size Distribution**

Monday, 14 December 2015

Poster Hall (Moscone South)

Guillermo José Arce¹, **Pablo Pasten**^{1,2}, **Marina Coquery**³, **María Ignacia Abarca**¹ and **Mauricio Montecinos**¹, (1)Pontificia Universidad Católica de Chile, Santiago, Chile, (2)Centro de Desarrollo Urbano Sustentable, Santiago, Chile, (3)IRSTEA, UR MALY (Freshwater Systems Ecology and Pollution), F-69626 Villeurbanne Cedex, France

Abstract:

The presence of toxic metals in watersheds affected by acid drainage (AD) imposes a challenge for sustainable supply of water for cities, agriculture and industry. The formation and settling of metal-rich HFO (hydrous ferric oxides) and HAO (hydrous aluminum oxides) is a relevant mechanism for the attenuation of dissolved metals from AD, particularly for arsenic. Organic matter is known to alter the chemical speciation and key physical properties like particle size distribution (PSD) and settling velocity of HFO and HAO particle suspensions. However, available experimental studies focus either on chemical or physical properties.

We used a suite of analytical techniques to probe the effects of organic matter on particle suspensions formed in natural waters and in laboratory model systems. Dissolved organic matter was added at different concentrations (0, 0.1, 0.3, 0.6, 1 and 1.5 mg C/L) to synthetic AD before neutralization with alkaline solutions. PSD and average particle size were measured with laser scattering transmissometry (LISST), while organic matter was characterized by total organic carbon (TOC) and UV-VIS spectrometry.

Larger concentrations of organic matter lead to the formation of particle suspensions with larger particle volume and size. When 1.5 mg C/L were added, the total particle volume concentration increased from 0.67 to 29.74 µL/L, while the mean particle size increased from 102 to 476 µm. These results suggest that organic matter influences the fate of metals from AD. Undergoing measurements include total and dissolved metal analyses with total reflection X-ray fluorescence (TXRF) and ICP-OES to confirm increased removal of dissolved arsenic.

The results from this research are necessary to understand the processes governing natural attenuation of metal contamination in fluvial systems affected by AD and to serve as the basis for enhanced natural attenuation schemes.

The authors acknowledge funding from Fondecyt 1130936.

[<< Previous Abstract](#) | [Next Abstract >>](#)

SEARCH**BROWSE PROGRAM****BROWSE BY PERSON**

