



HAL
open science

Geochemical and hydrologic controls of copper-rich surface waters in the Yerba Loca-Mapocho system

M. Montecinos, P. Pasten, Marina Coquery, E.G. Pizarro, M. I. Abarca, G. J. Arce

► **To cite this version:**

M. Montecinos, P. Pasten, Marina Coquery, E.G. Pizarro, M. I. Abarca, et al.. Geochemical and hydrologic controls of copper-rich surface waters in the Yerba Loca-Mapocho system. American Geophysical Union (AGU), Dec 2015, San Francisco, United States. pp.1, 2015. hal-02604505

HAL Id: hal-02604505

<https://hal.inrae.fr/hal-02604505>

Submitted on 16 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

H13L-1766

Geochemical and Hydrologic Controls of Copper-Rich Surface Waters in the Yerba Loca-Mapocho System

Monday, 14 December 2015

Poster Hall (Moscone South)

Mauricio Montecinos¹, Pablo Pasten², Marina Coquery³, Gonzalo E Pizarro¹, María Ignacia Abarca⁴ and Guillermo José Arce⁴, (1)Pontificia Universidad Católica de Chile, Santiago, Chile, (2)Centro de Desarrollo Urbano Sustentable, Santiago, Chile, (3)IRSTEA, UR MALY (Freshwater Systems Ecology and Pollution), F-69626 Villeurbanne Cedex, France, (4)Pontifical Catholic University of Chile, Santiago, Chile

Abstract:

Andean watersheds in Northern and Central Chile are naturally enriched with metals, many of them associated to sulfide mineralizations related to copper mining districts. The natural and anthropogenic influx of toxic metals into drinking water sources pose a sustainability challenge for cities that need to provide safe water with the smallest footprint. This work presents our study of the transformations of copper in the Yerba Loca-Mapocho system. Our sampling campaign started from the headwaters at La Paloma Glacier and continues to the inlet of the San Enrique drinking water treatment plant, a system feeding municipalities in the Eastern area of Santiago, Chile. Depending on the season, total copper concentrations go as high as 22 mg/L for the upper sections, which become diluted to <5 mg/L downstream. pH ranged from 3 to 5.6 while suspended solids ranged from <10 to 100 mg/L. We used Geochemist Workbench to assess copper speciation and to evaluate the thermodynamic controls for the formation and dissolution of solid phases. A sediment trap was used to concentrate suspended particulate matter, which was analyzed with ICP-MS, TXRF (total reflection X ray fluorescence) and XRD (X-ray diffraction). Major elements detected in the precipitates were Al (200 g/kg), S (60 g/kg), and Cu (6 g/kg). Likely solid phases include hydrous amorphous phases of aluminum hydroxides and sulfates, and copper hydroxides/carbonates. Efforts are undergoing to find the optimal mixing ratios between the acidic stream and more alkaline streams to maximize attenuation of dissolved copper. The results of this research could be used for enhancing in-stream natural attenuation of copper and reducing treatment needs at the drinking water facility.

Acknowledgements to Fondecyt 1130936 and Conicyt Fondap 15110020

[<< Previous Abstract](#) | [Next Abstract >>](#)