



Towards hydrochemical PUB – stable vs. heterogeneous NO₃ and COD signatures across hydrographic structure and size

Stéphane Ecrepont, Christophe Cudennec, Anne Jaffrézic, Alban de Lavenne

► To cite this version:

Stéphane Ecrepont, Christophe Cudennec, Anne Jaffrézic, Alban de Lavenne. Towards hydrochemical PUB – stable vs. heterogeneous NO₃ and COD signatures across hydrographic structure and size. EGU General Assembly 2017, Apr 2017, Vienna, Austria. Geophysical Research Abstracts, 19, pp.1, 2017. hal-02606213

HAL Id: hal-02606213

<https://hal.inrae.fr/hal-02606213>

Submitted on 16 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Towards hydrochemical PUB – stable vs. heterogeneous NO₃ and COD signatures across hydrographic structure and size

Stephane Ecrepont (1), Christophe Cudennec (1), Anne Jaffrezic (1), and Alban de Lavenne (2)

(1) UMR SAS, INRA, AGROCAMPUS OUEST 35000 Rennes, France, (2) HBAN, Irstea, Antony, France

Towards hydrochemical PUB – stable vs. heterogeneous NO₃ and DOC signatures across hydrographic structure and size

Ecrepont, S.1Cudennec, C.1 Jaffrézic, A.1 de Lavenne, A.2

1UMR SAS, Agrocampus Ouest, Rennes, France

2 HBAN, Irstea, Antony, France

Intensive agriculture is a major disturbing factor for water quality in Brittany, France. Observations of chemical data from 350 catchments over a 15 year period show that the high variability of hydrochemical dynamics between catchments in relation to geographic characteristics and farming practices, decreases with an increase in the catchment size. A stable signature of nitrate and DOC dynamics does emerge for bigger catchments, and was evidenced statistically.

We adapted a modified version of the standard deviation formula to calculate an index on mean inter-annual winter nitrate and dissolved organic carbon concentrations to characterize each catchment. The method was applied to the whole sample of catchments, some of them nested, to investigate variation of our new index across scales and regions. Results show an increasing and non-linear relationship between the criterion and the surface, with threshold effects. The stability of the thresholds across river basins in Brittany, and across seasons and years is explored.

This emergence relates to the progressive connection of streams with heterogeneous characteristic chemical signatures into a mixing dominant effect. The better assessment of this relationship opens two major perspectives: i) to define a geomorphology-based PUB (Prediction in Ungauged Basins) approach for hydrochemistry; ii) to identify the most critical sub-catchments for mitigating actions in terms of farming and landscape practices towards water quality recovery.