



HAL
open science

A conceptual snow model with an analytic resolution of the heat and phase change equations

Philippe Riboust, Nicolas Le Moine, Guillaume Thirel, Pierre Ribstein

► To cite this version:

Philippe Riboust, Nicolas Le Moine, Guillaume Thirel, Pierre Ribstein. A conceptual snow model with an analytic resolution of the heat and phase change equations. EGU General Assembly 2017, Apr 2017, Vienna, Austria. Geophysical Research Abstracts, 19, pp.1, 2017. hal-02606216

HAL Id: hal-02606216

<https://hal.inrae.fr/hal-02606216>

Submitted on 16 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A conceptual snow model with an analytic resolution of the heat and phase change equations

Philippe Riboust (1,2), Nicolas Le Moine (1), Guillaume Thirel (2), and Pierre Ribstein (1)

(1) Sorbonne Universités, UPMC Univ. Paris 06, CNRS, EPHE, UMR 7619 Metis, 4 place Jussieu, 75005 PARIS, France, (2) IRSTEA, HBAN Hydro, Antony, France (philippe.riboust@upmc.fr)

Compared to degree-day snow models, physically-based snow models resolve more processes in an attempt to achieve a better representation of reality. Often these physically-based models resolve the heat transport equations in snow using a vertical discretization of the snowpack. The snowpack is decomposed into several layers in which the mechanical and thermal states of the snow are calculated. A higher number of layers in the snowpack allow for better accuracy but it also tends to increase the computational costs.

In order to develop a snow model that estimates the temperature profile of snow with a lower computational cost, we used an analytical decomposition of the vertical profile using eigenfunctions (i.e. trigonometric functions adapted to the specific boundary conditions). The mass transfer of snow melt has also been estimated using an analytical conceptualization of runoff fingering and matrix flow.

As external meteorological forcing, the model uses solar and atmospheric radiation, air temperature, atmospheric humidity and precipitations. It has been tested and calibrated at point scale at two different stations in the Alps: Col de Porte (France, 1325 m) and Weissfluhjoch (Switzerland, 2540 m). A sensitivity analysis of model parameters and model inputs will be presented together with a comparison with measured snow surface temperature, SWE, snow depth, temperature profile and snow melt data.

The snow model is created in order to be ultimately coupled with hydrological models for rainfall-runoff modeling in mountainous areas. We hope to create a model faster than physically-based models but capable to estimate more physical processes than degree-day snow models. This should help to build a more reliable snow model capable of being easily calibrated by remote sensing and in situ observation or to assimilate these data for forecasting purposes.