



HAL
open science

On the impact of bias correcting and conditioning precipitation inputs on seasonal streamflow forecast quality

Louise Crochemore, Maria-Helena Ramos, Florian Pappenberger, Charles Perrin

► To cite this version:

Louise Crochemore, Maria-Helena Ramos, Florian Pappenberger, Charles Perrin. On the impact of bias correcting and conditioning precipitation inputs on seasonal streamflow forecast quality. EGU General Assembly 2017, Apr 2017, Vienna, Austria. Geophysical Research Abstracts, 19, pp.1, 2017. hal-02606262

HAL Id: hal-02606262

<https://hal.inrae.fr/hal-02606262>

Submitted on 16 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On the impact of bias correcting and conditioning precipitation inputs on seasonal streamflow forecast quality

Louise Crochemore (1), Maria-Helena Ramos (2), Florian Pappenberger (3), and Charles Perrin (2)

(1) Swedish Meteorological and Hydrological Institute (SMHI), Norrköping, Sweden (louise.crochemore@smhi.se), (2) Irstea, Catchment Hydrology Research Group, UR HBAN, Antony, France, (3) European Centre for Medium-Range Weather Forecasts (ECMWF), Shinfield Park, Reading, UK

Skillful seasonal streamflow forecasts are increasingly requested for decision-making in areas such as drought risk assessment or reservoir management. Meteorological forcing can be the major source of uncertainty in seasonal forecasts as early as in the first month of the forecast period. The choice of the hydrological model inputs thus has a major impact on the quality of generated streamflow forecasts. In this study, we assess the impact of two types of precipitation forecast post-treatment: 1) bias correction and 2) conditioning, on streamflow forecast quality.

We first evaluated several bias correction approaches and conditioned precipitation scenarios in sixteen catchments in France, with the help of ECMWF System 4 seasonal precipitation forecasts and the GR6J hydrological model. The results show that, in most catchments, raw seasonal precipitation and streamflow forecasts are often sharper than the conventional ESP method. However, they are not significantly better in terms of reliability. Forecast skill is generally improved when applying bias correction. The empirical distribution mapping of daily values was successful in improving forecast reliability, but sometimes at the expense of forecast sharpness.

We also evaluated several conditioning methods based on ECMWF System 4 precipitation forecasts to generate seasonal streamflow forecasts in the same sixteen catchments. Four precipitation indices based on System 4 precipitation were used to condition historical streamflow or historical precipitations to be used as input to the GR6J model. Our results evaluate how the conditioning impacts the reliability and sharpness of streamflow forecasts, as well as forecasts of drought indices. We show that conditioning past observations based on the three-month Standardized Precipitation Index (SPI3) can improve the sharpness of ensemble forecasts based on historical data, but also often decrease reliability.

References:

Crochemore, L., Ramos, M.-H., and Pappenberger, F.: Bias correcting precipitation forecasts to improve the skill of seasonal streamflow forecasts, *Hydrol. Earth Syst. Sci.*, 20, 3601-3618, doi:10.5194/hess-20-3601-2016, 2016.

Crochemore, L., Ramos, M.-H., Pappenberger, F., and Perrin, C.: Seasonal streamflow forecasting by conditioning climatology with precipitation indices, *Hydrol. Earth Syst. Sci. Discuss.*, doi:10.5194/hess-2016-285, in review, 2016.