



**HAL**  
open science

## Continental mapping of forest ecosystem functions reveals a high but unrealised potential for forest multifunctionality

P. Fons van Der, S. Ratcliffe, P. Ruiz-Benito, Michael Scherer-Lorenzen, Kris Verheyen, C. Wirth, M.A. Zavala, E. Ampoorter, L. Baeten, Luc Barbaro, et al.

### ► To cite this version:

P. Fons van Der, S. Ratcliffe, P. Ruiz-Benito, Michael Scherer-Lorenzen, Kris Verheyen, et al.. Continental mapping of forest ecosystem functions reveals a high but unrealised potential for forest multifunctionality. *Ecology Letters*, 2018, 21 (1), pp.31-42. 10.1111/ele.12868 . hal-02606618

**HAL Id: hal-02606618**

**<https://hal.inrae.fr/hal-02606618>**

Submitted on 19 Jan 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

1 ***"This is the peer reviewed version of the following article: van der Plas, F.***  
2 ***et al. Continental mapping of forest ecosystem functions reveals a high but***  
3 ***unrealised potential for forest multifunctionality. Ecol. Lett. 31–42 (2017),***  
4 ***which has been published in final form at doi:10.1111/ele.12868. This***  
5 ***article may be used for non-commercial purposes in accordance with***  
6 ***Wiley Terms and Conditions for Self-Archiving."***

7  
8 **TITLE:** Continental mapping of forest ecosystem functions reveals a high but unrealized  
9 potential for forest multifunctionality

10

11 **AUTHORS:**

12 Fons van der Plas<sup>1,2,3,\*</sup>, Sophia Ratcliffe<sup>3</sup>, Paloma Ruiz-Benito<sup>4,5</sup>, Michael Scherer-Lorenzen<sup>6</sup>,  
13 Kris Verheyen<sup>7</sup>, Christian Wirth<sup>3,8</sup>, Miguel A. Zavala<sup>5</sup>, Evy Ampoorter<sup>7</sup>, Lander Baeten<sup>7</sup>, Luc  
14 Barbaro<sup>9,10</sup>, Cristina C. Bastias<sup>11</sup>, Jürgen Bauhus<sup>12</sup>, Raquel Benavides<sup>6</sup>, Adam Benneter<sup>12</sup>,  
15 Damien Bonal<sup>13</sup>, Olivier Bouriaud<sup>14</sup>, Helge Bruelheide<sup>8,15</sup>, Filippo Bussotti<sup>16</sup>, Monique Carnol<sup>17</sup>,  
16 Bastien Castagneyrol<sup>9</sup>, Yohan Charbonnier<sup>9</sup>, Johannes H. C. Cornelissen<sup>18</sup>, Jonas Dahlgren<sup>19</sup>,  
17 Ewa Checko<sup>20</sup>, Andrea Coppi<sup>21</sup>, Seid Muhie Dawud<sup>22</sup>, Marc Deconchat<sup>9,10</sup>, Pallieter De Smedt<sup>7</sup>,  
18 Hans De Wandeler<sup>23</sup>, Timo Domisch<sup>24</sup>, Leena Finér<sup>24</sup>, Mariangela Fotelli<sup>25</sup>, Arthur Gessler<sup>26</sup>,  
19 André Granier<sup>13</sup>, Charlotte Grossiord<sup>27</sup>, Virginie Guyot<sup>9,10</sup>, Josephine Haase<sup>6,28</sup>, Stephan  
20 Hättenschwiler<sup>29</sup>, Hervé Jactel<sup>9</sup>, Bogdan Jaroszewicz<sup>20</sup>, François-Xavier Joly<sup>29</sup>, Tommaso  
21 Jucker<sup>30</sup>, Stephan Kambach<sup>15,31</sup>, Gerald Kaendler<sup>32</sup>, Jens Kattge<sup>8,33</sup>, Julia Koricheva<sup>34</sup>, Georges  
22 Kunstler<sup>35</sup>, Aleksi Lehtonen<sup>36</sup>, Mario Liebergesell<sup>3,8</sup>, Peter Manning<sup>1,2</sup>, Harriet Milligan<sup>34</sup>,

23 Sandra Müller<sup>6</sup>, Bart Muys<sup>23</sup>, Diem Nguyen<sup>37</sup>, Charles Nock<sup>6</sup>, Bettina Ohse<sup>3</sup>, Alain Paquette<sup>38</sup>,  
24 Josep Peñuelas<sup>39,40</sup>, Martina Pollastrini<sup>16</sup>, Kalliopi Radoglou<sup>41</sup>, Karsten Raulund-Rasmussen<sup>22</sup>,  
25 Fabian Roger<sup>42</sup>, Rupert Seidl<sup>43</sup>, Federico Selvi<sup>16</sup>, Jan Stenlid<sup>37</sup>, Fernando Valladares<sup>11</sup>, Johan van  
26 Keer<sup>44</sup>, Lars Vesterdal<sup>22</sup>, Markus Fischer<sup>1,2</sup>, Lars Gamfeldt<sup>42</sup>, Eric Allan<sup>1</sup>.

27

## 28 **AUTHOR AFFILIATIONS**

29 <sup>1</sup>Institute of Plant Sciences, University of Bern, Altenbergrain 21, 3013 Bern, Switzerland.

30 <sup>2</sup>Senckenberg Gesellschaft für Naturforschung, Biodiversity and Climate Research Centre, Senckenberganlage 25,  
31 60325 Frankfurt, Germany.

32 <sup>3</sup>Department of Systematic Botany and Functional Biodiversity, University of Leipzig, Johannisallee 21-23, 04103  
33 Leipzig, Germany.

34 <sup>4</sup>Biological and Environmental Sciences, School of Natural Sciences, University of Stirling, FK9 4LA, Stirling,  
35 United Kingdom.

36 <sup>5</sup>Grupo de Ecología y Restauración Forestal, Departamento de Ciencias de la Vida, Universidad de Alcalá, Edificio  
37 de Ciencias, Campus Universitario, 28805 Alcalá de Henares, Madrid, Spain.

38 <sup>6</sup>University of Freiburg, Faculty of Biology, Geobotany, Schänzlestr. 1, 79104 Freiburg, Germany.

39 <sup>7</sup>Forest & Nature Lab, Ghent University, Geraardsbergsesteenweg 267, B-9090 Melle-Gontrode, Belgium.

40 <sup>8</sup>German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5E, 04103  
41 Leipzig, Germany.

42 <sup>9</sup>BIOGECO, INRA, Univ. Bordeaux, 33610 Cestas, France.

43 <sup>10</sup>Dynafor, INPT-EI Purpan, INRA, Univ. Toulouse, 31320, Auzeville, France

44 <sup>11</sup>MNCN-CSIC, Serrano 115 bis 28006 Madrid, Spain.

45 <sup>12</sup>Faculty of Environment and Natural Resources, Chair of Silviculture, University of Freiburg, Fahrenbergplatz,  
46 79085 Freiburg, Germany.

47 <sup>13</sup>INRA, UMR EEF, 54280 Champenoux, France.

48 <sup>14</sup>Faculty of Forestry, Stefan cel Mare University of Suceava, Universitatii Street 13, Suceava 720229, Romania.

49 <sup>15</sup>Institute of Biology / Geobotany and Botanical Garden, Martin Luther University Halle-Wittenberg, Am Kirchtor  
50 1, 06108 Halle (Saale), Germany.

51 <sup>16</sup>Department of Agrifood Production and Environmental Sciences, Laboratory of Applied and Environmental  
52 Botany, University of Firenze, P.le Cascine 28, 50144 Firenze, Italy.

53 <sup>17</sup>Laboratory of Plant and Microbial Ecology, University of Liege, Botany B22, Chemin de la Vallee 4, 4000 Liege,  
54 Belgium.

55 <sup>18</sup>Systems Ecology, Department of Ecological Science, Faculty of Earth and Life Sciences, Vrije Universiteit  
56 Amsterdam, De Boelelaan 1085, 1081 HV, Amsterdam, The Netherlands.

57 <sup>19</sup>Swedish University of Agricultural Sciences, Skogsmarksgränd, 90183 Umeå, Sweden.

58 <sup>20</sup>Białowieża Geobotanical Station, Faculty of Biology, University of Warsaw, 17-230 Białowieża, Poland.

59 <sup>21</sup>University of Firenze, Department of Biology, Botanical Laboratories, Via G. La Pira 4, 50121 Firenze, Italy.

60 <sup>22</sup>Department of Geosciences and Natural Resource Management, University of Copenhagen, Rolighedsvej 23, 1958  
61 Frederiksberg C, Denmark.

62 <sup>23</sup>Department of Earth and Environmental Sciences, KU Leuven, University of Leuven, Celestijnenlaan 200E Box  
63 2411, BE-3001 Leuven, Belgium.

64 <sup>24</sup>Natural Resources Institute Finland (Luke), Yliopistokatu 6, FI-80100 Joensuu, Finland.

65 <sup>25</sup>Forest Research Institute of Thessaloniki, Greek Agricultural Organization-Dimitra, 57006 Vassilika,  
66 Thessaloniki, Greece.

67 <sup>26</sup>Swiss Federal Research Institute WSL, Research Unit Forest Dynamics, Zuercherstr, 111, 8903 Birmensdorf,  
68 Switzerland.

69 <sup>27</sup>Earth and Environmental Sciences Division, Los Alamos National Laboratory, Los Alamos, NM 87545, USA.

70 <sup>28</sup>Institute for Terrestrial Ecosystems, Department of Environmental Systems Science, ETH Zurich,  
71 Universitaetsstrasse 16, 8092 Zurich, Switzerland.

72 <sup>29</sup>Centre of Evolutionary and Functional Ecology (CEFE UMR 5175, CNRS – University of Montpellier –  
73 University Paul-Valery Montpellier – EPHE), 1919 route de Mende, 34293 Montpellier, France.

74 <sup>30</sup>Forest Ecology and Conservation, Department of Plant Sciences, University of Cambridge, Downing Street,  
75 Cambridge CB2 3EA, UK.

76 <sup>31</sup>UFZ – Helmholtz Centre for Environmental Research, Department Community Ecology, 06120 Halle (Saale),  
77 Germany.

78 <sup>32</sup>Forest Research Institute Baden-Wurttemberg, Wonnhaldestrase 4, 79100 Freiburg, Germany.

79 <sup>33</sup>Max Planck Institute for Biogeochemistry, Hans-Knöll-Straße 10, 07745 Jena, Germany.

80 <sup>34</sup>School of Biological Sciences, Royal Holloway University of London, Egham, Surrey TW20 0EX, UK.

81 <sup>35</sup>Univ. Grenoble Alpes, Irstea, UR EMGR, Centre de Grenoble, 2 rue de la Papeterie-BP 76, F-38402 Saint-Martin-  
82 d’Hères, France.

83 <sup>36</sup>Natural Resources Institute Finland (Luke), Jokiniemenkuja 1, FI-01370 Vantaa, Finland.

84 <sup>37</sup>Department of Forest Mycology and Plant Pathology, Swedish University of Agricultural Sciences, PO Box 7026,  
85 SE-750 07 Uppsala, Sweden.

86 <sup>38</sup>Centre for Forest Research (CFR), Université du Québec à Montréal, Montréal (Québec), Canada.

87 <sup>39</sup>CREAF, Cerdanyola del Vallès, 08913 Catalonia, Spain.

88 <sup>40</sup>CSIC, Global Ecology Unit CREAM-CSIC-UB-UAB, Bellaterra, 08913 Catalonia, Spain.

89 <sup>41</sup>Democritus University of Thrace (DUTH), Department of Forestry and Management of the Environment and  
90 Natural Resources, Pantazidou 193, 68200, Nea Orestiada, Greece.

91 <sup>42</sup>Department of Marine Sciences, University of Gothenburg, Carl Skottsbergs gata 22B, 41319 Göteborg, Sweden.

92 <sup>43</sup>University of Natural Resources and Life Sciences (BOKU), Institute of Silviculture, Vienna, Austria.

93 <sup>44</sup>Bormstraat 204 bus 3, 1880 Kapelle-op-den-Bos, Belgium.

94

95 \*corresponding author: [Fonsvanderplas@gmail.com](mailto:Fonsvanderplas@gmail.com); Senckenberganlage 25, D-60325 Frankfurt am Main, Germany;  
96 Tel.: +49 69 7542 1820; Fax: + 49 69 7542 7904

## 98 **AUTHOR CONTRIBUTIONS**

99 FvdP, EA, LG, MF, SR, PRB, MSL, CW, LB, RB, HB, TJ, SK, GK, CN, BO, AP and FR  
100 developed the ideas of this study at a workshop organized by EA and LG. FvdP, SR and PRB  
101 analysed the data. All authors, except FvdP, EA, MF, SKa, PM, BO, AP and FR contributed to  
102 the data collection. FvdP wrote the manuscript. All authors contributed in editing the manuscript.

103

104 **DATA ACCESSIBILITY STATEMENT**

105 Should the manuscript be accepted, the data supporting the results will be published in Dryad  
106 and the data DOI will be included at the end of the article.

107

108 **RUNNING TITLE:** continental mapping of multifunctionality

109

110 **KEYWORDS:** biodiversity, climate, ecosystem multifunctionality, ecosystem services, forest,  
111 FunDivEUROPE, large-scale, phylogenetic diversity, tree communities, upscaling

112

113 **TYPE OF ARTICLE:** letter

114

115 **NUMBER OF WORDS**

116 Abstract: 150

117 Main text: 4999

118

119 **NUMBER OF REFERENCES:** 51

120 **NUMBER OF FIGURES:** 4

121 **NUMBER OF TABLES:** 1

122 **NUMBER OF TEXT BOXES:** 0

123 **ABSTRACT**

124           Humans require multiple services from ecosystems, but it is largely unknown whether  
125 trade-offs between ecosystem functions prevent the realization of high ecosystem  
126 multifunctionality across spatial scales. Here, we combined a comprehensive dataset (28  
127 ecosystem functions measured on 209 forest plots) with a forest inventory dataset (105,316 plots)  
128 to extrapolate and map relationships between various ecosystem multifunctionality measures  
129 across Europe. These multifunctionality measures reflected different management objectives,  
130 related to timber production, climate regulation and biodiversity conservation/recreation. We  
131 found that trade-offs among them were rare across Europe, at both local and continental scales.  
132 This suggests a high potential for "win-win" forest management strategies, where overall  
133 multifunctionality is maximized. However, across sites, multifunctionality was on average 45.8-  
134 49.8% below maximum levels and not necessarily highest in protected areas. Therefore, using  
135 one of the most comprehensive assessments so far, our study suggests a high but largely  
136 unrealized potential for management to promote multifunctional forests.

137

138 **INTRODUCTION**

139           One of the greatest challenges in ecology is to understand the effects of global change  
140 and nature management on the multiple ecosystem functions on which humans depend (MEA  
141 2005). Such an understanding would help predicting the circumstances under which trade-offs  
142 between different ecosystem functions are minimal and therefore when their simultaneous  
143 provisioning, i.e. ecosystem multifunctionality (Hector & Bagchi 2007; Gamfeldt *et al.* 2008), is  
144 maximised. Previous studies have identified conditions promoting local-scale ecosystem  
145 multifunctionality, e.g. through the maximization of biodiversity (Lefcheck *et al.* 2015).  
146 However, whether such relationships also exist at large spatial scales, and how they vary in  
147 space, is less clear (Isbell *et al.* 2017). Understanding this is essential if ecosystem-functioning  
148 studies are to provide policy-relevant advice, because most policy focuses on large scales.

149           Forests provide a number of functions related to key services such as timber production,  
150 climate regulation and recreation (Gamfeldt *et al.* 2013), and are important for the conservation  
151 of many plant and animal species (FAO 2015). Understanding large-scale relationships between  
152 different functions is therefore important if we are to find “win-win” management scenarios,  
153 which meet different forest management objectives and promote forest multifunctionality.

154           Quantifying many ecosystem functions at large scales has so far proven challenging.  
155 Studies have used exhaustive remote sensing or ground-based measurements (e.g. Prince &  
156 Goward 1995; Ratcliffe *et al.* 2016), mechanistic models (e.g. McGuire *et al.* 2001), indirect  
157 measures (e.g. where certain habitat types are assumed to promote certain functions; Maskell *et*  
158 *al.* 2013) or a combination of these (Maes *et al.* 2012; Mouchet *et al.* 2017) to quantify single or  
159 multiple functions at large spatial extents. However, for some important functions, such as  
160 biological pest control or timber quality, large scale maps have not yet been developed, limiting



161 our understanding of ecosystem functioning synergies and trade-offs. In contrast, many local-  
162 scale studies, such as biodiversity experiments (e.g. Hector & Bagchi 2007; Zavaleta *et al.* 2009)  
163 or comparative studies (Lavorel *et al.* 2011), have accurately quantified a large number of  
164 functions. Extrapolating these small-scale observations to larger scales could increase our  
165 understanding of the drivers of ecosystem functioning trade-offs and the resulting provision of  
166 ecosystem multifunctionality.

167         Forests are often managed for a particular subset of functions related to certain ecosystem  
168 services (e.g. timber production, climate regulation or nature conservation) that are prioritized by  
169 a specific stakeholder group. We aimed to identify areas where functions of all these different  
170 sets are high and where trade-offs are weakest. To this end, we combined a multi-site dataset,  
171 containing accurate measures of multiple ecosystem functions, with a continental-scale  
172 inventory-based dataset with high spatial plot coverage. We extrapolated regional scale  
173 relationships between ecosystem functions and their drivers (e.g. forest community composition  
174 and climate) to larger spatial scales (Fig. S1) to map both individual ecosystem functions and  
175 ecosystem multifunctionality across Europe, in forests without recent intensive management. We  
176 then tested for potential trade-offs between sets of functions, at scales relevant for policymakers.

177         To do this, we developed different measures of multifunctionality corresponding to  
178 different management scenarios (Fig. 1). In these, functions related to (sustainable) timber  
179 production, climate regulation or biodiversity conservation/recreation were prioritized (Fig. 1).  
180 We also considered a scenario where all functions were valued equally. Our objectives were  
181 firstly, to identify "multifunctionality hotspots", i.e. areas with highest multifunctionality.  
182 Secondly, we investigated whether there are synergies (allowing for win-win management) or  
183 trade-offs between different multifunctionality measures at both continental and local scales, and

184 how these varied in space. Finally, we investigated whether forest protection status is associated  
185 with high multifunctionality, and thus whether potential win-win policies are realized in  
186 (protected) forests.

187

## 188 **MATERIALS AND METHODS**

189 Our approach to extrapolate ecosystem functioning relationships from regional to  
190 continental scales consisted of two main steps (Fig. S1). Firstly, statistical models were fitted to a  
191 comprehensive (many ecosystem functions), multi-site dataset ('fitting dataset'). Secondly, these  
192 models were extrapolated to a continental-scale dataset containing forest plots distributed across  
193 Western Europe ('inventory dataset'). These two datasets share variables related to climate, soils  
194 and tree composition, all potential drivers of ecosystem functioning. For three ecosystem  
195 functions which were independently measured in the inventory dataset, we cross-validated  
196 predicted ecosystem function values. Our approach allowed testing for trade-offs and synergies  
197 between individual ecosystem functions and between different multifunctionality measures, at  
198 different scales: 1) using all plots (thus including both local and large-scale variation in  
199 functions) and 2) within 20×20km localities.

200

### 201 *Fitting dataset: design*

202 As part of the EU-FP7 FunDivEUROPE project ([www.fundiveurope.eu](http://www.fundiveurope.eu)), which  
203 investigates how tree species composition and diversity drive forest ecosystem functioning, 209  
204 30×30 meter plots (Fig. S2) were established. The plots covered six major regions/countries,  
205 representing different forest types: 28 boreal forest (Finland), 43 temperate mixed forest  
206 (Poland), 38 temperate deciduous forest (Germany), 28 mountainous deciduous forest

207 (Romania), 36 thermophilous deciduous forest (Italy) and 36 Mediterranean mixed forest plots  
208 (Spain). These plots covered a broad climatic gradient: mean annual precipitation ranged from  
209 484 to 819mm, mean annual temperature from 1.4 to 14.1°C (WorldClim; Hijmans *et al.* 2005)  
210 and altitude from 87 to 1404m. Within regions, plots differed in the composition and diversity of  
211 regionally common tree species, while site-related factors were similar. Management was either  
212 at low intensity or absent (Baeten *et al.* 2013).

213

#### 214 *Measurement and collation of fitting data*

215 In all plots, we measured 28 different ecosystem characteristics/processes ('ecosystem  
216 functions' hereafter) linked to various ecosystem services (see overview in Fig. 1 and  
217 methodology in Supplementary Material). For each plot we compiled data on tree species  
218 composition (to derive measures of functional and phylogenetic diversity), stand structure, soil  
219 pH, altitude and 18 climatic variables. Previous studies demonstrated that climate (Cramer *et al.*  
220 2001), soil pH (Foy 1992), functional community composition (Diaz *et al.* 2004) and tree  
221 diversity (van der Plas *et al.* 2016; Liang *et al.* 2016) can all drive (forest) functioning.

222 In each plot, we identified all tree stems  $\geq 7.5$ cm in diameter at breast height (dbh) to  
223 species level. With these data, we calculated total and average tree basal area. In addition, by  
224 combining these observations with (1) published trait data (Kattge *et al.* 2011; Royal Botanic  
225 Gardens Kew 2015; see Table S1) representing key life-history strategies (Westoby *et al.* 2002),  
226 and (2) a phylogeny (Zanne *et al.* 2014), we calculated several metrics describing the functional  
227 identity, functional diversity and phylogenetic diversity of the tree communities. Firstly, we  
228 calculated Community Weighted Means (Garnier *et al.* 2004), reflecting functional identities of  
229 communities, based on species values for specific leaf area ( $\text{cm}^2 \text{g}^{-1}$ ), maximum life span (log-

230 transformed; yrs), maximum height (m), wood density ( $\text{g cm}^{-3}$ ), seed mass (mg), conifer  
231 (proportion) and evergreen (proportion). Secondly, we calculated the functional (trait) diversity  
232 within communities as Rao's Quadratic Entropy (Botta-Dukát 2005), for each trait separately  
233 and for all traits combined. Finally, we calculated several phylogenetic diversity metrics:  
234 Phylogenetic Species Variability, Phylogenetic Species Evenness (Helmus *et al.* 2007), Faith's  
235 Phylogenetic Diversity (Faith 1992) and (abundance-weighted) Mean Phylogenetic Distance  
236 (Webb *et al.* 2002). As inventory plots differed in size, tree species richness was not  
237 investigated, and we selected functional and phylogenetic diversity metrics uncorrelated with  
238 species richness.

239 To represent soil conditions we used pH (methods in Supplementary Materials), as it  
240 drives many functions and was the only soil variable available for the inventory dataset. Eighteen  
241 variables (see Table S2) related to climate (worldclim data; Hijmans *et al.* 2005) were collated at  
242 a 30 seconds spatial resolution. Altitude data were collated from [srtm.csi.cgiar.org](http://srtm.csi.cgiar.org).

243

#### 244 *Analysis of the drivers of ecosystem functioning*

245 We used the Random Forest (Breiman 2001) algorithm to explain ecosystem function  
246 variation in the fitting dataset. Random Forest is a machine-learning algorithm, powerful for  
247 making predictions (but less suitable in explaining mechanisms) and incorporating both linear  
248 and non-linear relationships, as well as interaction effects (Strobl *et al.* 2007). It is relatively  
249 insensitive to multicollinearity and overfitting (Hastie *et al.* 2008), allowing for the inclusion of  
250 many predictors. Initially, we included the 42 predictor variables described above (see also Table  
251 S2), describing abiotic conditions, climate, stand structure, functional identity, and functional  
252 and phylogenetic diversity. Random Forests were run in R (R Core Team 2013) with the

253 ‘randomForest’ library (Liaw & Wiener 2012). Following Seidl *et al.* (2011), we iteratively  
254 removed those variables not reducing the mean square error over random permutations of the  
255 same variable. For final Random Forests, we identified, using the ‘importance’ function, the  
256 degree to which the inclusion of each predictor decreases residual model variance.

257

### 258 *Forest inventory data*

259 We combined data from 163,451 plots of the National Forest Inventories (NFIs) of Spain  
260 (59,048 plots), France (40,844), Wallonia (Belgium, 1,238), Germany (47,832), Sweden (11,212)  
261 and Finland (2,456). NFIs contained data on individual trees in each plot, including species  
262 identity, dbh and basal area. Furthermore, estimates of timber production (increase in tree basal  
263 area per hectare per year), tree biomass and tree recruitment (tree saplings per hectare) were  
264 available for many plots. To ensure that data from different NFIs were comparable to the fitting  
265 dataset plots, we only included trees with dbh  $\geq 7.5$ cm. Furthermore, we only included the  
266 105,316 plots that were at low to mid-altitudes (<1500m), without indication of recent logging,  
267 and dominated by one of the ‘target’ species of the fitting dataset (Baeten *et al.* 2013).

268 We calculated the same climate, functional identity and functional and phylogenetic  
269 diversity variables for the NFI dataset as for the fitting dataset. Soil pH, calculated for the top  
270 10cm of the soil at 1km<sup>2</sup> resolution, was obtained from the ESDAC database (Panagos *et al.*  
271 2012). These variables had similar ranges as in the fitting dataset (Table S3).

272

### 273 *Extrapolating and mapping ecosystem functions across Europe*

274 We used the ‘predict’ function in R to predict values of each ecosystem function in  
275 inventory plots, based on the Random Forests (built using the fitting dataset with independently

276 collected FunDivEUROPE data; Baeten et al. 2013) and the climate, functional identity,  
277 diversity (of the most recent survey) and abiotic conditions in the inventory plots. To determine  
278 the accuracy of our predictions, we correlated the three ecosystem functions (timber production,  
279 tree biomass and tree recruitment) that were measured in inventory plots with the values  
280 predicted by the Random Forests. We did the validations across all plots at continental scale  
281 (local and large scale variation) and within (only local variation) and among (only large-scale  
282 variation) 20×20km grid cells ('localities') containing  $\geq 20$  plots. In addition, we compared  
283 observed correlations between ecosystem functions with extrapolated ones. We also compared  
284 the average values for tree biomass and recruitment between fitting and inventory datasets  
285 (productivity was not comparable as it was measured in different units). To investigate how  
286 mapped functions changed across latitude, we fitted linear models with linear and quadratic  
287 effects of latitude as predictors.

288

### 289 *Calculating multifunctionality and quantifying trade-offs*

290 We used the 'threshold-approach' (Gamfeldt *et al.* 2008) to calculate ecosystem  
291 multifunctionality for each inventory plot, based on the predicted values of individual ecosystem  
292 functions. Ecosystem multifunctionality was measured at both local and continental scales and  
293 defined as the number of functions exceeding a threshold. The threshold was defined as the  
294 proportion (25%, 50% (default threshold reported in main results), 75% or 90%) of the  
295 'maximum' value observed for that function, either within a 20×20km locality (local scale) or  
296 across Europe (continental scale). The maximum was defined as the 97.5<sup>th</sup> percentile of  
297 observed functioning across plots, thus removing extreme outliers. For a concrete example on  
298 quantifying multifunctionality, we refer to Fig. S3. We excluded ecosystem functions that (a) had

299 poor Random Forest fit, with  $R^2$  (correlation between observed and predicted) values <0.20  
300 (default analysis; Fig. 1C) and (b), as a sensitivity analysis, also those which had a low validation  
301  $R^2$  (see results: tree recruitment and the related function of seedling growth). As a further  
302 sensitivity analysis, we calculated ecosystem multifunctionality using Random Forest  $R^2$  values  
303 as weights.

304 We also calculated multifunctionality according to various management objectives,  
305 following Allan *et al.* (2015). In these measures, we gave different weightings to the various  
306 ecosystem functions, according to their presumed importance (based on a consensus of expert  
307 opinions of all authors) for delivering the ecosystem services required for the given objective  
308 (Fig. 1). The equal weights measure described above corresponds with most previous studies  
309 (e.g. Lefcheck *et al.* 2015). In the measures representing management objectives, functions were  
310 weighted with loadings ranging from 0 (unimportant) to 1 (high importance). Functions directly  
311 related to the objective received a weight of 1, i.e. timber production and quality for ‘timber  
312 production multifunctionality’, carbon sequestration-related functions for ‘climate regulation’  
313 and functions directly measuring biodiversity (e.g. bird/understory diversity) for ‘biodiversity  
314 conservation/recreation’. Other functions were weighted 0.25; 0.50 or 0.75, depending on their  
315 relevance (Fig. 1). We also quantified a ‘narrow-sense’ biodiversity conservation measure,  
316 where only functions directly measuring biodiversity were included, with weights of 1 (Fig. 1).

317 Relationships between multifunctionality measures can either be caused by large-scale  
318 climatic/biogeographical factors (e.g. temperature gradients) or local-scale factors (e.g.  
319 management, soil conditions). Therefore, using Pearson correlations, we tested for trade-offs and  
320 synergies, at both continental (all plots) and local scales (within localities with >10 plots). With  
321 *t*-tests we investigated whether local-scale correlations, differed from zero.

322 Several functions had high weights in multiple multifunctionality measures, reflecting their  
323 relevance for different ecosystem services (Fig. 1B). Raw correlation coefficients between  
324 multifunctionality measures are therefore inflated by this overlap. To remove this effect, we  
325 calculated a null expectation for the correlation-coefficients by reshuffling ecosystem function  
326 values, without replacement, across plots 100 times. This eliminated any correlations among  
327 functions, while maintaining the original distribution of values. With these resampled ecosystem  
328 functions, we again calculated the different multifunctionality measures, and the average and  
329 95% confidence intervals of the correlations between them. We calculated correlation-  
330 coefficients corrected for overlap in functions by subtracting expected values (in the absence of  
331 correlations among functions) from observed ones. As a sensitivity analysis, we repeated these  
332 analyses only including plots located within those 150 localities in which validations of both  
333 timber production and tree biomass were adequate (both  $r > 0.1$ ).

334

### 335 *Comparing multifunctionality between protected versus non-protected forests*

336 In total, 11.8% of the inventory plots were within protected areas which, depending on  
337 the NFI, indicated either that forestry activities were restricted (Germany, Sweden) or that the  
338 plot was in a National Park or nature reserve (Finland, France, Spain, Wallonia), see  
339 Supplementary Material for more detailed information. Within each country, we investigated, for  
340 each measure, whether local-scale multifunctionality was higher inside versus outside protected  
341 areas, using Welch's  $t$ -tests.

342

## 343 **RESULTS**

### 344 *Explaining variation in ecosystem functioning*



345 On average, across the different ecosystem functions in our fitting dataset, Random  
346 Forests explained 40.7% of the total variation. The explained variation in ecosystem functions  
347 ranged from high (timber production: 72.5%; resistance to insect herbivory: 67.6%) to low  
348 (browsing resistance: 2.4%, Fig. 1C). The single most important explanatory factor (i.e. with  
349 lowest residual variance) varied between the functions. For sixteen functions it was a climate  
350 variable, for six a functional identity variable, for two altitude, for two a functional diversity  
351 variable and soil pH and average stem diameter for one each (Fig. 1C; Table S4).

352 Three ecosystem functions allowed for validation of predicted values in inventory plots.  
353 For timber production and tree biomass, across all plots, predicted values correlated reasonably  
354 well with observed values, with ‘extrapolation’  $R^2$  values (correlation between predicted and  
355 observed values in inventory plots) of 0.219 and 0.280, respectively. For tree recruitment the  $R^2$   
356 was only 0.040; Fig. S4. Validations generally worked best at large spatial scales and less well at  
357 local scales. Correlations between predicted and observed values of timber production, tree  
358 biomass and tree recruitment were, respectively, 0.390; 0.472 and 0.027 *across* 20×20km  
359 localities, and on average 0.127 (range: 0-0.976); 0.124 (range: 0-0.971) and 0.091 (range: 0-  
360 0.967) *within* localities. Absolute values of tree biomass were similar between NFI observations  
361 and Random Forest predictions, but for tree recruitment the values differed (Fig. S5). For more  
362 information on model validations, see Supplementary Material (S3).

363

### 364 *Levels of ecosystem functioning and multifunctionality throughout Western Europe*

365 After removing ecosystem functions poorly explained by the Random Forests ( $R^2 < 0.2$ ;  
366 see Fig. 1C), we predicted levels of 22 ecosystem functions for the inventory plots (Fig. S6).  
367 Many of the mapped functions showed clear continental trends. For example, some (e.g. timber

368 production) had highest levels in central Western Europe, while others had highest values in  
369 boreal (e.g. timber quality) or Mediterranean (e.g. bat diversity) regions (Fig. S6; Table S5).  
370 Most functions tended to be highest at mid-latitudes. Consequently, most continental-scale  
371 multifunctionality measures were highest in central Western Europe (multifunctionality hotspots)  
372 and lowest in southern Europe (Fig. 2). When only diversity measures were considered (narrow-  
373 sense biodiversity conservation), multifunctionality was also high in southern/central Spain and  
374 parts of Scandinavia. These patterns were broadly similar when functions with a high proportion  
375 of explained variance were weighted more heavily (Fig. S7). As expected, local-scale  
376 multifunctionality values did not show any large-scale spatial patterns (Fig. S8). Local  
377 multifunctionality scores were on average 45.8%, 47.1%, 49.2%, 49.8% and 47.8% below their  
378 maximum possible score (i.e. all functions above the 50% threshold) in the timber production,  
379 climate regulation, broad-sense and narrow-sense biodiversity conservation and overall  
380 multifunctionality scenario, respectively, and higher than 90% of the maximum possible score in  
381 97, 49, 49 and 11,625 plots (out of 105,316 plots) in the timber production, climate regulation,  
382 broad-sense and narrow-sense biodiversity conservation scenario, respectively, whereas it  
383 exceeded 90% and 80% of maximum overall multifunctionality in only 3 and 446 plots  
384 respectively (Fig. 2B). Importantly, while ecosystem functions varied strongly at the continental  
385 scale (with 97.5 percentile values being on average 42.8% higher than mean values), there was  
386 also substantial variation within localities, with 97.5 percentile values being on average 12.6%  
387 higher than mean values (Table S6).

388

389 *Trade-offs and synergies*

390 Pairwise correlations between individual functions were positive on average at both  
391 scales, although correlations were weaker at local ( $\bar{r} = 0.012$ ) than at continental scales ( $\bar{r} =$   
392  $0.021$ ), probably due to lower variation in functioning within localities (Table S6). Moderately to  
393 strongly positive correlations ( $r > 0.3$ ;  $n = 57$  (continental-scale) and  $22$  (local scale))  
394 outnumbered negative ( $r < -0.3$ ;  $n = 45$  (continental-scale) and  $14$  (local-scale)) correlations  
395 (Table S7,8). At the continental scale, correlations between timber production and tree biomass  
396 were similar for observed ( $r = 0.55$ ) and extrapolated ( $r = 0.65$ ) values. However, within  
397 localities this match was weaker ( $\bar{r} = 0.63$  observed and  $0.24$  predicted), with fits generally best  
398 in France and central/southern Spain, and weaker in Germany and northeast Spain (Fig. S9).

399 As different multifunctionality variables had similar continental-scale patterns (Fig. 2),  
400 continental-scale correlations between most measures were positive (Table 1). Only correlations  
401 between narrow-sense biodiversity conservation and both timber production ( $r = -0.13$ ) and  
402 climate regulation multifunctionality ( $r = 0.01$ ) were not. These correlations became more  
403 positive at more extreme (25 and 90%) multifunctionality thresholds (Table S9-S11).

404 Within localities, similar patterns were found. Relationships between timber production,  
405 climate regulation and broad-sense biodiversity conservation/recreation were positive, whereas  
406 relationships between narrow-sense biodiversity conservation and other multifunctionality  
407 variables were close to zero, or negative, on average (Fig. 3, Table 1). Negative relationships  
408 largely disappeared when multifunctionality was based on 25% or 90% thresholds (Table S9-  
409 S11). Importantly, positive relationships between timber production and climate regulation  
410 multifunctionality, and to a lesser extent between timber production/climate regulation  
411 multifunctionality and broad-sense biodiversity conservation/recreation multifunctionality, were  
412 very widespread across Europe (Fig. 3).

413 We used null models to investigate whether observed correlations between  
414 multifunctionality variables were larger than expected. Relationships between multifunctionality  
415 variables were to a large extent driven by functions contributing to multiple multifunctionality  
416 variables, as observed minus expected correlation-coefficients were often close to zero (Fig. 3,  
417 Table 1). Nevertheless, at a continental scale, relationships between timber production, climate  
418 regulation and broad-sense biodiversity conservation multifunctionality remained significantly  
419 positive (all  $P < 0.05$ ). At the local scale, relationships between timber production and climate  
420 regulation multifunctionality also remained significantly (although weakly) positive, whereas  
421 relationships between timber production and the biodiversity conservation measures became  
422 significantly, weakly, negative. In sensitivity analyses these patterns hardly changed when (i)  
423 recruitment-related functions were omitted from multifunctionality measures, (ii) ecosystem  
424 functions with a high Random Forest fit had proportionally higher loadings in multifunctionality  
425 measures, or (iii) only plots from localities with high validation  $R^2$  values of Random Forests  
426 explaining timber production and tree biomass were included (Table 1). Negative relationships  
427 largely disappeared when multifunctionality was quantified based on 25% or 90% thresholds  
428 (Table S9-S11). Importantly, functional overlap-corrected correlation-coefficients between  
429 different ecosystem multifunctionality scenarios varied greatly, from positive to negative,  
430 throughout localities (Fig. 3).

431

#### 432 *Multifunctionality inside versus outside protected areas*

433 Local-scale associations between values of multifunctionality and protection status  
434 differed widely between countries and scenarios (Fig. 4). In Spain and Germany, timber  
435 production and climate regulation multifunctionality were lower inside protected areas, whereas

436 the opposite was observed in France. In Germany, biodiversity conservation-related  
437 multifunctionality was highest inside protected areas, whereas in France the opposite was found.  
438 These results were largely insensitive to the way in which multifunctionality was quantified  
439 (Table S12).

440

## 441 **DISCUSSION**

442 In our study trade-offs between groups of functions were rare in European forests, at both  
443 continental and local scales. We found synergies between individual ecosystem functions and  
444 few trade-offs between multifunctionality measures focused on timber production, climate  
445 regulation and biodiversity conservation/recreation. When corrected for overlap in functions  
446 among scenarios, some relationships were weakly positive throughout most of Europe (timber  
447 production versus climate regulation), some were weakly negative (timber production versus  
448 biodiversity conservation/recreation) and some were close to zero (climate regulation versus  
449 biodiversity conservation/recreation). The lack of strong trade-offs indicates that functions related  
450 to (sustainable) timber production can go hand in hand with functions related to services such as  
451 biodiversity conservation. Mapping local trade-offs and synergies across Europe revealed  
452 substantial variation in these relationships, showing that strong synergies are realized in a few  
453 environments. While biodiversity and timber production are currently maximised in some  
454 forests, suggesting a "win-win" for conservation and commercial forestry, across plots, average  
455 multifunctionality values were almost 50% below maximum possible levels, and the proportion  
456 of forest plots providing high levels of 'overall multifunctionality' (where timber production,  
457 climate regulation and biodiversity conservation are all maximized) was very small. Hence,  
458 while forest management has the potential to realize high multifunctionality, this is currently not

459 common. Most multifunctionality measures had many ecosystem functions in common, as some  
460 ecosystem functions are valued under a range of different management objectives (e.g. Chan *et*  
461 *al.* 2006; Allan *et al.* 2015). Relationships between different multifunctionality measures were  
462 generally much more strongly positive if not corrected for this functional overlap. While these  
463 raw correlations are statistically spurious (as the different measures partly contain the same data),  
464 they can be highly relevant for management. For instance, tree growth is important for both  
465 timber production and climate regulation, which suggests that forest management promoting tree  
466 growth will maximize both services. Our results therefore suggest many possibilities for win-win  
467 forest management strategies.

468 Our multifunctionality variables were intended to represent the bundle of functions  
469 needed to meet certain forest management objectives (following Allan *et al.* 2015). They should  
470 therefore be more useful to managers than traditional multifunctionality metrics that assume  
471 equal importance of each ecosystem function. However, they could be further improved to  
472 consider how multiple functions are related to final ecosystem services, using production  
473 functions, and then services can be valued in monetary or other units to calculate the overall  
474 benefits supplied by different management scenarios (e.g. Nelson *et al.* 2009; Bateman *et al.*  
475 2013). Ultimately, sustainable ecosystem management needs to minimize trade-offs between  
476 ecosystem benefits for different stakeholders (Díaz *et al.* 2015) and our targeted  
477 multifunctionality metrics represent a step towards quantifying and mapping these trade-offs at  
478 large scales.

479 Other studies, performed in grasslands (e.g. Lavorel *et al.* 2011) or across different  
480 ecosystems or land-use types (Chan *et al.* 2006) have documented strong trade-offs between  
481 ecosystem functions and services, especially between productivity-related functions and those

482 associated with biodiversity conservation or recreation. However, in forests, relationships  
483 between tree biomass and the biodiversity of associated taxa often show more mixed patterns  
484 (Jukes *et al.* 2007). For example, the positive relationship between tree productivity and bird  
485 diversity in our data could be due to the strong dependence of specialist species on forests with  
486 many old trees (Gil-Tena *et al.* 2007), while the trade-off between productivity and understorey  
487 biomass may be driven by light competition between trees and understorey plants. When  
488 biodiversity conservation multifunctionality was quantified using only the four direct measures  
489 of biodiversity, weakly negative relationships with timber production and climate regulation  
490 multifunctionality were found. Their approximately equal strength at continental and local scales  
491 (Table 1) suggests that the relationship was primarily driven by local-scale factors, such as stand  
492 composition. The negative response of understorey plants to tree growth is likely responsible for  
493 this trade-off, as it is difficult to maximize timber production whilst maintaining an open canopy.

494 We also found that protected forests were not necessarily associated with high local-scale  
495 ecosystem multifunctionality. In Spain, several multifunctionality measures were in fact lower  
496 inside protected areas. In other countries, patterns were more mixed, but overall  
497 multifunctionality was never highest inside protected areas. Importantly, associations between  
498 forest protection status and multifunctionality were unlikely to be driven by climate, as local-  
499 scale climatic variation is low within our 20×20km regions. Associations between local-scale  
500 multifunctionality and protection status seem therefore to be driven by local factors, such as tree  
501 diversity or composition. However, it is uncertain whether these observed relationships are  
502 causal, as forests were likely not designated to be protected at random. For example, they may  
503 have had low productivity and particular tree compositions before they were protected.  
504 Furthermore, services such as the conservation of forest specialist species were not quantified,

505 but these could be high inside protected areas. Many protected areas were only established  
506 relatively recently (Paillet *et al.* 2015), so protected forests may still be recovering from past  
507 management. Finally, we only investigated forests without evidence of recent logging activity,  
508 which may have reduced the contrast between protected and non-protected areas. Regardless,  
509 although our results suggest a high potential for win-win forest management scenarios, the  
510 simultaneous maximization of timber production, climate regulation and biodiversity has not yet  
511 been realized within protected areas.

512 Our results also provide evidence that climate drives large-scale variation in many  
513 ecosystem functions and the synergies between them. Many functions, such as tree biomass or  
514 litter production, had highest levels in central Western Europe (Fig. S6) and some synergies  
515 between multifunctionality scenarios were stronger at continental than at local scales. A strong  
516 continental-scale synergy between earthworm biomass and litter decomposition (Table S7) may  
517 have arisen because they were both strongly associated with climate (Table S4). The correlation  
518 was also present at the local scale (Table S8), suggesting additional direct links between them.  
519 While earlier studies have already shown the importance of climate for functions such as primary  
520 production and carbon sequestration (e.g. Cramer *et al.* 2001), our more comprehensive study  
521 shows that climate may be a driver of many more ecosystem functions, such as earthworm or  
522 microbial biomass. The fact that so many functions appear related to climate, especially to wet  
523 season precipitation (Table S4), may have important implications. For example, timber  
524 production multifunctionality was lower in dry climates, suggesting detrimental effects of  
525 projected future decreases in precipitation (IPCC 2014). However, while our approach is  
526 powerful in describing patterns, it is not suited to identify underlying processes. Therefore, more



527 research on the causality of climate-ecosystem functioning relationships (e.g. De Boeck *et al.*  
528 2008; Šímová & Storch 2017) is needed to predict ecosystem responses to climate change.

529 Extrapolations are still relatively rare in ecosystem functioning studies (but see Lee *et al.*  
530 2000; Isbell *et al.* 2014; Manning *et al.* 2015), although other subtopics of ecology, such as  
531 species distribution research (Elith & Leathwick 2009), have a much stronger tradition in this  
532 respect. Three ecosystem functions could be validated with independent observations, which  
533 showed that: (1) validations were generally adequate for timber production and tree biomass, but  
534 not for tree recruitment, (2) validations worked best at large spatial scales, whereas at local  
535 scales there was large variation in their accuracy but (3) relationships between different  
536 multifunctionality variables were insensitive to the inclusion of localities where the validation  
537 was less well supported. Our approach is therefore promising, but we emphasize that validations  
538 could only be carried out for those three ecosystem functions for which independent inventory  
539 data was available, so future validations of other functions are needed. Local-scale data related to  
540 soil fertility or management could thus further improve the accuracy of ecosystem function  
541 predictions.

542 Our study presents a new approach to quantify ecosystem functioning at scales relevant  
543 for policy makers. The increasing availability of large datasets on ecosystem functioning from  
544 integrated projects means our approach may become increasingly feasible for other systems and  
545 regions. A further possibility would be to combine local-scale ecosystem functioning datasets  
546 with remote sensing data to map services at large scales. Remote sensing approaches have  
547 successfully predicted some ecosystem functions, but have difficulties with other functions, such  
548 as soil processes (de Araujo *et al.* 2015). By combining data on forest and climate attributes with  
549 remotely sensed parameters, we could map ecosystem functions even more accurately in the

550 future. Our study is a first step in reaching the ultimate goal of predicting how future ecosystem  
551 functioning and service provision will be altered by ongoing global trends, such as climate  
552 change (IPCC 2014), eutrophication and acidification (Galloway *et al.* 2008) or land-use change  
553 (Newbold *et al.* 2015). Future studies could combine our approach with models on climate  
554 change (e.g. IPCC scenarios), biodiversity change (e.g. Isbell *et al.* 2014) or management  
555 scenarios to investigate the impacts of these global trends for the future functioning and service  
556 provisioning of forests and other ecosystems.

557         In conclusion, our study, among most comprehensive overviews of forest ecosystem  
558 functioning to date, showed that different measures of forest multifunctionality tend not to trade-  
559 off with each other, at both local and continental scales. Within some areas there were strong  
560 synergies between different multifunctionality measures, indicating that even though they are  
561 currently uncommon, "win-win" forest management strategies are possible and could be  
562 promoted in the future. However, we also found that multifunctionality is often not higher inside  
563 than outside protected areas. Our study therefore suggests a high but unrealized potential for  
564 multifunctionality in European forests.

565

## 566 **ACKNOWLEDGEMENTS**

567         This paper is a joint effort of the working group ‘Scaling biodiversity-ecosystem  
568 functioning relations: a synthesis based on the *FunDivEUROPE* research platforms’ on the 24<sup>th</sup>-  
569 26<sup>th</sup> November 2014 in Leipzig, Germany, kindly supported by sDiv, the Synthesis Centre of the  
570 German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, funded by the  
571 German Research Foundation (FZT 118). The FunDivEUROPE project received funding from  
572 the European Union’s Seventh Programme (FP7/2007–2013) under grant agreement No. 265171.

573 We thank the MAGRAMA for access to the Spanish Forest Inventory, the Johann Heinrich von  
574 Thünen-Institut for access to the German National Forest Inventories, the Natural Resources  
575 Institute Finland (LUKE) for making the Finnish NFI data available, the Swedish University of  
576 Agricultural Sciences for making the Swedish NFI data available, and Hugues Lecomte, from the  
577 Walloon Forest Inventory, for access to the Walloon NFI data. The study was supported by the  
578 TRY initiative on plant traits (<http://www.trydb.org>). The TRY initiative and database is hosted,  
579 developed and maintained at the Max Planck Institute for Biogeochemistry, Jena, Germany.  
580 TRY is/has been supported by DIVERSITAS, IGBP, the Global Land Project, the UK Natural  
581 Environment Research Council (NERC) through its program QUEST (Quantifying and  
582 Understanding the Earth System), the French Foundation for Biodiversity Research (FRB), and  
583 GIS "Climat, Environnement et Société" France.

584

## 585 REFERENCES

- 586 1. Baeten, L., Verheyen, K., Wirth, C., Bruelheide, H., Bussotti, F., Finér, L. *et al.* (2013). A  
587 novel comparative research platform designed to determine the functional significance of tree  
588 species diversity in European forests. *Perspect. Plant Ecol. Evol. Syst.*, 15, 281-291.
- 589 2. Bateman, I.J., Harwood, A.R., Mace, G.M., Watson, R.T., Abson, D.J., Ndrews, B. *et al.*  
590 (2013). Bringing economic services into economic decision-making: land-use in the United  
591 Kingdom. *Science*, 341, 45-50.
- 592 3. Botta-Dukát, Z. (2005). Rao's quadratic entropy as a measure of functional diversity based on  
593 multiple traits. *J. Veg. Sci.*, 16, 533-540.
- 594 4. Breiman, L. (2001). Random forests. *Mach. Learn.*, 45, 5-32.
- 595 5. Chan, K.M.A., Shaw, R., Cameron, D.R., Underwood, E.C. & Daily, G.C. (2006).  
596 Conservation planning for ecosystem services. *PLoS Biol.*, 4, 2138-2152.

- 597 6. Cramer, W. *et al.* (2001). Global response of the terrestrial ecosystem structure and function  
598 to CO<sub>2</sub> and climate change: results from six dynamic global vegetation models. *Global*  
599 *Change Biol.*, 7, 357-373.
- 600 7. De Araujo, C.C., Atkinson, P.M. & Deary, J.A. (2015). Remote sensing of ecosystem  
601 services: a systematic review. *Ecol. Indicators* 52, 430-443.
- 602 8. De Boeck, H.J., Lemmens, C.M.H.M., Zavalloni, C., Gielen, B., Mailchair, S., Carnol, M.,  
603 Merckx, R., Van den Berge, J., Geulemans, R. & Nijs, I. (2008). Biomass production in  
604 experimental grasslands of different species richness during three years of climate warming.  
605 *Biogeosciences*, 5, 585-594.
- 606 9. Díaz, S. *et al.* (2004). The plant traits that drive ecosystems: evidence from three continents.  
607 *J. Veg. Sci.*, 15, 295-304.
- 608 10. Díaz, S. *et al.* (2015). The IPBES conceptual framework - connecting nature and people.  
609 *Curr. Opin. Environ. Sustain*, 14, 1-16.
- 610 11. Elith, J. & Leathwick, J.R. (2009). Species distribution models: ecological explanation and  
611 prediction across space and time. *Ann. Rev. Ecol. Evol. Syst.*, 40, 677-697.
- 612 12. Faith, D.P. (1992). Conservation evaluation and phylogenetic diversity. *Biol. Cons.*, 61, 1-10.
- 613 13. FAO. (2015). *Global Forest Resources Assessment 2015—How are the world's forests*  
614 *changing?* (Food and Agriculture Organization of the United Nations).
- 615 14. Foy, C.D. (1992). Soil chemical factors limiting plant root growth. In: Hatfield, J.L. &  
616 Steward, B.A. (eds). *Limitations to plant growth*. Springer-Verlag, New York.
- 617 15. Galloway, J.N., Townsend, A.R., Erisman, J.W., Bekunda, M., Cai, Z.C., Freney, J.R. *et al.*  
618 (2008). Transformation of the nitrogen cycle: Recent trends, questions, and potential  
619 solutions. *Science*, 320, 889-892.

- 620 16. Gamfeldt, L. *et al.* (2013). Higher levels of multiple ecosystem services are found in forests  
621 with more tree species. *Nat. Comm.*, 4, 1340.
- 622 17. Gamfeldt, L., Hillebrand, H. & Jonsson, P.R. (2008). Multiple functions increase the  
623 importance of biodiversity for overall ecosystem functioning. *Ecology*, 89, 1223-1231.
- 624 18. Garnier, E. *et al.* (2004). Plant functional makers capture ecosystem properties during  
625 secondary succession. *Ecology*, 85, 2630-2637.
- 626 19. Gil-Tena, A., Saura, S. & Brotons, L. (2007). Effects of forest composition and structure on  
627 bird species richness in a Mediterranean context: Implications for forest ecosystem  
628 management. *Forest Ecol. Manag.* 242, 470-476.
- 629 20. Hastie, T., Tibshirani, R. & Friedman, J. (2008). The elements of statistical learning (2<sup>nd</sup>  
630 edn.). Springer ISBN 0-387-95285-5.
- 631 21. Hector, A. & Bagchi, R. (2007). Biodiversity and ecosystem multifunctionality. *Nature*, 448,  
632 188-190.
- 633 22. Helmus, M.R., Bland, T.J., Williams, C.K. & Ives, A.R. (2007). Phylogenetic measures of  
634 biodiversity. *Amer. Nat.*, 169, 3.
- 635 23. Hijmans, R.J., Cameron, S.E., Parra, J.L., Jones, P.G. & Jarvis, A. (2005). Very high  
636 resolution interpolated climate surfaces for global land areas. *Int. J. Climatol.*, 25, 1965-1978.
- 637 24. IPCC. (2014). *IPCC fifth assessment report*. Cambridge and New York.
- 638 25. Isbell, F., Gonzalez, A., Loreau, M., Cowles, J., Díaz, S., Hector, A. *et al.* (2017). Linking the  
639 influence and dependence of people on biodiversity across scales. *Nature* 546, 65-72.
- 640 26. Isbell, F., Tilman, D., Polasky, S. & Loreau, M. (2014). The biodiversity-dependent  
641 ecosystem service debt. *Ecol. Lett.*, 18, 119-134.


- 642 27. Jukes, M.R., Ferris, R. & Peace, A.J. (2007). The influence of stand structure and  
643 composition on diversity of canopy Coleoptera in coniferous plantations in Britain. *For. Ecol.*  
644 *Managem.*, 163, 27-41.
- 645 28. Kattge, J. *et al.* (2011). TRY – a global database of plant traits. *Global Change Biol.*, 17,  
646 2905-2935.
- 647 29. Klaus, V.H. *et al.* (2013). Direct and indirect associations between plant species richness and  
648 productivity in grasslands: regional differences preclude simple generalization of  
649 productivity-biodiversity relationships. *Preslia*, 85, 97-112.
- 650 30. Lavorel, S. *et al.* (2011). Using plant functional traits to understand the landscape distribution  
651 of multiple ecosystem services. *J. Ecol.*, 99, 135-147.
- 652 31. Lee, M., Manning, P., Rist, J., Power, S.A. & Marsh, C. (2010). A global comparison of  
653 grassland biomass responses to CO<sub>2</sub> and nitrogen enrichment. *Phil. Trans. R. Soc. B*, 365,  
654 2047-2056.
- 655 32. Lefcheck, J.S. *et al.* (2015). Biodiversity enhances ecosystem multifunctionality across  
656 trophic levels and habitats. *Nat. Comm.*, 6, 6936.
- 657 33. Liang, J. *et al.* (2016). Positive biodiversity-productivity relationships predominant in global  
658 forests. *Science*, 354, aaf8957.
- 659 34. Liaw, A. & Wiener, M. (2012). randomForest: Breiman and Cutler's random forests for  
660 classification and regression. Version 4.6.7. [http://cran.r-](http://cran.r-project.org/web/packages/randomForest/index.html)  
661 [project.org/web/packages/randomForest/index.html](http://cran.r-project.org/web/packages/randomForest/index.html)
- 662 35. Manning, P. *et al.* (2015). Simple measures of climate, soil properties and plant traits predict  
663 nation-scale grassland soil carbon stocks. *J. Appl. Ecol.*, 52, 1188-1196.

- 664 36. Maskell, L.C. *et al.* (2013). Exploring the ecological constraints to multiple ecosystem  
665 service delivery and biodiversity. *J. Appl. Ecol.*, 50, 561-571.
- 666 37. McGuire *et al.* (2001). Carbon balance of the terrestrial biosphere in the twentieth century:  
667 analyses of CO<sub>2</sub>, climate and land use effects with four process-based ecosystem models.  
668 *Global Biochem. Cy.*, 15, 183-206.
- 669 38. Millenium Ecosystem Assessment. (2005). *Ecosystems and human well-being: synthesis*.  
670 Island Press, Washington.
- 671 39. Mouchet *et al.* (2017). Bundles of ecosystem (dis)services and multifunctionality across  
672 Europe landscapes. *Ecol. Ind.*, 73, 23-28.
- 673 40. Naidoo, R. *et al.* (2008). Global mapping of ecosystem services and conservation priorities.  
674 *Proc. Nat. Acad. Sci. USA*, 105, 9495-9500.
- 675 41. Nelson, E., Mendoza, G., Regetz, J., Polasky, S., Tallis, H., Cameron, D.R. *et al.* Monitoring  
676 multiple ecosystem services, biodiversity conservation, commodity production, and tradeoffs  
677 at landscape scales. *Front. Ecol. Environ.*, 7, 4-11.
- 678 42. Newbold, T. *et al.* (2015). Global effects of land use on local terrestrial biodiversity. *Nature*,  
679 520, 45-50.
- 680 43. Paillet, Y. *et al.* (2015). Quantifying the recovery of old-growth attributes in forest reserves: a  
681 first reference for France. *Forest Ecol. Managem.* 346, 51-64.
- 682 44. Panagos, P., Van Liedekerke, M., Jones, A. & Montanarella, L. (2012). European Soil Data  
683 Centre: Response to European policy support and public data requirements. *Land Use Policy*,  
684 29, 329-338.
- 685 45. Prince, S.D. & Goward, S.N. (1995). Global primary production: a remote sensing approach.  
686 *J. Biogeogr.*, 22, 815-835.

- 687 46. R Core Team. (2013). R: A language and environment for statistical computing. - R  
688 Foundation for Statistical Computing.
- 689 47. Ratcliffe, S. *et al.* (2016). Modes of functional diversity control on tree productivity across  
690 the European continent. *Global Ecol. Biogeogr.*, 25, 251-261.
- 691 48. Royal Botanic Gardens Kew. (2015). *Seed Information Database (SID)*. Version 7.1.  
692 Available from: <http://data.kew.org/sid/>
- 693 49. Seidl, R., Schelhaas, M.J. & Lexer, M.J. (2011). Unraveling the drivers of intensifying forest  
694 disturbance regimes in Europe. *Glob. Change Biol.*, 17, 2842-2852.
- 695 50. Šímová, I. & Storch, D. (2017). The enigma of terrestrial primary productivity:  
696 measurements, scales, models and the diversity-productivity relationship. *Ecography*, 40,  
697 239-252.
- 698 51. Strobl, C., Boulesteix, A.L., Zeileis, A. & Hothorn, T. (2007). Bias in random forest variable  
699 importance measures: illustrations, sources and a solution. *BMC Bioinformatics*, 8, 25.
- 700 52. Van der Plas, F. *et al.* (2016). Jack-of-all-trades effects drive biodiversity-ecosystem  
701 multifunctionality relationships in European forests. *Nat. Comm.*, 7, 11109.
- 702 53. Webb, C.O., Ackerly, D.D., McPeck, M.A. & Donoghue, M.J. (2002). Phylogenies and  
703 Community Ecology. *Annu. Rev. Ecol. Syst.*, 33, 475-505.
- 704 54. Westoby, M., Falster, D.S., Moles, A.T., Vesk, P.A. & Wright, I.J. (2002). Plant ecological  
705 strategies: some leading dimensions of variation between species. *Annu. Rev. Ecol. Syst.*, 33,  
706 125-159.
- 707 55. Zanne, A.E. *et al.* (2014). Three keys to the radiation of angiosperms into freezing  
708 environments. *Nature*, 506, 89-92.



709 56. Zavaleta, E.S., Pasari, J.R., Hulvey, K.B. & Tilman, D. (2010). Sustaining multiple  
710 ecosystem functions in grassland communities requires higher biodiversity. *Proc. Nat. Acad.*  
711 *Sci. USA*, 107, 1443-1446.



713

714 Figure 1. A: Ecosystem functions included in this study, with the colours and numbers referring

715 to the bars/circles representing them in B and C. B: Weightings used to produce five ecosystem


716 multifunctionality measures, reflecting different management scenarios. From left to right, the

717 ‘equal-weights’, ‘timber production’, ‘climate regulation’, the ‘broad-sense biodiversity

718 conservation/recreation’ and the ‘strict-sense biodiversity conservation’ measure. In the equal


719 weights measure, all ecosystem functions are valued equally. In other measures, function

720 weightings reflect their importance for the management objective. Note that in the climate  
721 regulation scenario, loadings of the decomposition variables are negative. C: Proportion of  
722 variance of ecosystem functions explained by Random Forests. Letters above the bars indicate  
723 which type of predictor was most important in explaining variation: C = climate-related; I =  
724 functional identity-related; P = pH; A = altitude; D = biodiversity-related; S = stand structure  
725 related. In further analyses, only those functions with  $R^2$  values above 0.2 (dashed horizontal  
726 line) were included.



727

728 Figure 2. While high values of continental-scale multifunctionality (A, C-F) in central Europe  
 729 across a range of scenarios indicate large scale synergies, at local scales (B) high overall  
 730 multifunctionality is realized in only a few sites. Mapped levels of predicted large-scale  
 731 multifunctionality are rescaled as the proportion of functions above a 50% threshold. Green  
 732 values indicate relatively high functioning, while brown values indicate relatively low  
 733 functioning. In A), locations of fitting dataset plot are shown in red. In B, where overall, local-  
 734 scale multifunctionality is shown, the histogram indicates that in only a few plots, levels exceed  
 735 0.8.



736

737

738

739

740


741

742

743

744

Figure 3. Substantial variation in the degree of local scale synergies and trade-offs exists across Europe. Observed and observed minus expected correlation coefficients between multifunctionality measures, within 20×20 km grid cells. Top: Values of all observed multifunctionality measures, except for the narrow-sense biodiversity conservation measure, correlate positively at local scales. Bottom: these correlations are largely driven by overlap in ecosystem functions, as observed minus expected correlation-coefficients are close to zero. Average correlations that deviate significantly from zero are indicated with an asterisk (\*).



745

746 Figure 4. Local-scale ecosystem multifunctionality is generally not higher inside protected areas,

747 for different multifunctionality measures and countries. Bars above zero indicate that

748 multifunctionality is higher inside than outside protected areas, while bars below zero indicate

749 the opposite. A: Equal-weight multifunctionality. B: timber production multifunctionality. C:

750 climate regulation multifunctionality. D: broad-sense biodiversity conservation/recreation

751 multifunctionality. E: narrow-sense biodiversity conservation/recreation multifunctionality.

752

753 Table 1. Correlations between values of different multifunctionality measures at both continental  
754 and local scales and both across all plots and within countries. Here, multifunctionality was  
755 based on a 50% threshold level. Correlations were also quantified after correcting for the overlap  
756 in ecosystem functions between multifunctionality measures. This is indicated as ‘no functional  
757 overlap’ or ‘no FO’ in the table. As sensitivity analyses, correlations were also calculated based  
758 on (a) multifunctionality measures in which recruitment-related functions were excluded, (b)  
759 multifunctionality measures in which loadings of ecosystem functions was proportional to  
760 Random Forest R<sup>2</sup> values and (c) only those plots within 20x20 km grid cells with a high  
761 validation R<sup>2</sup> (>0.10) for timber production and tree biomass. Significant correlations are shown  
762 in bold. TP = timber production, CR = climate regulation, BCB = broad-sense biodiversity  
763 conservation and BCN = narrow-sense biodiversity conservation.

	TP-CR	TP-BCB	TP-BCN	CR-BCB	CR-BCN
Continental scale, raw	<b>0.81</b>	<b>0.57</b>	<b>-0.13</b>	<b>0.63</b>	0.01
Continental scale, no FO	<b>0.06</b>	<b>0.15</b>	<b>-0.13</b>	<b>0.16</b>	0.01
Continental scale, no FO, no recruitment-related EFs	<b>0.07</b>	<b>0.16</b>	<b>-0.09</b>	<b>0.20</b>	<b>0.08</b>
Continental scale, no FO, corrected for EF R <sup>2</sup> values	<b>0.10</b>	<b>0.18</b>	<b>-0.17</b>	<b>0.12</b>	<b>-0.10</b>
Continental scale, no FO, only plots with high validation	<b>0.05</b>	<b>0.12</b>	<b>-0.35</b>	<b>0.11</b>	<b>-0.17</b>
Local scale	<b>0.79</b>	<b>0.31</b>	<b>-0.12</b>	<b>0.44</b>	-0.01
Local scale, Spain only	<b>0.79</b>	<b>0.32</b>	<b>-0.11</b>	<b>0.46</b>	0.02
Local scale, France only	<b>0.80</b>	<b>0.30</b>	<b>-0.12</b>	<b>0.42</b>	<b>-0.03</b>
Local scale, Wallonia only	<b>0.78</b>	<b>0.12</b>	<b>-0.31</b>	<b>0.38</b>	<b>-0.07</b>
Local scale, Germany only	<b>0.80</b>	<b>0.31</b>	<b>-0.16</b>	<b>0.47</b>	-0.01
Local scale, Sweden only	<b>0.73</b>	<b>0.30</b>	-0.03	<b>0.33</b>	-0.03
Local scale, Finland only	<b>0.77</b>	<b>0.34</b>	-0.08	<b>0.44</b>	-0.02
Local scale, no FO	<b>0.01</b>	<b>-0.08</b>	<b>-0.13</b>	<b>0.03</b>	<b>-0.01</b>
Local scale, no FO, Spain only	<b>0.01</b>	<b>-0.08</b>	<b>-0.11</b>	<b>0.05</b>	<b>0.02</b>
Local scale, no FO, France only	<b>0.01</b>	<b>-0.09</b>	<b>-0.13</b>	<b>0.01</b>	<b>-0.03</b>
Local scale, no FO, Wallonia only	0.00	<b>-0.26</b>	<b>-0.31</b>	<b>-0.03</b>	<b>-0.07</b>
Local scale, no FO, Germany only	<b>0.02</b>	<b>-0.08</b>	<b>-0.16</b>	<b>0.06</b>	<b>-0.01</b>
Local scale, no FO, Sweden only	<b>-0.05</b>	<b>-0.09</b>	<b>-0.03</b>	<b>-0.08</b>	<b>-0.04</b>
Local scale, no FO, Finland only	<b>-0.01</b>	<b>-0.05</b>	<b>-0.08</b>	<b>0.03</b>	<b>-0.02</b>
Local scale, no FO, no recruitment-related EFs	<b>0.03</b>	<b>-0.12</b>	<b>-0.14</b>	<b>-0.04</b>	<b>-0.02</b>
Local scale, no FO, corrected for EF R <sup>2</sup> values	<b>0.09</b>	<b>-0.07</b>	<b>-0.17</b>	<b>-0.04</b>	<b>-0.08</b>
Local scale, no FO, only plots with high validation	<b>0.10</b>	<b>-0.15</b>	<b>-0.29</b>	-0.06	-0.13

764