



HAL
open science

In situ measurement of molecule mobility in mucilage polysaccharide gels from different species

A. Berger, Olivier Grandjean, Corinne C. Rondeau-Mouro, H. North

► **To cite this version:**

A. Berger, Olivier Grandjean, Corinne C. Rondeau-Mouro, H. North. In situ measurement of molecule mobility in mucilage polysaccharide gels from different species. 11e journées du Réseau Français des Parois, Jun 2017, Orléans, France. 2017. hal-02606626

HAL Id: hal-02606626

<https://hal.inrae.fr/hal-02606626>

Submitted on 16 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Adeline Berger¹; Olivier Grandjean¹; Corinne Rondeau-Mouro²; Helen North¹

¹INRA, Institut Jean-Pierre Bourgin, INRA, Agro Paris-Tech, CNRS, Université Paris Saclay, 78026 Versailles, France,
²IRSTEA, UR OPAALE, 17 avenue de Cucillé, CS 64427, 35044 Rennes, France

Introduction

On imbibition, the seeds of certain species form a polysaccharide hydrogel, termed mucilage. In *Arabidopsis thaliana* and *Camelina sativa*, this mucilage is composed of 2 layers, the outer being water-soluble while the inner is tightly attached to the seed surface. Determining the macromolecular properties of mucilage polymers in the inner mucilage layer is usually achieved by detaching the polysaccharides from the tegument either by physical or enzymatic methods, which alters or eliminates the structure. In order to characterize these properties, we have developed three non-destructive microscopy techniques. Based on fluorescence excitation or nuclear magnetic resonance, these in situ methods aim at investigating the physicochemical properties of mucilage polysaccharides that could affect accessibility and mobility of molecules within the hydrogel. This work presents results obtained on *Arabidopsis* and *Camelina* seeds comparing their inner mucilage properties.




Figure 1. The organisation of mucilage constituents in *Arabidopsis* (A, B, E and F) and *Camelina* (C, D, G and H) seeds. (A) and (C) showing seed coat features, (B) and (D) after cellulose staining of inner mucilage with 0.01% pontamine fast scarlet 4b in 50mM NaCl. Immunolabelling indicates a difference in the heteromannan localisation (green in pictures (E) and (G), magenta showing cellulose staining). FITC polysucrose 40kDa shows differences in the permeability of the inner mucilage. Scale bar: 100µm (A to D), 50µm (E to H).

Fluorescence Recovery After Photobleaching

Material: Seeds imbibed in water and then stained with fluorescent probes (FITC-polysucrose) of different sizes (20kD, 40kDa and 400kDa).




Figure 2. FRAP acquisition sequence. A Leica SP5 confocal microscope was used to achieve pre-bleach, bleach and post-bleach.




Figure 3. *Camelina* mucilage traps some molecules compared to this of *Arabidopsis*. Results of FRAP data analysis³ after normalization and fitting the raw data using a biexponential algorithm shown in (A) for *Arabidopsis* and (B) for *Camelina*.




Figure 4. Differences in *Arabidopsis* and *Camelina* molecule mobility. (A) represents the mobile fraction of different sizes of fluorescent probe (pink: 20kDa, green: 40kDa and blue: 400kDa) in *Arabidopsis* or *Camelina* mucilage and (B) the t-half.

Conclusions: - FRAP can be to obtain information about molecule mobility; if molecule moves freely or is subject to steric hindrance.
- The physical properties of *Camelina* and *Arabidopsis* inner mucilage differ.

Fluorescence anisotropy

Material: *Arabidopsis* seeds imbibed in water and then stained with fluorescent probes (FITC-polysucrose) of different sizes (20kD, 40kDa and 400kDa).




Figure 5. Anisotropy measurement⁴. Fluorescence emission intensity (A) perpendicular and (B) parallel to excitation plane, is analysed with imageJ to obtain (C) the anisotropy and (D) the maximum intensity threshold of the (C) image.




Figure 6. Anisotropy values obtained with different fluorophore sizes (pink: 20kDa, green: 40kDa and blue: 400kDa).

Conclusions: - Anisotropy measurement of molecule mobility provides information about hydrocolloid viscosity and molecular crowding
- Data acquisition is fast (< 5 min) allowing different samples to be compared in a single experiment
- This technique is independent of fluorophore concentration.

Magnetic resonance micro-imaging

Material: *Camelina* seeds imbibed in water without a probe molecule. Water mobility is measured through spatially localized T2 relaxation time.




Figure 7. T2 maps of *Camelina* seeds imbibed in water. Data was acquired using the MSME sequence (Bruker, 11.7 T) with 128 echos, TE=2.96 ms, TR=1s. Resolution = 55µm²x1mm with a matrix of 256x256 voxels. (A) : distribution of the shorter T2 water component (0-20 ms) within seeds. (B), (C) : distribution of water and fat with T2 between 20-40 ms, surrounding ring with T2 around 60 ms is assigned to water in the inner mucilage layer. T2 of soluble mucilage in the water phase outside seeds exceed 120-140 ms. Scale bar: 1mm.

Conclusions: - MRµl allows quantification of water and fat in or outside seeds of diameter > 1mm².
- Rapid image acquisition (15 min) enable studies of the rate and evolution over time of water distribution and mobility in the inner mucilage layer.
- The first MRµ-images of *Camelina* indicate that measure of the water mobility and its change with time is possible with a resolution of 55 µm² radial slice.

Perspectives

Three complementary, non-destructive *in situ* methods have been developed which provide information about hydrocolloid organization and the rheological properties of seed mucilage. Each of them is rapid enough for the analysis of several different seeds in a single experiment.

Contact

Adeline Berger
Jean-Pierre Bourgin institute
INRA Versailles - Grignon
@ : adeline.berger@inra.fr
<http://www-ijpb.verailles.inra.fr/en/index.htm>
https://www6.inra.fr/observatoire-vegetal_eng/
☎ (33) (0) 1 30 83 37 07



References

- Voiniciuc C., Schmidt M.H., Berger A., Yang B., Ebert B., Scheller H.V., North H.M., Usadel B., Günl M. (2015) MUCILAGE-RELATED10 Produces Galactoglucomannan That Maintains Pectin and Cellulose Architecture in *Arabidopsis* Seed Mucilage. *Plant Physiol.* 169(1):403-20.
- Saez-Aguayo S., Rondeau-Mouro C., Macquet A., Kronholm I., Ralet M.-C., Berger A., Sallé C., Poulain D., Granier F., Botran L., Loudet O., de Meaux J., Marion-Poll A., North H.M. (2014) Local Evolution of Seed Flotation in *Arabidopsis*. *PLoS Genet.* 10(3): e1004221.
- Rapsomaniki M.A., Kotsantzi P., Symeonidou I.E., Giakoumakis N.N., Taraviras S and Lygerou Z (2012). easyFRAP: an interactive, easy-to-use tool for qualitative and quantitative analysis of Fluorescence Recovery After Photobleaching data. *Bioinformatics.* 28(13):1800-1801
- Marquer C., Fruchart-Gaillard C., Mourier G., Grandjean O., Girard E., le Maire M., Brown S., Servent D. (2010) Influence of MT7 toxin on the oligomerization state of the M1 muscarinic receptor. *Biol Cell.* 102(7):409-20.