



**HAL**  
open science

## Ecologically relevant flow metrics for intermittent rivers and ephemeral streams

G. Laaha, T. Gauster, Eric Sauquet, C. Sefton, R. Stubbington, S. M. Parry

► **To cite this version:**

G. Laaha, T. Gauster, Eric Sauquet, C. Sefton, R. Stubbington, et al.. Ecologically relevant flow metrics for intermittent rivers and ephemeral streams. EGU, Apr 2018, Vienna, Austria. pp.1, 2018. hal-02607367

**HAL Id: hal-02607367**

**<https://hal.inrae.fr/hal-02607367>**

Submitted on 16 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Ecologically relevant flow metrics for intermittent rivers and ephemeral streams



Gregor Laaha<sup>1</sup>, Tobias Gauster<sup>1</sup>, Eric Sauquet<sup>2</sup>, Cath Senfton<sup>3</sup>, Rachel Stubbington<sup>4</sup>, Simon Parry<sup>3</sup>, and the COST-Action SMIREs author team



1) University of Natural Resources and Life Sciences Vienna (BOKU), Austria; 2) IRSTEA Lyon, France; 3) CEH Wallingford, UK; 4) Nottingham Trent University, UK

## Background


The COST-Action SMIREs brings together scientists from various disciplines to foster a common understanding of hydrology, hydrochemistry and ecology of intermittent rivers and ephemeral streams. Such rivers (IRES) are characterized by episodes of stagnant waters or drying-off, occurring either annually during the drought season or exceptionally during extreme drought years. Since IRES can be seen as a special, ultimate, case of low-flow regimes when discharge reaches zero-values, relevant low-flow characteristics exist but need to be adapted to characterise ecologically relevant features of the particular flow regime.

## Objectives

We present a set of metrics and methods for characterizing flow intermittency in an ecologically relevant way, stemming from interdisciplinary discussions among SMIREs experts of various disciplines. The indices are compatible with standard low-flow indices of the WMO manual on low-flow estimation and prediction (Gustard and Demuth, 2008) and the handbook Hydrological Drought (Tallaksen and Van Lanen, 2004), extending these indices to no-flow conditions.

## 1. General definitions

- All indices derived from daily discharge hydrographs  $Q$
- No-flow spell: period of time when discharge is less or equal some no-flow threshold (= detection limit):  $Q \leq Q_0$  (default 1 l/s)
- No-flow event: a contiguous period of days with no-flow occurrence, terminated by rewetting period. Some pooling is usually applied to successive, dependent events in order to define an independent sequence of events.
- All annual characteristic may be calculated based on calendar years or hydrological years (e.g. for European climates usually starting with 1 April)
- All annual characteristics can be calculated from the whole time series (all-season characteristics) or stratified by seasons (seasonal characteristics)



## 2. No-flow characteristics

### 2.1 Proportion of no-flow years

Name	Measure	Acronym	Function	Description
Proportion of no-flow years	$f_0$	$no\_flow\_years(x)$		Number of years with no-flow occurrence (when flow is at least for one day below the no-flow threshold) divided by the study period in years

### 2.2 Number of no-flow days

Name	Measure	Acronym	Function	Description
Mean annual number of no-flow days	Location	$MAN$	$MAN(x)$	Mean annual number of days with no-flow occurrence (when flow is below the no-flow threshold)
CV of the annual number of no-flow days	Variability	$CVAN$	$CVAN(x)$	Coefficient of variation of the annual number of days with no-flow occurrence (when flow is below the no-flow threshold)
Distribution of the annual number of no-flow days	Distribution	$FAN$	$FAN(x)$	Statistical distribution (cdf) of the annual number of days with no-flow occurrence (when flow is below the no-flow threshold)

### 2.3 Duration

Name	Measure	Acronym	Function	Description
Mean annual maximum duration	Location	$MAMD$	$MAMD(x)$	Mean duration of the longest annual no-flow event (during which flow remains below the no-flow threshold)
CV of annual maximum duration	Variability	$CVAMD$	$CVAMD(x)$	Coefficient of variation of the duration of the longest annual no-flow event (during which flow remains below the no-flow threshold)
Distribution of annual maximum duration	Distribution	$FAMD$	$FAMD(x)$	Statistical distribution (cdf) of the duration of the longest annual no-flow event (during which flow remains below the no-flow threshold)

### 2.4 Timing and seasonality

Name	Measure	Acronym	Function	Description
Mean onset	Location	$\tau_0$	$\tau_{au0}(x)$	Mean Julian date (day-of-year) of the first annual no-flow day (using circular statistics)
Variability of onset	Variability	$\tau_{0v}$	$\tau_{au0r}(x)$	Circular variability index $r$ (between 0 and 1) of the onset
Mean termination	Location	$\tau_E$	$\tau_{auE}(x)$	Mean Julian date (day-of-year) of the end of the first no-flow spell (using circular statistics)
Variability of termination	Variability	$\tau_{Ev}$	$\tau_{auEr}(x)$	Circular variability index $r$ (between 0 and 1) of the termination
Mean seasonality	Location	$\tau$	$\tau_{au}(x)$	Mean Julian date of all no-flow days (using circular statistics)
Strength of seasonality	Variability	$\tau_r$	$\tau_{aur}(x)$	Circular variability index $r$ (between 0 and 1) of all no-flow days

## 3. Rate of change before/after no-flow event

Name	Measure	Acronym	Function	Description
Mean recession rate	Location	$k$	$k(x)$	Mean rate of decay of the hydrograph during flow recession periods
Standard deviation of recession rate	Variability	$k_{std}$	$k_{std}(x)$	Standard deviation of rate of decay of the hydrograph during flow recession periods
Mean annual number of reversals in flow magnitude	Location	$nrv$	$nrv(x)$	Mean annual number of reversals in flow magnitude


Similar indices for recovery rates are needed, but would require a clear definition of the termination of the recovery period from drought. We here suggest to define the recovery period consistently with process-based pooling criteria (e.g. used in Laaha et al. 2017): events are pooled unless the day when the catchment has completely recovered from a negative water balance, marking the end of recovery period.

## 4. Low-flow frequency analysis

Annual minima series of IRES contain zero values, hence a conditional probability model is required. First, a distribution model  $G(x)$  is fitted to the censored series (i.e., the annual minima for all years except for no-flow years). In a second step, the distribution  $F(x)$  of the entire annual minima series containing no-flow years is obtain by

$$F(x) = f_0 + (1 - f_0)G(x),$$

using  $f_0$  (section 2.1) as an estimator of the annual probability that the river falls dry. The figure shows an example of a river with 30% no-flow years.



## Conclusions

R-Software package *smires* (Gauster et al. 2017) has been compiled to foster the use of statistically sound no-flow indices that are consistent with earlier definitions in low-flow hydrology. In addition, more qualitative characteristics about flow states are needed, which are subject of ongoing SMIREs research.