



HAL
open science

Copper bioaccumulation and internalization during freshwater biofilm development using stable Cu isotope experiment

Aymeric Dabrin, J. Gahou, M. Masson, Christopher Brosse, B. Volat, C. Bonnineau, Stéphane Pesce, Marina Coquery

► To cite this version:

Aymeric Dabrin, J. Gahou, M. Masson, Christopher Brosse, B. Volat, et al.. Copper bioaccumulation and internalization during freshwater biofilm development using stable Cu isotope experiment. SETAC Europe 28th annual meeting, May 2018, Rome, Italy. pp.1, 2018. hal-02607501

HAL Id: hal-02607501

<https://hal.inrae.fr/hal-02607501>

Submitted on 16 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Following copper bioaccumulation and internalization during freshwater biofilm development using stable Cu isotope

Dabrin*, A., Gahou, J., Masson, M., Brosse, C., Volat, B., Bonnineau, C., Pesce, S., Coquery, M.

Irstea, UR RIVERLY, centre de Lyon-Villeurbanne, 5 Rue de la Doua, CS 20244, F-69625 Villeurbanne Cedex, France.

* aymeric.dabrin@irstea.fr

Context and aim of the study


In small streams, microbial communities form river biofilms attached to solid substrates by producing extracellular polymeric substances (EPS). This matrix may act as a protective layer by limiting cellular contact with surface water contaminants. Thus, several studies (i.e. Ivorra et al. 2000) have suggested that during biofilm growth, biofilm and EPS matrix thickness could limit cellular bioaccumulation.

Aim of the study


To test the hypothesis that biofilm is more exposed to Cu in early colonization stage than a mature biofilm, we conducted an experiment under controlled conditions to follow the bioaccumulation of two Cu isotopes in the different biofilm fractions throughout biofilm growth and maturation. After a first period of development (0 to 20 days) in natural dissolved copper medium, biofilm was transferred to a mono-isotopic (^{65}Cu) copper-enriched medium for 20 additional days. After 20 and 40 days, a sequential extraction was applied to recover Cu from the intracellular fraction, and from the colloidal and capsular EPS fractions. Copper concentrations and isotopic ratios were determined by ICP-MS in water collected at various times of the experiment and after 20 and 40 days in the different fractions of the biofilm.

Material and methods

Experimental procedure



Sequential extraction of EPS on biofilms



Cu analyses

- Analysis of ^{63}Cu and ^{65}Cu by ICP-MS with CCT (Thermo X7 Series II)
- Mass bias correction for isotopic ratio measurement ($^{64}\text{Zn}/^{66}\text{Zn}$)
- External calibration for Cu concentrations measurements
- LQ = 0.05 µg/L for water samples
- LQ = 2 µg/g d.w. for biofilms samples




EPS Characterization

- Polysacc
- Prot
- HPSEC


Biofilm growth

XXX

Cu in dissolved medium




Biofilm growth




EPS characterization

Polysaccharids




Proteins




- Polysaccharids and proteins concentrations are 2.7 and 2.0 times higher in colloidal fraction than in capsular fraction.
- No change of polysaccharids and proteins concentrations between the early stage of biofilm development and the second phase of growing (AQ 1 and 2).
- Decrease of polysaccharids and proteins concentrations in the third aquarium.

Bioaccumulation of Cu in biofilm


1 Total concentration of Cu in biofilm



2 Cu concentrations in the fractions of the biofilm



3 Isotopic ratios of $^{63}\text{Cu}/^{65}\text{Cu}$ in the fractions of the biofilm



Conclusions

- According to analytical procedure (total Cu concentrations, Cu concentrations in the different fraction of the biofilm or isotopic tracing), informations on Cu accumulation kinetics can be very different.

- While a total Cu concentration approach seems to indicate a Cu saturation over time, isotopic tracing approach suggests that the entire biofilm (EPS + cells) is continuously in intense renewal as well as the bioaccumulated Cu.

- In aquarium n°3, the increase of dissolved Cu during the second period of exposure has led to a less Cu accumulation. In that case, we noticed an isotopic ratio slightly higher in cells than in EPS fractions, suggesting a potential protective effect of EPS matrix

Perspectives

- Refine kinetics of accumulation during the second period of exposure (biofilm extraction every day to follow isotopic ratios in the different fractions)
- Coupling tracing approaches with microscopic/analysis approaches (LA-ICP-MS, µ-SXRF)

