



Interaction between climate and management on beta diversity components of vegetation in relation to soil properties in arid and semi-arid oak forests, Iran

M. Heydari, F. Aazami, M. Faramarzi, R. Omidipour, M. Bazgir, D. Pothier, Bernard Prévosto

► To cite this version:

M. Heydari, F. Aazami, M. Faramarzi, R. Omidipour, M. Bazgir, et al.. Interaction between climate and management on beta diversity components of vegetation in relation to soil properties in arid and semi-arid oak forests, Iran. *Journal of Arid Land*, 2019, 11 (1), pp.43-57. 10.1007/s40333-018-0024-z . hal-02607590

HAL Id: hal-02607590

<https://hal.inrae.fr/hal-02607590>

Submitted on 16 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Interaction between climate and management on beta diversity components of vegetation in relation to soil properties in arid and semi-arid oak forests

Mehdi Heydari, Fatemeh Aazami, Marzban Faramarzi, Reza Omidipour, Masoud Bazgir, David Pothier Bernard Prévosto

Abstract

This study aimed to investigate the interaction between regions with different climatic conditions (arid vs. semi-arid) and management (protected vs. unprotected) on the turnover and nestedness of vegetation in relation to physical, chemical and biological properties of soils in the Ilam province of Iran. In each of the two regions, we sampled eight sites (4 managed and 4 unmanaged sites) within each of which we established four circular plots (1000 m²) that were used to inventory woody species, while two micro-plots (1×1 m²) were established in each 1000-m² plot to inventory herbaceous species. In each sample unit, we also extracted three soil samples (0–20 cm depth) for measuring soil properties. The results indicated that the interaction between region and conservation management significantly affected the percent canopy cover of Persian oak (*Quercus brantii* Linddl), soil respiration, substrate-induced respiration, as well as beta and gamma diversities and turnover. The percent canopy cover of oak was positively correlated with soil silt, electrical conductivity, available potassium, and alpha diversity, whereas it was negatively correlated with plant turnover. In addition, plant turnover was positively related to available phosphorus while nestedness was positively related to organic carbon and total N. According to these results, we conclude that physical, chemical and biological characteristics of limited ecological niche generally influence plant diversity. Also, this study has indicated the major contribution of the β -diversity on γ -diversity, especially in the semi-arid region, because of a higher heterogeneity in this area.

Keywords: climatic conditions; conservation; Beta diversity; oak forests

1. Introduction

32 Forest ecosystems have a critical role to mitigate soil loss and erosion, regulate weather and
33 maintain habitats and biodiversity (Fathizadeh et al., 2017). Forest degradation and/or destruction had
34 many negative effects on plant and soil microorganism diversity and composition (Yacht et al., 2017;
35 Moradi Behbahani et al., 2017) and soil properties (Feng et al., 2017). Conservation management can
36 be a key factor in preventing worldwide destruction of forest ecosystems and promoting the
37 sustainability of these valuable resources.

38 Biodiversity indices are generally used to assess and evaluate ecosystem degradation (Wilson and
39 Tilman, 2002). However, with the increasing importance of biodiversity conservation, the use of new
40 and effective methods to investigate plant biodiversity has become increasingly popular. Diversity
41 partitioning (additive vs. multiplicative) is one of these methods that was frequently used during the
42 two recent decades by ecologists devoted to biodiversity conservation (Erfanzadeh et al., 2015).
43 According to this method, total regional diversity can be partitioned additively ($\gamma = \alpha +$
44 β ; Lande, 1996) or multiplicatively ($\gamma = \alpha \times \beta$) into within (α diversity) and
45 among (β diversity) components across different spatial/temporal scales (Crist et al., 2003). The
46 additive partitioning method has been increasingly used in rangeland and forest studies in relation to
47 plant species diversity (i.e. Chávez and Macdonald, 2012; Zhang, et al., 2014; Erfanzadeh et al.,
48 2015; Schulze, et al., 2016).

49 Beta diversity is an important biodiversity index that is defined as the variation in community
50 composition from one place to another. Baselga (2010) demonstrated mathematically that beta
51 diversity can be decomposed in two main processes: turnover and nestedness. Spatial turnover is
52 defined as the continuous or sudden replacement of species along an environmental gradient due to
53 environmental pressure, climate change or competition (Baselga, 2010; Lafage et al., 2015). For its
54 part, nestedness refers to communities in which species composing a low-diversity community are a
55 subset of a high-diversity community that was subjected to species migration or extinction (Calderón-
56 Patrón et al., 2013). By analyzing beta diversity components and their respective contribution to the
57 total beta diversity, we can detect the main mechanism involved in changes in community
58 composition.

59 Semi-arid forests of the Zagros Mountains are dominated by Persian oak (*Quercus brantii* Linddl)
60 and cover an area of about 5 million ha (40% of Iranian forests; Sagheb-Talebi et al., 2014). They
61 form one of the most important ecosystems of Iran due to their high level of biodiversity and the
62 presence of endemic species (Marvi Mohadjer, 2005). The Zagros region has a great impact on
63 people's livelihood due to the dependency of people on local resources (Salehi et al., 2013).

64 Unfortunately, during the three last decades, the vegetation of the Zagros forests was highly degraded
65 due to incorrect management practices and human activities such as human-caused fire, over-grazing,
66 excessive cutting, firewood harvesting and land use changes (Heydari et al., 2016).

67 While forest degradation could create important changes in vegetation composition and forest
68 structure, forest soils can also be impacted. Soils are an important constituent of ecosystems and play
69 a critical role in the development of forest vegetation, which, in turn, have noticeable effects on the
70 development of physical, chemical and biological soil properties (Kooch et al., 2007; Onyekwelu et
71 al., 2006; Wang, 2007).

72 Biodiversity conservation in Zagros forests should take advantage of new management practices to
73 help reduce the progression of its degradation caused by drought-induced oak mortality. Conservation
74 plan including exclosures is one management solution that was effective in Zagros forests (Heydari et
75 al., 2013a). Tárrega et al. (2009) reported that different management regimes in dehesa ecosystems
76 (grazed, ungrazed, ungrazed with shrub cutting) had a critical effect on soil properties and vegetation
77 in oak (*Q. pyrenaica*) stands, and that conservational practices led to improvement in soil fertility.
78 This is in accordance with results indicating that conservational practices (grazed vs. no grazed) had
79 positive effects on soil properties and vegetation (Strandberg et al., 2005). In addition, Sheklabadi et
80 al. (2007) observed that in areas degraded by grazing, the amount of organic carbon, total nitrogen
81 and microbial respiration were lower than in non-degraded areas.

82 Climate is another factor that has profound effects on vegetation diversity and composition as well
83 as on soil properties (Kardol et al., 2010). Climatic factors, particularly temperature and
84 precipitation, affect the formation and evolution of soils (Binkley and Fisher, 2012), the
85 accumulation of organic carbon (Wang et al., 2013), the cycling rate of nutrients (Auyeung et al.,
86 2013) and the soil biological activity (Sardans and Peñuelas, 2005; Steinweg et al., 2013).

87 Due to the interaction between [soil characteristics, climatic conditions and plant species](#) (von
88 [Lutzow & Kogel-Knabner, 2009](#); [Mirzaei and Moradi, 2017](#); Toure et al., 2015), changes in plant
89 species diversity and composition between regions with different climatic conditions are expected.
90 On the other hand, plant communities and their characteristics such as life form spectra (Raunkiaer,
91 1934) gradually reach an equilibrium with climatic conditions to form so-called climatic climax
92 communities.

93 The effects of forest degradation and management strategies on plant diversity and soil attributes
94 have been recently investigated in the Zagros forest of Iran (Parma & Shataee Jouybari, 2010;

Heydari et al., 2017). However, no study has addressed the effects of climate with or without interaction with management practices on diversity components (alpha and beta diversity) as well as on beta diversity components (turnover and nestedness). This study aims to fill this gap by evaluating the effects of climate, management strategies and their interaction on soil physical, chemical and biological properties as well as on biodiversity components.

100

101


102 2. Material and methods

103 2.1. Study area

104 This study took place in managed and unmanaged (see details below) forests dominated by Persian oak (*Quercus brantii* Linddl) in western Iran. These forests were located in two regions (Fig. 1) with 105 different climatic conditions. According to the classification system of de Martonne (1925), the 106 climate of these regions are both Mediterranean, but the semi-arid Dalab region is characterized by 107 annual precipitation of 590 mm and mean annual temperature of 16.7 °C, while the mean annual 108 precipitation and temperature of the arid Koshk region are 408 mm and 18.5 °C, respectively. These 109 regions are both part of the Zagros mountains, and the geology and the pedogenesis of the two 110 regions are similar. Calcareous soils with a clay loamy texture, dominate the soils of these regions.

112 In each region, we selected four managed and four unmanaged sites that were described by 113 Heydari et al. (2013 b). In managed sites, a 20-year period of strict conservation was applied 114 following long-term human disturbances. For their part, unmanaged sites were never protected 115 against long-term anthropogenic land degradation such as tree and shrub cutting for firewood and 116 development of arable lands as well as animal husbandry (Heydari et al., 2013 b). The vegetation of 117 the managed sites of Dalab (MD) was dominated by *Cerasus microcarpa*, *Daphne mucronata*, 118 *Acantholimon bromoifolium*, *Teucrium polium*, *Alyssum marginatum*, *Centaurea depressa*, *Centaurea* 119 *amadanensis*, *Centaurea koeieana*. In the unmanaged sites of Dalab (UMD), the dominant 120 vegetation was composed of *Capparis parviflora*, *Astragalus fasciculifolius*, *Astragalus* 121 (*Leucocercis*) *curviflorus*, *Bromus tomentellus*, *Gundelia Tournefortii*, *Erodium cicutarium*, 122 *Stellaria media*, *Valerianella dactylophylla*. In the managed sites of Koshk (MK), the dominant 123 species were *Pistacia khinjuk*, *Amygdalus lycioides*, *Populus euphratica*, *Medicago rigidula*, *Sinapis* 124 *arvensis*, *Trifolium purpureum*, *Ziziphora capitata*, *Anthemis pseudocotula*. Finally, in the 125 unmanaged sites of Koshk (UMK), the vegetation was dominated by *Nerium oleander*, *Vitex*

126 *pseudo-Negundo Medicago rigidula*, *Sinapis arvensis*, *Phlomis olivieriv*, *Crepis katschyana*,
 127 *Scabiosa leucactic* (Aazami, 2016).
 128



129
 130 Fig. 1 Study area location and sampling method; ■ : managed and ▲ : unmanaged sites;
 131 ○ : Marco-plot and □ : Micro-plot
 132

133 2.2. Experimental design and data collection

134 The combination of our two regions (D=Dalab, K=Koshk) with the two types of management
 135 (M=managed, UM=unmanaged) results in four treatments (MD, UMD, MK, UMK) and with
 136 different climatic conditions (semi-arid Dalab region vs. arid Koshk region). In each treatment, we
 137 selected four independent sites that were considered as replicates. However, to decrease variability,
 138 geographically adjacent sites were selected in each region. In the center of each site, we used a
 139 systematic sampling method to establish a cluster of four circular, 1000-m² plots located 100 m
 140 apart (Fig. 1). These plots were used to estimate the percent cover of all woody species while two
 141 squared 1-m² microplots, located within each main plot, were randomly established to inventory
 142 herbaceous species (each species was given a value of abundance) in May 2015. In total, we have 16
 143 sampling sites (8 in Koshk and 8 in Dalab), 64 macroplots (1000-m²) and 128 microplots.

144

145 2.3. Soil sampling and laboratory methods

146 We extracted soil samples at 0-20 cm depth in May 2015. Physical, chemical and biological soil
147 properties were determined from composite samples, each consisting of three sub-samples that were
148 gathered from three randomly selected points in each main plot (for a total of 64 composite soil
149 samples). Soils were sieved (2-mm mesh) to remove roots and debris prior to laboratory analyses
150 and divided into two subsamples.

151 The first subsample was air-dried for measuring physico-chemical properties. Soil texture was
152 measured using a hydrometer (Bouyoucos, 1962). Soil organic carbon (OC) was determined by
153 dichromate oxidation using the Walkley-Black method (Nelson and Sommers, 1982). Soil pH and
154 electrical conductivity (EC) were determined electrometrically (in H₂O, 2:1 v/m) with a
155 conductivity probe in filtered extracts (Kalra and Maynard, 1991), respectively. Soil cation
156 exchange capacity (CEC) was determined following extraction in buffered sodium acetate (NaOAc,
157 at pH 8.2; [Sumner and Miller, 1996](#)). Total nitrogen was measured by Kjeldahl digestion (Bremner,
158 1996). Available phosphorus (P_{ava}), as ortho-PO₄⁻², was determined using the method of Bray and
159 Kurtz (1945). Available potassium (K_{ava}) was determined following ammonium acetate (pH 7)
160 extraction and quantified by flame photometry (Black, 1986). Lime percentage, expressed using the
161 Total Neutralizing Value (T.N.V), was determined using the NaOH titration method.

162 The second subsample was maintained at field moisture and stored at 4 °C for subsequent
163 measurements of soil microbial activity. Soil basal respiration (BR) was measured by trapping and
164 quantifying CO₂ that was emitted from soil samples over a five-day period (Alef and Nannipieri,
165 1995). Substrate-induced respiration (SIR) was determined using glucose (1 %) as the substrate and
166 the evolved CO₂ was determined after 8 h incubation. Evolved CO₂ was adsorbed in 1 M NaOH and
167 measured by 0.1 M HCl titration (Anderson and Domsch, 1978).

168 Additional undisturbed soil cores distributed in the four treatments were collected for the
169 determination of bulk density (BD) in the 0-15 cm mineral layer (Blake and Hartge, 1986). Soils
170 were immediately sieved (2 mm) and kept in a plastic box to avoid evaporation. Soil moisture
171 content or WC (water mass/soil dry mass) was measured gravimetrically (oven-drying at 105 °C for
172 24 h; Famiglietti et al., 1998)

173

174 2.4. Diversity partitioning

175 To estimate the diversity components, we used the additive partitioning method as shown in
176 equation (1) (Crist et al., 2003):

$$177 \gamma = \alpha_1 + \sum_{i=1}^m \beta_i \quad [1]$$

178

179

180 where γ is the total species diversity, m is the number of scales, α_1 is the average diversity within
181 sample units in each site, and β_i is the beta diversity at each scale i . This method enabled us to
182 estimate the total species richness in each region (γ), which was partitioned into α_1 (the mean
183 number of species found per 1 m² plot) and β_1 (the difference between α_1 and γ_{site} , in which γ_{site}
184 was the total species richness of each site).

185 Then, we additively partitioned the β -diversity into two components, i.e. the spatial turnover and
186 nestedness. To do this, we used the method suggested by Baselga (2010), which is based on the
187 multiple dissimilarity derived from the Sørensen coefficient of dissimilarity (Baselga 2010). This
188 analysis was performed using the betapart package (Baselga and Orme 2012) within the R software
189 (R Development Core Team 2013).

190

191 2.5. Statistical analysis

192 First, the normality of each variable was verified using the Kolmogorov–Smirnov test as well as
193 variance homogeneity (Levene Test). Mathematical transformations of data were used when
194 necessary to correct deviations from normality and heterogeneity of variance. Then we used two-
195 way ANOVAs to detect significant differences between variable means associated with treatments,
196 regions, and the interaction between treatments and regions. If the two-way ANOVAs detected
197 significant differences, we applied a Duncan test for pairwise comparisons. Pearson correlation
198 coefficients were calculated to investigate possible relationships between oak cover, soil properties
199 and diversity components. Finally, relationships between beta diversity components and soil factors
200 were analyzed using linear regressions. All analyses were performed with the R statistical software
201 (R Core Team 2013).

202

203

204

205 **3. Results**

206 3.1. Effects of management measures and climate

207 The results showed that management (protected vs. unprotected), climatic conditions (arid vs. semi-
 208 arid) and their interaction had a significant (P -value < 0.05) effects on oak canopy cover (Table 1).
 209 These variables also produced significant effects on water content, silt, BR and SIR. However, for
 210 T.N.V, clay, Pava, Kava, total N and OC, only management and climatic conditions were significant
 211 and their interaction was not significant (Table 1). The percent of sand and BD were solely affected
 212 by management, while EC and pH were affected only by climatic conditions.

213 Diversity components (alpha, beta and gamma) and beta diversity components (turnover and
 214 nestedness) showed different trends in relation to management, climatic conditions and their
 215 interaction (Table 1). We detected significant effects of management and climatic conditions for all
 216 diversity components but a significant influence of the interaction was revealed only for beta and
 217 gamma diversity as well as turnover. Therefore, the effects of management on alpha diversity and
 218 nestedness were similar across the two regions (Table 1).

219 Table 1 Results of two way-ANOVA (F-value and significance level) testing for regions with different
 220 climatic conditions and management on oak canopy cover percentage, soil properties, diversity components
 221 (alpha, beta and gamma) and beta diversity components (spatial turnover and nestedness); (EC: electrical
 222 conductivity; OC: organic carbon; BD: Bulk density; WC: Water content; T.N.V: total neutralizing Value;
 223 N_{tot} : total nitrogen; K_{ava} : available potassium; P_{ava} : available phosphorus; BR: basal respiration; SIR:
 224 substrate induced respiration)

	Variable	Management	Climate condition	Management \times Climate condition
Oak canopy cover and soil physical properties	Oak canopy cover (%)	11.662**	8.396**	6.04*
	BD (g/cm ³)	1.037*	2.846	2.732
	Water content (%)	15.859**	11.017**	5.475*
	Sand (%)	5.988*	0.828	0.198
	Clay (%)	3.562*	0.608*	0.007
	Silt (%)	0.463*	3.795**	0.372*
Soil chemical properties	T.N.V (%)	26.447***	14.393**	2.084
	OC (%)	62.331***	43.721***	0.388
	Pava (ppm)	13.529	394.731***	0.438
	EC (dS/m)	1.752	6.209*	5.1*
	pH	4.134	62.152***	1.145
	N_{tot} (%)	62.331***	43.721***	0.388
	K_{ava} (ppm)	0.219	43.522***	0.005
Soil biological	BR (mg CO ₂ - C/kg soil/day)	177.92***	368.656***	112.515***

variables		105.622***	162.689***	33.996***
SIR (mg CO ₂ -C/kg soil/day)				
Diversity	Alpha diversity	160.163***	4.379*	0.054
	Beta diversity	136.289***	39.472***	37.768***
	Gamma diversity	370.559***	20.931***	35.477***
	Turnover	11.626**	183.214***	42.374***
	Nestedness	13.636**	4.909*	2.182

*** Significant at $P = 0.001$; ** Significant at $P = 0.01$ and * significant at $P = 0.05$

3. 2. Oak canopy cover and soil physico-chemical properties

Oak canopy cover was significantly higher in the Dalab region than in the Koshk region. This variable was also significantly higher in managed than in unmanaged plots for both Dalab and Koshk regions (Table 2). Available K was higher in the Dalab than in the Koshk region while the reverse was true for P_{ava} .

EC was higher in the unmanaged than in the managed plots of the Koshk region while it was lower in the Dalab region within which no significant differences were found between managed and unmanaged plots. pH was unaffected by management in both regions but was higher in the Koshk than in the Dalab region. Clay content was significantly higher in MD than in the other treatments, whereas silt content was higher in MK. Sand content was higher in unmanaged plots but the difference was significant in the Koshk region only. The treatment order for total nitrogen and organic matter contents was MD > UMD = MK > UMK whereas that of T.N.V was UMK > UMD > MK > MD (Table 2). Water content and bulk density were highest in the managed plots of the Dalab region and in the unmanaged plots of the Koshk region, respectively.

3. 3. Soil biological variables

The mean values of BR and SIR in the Dalab region were higher than those of the Koshk region but the influence of management was significant in both regions with higher values of BR and SIR in managed than in unmanaged plots (Table 2).

Table 2 Oak canopy cover, soil physico-chemical and microbial properties (mean \pm SE) measured in the different regions; (EC: electrical conductivity; OC: organic carbon; BD: Bulk density; WC: Water content; T.N.V: total neutralizing Value; N_{tot}:total nitrogen; K_{ava}: available potassium; P_{ava}: available phosphorus; BR: basal respiration; SIR: substrate induced respiration)

Variable	Koshk		Dalab	
	Managed	Unmanaged	Managed	Unmanaged
Oak canopy cover (%)	10.18 \pm 2.33 b	7.52 \pm 1.32b	20.13 \pm 3.92a	8.94 \pm 1.22b
BD (g/cm ³)	1.37 \pm 0.02b	1.48 \pm 0.03a	1.36 \pm 0.01b	1.38 \pm 0.02b
Sand (%)	47.87 \pm 1.61b	55.65 \pm 2.45a	48.58 \pm 1.24b	52.71 \pm 2.34ab
Clay (%)	27.71 \pm 1.51b	24.43 \pm 0.51b	31.04 \pm 2.16a	27.04 \pm 3.2b
Silt (%)	24.42 \pm 1.64a	19.92 \pm 2.04b	20.38 \pm 1.24b	20.25 \pm 0.91b
WC (%)	40.87 \pm 0.59b	38.06 \pm 0.85b	50.54 \pm 3.09a	39.74 \pm 0.99b
pH	7.38 \pm 0.01a	7.37 \pm 0.01a	7.14 \pm 0.04b	7.20 \pm 0.01b
EC (ds/m)	0.38 \pm 0.03 b	0.52 \pm 0.03a	0.29 \pm 0.01c	0.32 \pm 0.01c
Total nitrogen (%)	0.23 \pm 0.010b	0.18 \pm 0.010c	0.33 \pm 0.010a	0.21 \pm 0.006b
OC (%)	3.37 \pm 0.31b	2.81 \pm 0.55c	5.7151 \pm 0.32a	3.55 \pm 0.11b
T.N.V (%)	43.93 \pm 0.91b	52.40 \pm 0.47a	35.83 \pm 3.39c	48.85 \pm 1.87ab
K _{ava} (ppm)	263.09 \pm 15.00bc	169.80 \pm 8.47c	400.79 \pm 17.56a	387.67 \pm 39.55a
P _{ava} (ppm)	9.95 \pm 0.07a	9.09 \pm 0.13a	7.60 \pm 0.21b	7.07 \pm 0.05b
BR (mg CO ₂ - C/kg soil/day)	12.21 \pm 0.35c	10.49 \pm 0.21c	34.01 \pm 0.41a	16.40 \pm 1.34b
SIR (mg CO ₂ -C/kg soil/day)	23.61 \pm 0.62b	21.84 \pm 1.48c	49.40 \pm 1.95a	24.58 \pm 1.46b

Within rows, means with the same letters do not significantly differ ($p > 0.05$)

3.4. Alpha, Beta and Gama diversity and plant beta diversity components

Alpha, beta and gamma diversity values were higher in managed than in unmanaged plots in both regions. Diversity values were significantly higher (beta and gamma) or similar (alpha) between managed Dalab and managed Koshk whereas no significant differences were detected between the two regions in the unmanaged plots (Fig. 2 a).

Between the two components of the beta diversity, spatial turnover was largely higher than nestedness in each treatment (Fig. 2 b). Turnover significantly decreased according to the following order MD > UMD > UMK > MK. By contrast, nestedness was lowest in MD while there were no significant differences among the other treatments.





Fig. 2 Variations of the diversity components (mean \pm SE) according to the treatments for a) Alpha, Beta and Gamma components and b) Turnover and Nestedness. Means with the same letters for a given diversity component are not significantly different based on Duncan's multiple range test ($p > 0.05$)

3.5. Relationships between oak canopy cover, soil properties and diversity components







The oak canopy cover was positively correlated with silt, pH, K_{ava} and alpha diversity, while it was negatively correlated with spatial turnover (Table 3).


Table 3 Pearson correlation coefficients between the oak canopy cover and soil properties and components of diversity; (EC: electrical conductivity; OC: organic carbon; BD: Bulk density; WC: Water content; T.N.V: total neutralizing Value; N_{tot} :total nitrogen; K_{ava}: available potassium; P_{ava}: available phosphorus; BR: basal respiration; SIR: substrate induced respiration)

Variables	oak canopy cover	Variables	oak canopy cover	Variables	oak canopy cover
BD (g/cm ³)	- 0.14 ^{ns}	P _{ava} (ppm)	0.14 ^{ns}	Total nitrogen (%)	- 0.02 ^{ns}
WC (%)	- 0.08 ^{ns}	K _{ava} (ppm)	0.36 [*]	Alpha	0.36 [*]
Sand (%)	- 0.23 ^{ns}	TNV (%)	0.09 ^{ns}	Beta	- 0.28 ^{ns}
Clay (%)	- 0.03 ^{ns}	BR (mg CO ₂ - C/kg/day)	- 0.25 ^{ns}	Gamma	- 0.09 ^{ns}
Silt (%)	0.14 [*]	SIR (mg CO ₂ -C/kg soil/day)	- 0.28 ^{ns}	Spatial turnover	- 0.62 [*]
pH	0.32 [*]	EC (dS/m)	0.13 ^{ns}	Nestedness	0.12 ^{ns}

3.6. Linear regressions between beta diversity components and soil properties

There were strong negative relationships between spatial turnover and BD, EC, pH and P_{ava}. By contrast, spatial turnover was strongly and positively related to K_{ava}, BR and SIR (Fig. 3).





292

293 Fig. 3 Simple linear regressions between spatial turnover of plant species and soil properties;


294 * p-value < 0.05 and ** p-value < 0.01

295

296 The nestedness of plant species was significantly and positively related to T.N.V., and negatively
 297 related to WC, OC, N_{tot}, BR, and SIR (Fig.4).

298

299



300

301

302

303 Fig. 4 Simple linear regressions between nestedness of plant species and soil properties; * p-
 304 value < 0.05 and ** p-value < 0.01

305

306

307

308 5. Discussion

309 5.1. Effect of management and climatic conditions on soil properties

310 We found that some soil attributes were influenced by climatic conditions while others responded to
311 conservation measures. For instance, pH was probably mainly influenced by climatic conditions and
312 showed significantly lower values in the more humid Dalab region than in Koshk. However, we cannot
313 exclude that the higher organic matter content in soils of Dalab has stimulated the microbial activity, in
314 particular respiration, which has led to enhanced carbon dioxide production and the formation of
315 carbonic acid forms reducing soil pH (Brady and Weil, 2008).

316 Similarly, soil K_{ava} was higher in the semi-arid Dalab region than in the arid Koshk region whereas the
317 reverse was observed for P_{ava} . In accordance, Sardans and Peñuelas (2007) emphasized the influence of
318 climatic conditions, especially drought, in modifying soil phosphorus and potassium accumulation
319 patterns in Mediterranean woodlands.

320 Total nitrogen and organic carbon were higher in the more humid Dalab region than in the arid Koshk
321 region. Accordingly, climatic conditions influence soil organism activities, which play an important role
322 in improving the soil nutrient cycle (Frossard et al., 2000). For example, temperature affects the
323 decomposition of organic matter and thus influences the physical, chemical and biological soil
324 characteristics (von Lutzow & Kogel-Knabner, 2009). Therefore, the higher organic carbon content in
325 the Dalab region is likely related to soil nutrient availability and thus plays an important role in soil
326 fertility (Bationo et al., 2006). The decrease in nitrogen and organic carbon in unmanaged sites was
327 consistent with the result reported by Dahlgren et al. (2003) and Heydari et al. (2014) who showed that
328 human disturbances and site degradation decreased the amount of organic matter and soil nutrients.
329 Higher cover of woody and herbaceous species in managed sites protects soils from erosion (Mekuria
330 and Aynekulu, 2013) while degradation agents such as livestock grazing in unmanaged Koshk decrease
331 plant biomass accumulation, which in turn affects soil fertility (Savadogo et al., 2007).

332 EC was higher in the Koshk than in the Dalab region and unmanaged plots were associated with higher
333 EC values in Koshk. Some disturbances, such as overgrazing or tree cutting, reduce the protection of
334 soil by vegetation and tend to dry out the soil surface and increase [electrical conductivity](#) (Binkley and
335 Fisher, 2012), a process that is especially pronounced in arid and semi-arid regions (Pan et al., 2014).
336 This explanation also holds for the soil water content, which is higher in managed than in unmanaged
337 areas (Kidron and Gutschick, 2013).

Bulk density was higher in unmanaged than in managed areas although the difference was only significant in the Koshk region. It is well known that soil compaction that is induced by overgrazing and low soil organic carbon increases bulk density (Li et al., 2011, Chaudhari et al., 2013) as observed in the Koshk region.

Finer soil particles (i.e. clay and silt contents) were significantly higher in managed than in unmanaged sites, while sand was higher in unmanaged ones. Due to a higher percentage of bare soil and a lower vegetation cover, soils of unmanaged sites are more exposed to run off and erosion favouring the loss of silt and clay particles (Tessema et al., 2011).

Lime percentage (T.N.V) was higher in unmanaged sites than in managed ones in both arid and semi-arid climates. Accordingly, Celik (2005) showed that the amount of lime increased with forest degradation and land-use change, particularly from forestland to agriculture. Changes in the content of soil lime can also be influenced by other factors such as soil organic carbon, biological activities (soil respiration) and CO₂ pressure (pCO₂). Indeed, higher amounts of soil organic matter enhance CO₂ pressure due to higher heterotrophic and autotrophic respiration and then increase carbonate dissolution (Brady and Weil, 2008).

5.2. Soil biological variables

Basal respiration (BR) and substrate-induced respiration (SIR) in the the Dalab region were higher than those obtained from the Koshk region. **More abundant precipitations and lower temperatures in the semi-arid Dalab region compared to the arid Koshk region** are favourable to vegetation development and thus increase soil organic matter (Binkley and Fisher, 2012). This process promotes microorganism activities, which can explain the higher BR and SIR values. This result is consistent with the findings of Sardans and Peñuelas (2005) who showed that drought reduces soil enzyme activity in a Mediterranean holm oak forest while more generally, mesic climatic conditions are favourable to the activity of soil organisms (e.g. Qiu et al., 2005; Mansourzadeh & Raiesi, 2012). The same process applies to explain the higher BR and SIR values measured in managed plots compared to unmanaged plots. Indeed, the lower vegetation cover in unmanaged areas caused by human disturbances led to lower organic matter content and reduced respiration values (Cheng et al., 2013).

5.3. Alpha, Beta and Gama diversity and beta diversity components

Alpha and gamma diversity values as well as both components of beta diversity (turnover and nestedness) were significantly affected by management type and climatic conditions and, to a lesser degree, by the interaction between these two variables. This is in accordance with the observations that changes in climatic and management conditions influence species occurrence, diversity and distribution in various ecosystems (Adler & Levine, 2007; van der Putten, 2013), particularly in arid and semi-arid rangelands (Erfanzadeh et al. 2015) .

However, our results clearly indicated that management had a major influence on alpha, beta and gamma diversity (Fig 2a) whereas climatic conditions had no significant effect in our unmanaged Mediterranean ecosystems contrary to the results of some previous studies (e.g. Rodriguez et al., 2017). Conservation management positively affected the different components of species diversity in our study areas likely because it provided sufficient opportunities for vegetation rehabilitation and plant community reconstruction (Miura et al., 2003; Coetzee et al., 2014) as well as preservation of native plant species diversity (Cox and Underwood, 2011).

The fact that beta and gamma diversities were not influenced by climatic conditions when there are no conservation measures indicate that the role of climate on vegetation composition and diversity is considerably reduced compared to the influence of anthropogenic disturbances. In fact, gamma diversity, which corresponds to the number of species in each region, peaked at 65 and 50 in the managed Dalab and Koshk regions, respectively, whereas alpha diversity was approximately the same (24 and 25 respectively in MD and MK). Beta diversity, which represents the difference between the plots diversity (alpha), is therefore of 41 species for Dalab and 25 for Koshk. This indicates that the beta diversity in managed areas of both regions has a similar (Korshk) or a higher (Dalab) contribution to total diversity than alpha diversity. The reduction of the negative impacts of degradation agents by the conservation management strategy resulted not only in a higher diversity at the plot level (alpha diversity; Erfanzadeh et al., 2015) but also increased species composition at larger scales (Hermy and Verheyen, 2007). Our results are consistent with the findings of Tang et al. (2011) who observed a significant positive effect of conservation management on plant species diversity and those of Erfanzadeh et al. (2015) who showed that conservation management can increase plant diversity at the regional scale in arid regions.

Our results emphasize the major contribution of spatial turnover in beta diversity in comparison to nestedness whose contribution was limited in all treatments. These results imply that the variation in species assemblages in the studied regions is less explained by the presence of subsets of species from

richer sites than by the high degree of species replacement from one location to the other (Baselga, 2010). First, the more favourable environmental conditions of the Dalab region can explain its higher turnover rate compared to the Koshk region because better soil qualities can positively affect plant species distribution and migration (Harrison et al., 2006; LaManna et al., 2017) and result in increased turnover. Second, the clear influence of conservation measures in both areas suggest that they should be applied more extensively rather than limited to the richest sites (Balsega, 2010).

405

5.4. Relationships between percent oak canopy cover, soil properties and diversity components

Overstory canopy cover is an important ecological characteristic of Zagros oak forests, which was significantly reduced in disturbed areas (Heydari et al., 2014). Previous studies indicated that canopy cover controls the biological, physical and chemical soil properties as well as the herbaceous coverage, which in turn, influence the seedling recruitment and development (Moradi et al., 2017; Caldeira et al., 2014).

Our results show that oak canopy cover was positively correlated with silt, pH, Kava and alpha diversity while it was negatively correlated with turnover (Table 3). The higher organic matter content in soils under a more developed canopy cover can improve the cation exchange capacity and thus increase the soil pH (Dahlgren et al., 2003). Oak trees, thanks to their deep rooting system, can bring up nutrients from lower soil layers where they can be recycled via leaf litter, a process that can explain higher soil K content with increasing canopy cover (Moreno et al., 2007).

The canopy cover and litter, by creating a suitable microclimate and by limiting leaching, plays an important role in seed storage of plant species and thus increase the diversity of plant species (Arriaga & Mercado, 2004; Heydari et al. 2013 a). This can explain the positive relationship between oak canopy cover and the alpha diversity of plant species. As already observed by Heydari et al. (2015) in the same region, human degradation and lack of protective measures significantly and negatively affect overstory canopy, which, in turn, decrease species diversity.

In contrast, the negative correlation observed between oak canopy cover and turnover of plant species could reflect the modification of the microclimatic conditions below canopy, particularly light availability, which is beneficial to some shade-tolerant species but detrimental to many light-demanding plants (Benítez et al., 2015). Sabatini et al (2014) proposed that the ecological factors driving species

composition and turnover, like canopy cover, forest structure or topography, are system-specific and thus vary according to forest types.

Species turnover is negatively correlated with BD, EC, pH and Pava, but positively with Kava, BR and SIR. Turnover means the continuous or sudden replacement of species along an environmental gradient, and that removal or replacement is controlled by environmental variables and competition (Lafage *et al*, 2015; Baselga, 2010). Past studies have shown that the reduction of habitat quality could lead to the elimination of the most sensitive and non-resistant species (extinction) or species with narrow ecological niche range (Tilman and Lehman, 2001). In contrast, the improvement of soil conditions linked to conservation management efforts, can cause increased migration, development and distribution of species (Harrison *et al.*, 2006; LaManna *et al.*, 2017) and may increase the turnover.

Nestedness is observed when species of a specific habitat are a subset of species of a richer habitat (in term of species), and is usually explained by species extinction or selective migration (Calderón-Patrón, *et al.*, 2013; Baselga, 2010). In our study, nestedness is significantly and positively related to TNV, and negatively related to WC, OC, N_{tot}, BR, and SIR. Changes in these factors can lead to limitation in ecological niches -for instance, highly calcareous soils are not adapted to lime-intolerant species- and can increase the nestedness due to extinction of species (Barnes *et al.*, 1998). In contrast, more fertile soils (e.g. with higher NPK or OC contents) may increase the competition and thus species replacement, which can lead to a higher turnover.

Given that turnover was by far a greater contributor to total beta diversity than nestedness, our results support the hypothesis that site productivity (Harrison *et al.*, 2006; LaManna *et al.*, 2017 but see Zemunik *et al.*, 2016) and environmental heterogeneity (Chase and Leibold, 2002; Veech and Crist, 2007; Stegen *et al.*, 2013) promote beta diversity.

6. Conclusion

Our results indicate that conservation management results in positive changes in the physical, chemical and biological soil properties as well as in canopy cover of oak trees, which improve plant species diversity in both studied regions.

This confirms the effectiveness of conservation management to increase plant species diversity independently of the climatic conditions although we have noted a more pronounced effect in the less arid area. More specifically, this study has shown the major contribution of the β -diversity on γ -diversity, especially in the semi-arid region, because of a higher heterogeneity in this area.

458 Knowledge of the contribution of the components of beta diversity can be a valuable guide for
459 managers in adopting appropriate management decisions. The high spatial turnover and the low
460 importance of nestedness observed in this study imply that conservation measures should be applied
461 to a large number of sites of various qualities rather than to a limited number of the richest sites.
462 Besides, the negative correlation between oak cover and turnover suggests that oak forests could
463 benefit from a silviculture aiming at enhancing micro-environmental heterogeneity, such as the
464 creation of gaps, in order to promote species conservation and diversity.

465

466 **Acknowledgements**

467 **References**

- 468 Aazami F. 2016. Additive partitioning of plant species diversity in relation to management and
469 environmental variables in semi-arid area in western Iran. MSc thesis. Forest science group. Ilam
470 university, Ilam. 86 p.
- 471 Adler PB, Levine JM. 2007. Contrasting relationships between precipitation and species richness in
472 space and time. *Oikos*, 116: 221-232.
- 473 Alef Z, Nannipieri KP. 1995. Methods in applied soil microbiology and biochemistry. Academic,
474 London p 576.
- 475 Anderson JPE, Domsch KH. 1978. Physiological method for quantitative measurement of
476 microbial biomass in soils. *Soil Biological and Biochemistry*, 10:215–221.
- 477 Arriaga L, Mercado C. 2004. Seed bank dynamics and tree- fall gaps in a northwestern Mexican
478 *Quercus- Pinus* forest. *Journal of Vegetation Science*, 15: 661- 668.
- 479 Auyeung DSN, Suseela V, Dukes JS. 2013. Warming and drought reduce temperature sensitivity
480 of nitrogen transformations. *Global Change Biology*, 19 (2): 662-676.
- 481 Barnes BV, Zak DR, Spurr SH. 1998. *Forest Ecology*. John Wiley and Sons Inc, New York.
- 482 Baselga A, Orme CDL. 2012. Beta part: an R package for the study of beta diversity. *Methods in*
483 *Ecology and Evolution*, 3: 808–812.
- 484 Baselga A. 2010. Partitioning the turnover and nestedness components of beta diversity. *Global*
485 *Ecology and Biogeography*, 19:134–143.
- 486 Bationo A, Kihara J, Vanlauwe B, et al. 2007. Soil organic carbon dynamics, functions and
487 management in West African agro-ecosystems. *Agricultural Systems*, 94 (1): 13-25.
- 488
- 489 Benítez AR, Prieto M, Aragon G. 2015. Large trees and dense canopies: key factors for
490 maintaining high epiphytic diversity on trunk bases (bryophytes and lichens) in tropical montane
491 forests. *Forestry*, 88: 521–527.

492 Binkley D, Fisher RF. 2012. Ecology and management of forest soils, 4rd edn. Wiley, New York.
493 362 p.

494 Black CA. 1986. Methods of soil analysis, Part 1. ASA, Madison, WI, 9. P: 545- 566.

495 Blake GR, Hartge KH. 1986. Bulk density. In: Klute A (ed) Methods of soil analysis: part I—
496 physical and mineralogy methods, 2nded, Agronomy monograph no. 9, American Society of
497 Agronomy, Soil Science Society of America, Madison pp 363–376

498 Bouyoucos GJ. 1962. Hydrometer method improved for making particle size analysis of soils.
499 American Society of Agronomy Journal, 54: 44-46.

500 Brady NC., Weil RR. 2008. The Nature and Properties of Soils. 14th Edition. Prentice Hall, Upper
501 Saddle River, New Jersey. 975 p.

502 Bray RH, Kurtz LT. 1945. Determination of total organic and available forms of phosphorus in
503 soils. Soil Science, 59: 39-45.

504 Bremner JM. 1996. Nitrogen—total. In: Sparks DL (ed) Methods of soil analysis: part 3—chemical
505 methods. SSSA Book Series 5, Soil Science Society of America, Madison pp 1085–1122.

506 Caldeira MC, Ibañez I, Nogueira C, et al. 2014. Direct and indirect effects of tree canopy
507 facilitation in the recruitment of Mediterranean oaks. Journal of Applied Ecology, 51: 349–358

508 Calderón-Patrón JM, Moreno CE, Pineda-López R, et al. 2013. Vertebrate dissimilarity due to
509 turnover and richness differences in a highly beta-diverse region: the role of spatial grain size,
510 dispersal ability and distance. PLoS ONE, 8 (12): 1–10.

511 Celik I. 2005. Land-use effects on organic matter and physical properties of soil in a southern
512 mediterranean highland of Turkey. Soil Tillage Research, 83: 270-277.

513 Chase JM, Leibold MA. 2002. Spatial scale dictates the productivity– biodiversity relationship.
514 Nature, 416: 427–430.

515 Chaudhari PR, Ahire DV, Ahire VD, et al. 2013. Soil Bulk Density as related to Soil Texture,
516 Organic Matter Content and available total Nutrients of Coimbatore Soil. International Journal of
517 Scientific and Research Publications.

518 Chávez V, Macdonald SE. 2012. Partitioning Vascular Understory Diversity in Mixedwood
519 Boreal Forests, the Importance of Mixed Canopies for Diversity Conservation, Forest Ecology and
520 Management, 271: 19-26.

521 Cheng F, Peng X, Zhao P, et al. 2013. Soil Microbial Biomass, Basal Respiration and Enzyme
522 Activity of Main Forest Types in the Qinling Mountains. PLoS ONE, 8 (6): e67353.

523 Coetzee BWT, Gaston KJ, Chown SL. 2014. Local Scale Comparisons of Biodiversity as a Test
524 for Global Protected Area Ecological Performance: A Meta-Analysis. PLoS ONE, 9(8): e105824.

525 Cox RL, Underwood EC. 2011. The Importance of Conserving Biodiversity Outside of Protected
526 Areas in Mediterranean Ecosystems. PLoS ONE, 6 (1): e14508.

527 Crist TO, Veech JA, Gering JC, et al. 2003. Partitioning species diversity across landscapes and
528 regions: a hierarchical analysis of α , β , and γ diversity. *The American Naturalist* 162: 734–743.

529 Dahlgren RA, Horwath WR, Tate KW, et al. 2003. Blue oak enhance soil quality in California oak
530 woodlands. *California Agriculture*, 57 (2): 42-47.

531 Erfanzadeh R, Omidipour R, Faramarzi M. 2015. Variation of plant diversity components in
532 different scales in relation to grazing and climatic conditions. *Plant Ecology and Diversity*, 8(4):
533 537-545.

534 Famiglietti J, Rudnicki J, Rodell M. 1998. Variability in surface moisture content along a hill slope
535 transect: rattlesnake Hill, Texas. *Journal of Hydrology*, 210: 259–281.

536 Fathizadeh O, Hosseini SM, Zimmermann A, et al. 2017. Estimating linkages between forest
537 structural variables and rainfall interception parameters in semi-arid deciduous oak forest stands.
538 *Science of the Total Environment*, 601-602:1824-1837.

539 Feng C, Ma Y, Fu S, et al. 2017. Soil Carbon and Nutrient Dynamics Following Cessation of
540 Anthropogenic Disturbances in Degraded Subtropical Forests. *Land Degradation and Development*,
541 28 (8): v 2457-2467.

542 Frossard E, Condorn LM, Oberson A, et al. 2000. Processes governing phosphorus availability in
543 temperate soils. *Journal of Environmental Quality*, 29: 15–23.

544 Zemunik G, Turner BL, Lambers H, et al. 2016. Increasing plant species diversity and extreme
545 species turnover accompany declining soil fertility along a long-term chronosequence in a
546 biodiversity hotspot. *Journal of Ecology*, 104 (3): 792-805.

547 Harrison S, Davies KF, Saford HD, et al. 2006. Beta diversity and the scale-dependence of the
548 productivity–diversity relationship: a test in the Californian serpentine flora. *Journal of Ecology*,
549 94:110–117.

550 Hermy M, Verheyen K. 2007. Legacies of the past in the present-day forest biodiversity: a review
551 of past land-use effects on forest plant species composition and diversity. *Ecological Research*, 22:
552 361–371.

553 Heydari M, Omidipour R, Abedi M, et al. 2017. Effects of fire disturbance on alpha and beta
554 diversity and on beta diversity components of soil seed banks and aboveground vegetation. *Plant*
555 *Ecology and Evolution*, 150 (3): 1–10.

556 Heydari M, Faramarzi M, Pothier D. 2016. Post-fire recovery of herbaceous species composition
557 and diversity, and soil quality indicators one year after wildfire in a semi-arid oak woodland.
558 *Ecological Engineering*, 94: 688-697.

559 Heydari M, Pourbabaei H, Esmailzadeh O. 2015. The effects of habitat characteristics and human
560 destructions on understory plant species biodiversity and soil in Zagros forest ecosystem. *Journal of*
561 *plant researches*, 28 (3): 535-548.

562 Heydari M, Poorbabaei H, Bazgir M, et al. 2014. Earthworms as indicators of different forest
563 management and human disturbance in Ilam Oak foresto Iran. *Folia Forestalia Polonica, series A* 56
564 (3): 121–134.

565 Heydari M, Poorbabaie H, Esmaelzade O, et al. 2013 a. Germination characteristics and diversity
566 of soil seed bank and aboveground vegetation in disturbed and undisturbed oak forests. *Forest*
567 *Ecosystems (Forest Science and Practice)*, 15(4): 286–301.

568 Heydari M, Poorbabaie H, Salehi A, et al. 2013 b. Application of two-step clustering methods to
569 investigate effects of oak forests conservative management of Ilam city on soil properties. *Iranian*
570 *Journal of Forest and Poplar Research*, 21 (2): 329-343.

571 Kalra YP, Maynard DG. 1991. Methods manual for forest soil and plant analysis. For. Can.,
572 Northwest Reg., North. For. Cen., Edmonton, AB. Inf. Rep. NOR-X-311.

573 Kardol P, Cregger MA, Campany CE, et al. 2010. Soil ecosystem functioning under climate
574 change: plant species and community effects. *Ecology*, 91: 767–781.

575 Kidron GJ, Gutschick VP. 2013. Soil moisture correlates with shrub-grass association in the
576 Chihuahuan Desert. *Catena*, 107: 71–79.

577 Kooch Y, Jalilvand H, Bahmanyar MA, et al. 2007. Ecological distribution of Indicator species
578 and effective edaphical factors on the northern Iran lowland forests. *Journal of Applied Science*, 7:
579 1475 – 1483.

580 Kouba Y, Martí'nez-García F, Frutos Ad, et al. 2014. Plant β -diversity in human-altered forest
581 ecosystems: the importance of the structural, spatial, and topographical characteristics of stands in
582 patterning plant species assemblages. *European Journal of Forest Research*, 133 (6): 1057-1072.

583 Lafage D, Maugenest S, Bouzille JB, et al. 2015. Disentangling the influence of local and
584 landscape factors on alpha and beta diversities: opposite response of plants and ground-dwelling
585 arthropods in wet meadows. *Ecological Research*, 30 (6):1025–1035.

586 LaManna JA, Blote RT, Burkle LA, et al. 2017. Negative density dependence mediates
587 biodiversity–productivity relationships across scales. *Nature Ecology and Evolution*, 1:
588 1, pages1107–1115.

589 Lande R. 1996. Statistics and partitioning of species diversity, and similarity among multiple
590 communities. *Oikos*, 76: 5–13.

591 Li Y, Zhao H, Zhao X, et al. 2011. Effects of grazing and livestock exclusion on soil physical and
592 chemical properties in desertified sandy grassland, Inner Mongolia, northern China. *Environmental*
593 *Earth Sciences*, 63(4): 771-783.

594 Marvi Mohadjer M. 2005. *Silviculture*. Tehran University Press.No 2709. 387 p.

595 Mekuria W, Aynekulu E. 2013. Exclosure land management for restoration of the soils in degraded
596 communal grazing lands in northern Ethiopia. *Land Degradation & Development*, 24: 528–538.

597 [Mirzaei J, Moradi M. 2017. Relationships between flora biodiversity, soil physiochemical](#)
598 [properties, and arbuscular mycorrhizal fungi \(AMF\) diversity in a semi-arid forest. *Plant Ecology*](#)
599 [and Evolution](#), 150 (2): 151-159.

600 Miura S, Yoshinaga S, Yamada T. 2003. Protective effect of floor cover against soil erosion on
601 steep slopes forested with *Chamaecyparis obtusa* (hinoki) and other species. *Journal of Forest*
602 *Research*, 8 (1): 27–35.

603 Moradi Behbahani S, Moradi M, Basiri R, Mirzaei J. 2017. Sand mining disturbances and their effects on
604 the diversity of arbuscular mycorrhizal fungi in a riparian forest of Iran. *Journal of Arid Land*, 9 (6): 837–
605 849.

606 Moreno G, Obrador JJ, Garcia A. 2007. Impact of evergreen oaks on soil fertility and crop
607 production in intercropped dehesas. *Agriculture, Ecosystems and Environment*, 119: 270–280.

608 Moradi M, Imani F, Naji HR, Moradi Behbahani, S, Ahmadi MT. 2017. Variation in soil carbon
609 stock and nutrient content in sand dunes after afforestation by *Prosopis juliflora* in the Khuzestan
610 province (Iran). *iForest-Biogeosciences and Forestry*, 10 (3): 585-589.

611 Nelson DW, Sommers LE. 1982. Total carbon, organic carbon and organic matter. In: Page AL
612 (ed) *Methods of soil analysis: part 3—chemical methods*. Book series no. 5, Soil Science Society of
613 America, Madison pp 961–1010.

614 Onyekwelu JC, Mosandl R, Stimm B. 2006. Productivity, site evaluation and state of nutrition of
615 Gmelina arborea plantations in Oluwa and Omo forest reserves. *Nigeria Forest Ecology and*
616 *Management*, 229: 214-227.

617 Pan YX, Wang XP, Li XR, et al. 2014. The influence of *Caragana korshinskii* shrub on soil and
618 hydrological properties in a revegetation-stabilized desert ecosystem. *Hydrological Sciences Journal*,
619 59 (10): 1925-1934.

620 Parma R, Shataee Jouybari Sh. 2010. Impact of physiographic and human factors on crown cover
621 and diversity of Woody species in the Zagros forests (Case study: Ghalajeh forests, Kermanshah
622 province). *Iranian Journal of Forest and Poplar Research*, 18 (4): 539-555.

623 Qiu S, McComb AJ, Bell RW, et al. 2005. Response of soil microbial activity to temperature,
624 moisture, and litter leaching on a wetland transect during seasonal refilling. *Wetlands Ecology and*
625 *Management*, 13 (1): 43-54.

626 R Core Team. 2013. *R: A Language and Environment for Statistical Computing*. Foundation for
627 Statistical Computing, Vienna, Austria.

628 Raunkiaer C. 1934. *The life forms of plant and statistical plant geography*. Oxford, Clarendon
629 Press. de Martonne E. 1925. *Traite´ de ge´ographie physique*. 3 tomes, Paris.

630 Rodriguez N, Santonja M, Baldy V, et al. 2017. Ecological patterns of plant diversity in a
631 plantation forest managed by clearfelling. *Journal of Applied Ecology*, 43 (6):1160 – 1171.

632 Sabatini FM, Burrascano S, Tuomisto H, et al. 2014. Ground Layer Plant Species Turnover and
633 Beta Diversity in Southern-European Old-Growth Forests. *PLoS ONE*, 9 (4): e95244.

634 Sagheb-Talebi K, Sajedi T, Pourhashemi M. 2014. *Forests of Iran*. Springer Dordrecht Heidelberg
635 New York London.

636 Salehi A, Söderberg U, Eriksson LO, et al. 2013. Impacts of Forest-Based Activities on Woodland
637 Characteristics in a Forested Watershed of Southern Zagros, Iran. *Caspian Journal of Environmental*
638 *Sciences*, 11 (2): 161-176.

639 Sardans J, Peñuelas J. 2005. Drought decreases soil enzyme activity in a Mediterranean holm oak
640 forest. *Soil Biology and Biochemistry*, 37: 455–461.

641 Sardans J, Peñuelas J. 2007. Drought changes phosphorus and potassium accumulation patterns in
642 an evergreen Mediterranean forest. *Functional Ecology*, 21: 191–201.

643 Savadogo P, Sawadogo L, Tiveau T. 2007. Effects of grazing intensity and prescribed fire on soil
644 physical and hydrological properties and pasture yield in the savanna woodlands of Burkina Faso
645 Patrice. *Agriculture, Ecosystems and Environments*, 118: 80-92.

646 Schulze ED, Aas D, Grimm GW, et al. 2016. A review on plant diversity and forest management
647 of European beech forests. *European Journal of Forestry Research*, 135:51–67.

648 Sheklabadi M, Khademi H, Karimiyan Iqbal M, et al. 2007. Effects of Climate and long-term
649 grazing on some biological indicators of soil quality in Rangelands of Central Zagros. *Journal of*
650 *Water and Soil Science*, 11 (41): 103-116.

651 Stegen JC, Freestone AL, Crist TO, et al. 2013. Stochastic and deterministic drivers of spatial and
652 temporal turnover in breeding bird communities. *Global Ecology and Biogeography*, 22 (2): 202–
653 212.

654 Steinweg JM, Dukes JS, Paul EA, et al. 2013. Microbial responses to multi-factor climate change:
655 effects on soil enzymes. *Frontier in Microbiology*, 146: 1-11.

656 Strandberg B, Kristiansen SM, Tybirk K. 2005. Dynamic oak-scrub to forest succession: effects of
657 management on understory vegetation, humus forms and soils. *Forest Ecology and Management*,
658 211: 128–318.

659 Sumner ME, Miller WP. 1996. Cation exchange capacity and exchange coefficients. In Sparks DL
660 (ed.) *Methods of soil analysis: part 3—chemical properties*. SSSA book series 5, Soil Science
661 Society of America, Madison pp 1085–1122.

662 Tang Z, Fang J, Sun J, et al. 2011. Effectiveness of Protected Areas in Maintaining Plant
663 Production. *PLoS ONE* 6:e19116.

664 Tárrega R, Calvo L, Taboada A, et al. 2009. Abandonment and management in Spanish dehesa
665 systems: Effects on soil features and plant species richness and composition. *Forest Ecology and*
666 *Management*, 257: 731–738.

667 Tessema ZK, de Boer WF, Baars RMT, et al. 2011. Changes in soil nutrients, vegetation structure
668 and herbaceous biomass in response to grazing in a semi-arid savanna of Ethiopia. *Journal of Arid*
669 *Environments*, 75: 662-670.

670 Tilman D, Lehman C. 2001. Human-caused environmental change: impacts on plant diversity and
671 evolution. *Proceedings of the National Academy of Sciences*, 98: 5433–5440.

672 Toure T, Ge JW, Zhou JW. 2015. Interactions between soil characteristics, environmental factors,
673 and plant species abundance: A case study in the Karst mountains of Longhushan Nature Reserve,
674 southwest China. *Journal of Mountain Science*, 12 (4): 943–960.

675 van der Putten WH, Bardgett RD, Bever JD, et al. 2013. Plant–soil feedbacks: the past, the present
676 and future challenges. *Journal of Ecology* 101: 265–276.

677 Veech JA, Crist TO. 2007. Habitat and climate heterogeneity maintain beta-diversity of birds
678 among landscapes within ecoregions. *Global Ecology and Biogeography*, 16: 650–656

679 von Lutzow M, Kogel-Knabner I. 2009. Temperature sensitivity of soil organic matter
680 decomposition: What do we know?. *Biology and Fertility of Soils*, 46:1–15.

681 Wang G, Zhou Y, Xu X, et al. 2013. Temperature Sensitivity of Soil Organic Carbon
682 Mineralization along an Elevation Gradient in the Wuyi Mountains, China. *PLoS ONE*, 8(1):
683 e53914.

684 Wang P. 2007. Measurements and simulation of forest leaf area index and net primary productivity
685 in Northern china. *Journal of Environmental Management*, 85: 607-615.

686 Wilson S, Tilman DD. 2002. Quadratic Variation in Old-Field Species Richness along Gradient of
687 Disturbance and Nitrogen. *Ecology*, 83 (2): 492-504.

688 Yacht ALV, Barrioz SA, Keyser PD, et al. 2017. Vegetation response to canopy *disturbance* and
689 season of burn during oak woodland and savanna restoration in Tennessee. [Forest Ecology and](#)
690 [Management](#), 390: 187-202.

691 Zhang Y, Zhang S, Ma K, et al. 2014. Woody Species Diversity in Forest Plantations in a
692 Mountainous Region of Beijing, China: Effects of Sampling Scale and Species Selection. *PLoS*
693 *ONE*, 9 (12): e115038.