



HAL
open science

Habitat use at multiple scales as indicator of fish sensitivity to the artificialisation of large rivers

Hervé Capra, Hervé Pella, Laura Plichard, Nicolas Lamouroux, Michaël Ovidio

► **To cite this version:**

Hervé Capra, Hervé Pella, Laura Plichard, Nicolas Lamouroux, Michaël Ovidio. Habitat use at multiple scales as indicator of fish sensitivity to the artificialisation of large rivers. ISE2018 - Ecohydraulics, Aug 2018, Tokyo, Japan. pp.4. hal-02608150

HAL Id: hal-02608150

<https://hal.inrae.fr/hal-02608150v1>

Submitted on 5 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

HABITAT USE AT MULTIPLE SCALES AS INDICATOR OF FISH SENSITIVITY TO THE ARTIFICIALISATION OF LARGE RIVERS

HERVE CAPRA, HERVE PELLA, LAURA PLICHARD, NICOLAS LAMOUREUX
IRSTEA, UR RiverLy, 5 Rue de la Doua, CS 20244,
69625 Villeurbanne Cedex, France

MICHAEL OVIDIO
University of Liège, UR-FOCUS. Biology of behaviour Unit, Laboratory of Fish Demography and
Hydroecology, 22 Quai Van Beneden, 4020 Liège, Belgium

Habitat selection and seasonal mobility are major components of the biology of fish populations. The aim of this study is to characterise fish habitat use templates at two different spatial scales (over reaches of 2 vs. 35 km) for two native rheophilic cyprinids (*Barbus barbus* and *Squalius cephalus*) and for the non-native catfish (*Silurus glanis*). The study was carried out in the Upper Rhône River characterised by strong and rapid variations of flow and thermal regimes (caused by hydroelectric and nuclear power plants). Results reveal contrasted mobility patterns, habitat uses and home-range sizes between native and non-native fish species, but also a high inter-individual variability. At a local scale, fish modify their habitat selection when discharge changes in selecting "least constraining" conditions. At a larger scale they are able to travel rapidly between their favoured habitats.

1 INTRODUCTION

Many rivers of the northern hemisphere are heavily affected by human activity causing major changes to the rivers' continuity, morphology, flow and thermal regimes [1]. Habitat selection, mobility, spatial and temporal distribution of fish are major elements of their population dynamics [2, 3, 4, 5]. However, little information is available about the behavioral ecology of holobiotic fish in large European rivers at local as well as at large scale.

We evaluated, at different spatial and time scales, the habitat selection and displacement strategies of two native species of cyprinids (barbel and chub) and of one exotic fish species (catfish) in the artificial hydrological and thermal conditions of the Upper Rhône River at Bugey. The hydrological regime is influenced by peak power generation that generates artificial daily flow variations (hydropeaks). The thermal regime is modified by the warmed water discharges from a nuclear power plant (The Bugey NPP of Electricité de France).

2 MATERIAL AND METHODS

The study reach is a 35.5-km undiverted section of the French Upper Rhône River located north-east of Lyon (Figure 1) between the Sault-Brénaz hydroelectric facility (located upstream; Compagnie Nationale du Rhône - CNR) and the Jons-Cusset hydroelectric facility (located downstream; Électricité De France - EDF). The upstream limit (river kilometer Km0) of the study reach is distinguished by an artificial riprap weir (which cannot be passed by boat). The downstream limit (Km35.5) is identified by the dam of Jons. Sault-Brénaz's and Jons-Cusset's hydroelectric facilities cannot be passed by fish moving upstream and downstream passages remains possible through the turbines or spillways. A nuclear power plant located on the right bank of the study reach at Km 17.5 pumps ≈ 100 m³/s to cool its four reactors and discharges warmed-up water (between 7 and 10 °C warmer than the upstream water). The warm water discharge creates a strong transversal temperature difference between the left bank and the right bank [6, 7].

We assessed micro-habitat selection and mobility patterns of barbel, chub and catfish using acoustic telemetry. In 2009, n=5 barbel, 7 chub and 6 catfish were intensively tracked with fixed acoustic telemetry equipment to study their microhabitat (hydraulics, temperature and substrate) selection [8]. The fish were located every 3-4 s over 3 months in a 2 km-reach (Km16 – 18; Figure 1). In 2010, we used an active acoustic equipment to track n=37 barbels, 23 chubs and 13 catfishes for 7 months on a weekly basis for assessing longitudinal home ranges and movements patterns between preferred residence areas in a 35.5 km-reach divided in 71 500 m-sections [9] (Figure 1). Seven of the individuals tagged in 2009 were also tracked in 2010.

Local [7] and global [10] habitat availability in the study reaches were simulated by a two-dimensional hydrodynamic model (average water column velocity) calibrated and validated at the study reach scale.




Figure 1. The 2010 study reach of the upper Rhône River was located upstream of Lyon, between Sault-Brénaz (Km0; 45°51'19.74"N; 5°24'23.40"E) and Jons's (Km35.5; 45°46'4.23"N; 4°54'49.90"E) hydroelectric facilities. The main riverbed is shown within a digital elevation model (elevations in m) in order to illustrate the section's 15-m level difference [9]. The 2009 study reach was located from Km16 to Km18, in front of the nuclear power plant.

3 RESULTS AND DISCUSSION

At the local scale, the fish tracked in 2009 modified their microhabitat selection when the flow changed by selecting hydrodynamic conditions that were the least constraining. Avoiding fast-flowing midstream habitat, fish generally live along the banks in areas where the dewatering risk is high. When discharge decreased, however, they selected higher velocities but avoid both dewatering areas and very fast-flowing midstream habitat [11].

At large-scale the fish tracked in 2010 moved rapidly between specific (and thus probably favoured) habitats (Figure 3). In comparison with catfish, barbel and chub demonstrated wider longitudinal home ranges, move more and over longer distances with greater inter-individual variability of behaviours [9]. Catfish selected habitats with a morphology that was little diversified, less sensitive to variations in flow, and with warmer water temperature (i.e. downstream from the discharges of the Bugey nuclear power plant). Hydrodynamic description at 500 m-section scale showed that barbel selected mainly shallow sections characterized by high wetted surface area variability and high range of flow velocity. Chub selected mainly sections with high range of depth and flow velocity. Catfish mainly selected sections characterized by low wetted surface area variability, warmer temperature and high range of depth and flow velocity.

In the Rhône River, the observations of movements along potential preferential routes suggest that fish can memorize the bathymetry (Figure 2) but also the hydraulic and thermal variations of their environment. The results suggest that habitat degradation is more damaging to cyprinids than catfish in large anthropogenic rivers.




Figure 2. Movements of the catfish-#3856 during four days (3, 6, 15 and 25 from left to right) in August 2009 between Km17 and Km18. This fish always arrived from upstream (red dots) moving to the main release of warm water (thick red arrow; thin red arrow indicates the secondary release of warmed water), stayed there for a few hours (grey dots) and went back in upstream direction (yellow dots). Time corresponding to the different steps of the daily movements are indicated in the upper left corner for each day. The bathymetry of the study reach is indicated in blue (184.7 m, dark blue to 188.5 m, white). Flow direction is indicated with the blue arrow.




Figure 3. Movements of the barbel #3289 (blue) and #3548 (red) in 2010 in the Rhône River at large scale in time (top) and in space (bottom; see Figure 1 for 500 m- sections description). Color of each 500m- section corresponds to the number of contact of fish in this section. Arrows indicate the main steps (with the same numbers in top and bottom figures) of fish's movements.

4 ACKNOWLEDGMENTS

The authors thank the Agence de l'Eau Rhône Méditerranée et Corse, Electricité de France (EDF-DTG), the European Union/FEDER and the Région Aquitaine for their financial support. We gratefully acknowledge the HTI-Sonar engineers (Tracey Steig, Patrick Neelson, David Ouellette & Samuel Johnston) for their valuable assistance in the processing of acoustic data. Finally, we wish to thank Julien Bergé, Pascal Roger, Raphael Mons, Frédérique Bau, Olivier Croze and the numerous people who participated in the field work and more specifically Cédric Giroud (professional fisherman) for net fishing.

REFERENCES

- [1] Fullerton A. H., Burnett K. M., Steel E. A., Flitcroft R. L., Pess G. R., Feist B. E., Torgersen C. E., Miller D. J. and Sanderson, B. I., "Hydrological connectivity for riverine fish: measurement challenges and research opportunities", *Freshwater Biology*, 55, (2010), pp 2215 2237.
- [2] Benitez J.-P., Matondo B.N., Dierckx A. and Ovidio M., "An overview of potamodromous fish upstream movements in medium-sized rivers, by means of fish passes monitoring", *Aquatic Ecology*, 49, (2015), pp 481–497.
- [3] Fredrich F., Ohmann S., Curio B. and Kirschbaum, F., "Spawning migrations of the chub in the River Spree, Germany", *Journal of Fish Biology*, 63, (2003), pp 710 723.
- [4] Gardner C. J., Deeming D. C. and Eady P. E., "Seasonal Water Level Manipulation for Flood Risk Management Influences Home-Range Size of Common Bream *Abramis brama* L. in a Lowland River", *River Research and Applications*, 31, (2015), pp 165 172.
- [5] Radinger J. and Wolter C., "Patterns and predictors of fish dispersal in rivers", *Fish and Fisheries*, 15, (2014), pp 456 473.
- [6] Ginot V., Souchon Y. and Roger P., "Impact de l'élévation artificielle de température induite par le fonctionnement du Centre Nucléaire de Production Electrique du Bugey (fleuve Rhône) sur les communautés de poissons", *Hydroécologie Appliquée*, 8, (1996), pp 1 33.
- [7] Capra H., McNeil E., Bouillon M.-C., Pella H. and Alfaro C., "Intérêt d'un modèle hydrodynamique en deux dimensions pour interpréter le comportement des poissons dans les grands cours d'eau", *La Houille Blanche*, (2011), pp 28–33.
- [8] Bergé J., Capra H., Pella H., Steig T., Ovidio M., Bultel E. and Lamouroux N., "Probability of detection and positioning error of a hydro acoustic telemetry system in a fast-flowing river: Intrinsic and environmental determinants", *Fisheries Research*, 125–126, (2012), pp 1–13.
- [9] Capra H., Pella H. and Ovidio M., "Individual movements, home-ranges and habitat use by native rheophilic cyprinids and non-native catfish in a large regulated river", *Fisheries Management and Ecology*, (in press).
- [10] Capra H., Pella H., Morin J., LePichon C., Perraud C., Detry T., Secretan Y., Jouve P. and Matte P., "Conséquences de l'artificialisation de l'hydrologie du Rhône sur la structuration des communautés d'invertébrés et de poissons", *Rapport Final. Rapport Cemagref – Agence de l'Eau RM&C*, (2011), 109 p.
- [11] Capra H., Plichard L., Bergé J., Pella H., Ovidio M., McNeil E. and Lamouroux N., "Fish habitat selection in a large hydropeaking river: Strong individual and temporal variations revealed by telemetry", *Science of The Total Environment*, 578, (2017), pp 109 120.