



HAL
open science

Perfluorooctane sulfonate toxicokinetics at different temperatures using a rainbow trout physiologically-based toxicokinetic model

A. Vidal, Rémy Beaudouin, Jeanne Garric, Marc Babut

► To cite this version:

A. Vidal, Rémy Beaudouin, Jeanne Garric, Marc Babut. Perfluorooctane sulfonate toxicokinetics at different temperatures using a rainbow trout physiologically-based toxicokinetic model. SETAC: Environmental Risk Assessment of PFAS, Aug 2019, Durham, United States. pp.1, 2019. hal-02609660

HAL Id: hal-02609660

<https://hal.inrae.fr/hal-02609660v1>

Submitted on 16 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Alice VIDAL¹, Rémy BEAUDOUIN², Jeanne GARRIC¹, Marc BABUT^{1*}

¹ Irstea, UR RIVERLY, Villeurbanne, France

² INERIS, Models for Ecotoxicology and Toxicology (METO), Verneuil-en-Halatte, France

Study objectives

Project aiming to improve our understanding of the toxicokinetics of perfluoroalkyl substances (perfluorooctane sulfonate - PFOS, perfluorohexane sulfonate and perfluorononanoate - PFNA) in fish, using the rainbow trout (*Onchorynchus mykiss*) as model species.

Ultimately, the model developed should account for the influence of environmental factors, and to be predictive.

First steps (current status):

- physiologically-based toxicokinetic model (descriptive): uptake is assumed to occur through respiration and diet, elimination hypothetically occurs through feces, urine and gills.
- influence of water temperature on PFOS uptake and elimination.

Model design (developed on experimental data at 19°C)

Basic model: exposure through diet only.


Additional hypotheses:

– enterohepatic cycle

– absorption / elimination through the gills

Calibration at 19°C ⇒ cardiac output adjusted according to Arrhenius law.

Good fit to experimental data (below)



Entero-hepatic cycle: whatever the reabsorption rate, no improvement of model outputs:


- Bayesian Information Criterion (BIC) higher for the sub-models including this cycle [782 - 803, depending on reabsorption rate] than for the basic model [777].

Considering uptake through gills (diffusion): lower overall Relative Error to the mean (RE).

Elimination bile-feces > gills > urine (idem, lower RE).

Uptake through gills: need to account for ionization / active transport.

Same for elimination through gills?




Vidal A, Garric J, Babut M, Beaudouin R. (2019) Elucidating the fate of perfluorooctanoate sulfonate using a rainbow trout (*Onchorynchus mykiss*) physiologically-based toxicokinetic model. *Sci. Total Environ.* 691, 1297-1309

Approach

Combination of experiments and modelling.

Experiments at 7°C, 11°C and 19°C:

- sub-adult trout (15 months old, ≈300 g at the start of exposure)
- spiked food: PFOS, PFHxS and PFNA 500 µg kg⁻¹ dw each
- water temperature 7 °C, 11°C and 19 °C
- exposure for 28 to 42 days; depuration for 28 to 35 days.




Sampling and analysis:

- blood, liver, kidney, viscera, brain.
- food
- water
- analysis UHPLC/ MS-MS (MRM mode)

Modelling approach:

- trout physiology based on Nichols *et al.* (1990) and Grech *et al.* (2019);
- (.../...)

Vidal A., Lafay F., Daniele G., Vulliet E., Rochard E., Garric J., Babut M. (2019) Does water temperature influence the distribution and elimination of perfluorinated substances in rainbow trout (*Oncorhynchus mykiss*)? *Environ. Sci. Pollut. Res.* 26 (16) 16355-16365



(.../...)

- growth accounted for;
- *a priori* information on plasma:tissue partition coefficients derived from Goeritz *et al.* (2013);
- PFAS TK parameters obtained by Bayesian inference using our experimental data

Influence of water temperature modelled on the basis of Arrhenius law.

Processes concerned:

- fish growth
- cardiac output and ventilation rate fitted or based on literature data.
- organ perfusion rates
- absorption rates
- clearances
- blood:tissue partition coefficients

Nichols JW, McKim JM, Andersen ME, Gargas ML, Clewell Iii HJ, Erickson RJ. 1990. A physiologically based toxicokinetic model for the uptake and disposition of waterborne organic chemicals in fish. *Toxicol. Appl. Pharm.* 106:433-447
 Grech, A., Tebby, C., Brochet, C., Bois, F.Y., Bado-Nilles, A., Dorne, J.L., Quignot, N., Beaudouin, R., 2019. Generic physiologically-based toxicokinetic modelling for fish: Integration of environmental factors and species variability. *Sci. Tot. Environ.* 651, 516-531
 Goeritz J, Falk S, Stahl T, Schäfers C, Schleichriem C. 2013. Biomagnification and tissue distribution of perfluoroalkyl substances (PFASs) in market-size rainbow trout (*Oncorhynchus mykiss*). *Environ. Toxicol. Chem.* 32:2078-2088

Effect of water temperature on PFAS toxicokinetics

Arrhenius law

$$\dot{A}_T = \exp\left(\frac{T_A}{T_r} - \frac{T_A}{T}\right)$$

T_A : Arrhenius temperature (°K)
 T : water temperature (°K)
 T_r : temperature at which the physiological process was originally modelled (°K)

Variation of process rate (Arrhenius S., 1889)

$$\dot{k}_T = \dot{k}_r \times \dot{A}_T$$

\dot{k}_T : process rate at T °K
 \dot{k}_r : process rate at T_r °K
 Arrhenius, S. (1889) "On the reaction velocity of the inversion of cane sugar by acids" *Zeitschrift für Physikalische Chemie*, 4, 226

Basic model ⇒ influence of temperature on fish growth:

- model output depends of T°C, food quantity (α), growth rate (k), and maximal size (L_m);
- calibration on experimental data from Goeritz *et al.* (2013) assuming $\alpha = 1.0$
- respective α values at 7°C and 19 °C equal 0.56 and 0.58

Hypotheses: each process considered separately; additionally, all processes together.



Bayesian inference: adjustment to all experimental data (4 temperatures) simultaneously.

Bayesian Information Criterion (BIC) allows determining the best model.

Lower BIC values obtained when all processes (cardiac output, organ perfusion rates, partition coefficients, absorption rates, clearances) included.

Good agreement between observed and predicted concentrations in most organs at 7°C, 15° and 19°C.

- Predicted cardiac outputs and oxygen consumption in agreement with available experimental data.
- Predicted PFOS concentrations farther from observations in viscera and kidney than in muscle, blood and liver



Barron, M.G., Tarr, B.D., Hayton, W.L., 1987. Temperature-dependence of cardiac output and regional blood flow in rainbow trout, *Salmo gairdneri* Richardson. *J. Fish Biol.* 31, 735-744
 Myrick, C.A., Cech Jr, J.J., 2000. Temperature influences on California rainbow trout physiological performance. *Fish Physiol. Biochem.* 22, 245-254
 Elliott, J.W. 1969 The oxygen requirements of Chinook salmon. *The Progressive Fish-Culturist* 31, 67-73

Acknowledgements

The project was funded by the Rhone-Mediterranean and Corsica Water Agency. The Rhone-Alpes Region provided a doctoral fellowship grant for Alice VIDAL.

Contact: remy.beaudouin@ineris.fr; marc.babut@irstea.fr