



HAL
open science

A glyphosate-based herbicide induces sub-lethal effects in early life stages and liver cell line of rainbow trout, *Oncorhynchus mykiss*

Shannon Weeks Santos, Patrice Gonzalez, Bettie Cormier, Nicolas Mazzella, Bertille Bonnaud, Soizic Morin, Christelle Clerandeanu, Bénédicte Morin, Jérôme Cachot

► To cite this version:

Shannon Weeks Santos, Patrice Gonzalez, Bettie Cormier, Nicolas Mazzella, Bertille Bonnaud, et al.. A glyphosate-based herbicide induces sub-lethal effects in early life stages and liver cell line of rainbow trout, *Oncorhynchus mykiss*. *Aquatic Toxicology*, 2019, 216, pp.105291-10.1016/j.aquatox.2019.105291 . hal-02609717

HAL Id: hal-02609717

<https://hal.inrae.fr/hal-02609717v1>

Submitted on 5 Jan 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **A glyphosate-based herbicide induces sub-lethal effects in early life stages and liver**
2 **cell line of rainbow trout, *Oncorhynchus mykiss*.**

3 Shannon Weeks Santos¹, Patrice Gonzalez¹, Bettie Cormier¹, Nicolas Mazzella³, Bertille
4 Bonnaud³, Soizic Morin³, Christelle Clérandeau¹, Bénédicte Morin¹, Jérôme Cachot^{1,*}

5

6 ¹UMR CNRS 5805 EPOC, University of Bordeaux, Allée Geoffroy Saint-Hilaire, CS 50023,
7 33615 Pessac Cedex, France

8 ²UMR CNRS 5805 EPOC, University of Bordeaux, Place du Dr B. Peyneau, 33120
9 Arcachon, France

10 ³IRSTEA, UR EABX, 50 avenue de Verdun 33612 Cestas cedex, France

11 ***Corresponding author:** jerome.cachot@u-bordeaux.fr

12

13 **Abstract**

14 Most pesticides used in agriculture end up in the aquatic environment through runoff and
15 leaching of treated crops. One of the most commonly used herbicides is glyphosate. This
16 compound or its metabolites are frequently detected in surface water in Europe. In the present
17 study, *in vivo* and *in vitro* studies were carried out using the early life stages of rainbow trout
18 (*Oncorhynchus mykiss*) and the cell line RTL-W1 (a liver cell line from rainbow trout) to
19 characterize the toxic effects of glyphosate at environmentally-realistic concentrations. Both
20 studies were performed using the commercial formulation Roundup® GT Max, and technical-
21 grade glyphosate for the *in vitro* study. Eyed-stage embryos were exposed for 3-weeks to sub-
22 lethal concentrations (0.1 and 1 mg/L) of glyphosate using Roundup. Numerous toxicity
23 endpoints were recorded such as survival, hatching success, larval biometry, developmental
24 abnormalities, swimming activity, genotoxicity (formamidopyrimidine DNA-glycosylase Fpg-
25 modified comet assay), lipid peroxidation (TBARS), protein carbonyls and gene transcription.
26 Neither concentrations affected embryonic or larval survival, and no significant increases of
27 developmental abnormalities were observed. However, a significant decrease was observed
28 in the head size of larvae exposed to 1 mg/L of glyphosate. In addition, a significant increase
29 in mobility was observed for larvae exposed to the weakest concentration compared to control
30 larvae. Remarkably, TBARS levels were significantly decreased on larvae exposed to 1 mg/L
31 (a.i.), and *cat* and *cox1* genes were differently transcribed from controls. DNA damage was
32 detected by the Fpg-modified comet assay in RTL-W1 cell line exposed to the technical-grade
33 glyphosate and Roundup formulation. The results suggest that sub-chronic exposure to
34 glyphosate, at environmental concentrations, represent a potential risk for aquatic organisms.

35

36 **Keywords:** pesticide, fish embryos, liver cell line, cytotoxicity, embryotoxicity, teratogenicity,
37 genotoxicity, photomotor response

38

39 1. Introduction

40 One of the most commonly used pesticides are the glyphosate-based herbicides, usually
41 transported by agricultural runoff and frequently detected in surface water at high
42 concentrations (Peruzzo et al., 2008). Glyphosate is the active ingredient of Roundup®
43 herbicide; and is commonly used in the form of salt of isopropylamine glyphosate. Glyphosate
44 is a broad-spectrum, non-selective and systemic herbicide for the control of weeds and grass,
45 used in both agricultural and non-agricultural areas. The main degradation products of
46 glyphosate are aminomethyl phosphonic acid (AMPA) and CO₂ (Grandcoin et al., 2017).

47 Half-life of glyphosate has been determined in several studies, ranging from few days to 2
48 weeks in freshwater (Giesy et al., 2000). However, its dissipation depends on the local
49 conditions regulated by chemical, physical and biological factors (Giesy et al., 2000), where
50 half-life could last sometimes more than 60 days (Myers et al., 2016).

51 Environmental exposure to glyphosate is extensive, due to the vast quantities used annually
52 all over the world (Van Bruggen et al., 2018). Increased use of glyphosate is closely linked to
53 the endorsement of genetically modified glyphosate-resistant crops (Van Bruggen et al., 2018),
54 cultivated at about 100 million hectares in 22 countries (mostly soybean, maize, canola and
55 cotton) (<http://www.fao.org/docrep/015/i2490e/i2490e04d.pdf>). This is particularly true in North
56 and South America, where elevated glyphosate concentrations were reported in different
57 streams and lakes near agricultural basins. For instance, in the Pampa region (Argentina)
58 glyphosate residues were detected up to 4.52 µg/L in surface water (Castro Berman et al.,
59 2018). However, higher concentrations were detected in streams near transgenic soybean
60 cultivation in Pergamino-Arrecifes (North of Buenos Aires), where levels of glyphosate in water
61 varied from 100 to 700 µg/L (Peruzzo et al., 2008). Coupe et al., (2012) studied the fate of
62 glyphosate in different agricultural basins in North America and France, and maximum
63 concentrations of glyphosate were observed between 73 and 430 µg/L.

64 Several studies have documented the toxicity of glyphosate in various aquatic invertebrates,
65 and acute toxicity thresholds in fish are generally much higher than the concentrations found
66 in streams following applications of crops (Folmar et al., 1979). Commercial formulations of
67 glyphosate seems to be more toxic than the pure molecule, due to interference from
68 substances such as polyethoxilene amine surfactant (POEA) (Folmar et al., 1979; Giesy et al.,
69 2000; Navarro and Martinez, 2014; Tsui and Chu, 2003) which helps the active ingredient
70 penetrate the plant surface. Since fish are susceptible to glyphosate exposure by direct uptake
71 through their gills and via their diet (Giesy et al., 2000), there are several studies that have
72 demonstrated sublethal effects of glyphosate on fish. For example, effects on genotoxicity
73 (Çavaş and Könen, 2007; Guilherme et al., 2012, 2010), acetylcholinesterase (AChE) inhibition
74 (Salbego et al., 2010) swimming alterations (Bridi et al., 2017; Valéria D.G. Sinhorin et al.,
75 2014), reproduction (Uren Webster et al., 2014), and formation of reactive oxygen species (de
76 Moura et al., 2017; Gluszczak et al., 2007; Harayashiki et al., 2013; Üner et al., 2006) have
77 been observed in different fish species.

78 The use of rainbow trout fish (*Oncorhynchus mykiss*) in ecotoxicology is very well documented;
79 and a number of previous studies have looked at the toxicity of glyphosate in this species
80 (Hildebrand et al., 1982; Morgan and Kiceniuk, 1992; Tierney et al., 2007; Topal et al., 2015).
81 Studies have been performed using early life stages (ELS) of fish on the deleterious effects of
82 glyphosate (Sulukan et al., 2017; Yusof et al., 2014; Zebra et al., 2017; Zhang et al., 2017);
83 however, few studies have been done on ELS of rainbow trout. ELS of rainbow trout can be
84 easily raised under laboratory conditions, and because of its slow embryo-larval development,
85 toxicity tests allow longer sub-chronic exposure to toxicants.

86 On the other hand, the use of cell lines allows the screening of molecules, the study of the
87 mode of action of chemicals and the toxicity assessment of complex environmental samples
88 (Bols et al., 2005; Castaño et al., 2003). Several studies have been done studying the effects
89 of glyphosate on fish cell lines (Alvarez-Moya et al., 2014; Lopes et al., 2018; Qin et al., 2017).
90 For this work, a reference cell line of rainbow trout, RTL-W1 (Rainbow Trout Liver-Waterloo

91 1), was selected. This cell line, developed by Lee et al. (1993), is derived from untransformed
92 liver tissue of rainbow trout. The RTL-W1 line consists of adherent fibroblastic cells and has
93 the ability to metabolize xenobiotics.

94 The aim of this work was to study the effects and mechanisms of glyphosate toxicity, using a
95 commercially available product called Roundup on rainbow trout, focusing on ELS, cell cultures
96 and a wide range of endpoints. Exposure of rainbow trout embryos and larvae was conducted
97 using Roundup® for 3 weeks. Several endpoints were studied, such as viability, hatching
98 success, biometric changes, locomotion, genotoxicity, lipid and protein oxidation, and gene
99 transcription. In order to explain some of the observed effects, 10 genes were selected
100 according to their function in antioxidant defense (*cat* and *sod*), detoxification (*gst*),
101 mitochondrial metabolism (*cox1* and *12s*), DNA repair (*ogg1* and *rad51*), apoptosis (*bax*) and
102 reproduction (*er-b* and *cyp19a1*) on fish. In addition, cytotoxicity assays on the RTL-W1 cell
103 line were implemented to screen the toxicity of technical grade glyphosate and Roundup®.

104

105 **2. Materials and methods**

106 **2.1. *In vivo* study: rainbow trout**

107 **2.1.1. Test chemicals**

108 Preparation of glyphosate solutions was carried out using the commercial formulation of
109 Roundup® GT Max. The active substance is 480 g/L of glyphosate acid, which is equivalent
110 to 588 g/L of potassium salt of glyphosate. Two stock solutions were prepared at 0.1 and
111 1 g/L of glyphosate (active ingredient a.i.) with osmosis water. From these stock solutions,
112 exposure solutions at 0.1 and 1 mg/L of glyphosate (a.i.) were prepared.

113 **2.1.2. Exposure system**

114 Eyed stage embryos, at 288 °D (degree days), from rainbow trout (*Oncorhynchus mykiss*)
115 were obtained from INRA-PEIMA (Sizun, FR). Rainbow trout embryos were exposed to 0

116 (control), 0.1 and 1 mg/L of glyphosate (a.i.) in total darkness and with a temperature of 12°C
117 in a climate chamber for 3 weeks. Each studied condition consisted in 3 replicates with 100
118 embryos in 1 L aquaria. Exposure solutions was prepared in three 5 L tanks of spring water
119 from Laqueuille (4.7 mg/L Ca, 1.8 mg/L Mg, 5.9 mg/L Na, 2.8 mg/L K, 40.3 mg/L HCO₃⁻, 0.2
120 mg/L SO₄²⁻, 0.5 mg/L NO₃⁻, pH 7.5, <1.2 mg/L Cl⁻) and was renewed every two days. A
121 peristaltic pump (Watson Marlow, USA) was used to maintain a continuous flow rate of water
122 (9 mL/min) into the incubation aquaria. Dissolved oxygen concentration was measured each
123 day with a fiber-optic oxygen mini-sensor Fibox 3 (PreSens Precision Sensor, Regensburg,
124 Germany) and data was recorded with OxyView v6.02 software (PreSens Precision Sensor).

125

126 **2.1.3. Chemical analysis in water**

127 Concentrations of glyphosate and its main metabolite, amino methyl phosphonic acid (AMPA),
128 were analyzed in water samples. Water samples were collected at T₀ and T₄₈ (48 h after
129 exposure and before water was changed). Glyphosate and AMPA were measured by the
130 method described by Fauvelle et al. (2015). Briefly, 5 mL of each samples were spiked with
131 150 µL of glyphosate and AMPA 13C 15N at 20 ng mL⁻¹. Then, 325 µL of 50 mM borate-Na
132 solution and 200 µL EDTA-Na₂ 200 mM were added. After homogenization, solutions were left
133 for 5 minutes, 4.5 mL of acetonitrile and 600 µL of FMOC-Cl (50 mg mL⁻¹) were added and
134 samples were left in dark for 30 minutes in order to form FMOC derivates. Acetonitrile was
135 evaporated under nitrogen flow until the volume was below 5 mL. Then, a liquid-liquid
136 extraction with 1.5 mL of ethyl acetate was performed three times. Ethyl acetate was
137 evaporated under nitrogen flow for 15 minutes. One hundred µL of formic acid was added and
138 the sample volume was adjusted to 5 mL. A solid phase extraction was then performed on
139 OASIS HLB cartridges (3 mL, 60 mg, 30 µm particle size, Waters) conditioned with 1 mL of
140 MeOH and 1 mL of formic acid 0.1 %. Samples were loaded on cartridges, and the cartridges
141 were rinsed with 1 mL of formic acid 0.1 % and 1 mL deionized water. The cartridges were

142 dried under nitrogen flow before elution with 2 mL of ammonium hydroxide/deionized
143 water/MeOH 2/30/68. Extracts were evaporated under nitrogen flow until stabilization volume
144 (0.5 mL). The volume was adjusted to 1 mL with deionized water. Analyses was performed by
145 HPLC-ESI MSMS.

146

147 **2.1.4. Embryo-toxicity assay**

148 The viability of embryos and larvae was recorded daily and dead specimens were removed
149 immediately. Half-hatched embryos were considered when part of the body was inside the
150 **chorion**. Embryonic or larval mortality was the number of dead individuals compared to the
151 total number of embryos at the start of the experiment or total number of hatching larvae. The
152 half-hatched embryo rate was calculated by dividing the number of half-hatched embryos by
153 the total number of embryos at the beginning of the experiment. Hatching time expressed in
154 degree days (DD) was the duration of embryonic development from fertilization to hatching. At
155 the end of the experiment, yolk-sac larvae (540 °D) were placed in Petri dishes with carbonated
156 water and ice to sedate them, and photos were taken for each larva with a stereomicroscope
157 (MZ 7.5 Leica) coupled to a camera CCD (DFP420C Leica) and a cold light (Intralux® 4100,
158 Volpi AG, Schlieren, Switzerland). From the photos, total body length (**from the end of upper**
159 **jaw to the base of the caudal fin**) and head length (**from the end of the upper jaw to the end of**
160 **the pectoral fin attachment level**) were measured for each larva. The presence of
161 developmental anomalies - including edemas, yolk-sac absorption, spinal malformations,
162 craniofacial anomalies, presence of hemorrhages - was recorded in 15 larvae per replicate
163 randomly chosen.

164

165 **2.1.5. Swimming behavior analysis**

166 Analysis of swimming behaviour was carried out on 12 yolk-sac larvae per replicate at 528 DD.
167 The larvae were acclimated individually 30 minutes in the dark at 12°C in 6-well microplates
168 containing 8 mL of exposure solution. The microplates were placed in the recording chamber
169 (Daniovision Image Analysis System with Ethovision software version 12.0 Noldus) connected
170 to a thermoregulation system set at $12 \pm 0.5^\circ\text{C}$ (Pilot one®, Huber). Larvae were subjected to
171 a light/dark cycle of 30 minutes, divided into 10 minutes dark, 10 minutes light, and 10 minutes
172 dark. This cycle is designed to analyze the photomotor response of larvae in response to light
173 stimulation. An infrared camera in the recording chamber records the movement of each larva
174 focusing on their center of gravity. The average velocity of each larva was calculated over 30
175 seconds. The total distance traveled, time of mobility and the time spent in the peripheral area
176 of the wells were determined for each larva.

177

178 **2.1.6. Biochemical analyses**

179 *Preparation of supernatant*

180 At the end of the exposure, 4 pools of 2 yolk-sac larvae were frozen in liquid nitrogen and
181 stored at -80°C until analysis. Larvae (approximately 250 mg) were homogenized in a
182 phosphate buffer (0.1 M; pH 7.5; 4°C) using an UltraTurrax® tissue homogenizer fitted with a
183 potter at 3,000 rpm (4°C). Then, samples were centrifuged at 9,000 g for 25 min at 4°C . The
184 supernatant S9 fraction obtained were placed in different tubes for total protein, TBARS and
185 protein carbonyl measurements.

186 *Total protein*

187 The total protein concentration was determined using the method of Lowry et al. (1951) on S9
188 fraction. Bovine Serum Albumin (BSA) was used as a standard. Measurements were
189 performed using a spectrophotometer microplate reader (Synergy HT, BioTek).

190 *Lipid peroxidation (TBARS)*

191 Lipid peroxidation was assessed following the method of Buege and Aust (1978) adapted to a
192 microplate reader. Five hundred μL of S9 fraction were added to 500 μL of a solution
193 containing 20 % of butylated hydroxytoluene (BHT) and 20 % of trichloroacetic acid (TCA).
194 The mixture was then centrifuged for 10 min at 9,000 g. Afterwards, 600 μL of supernatant was
195 added to 480 μL of TRISbase (25 mM) - TBA (thiobarbituric acid – 100 mM) and 120 μL of
196 0.6N HCl and heated at 80°C for 15 min. Mixtures were subsequently cooled and mixed.
197 TBARS levels were read using a UV-spectrophotometer (Synergy HT, BioTek) in a microplate
198 at 530 nm. Results were expressed as nmoles of thiobarbituric acid reactive substance
199 (TBARS) equivalents/mg of protein.

200

201

202 *Carbonylated protein analysis*

203 Carbonylated protein content was measured using the method described in Augustyniak et al.
204 (2015). 50 μL of 11 % streptomycin sulfate – phosphate buffer (100 mM pH 7.4) was added to
205 500 μL of S9 fraction, mixed and incubated for 15 min at room temperature. Then, the mixture
206 was centrifuged for 10 min at 6,000 g. Afterwards, supernatant was divided into two tubes (200
207 μL each) where 200 μL of supernatant was added to 800 μL of HCl 2.5 M used as a control
208 tube, and 200 μL of supernatant was added to 800 μL of DNPH (2,4-dinitrophenylhydrazine
209 10 Mm) used as a sample tube. Subsequently, the mixture was incubated for 1 h at room
210 temperature with vortexing every 15 min. Proteins were precipitated with 1 mL of 20 % TCA
211 (trichloroacetic acid), vortexed and centrifuged for 10 min at 10,000 g. The pellets were rinsed
212 with 1 mL of ethanol-ethyl acetate (v:v), vortexed and centrifuged three times. Pellets were
213 then solubilized with 500 μL of 6 M guanidine HCl and centrifuged at 10,000 g for 10 min. The

214 carbonyl content was measured using a UV-spectrophotometer (Biotek Synergy HT) at
215 370 nm. Results were expressed as nmoles of DNPH incorporated/mg protein.

216

217 **2.1.7. Gene expression**

218 Six yolk-sac larvae per replicate were collected **individually** in a storage buffer (RNA later,
219 Qiagen). Samples were deep-frozen in liquid nitrogen and then stored at -80°C until analysis.

220 *RNA extraction*

221 Total RNA extraction from whole larvae was **done following** the kit “SV Total RNA Isolation
222 system” (Promega) **according to** the supplier’s recommendations. This kit included a DNaseI
223 treatment to avoid genomic DNA contamination of the samples. **All details of RNA extraction**
224 **are described in Weeks et al. (2019). For each exposure condition, samples were analyzed in**
225 **triplicate.**

226 *Retro-transcription of total RNA into cDNA*

227 The retro-transcription of total purified RNA was realized with the kit “GoScript Reverse
228 Transcription System” (Promega), **following the indications described at Weeks et al. (2019).**
229 The cDNA thus obtained were stored at -20°C pending analysis by quantitative real-time PCR
230 reaction.

231 *Quantitative real-time PCR*

232 Twelve genes were selected and specific primer-pairs were designed with primer3plus
233 software (Table 1). **All primer-pairs used in this study has an efficiency upper than 95 %.** Real-
234 time qPCR was carried out using GoTaq® qPCR Master Mix kit (Promega) **and was** performed
235 in a Mx3000P® qPCR System (Stratagene), **as fully described in Weeks et al. (2019).** **For each**
236 **reaction, specificity** of amplifications was determined from the dissociation curve of the PCR

237 products. This dissociation curve was obtained by following the SYBR Green fluorescence
238 level during a gradual heating of the PCR products from 60 to 95 °C.

239 Cycle thresholds (Ct) were obtained from MxPro™ qPCR software for each gene. Two different
240 housekeeping genes were used for standardization (*rpl7* and *ef1α*) and were found to be stable
241 in our conditions. Consequently, relative quantification of each gene expression level was
242 normalized according to the mean Ct value of these two reference genes and using the $2^{-\Delta\Delta Ct}$
243 methods (Livak and Schmittgen, 2001). The expression factor (induction if >2 and repression
244 if <0.5) of each gene was calculated for each condition by dividing the transcription level of
245 exposed individuals by that observed in control ones

246

247 **2.2. *In vitro* study using RTL-W1 cell line**

248 **2.2.1. Cell exposure**

249 The RTL-W1 cell line was obtained from rainbow trout liver (Lee et al., 1993). For cell culture,
250 L15 Leibovitz medium supplemented with 5 % FBS (Fetal Bovine Serum) and
251 1 % Penicillin/Streptomycin (100 IU/mL) was used. The cells were kept in polypropylene flasks
252 of 75 cm² (Cell start® cell culture Flask Greiner) at 20 °C. The analysis was carried out with
253 cells aged from passage from 65-72.

254 The cytotoxicity and genotoxicity test were carried out in 96- and 24-well polypropylene
255 microplates, respectively. For both MTT and comet assay, cell lines were seeded 24 h prior
256 glyphosate exposure in triplicate. Cell density was 200 000 cells/mL. For the MTT assay, cell
257 lines were **exposed** to concentrations from 0.05 to 1000 mg/L of glyphosate for 24 h, using
258 both technical and commercial formulation Roundup®. For the comet assay, the
259 concentrations tested were the same studied in our *in vivo* study (0.1 and 1mg/L of glyphosate)
260 using technical and commercial formulation of glyphosate.

261 **2.2.2. MTT assay for cytotoxicity evaluation**

262 The cytotoxicity test was performed using serum free L15 medium containing 10 % of 3(4,5-
263 dimethyl-2thiazholyl)-2,5-diphenyl-2H-tetrazoliumbromide (MTT). 24 h after chemical
264 exposure, the medium was removed and cells were rinsed with PBS. 100 µL of the MTT
265 solution was added to the wells. After incubation of 1 h in the dark (time to allow cells to reduce
266 tetrazolium to formazan), the MTT solution was removed and 100 µL of isopropanol solution
267 (4% 1N HCl) was added. Then the microplate was shaken horizontally in the dark for 15 min
268 to dissolve the formazan crystals. Following this step, the formazan coloration was quantified
269 in a Bio-Tek Synergy HT spectrophotometer at 570 nm.

270 **2.3. Genotoxicity test**

271 The alkaline comet assay was performed following Le Bihanic et al. (2014) and Weeks-Santos
272 et al. (2019) in blood cells of larvae, and RTL-W1 cell line following Pannetier et al., (2018).
273 Blood sampling was performed in 6 larvae per replicate (previously anesthetized with ice water
274 and few drops of carbonated water) by decapitation using a heparinized pipette. Samples were
275 stored in microtubes with 200 µL of cryopreservation solution (250 mM sucrose, 40 mM citrate
276 trisodique, 5 % DMSO, pH adjusted to 7.6 with nitric acid 1 M) and immediately frozen in liquid
277 nitrogen until analysis. The comet assay for the RTL-W1 cell line (1.5 to 2×10^5 cells/mL) was
278 performed 24 h after cell exposure to glyphosate. For each condition, 4 replicates were
279 prepared. The cells were rinsed, trypsinized and transferred into microtubes. The cells were
280 then centrifuged (5 min, 20 °C at 1000 rpm) and supernatant was removed. Cell pellets were
281 re-suspended in 100 µL of L15 medium (without FBS) before being mixed with 200 µL of low
282 melting point agarose (0.75 % LMPA). A 50 µL of cell suspension (blood and cell line) were
283 deposited on slides previously coated with NMPA (Normal Melting Point Agarose, 0.8% w/w)
284 and covered with an 18x18 mm coverslip. The slides were then immersed in a lysis solution
285 (10 mM Tris; 2.5 M NaCl; 100 mM EDTA; 1% Triton X-100; 10% DMSO; pH adjusted to 10
286 with NaOH) at 4°C for 90 min in the dark. At the end of the lysis, the slides were then rinsed 3
287 times for 5 min in an enzyme buffer at pH 8 (Biolabs, Evry, France). Then the slides were
288 immersed for 30 min in two hellendahls, the first one containing 60 mL of buffer with 12 µL of

289 the enzyme Fpg (Biolabs, Evry, France) diluted in 1/5000, and the second one with only buffer.
290 Following exposure to enzymes, the slides were incubated in an alkaline solution (0.3 M NaOH,
291 1 mM EDTA, pH > 13) at 4°C for 40 min for the RTL-W1 cells and 20 min for blood cells to
292 allow the DNA to unwind. Electrophoresis was then performed in the same solution at a voltage
293 of 25 V and 300 mA for 20 min. The slides were rinsed 3 times with neutralizing solution (0.4
294 M Tris, pH 7.5) for 5 min at 4°C. Afterwards, the slides were dehydrated in absolute ethanol
295 for 20 min and then allowed to dry at room temperature for at least 12 h. Slides were stained
296 with 20 µg/mL of ethidium bromide solution and covered with a 22x22 mm coverslip. Comet
297 analysis was carried out using an epifluorescence microscope (Olympus BX51) (zoom x20)
298 equipped with an Olympus U-RFL-T reflected fluorescence system lamp. The comets were
299 quantified using the Comet Assay IV software (Instrument Perspective Ltd). Results are
300 expressed as percentage of degradation of DNA tail for 100 randomly selected nuclei per slide.

301

302 **2.4. Statistics**

303 **Sampling of larvae (individuals and pools), from each exposure condition,** were performed in
304 triplicate and each replicate was considered as an independent sample. All data are expressed
305 by the mean ± SE (Standard Error). For the MTT test, the EC₅₀ was calculated by PRISM 5
306 software (GraphPad software, California, USA). Statistical analyzes were carried out using R
307 (<http://cran.r-projet.org/>). The Normality of data distribution was verified on the residues by the
308 Shapiro-Wilk test ($p < 0.01$) and the homogeneity of variances was evaluated by the Levene
309 test ($p < 0.05$). In the case of normal distribution, a one-way ANOVA analysis was used
310 ($p < 0.05$) followed by a Tukey post-hoc test. In the case that data was not normal,
311 comparisons were carried out by non-parametric tests of Kruskal-Wallis ($p < 0.05$).

312

313 **3. Results**

314 **3.1. Exposure conditions**

315 Table 2 shows the concentrations of glyphosate in water for each experimental conditions. The
316 analyses were carried out at 0 and 48 hours after exposure to estimate the possible losses of
317 the molecule. The measured concentrations of glyphosate were comparable ($\pm 20\%$) to the
318 nominal concentrations. No concentration variation was noted during 48h. The glyphosate's
319 metabolites, aminomethylphosphonic acid (AMPA) were also analyzed but not detected at T0
320 and T48.

321 **3.2. Embryonic and larval survival**

322 Dissolved oxygen in the exposure water varied between 83.8 and 93% throughout the duration
323 of this study. Exposure to 0.1 and 1 mg/L of glyphosate (a.i.) did not induce significant mortality
324 in trout embryos and larvae throughout the duration of exposure (table 3). Both, embryonic
325 and larval survival, were greater than 90 % in all studied conditions. All embryos hatched
326 successfully. The duration of development was slightly longer for both groups exposed to
327 glyphosate compared to control, however no significant differences were observed.

328 **3.3. Biometry**

329 No significant differences were observed in total larval length between the studied conditions
330 and the control (Figure 1-A). Nevertheless, measurement of larvae head size showed
331 significant decreases in larvae exposed to 1 mg/L of glyphosate compared to control (Figure
332 1-B). Head size in unexposed larvae was 4.76 ± 0.04 mm against 4.55 ± 0.11 mm for larvae
333 exposed to 0.1 mg/L of glyphosate, and 4.43 ± 0.14 mm on larvae exposed to
334 1 mg/L of glyphosate. The ratio between total length and head size (Figure 1-C) showed a
335 significant dose-dependent decrease from control ($24.79 \pm 0.14\%$) and larvae exposed to
336 0.1 mg/L of glyphosate ($24.14 \pm 0.27\%$) and 1 mg/L of glyphosate ($23.37 \pm 0.3\%$).

337 **3.4. Malformations**

338 Embryo–larval exposure to glyphosate did not result in significant induction of malformation
339 when compared to non-exposed larvae. Control condition presented **13.3 ± 6.7 %** of

340 malformed larvae. However, larvae exposed to 1 mg/L of glyphosate (a.i.) showed a significant
341 increase in developmental anomalies over larvae exposed to 0.1 mg/L of glyphosate (a.i.) with
342 $26.7 \pm 6.7 \%$ and $8.9 \pm 3.8 \%$ respectively (Figure 2).

343 **3.5. Swimming behavior**

344 Figure 3 (A and B) shows the responses of larvae to light stimulation. Results represent the
345 average speed of larvae exposed to glyphosate with alternating periods of luminosity. Under
346 each condition, the same tendency was observed with an increase in larval velocity during the
347 light period. No significant differences were observed at the first period of darkness when
348 comparing the different treatments (Figure 3-A and B). When the light was turned on, the stress
349 caused an increase in the average speed of the larvae exposed to 0.1 mg/L of glyphosate with
350 a pic of $29.2 \pm 2.3 \text{ cm/s}$ when compared to control and larvae exposed to 1 mg/L of glyphosate
351 (22.4 ± 1.5 and $23.1 \pm 3.1 \text{ cm/s}$ respectively) (Figure 3-B). However, after 4 min of light exposure,
352 this increase of velocity was no longer different for larvae exposed to 0.1 mg/L of glyphosate
353 when compared to other conditions (Figure 3-A). Likewise, no significant differences were
354 observed at the second dark period.

355 Figure 4 shows the average cumulative time of immobility, mobility and high mobility for larvae
356 exposed to both glyphosate conditions and control. Larvae exposed to 0.1 mg/L of glyphosate
357 were significantly highly mobile ($8.04 \pm 1.25 \text{ s}$) in the light period when it was compared to
358 control ($4.72 \pm 0.63 \text{ s}$) and larvae exposed to 1 mg/L of glyphosate ($4.19 \pm 0.38 \text{ s}$).

359 **3.6. Genotoxicity in blood cells**

360 The average level of DNA damage for each studied condition, with and without treatment by
361 Fpg is presented in Figure 5. No significant differences were observed in DNA damage in all
362 conditions when cells were not treated with Fpg enzyme ($6.85 \pm 2.11 \%$ for control,
363 $8.52 \pm 2.33 \%$ for 0.1 mg/L of glyphosate condition, and $7.28 \pm 1.69 \%$ for 1 mg/L of glyphosate
364 condition). A global increase of DNA damage was observed after Fpg treatment but no

365 significant differences were observed between conditions (20.86 ± 3.73 % for control
366 condition, 22.37 ± 2.12 % for larvae exposed to 0.1 mg/L of glyphosate and 19.88 ± 1.02 % for
367 larvae exposed to 1 mg/L of glyphosate).

368

369

370 **3.7. Lipid peroxidation (TBARS) and protein carbonyls**

371 TBARS levels showed a significant reduction in larvae exposed to 0.1 mg/L of glyphosate when
372 compared to control (figure 6-A). In the other hand, larvae exposed to glyphosate did not show
373 any significant changes in protein carbonyls (figure 6-B).

374 **3.8. Gene expression**

375 After 3-week exposure of rainbow trout to glyphosate, only a handful of significant changes
376 were observed on gene expression on larvae exposed to 1 mg/L. *Cox1* gene was significantly
377 down-regulated (0.22) when *cat* gene level was increased (2.13). The expression of *sod*, *gst*,
378 *ERb*, *12s*, *ogg1*, *rad51*, *bax* and *Arom* were not significantly differentially regulated following
379 glyphosate exposure (data not showed).

380 **3.9. Cytotoxicity**

381 The cytotoxicity data for glyphosate and Roundup® (a.i.) was obtained using the MTT assay
382 on RTL-W1 (Figure 7). Cytotoxicity was observed only at concentrations above 250 mg/L of
383 glyphosate, and 200 mg/L of Roundup® (a.i.). The EC_{50} calculated at 24 h was 730 and
384 710 mg/L for glyphosate and Roundup® (a.i.), respectively.

385 **3.10. Genotoxicity in RTL-W1 cell line**

386 With the standard comet assay, no genotoxic effect was detected after exposure to both
387 glyphosate and Roundup® whatever the tested concentrations. However, with the modified

388 Fpg assay, significant genotoxic were observed on RTL-W1 cell line exposed to 0.1 and 1 mg/L
389 of technical glyphosate with 33.6±3.1 and 33.5±3.2% of DNA damage, respectively, when
390 compared to control condition with 25.4 ± 2.9 % of DNA damage (Figure 8). The same was
391 observed using Roundup® formulation where significant DNA damage was at 26.8 ± 1.5 and
392 23.9 ± 2.3 % for cells exposed to 0.1 and 1 mg/L of Roundup® (a.i.) when compared to control
393 with 17.9 ± 2.1 % of DNA damage (Figure 8).

394

395 4. Discussion

396 According to the World Health Organization (WHO, 1996), the acute toxicity of Roundup is
397 considered to be low in vertebrates. Because of its widespread use, and its slow degradation,
398 this herbicide is often found in aquatic environments at relatively high concentrations (Vera et
399 al., 2010) and thus could represent a threat for pollutant-sensitive species or early life stages
400 (ELS). Several authors have studied the acute toxicity of glyphosate on ELS, fingerlings and
401 adults of rainbow trout (Folmar et al., 1979; Hildebrand et al., 1982; Morgan and Kiceniuk, 1992;
402 Anton et al., 1994). 96h LC₅₀ for rainbow trout embryos and larvae was estimated to 16 and
403 3.4 mg/L glyphosate (a.i.) respectively (Folmar et al., 1979). However, acute toxicity varies
404 according to the commercial formulation. For example, 96 h LC₅₀ on rainbow trout fingerlings
405 was estimated to be 54.8 mg/L using Roundup® formulation (Hildebrand et al., 1982); and
406 10.4 mg/L using Vision formulation (Morgan and Kiceniuk, 1992). The work of Yusof et al.
407 (2014) focused on glyphosate toxicity on Java medaka. Their results showed that 50 % of
408 embryos exposed to 100 mg/L of glyphosate died after 16 days of exposure, and a decrease
409 on hatching rate in a concentration-dependent manner from 100 to 500 mg/L of glyphosate.

410 The *in vitro* study analyzed the toxicity of glyphosate using the rainbow trout liver cell line (RTL-
411 W1) considering technical grade glyphosate and its commercial formulation Roundup. The
412 toxicity test carried out on trout liver cells may provide additional information about the toxicity
413 mechanistic of pollutants (Bols et al., 2005; Castaño et al., 2003). The RTL-W1 cell line can

414 be considered a suitable model, given that the liver is the main organ responsible for
415 metabolising pollutants (Belpaeme et al., 1998). The results obtained on RTL-W1 in this study
416 highlight the cytotoxic effects of glyphosate, but at high concentrations above 200 mg/L. Our
417 results also indicate that the commercial formulation is slightly more cytotoxic than the
418 technical grade compound, which could be related to the presence of additives, especially
419 surfactants (POEA) in the commercial formulation. Similar studies on human cell lines have
420 shown that glyphosate-based formulations are usually more cytotoxic than the technical grade
421 compound (Gasnier et al., 2009; Koller et al., 2012; Martínez et al., 2007; Mesnage et al., 2013;
422 Vanlaeys et al., 2018). In addition, the study of Gasnier et al. (2009) evidenced that the
423 concentration of glyphosate in the commercial formulation is not related to toxicity. Indeed, the
424 formulation containing 400 g/L of glyphosate (a.i.) (Grands Travaux®, homologation 8800425)
425 has a lower LC₅₀ than its homolog containing 450 g/L (Grands Travaux plus®, homologation
426 2020448) confirming that the nature and concentration of adjuvants have a real impact on the
427 toxicity of the mixture. Very few studies have been done on fish cell lines regarding the toxic
428 effects of glyphosate. The LC₅₀ of glyphosate on diploid and triploid fin cell lines from
429 *Misgurnus anguillicaudatus* (DIMF and TRMF) were 315.34 and 371.77 mg/L respectively (Qin
430 et al., 2017). Cytotoxicity of Roundup was also studied on zebrafish cell line ZF-L regarding
431 the integrity of the plasma membrane, mitochondrial activity and lysosomal integrity. The
432 authors reported a significant reduction of cell viability from 67.7 µg/L (a.i.) (Goulart et al.,
433 2015). LC₅₀ of mononuclear blood cells was determined at 56.4 mg/L for Roundup, and
434 1630 mg/L for technical grade glyphosate (Martínez et al., 2007). These differences of toxicity
435 might depend on the concentration of the active agent but also the nature and concentration
436 of its adjuvants, as well as the cell line used.

437 In the literature, there are few studies concerning the effects of glyphosate on fish growth and
438 the findings are often inconsistent. Rainbow trout fingerlings exposed up to 100 µg/L of
439 glyphosate (a.i.) using Vision formulation (Monsanto Co.) did not show significant effect on
440 length or weight after two months of exposure (Morgan and Kiceniuk, 1992). *Leporinus*

441 *obtusidens*, a South American fish species, was exposed to 1 and 5 mg/L of glyphosate (a.i.)
442 using Roundup for 90 days and exhibited a lower growth rate (with reductions between 10 and
443 15 % respectively) and a lower weight gain (between 44 and 65 % respectively) when
444 compared to control fish (Salbego et al., 2010). Similarly, Bridi et al. (2017) reported a reduced
445 body length in zebra fish larvae (*Danio rerio*) exposed from 0.01 to 0.5 mg/L of Roundup (a.i.)
446 for 96 h. Koakoski et al., (2014) also observed a reduction of the weight gain and biomass of
447 *Rhamdia quelen* fingerlings when exposed to 1.21 mg/L of Roundup for 96 h and after 180
448 days of depuration. ~~Another study using adult fishes (*Piaractus Mesopotamicus*) reported that~~
449 ~~glyphosate reduced food intake and therefore could have an impact on normal growth~~
450 ~~(Cardoso Giaquinto et al., 2017).~~ Furthermore, some authors have stated that glyphosate may
451 have an effect on growth hormones and cortisol levels in fish (Cericato et al., 2008; El-Shebly
452 and El-kady, 2008; Koakoski et al., 2014). Cericato et al. (2008) observed that cortisol levels
453 in fish exposed to glyphosate were higher than in non-exposed fish. Indeed, cortisol is released
454 in response to stress and contributes to restore homeostasis (De Boeck et al., 2001), and
455 some evidence suggest that elevation of cortisol might interfere with normal growth of fishes
456 by stimulating energy-consuming processes (Bernier et al., 2004; De Boeck et al., 2001). In
457 our study, a 3-week exposure of rainbow trout embryos to 0.1 and 1 mg/L glyphosate did not
458 induce significant reductions in total body length of larvae. However, head length of larvae was
459 significantly smaller for those exposed to the highest tested concentration, and the ratio of
460 head to total body length showed a significant decrease in a concentration-dependent manner.
461 Interestingly, Zebral et al. (2017) evaluated eye diameter and distance between eyes in
462 pejerrey embryos (*Odontesthes humensis*) exposed to this herbicide (0.36-5.43 mg/L) for 96 h
463 and observed that both parameters were significantly reduced in a concentration-dependent
464 manner in exposed groups. Similar results were found by Zhang et al. (2017) in zebra fish
465 embryo (*D. rerio*) but using higher concentrations (up to 400 mg/L) of glyphosate for 96 h.
466 Zebral et al. (2017) suggested that glyphosate might alter the retinoic acid pathway, which
467 plays a major role in growth and development. Paganelli et al. (2010) also indicated that
468 glyphosate produces teratogenic effects on vertebrates by impairing retinoic acid signaling.

469 ~~Our results showed a trend of increasing spinal deformities when rainbow trout embryos were~~
470 ~~exposed to 1 mg/L of glyphosate. Several studies have reported significant body~~
471 ~~malformations, spinal curvature, pericardial and yolk sac edemas on embryos of zebra fish~~
472 ~~(Sulukan et al., 2017; Zhang et al., 2017) and Java medaka (Yusof et al., 2014) using relatively~~
473 ~~high concentrations of Roundup® from 1 to 500 mg/L (a.i.).~~

474 Larvae exposed to 0.1 mg/L of Roundup® (a.i.) were more active under light stimulation.
475 Several previous studies have also examined the effects of glyphosate on fish swimming
476 behaviour. In concordance with our results, Morgan et al., (1991) observed that after one-
477 month exposure to 45.75 µg/L of glyphosate, under Vision's commercial formulation, fry
478 rainbow trout presented erratic and agitated behaviour compared to unexposed fish. Similar
479 abnormal behaviours and hyperactivity were also reported in Nile tilapia (Ayoola, 2008) and
480 *Tilapia zillii* (Nwani et al., 2013) exposed from 2 to 310 mg/L for 4 days and from 216 to
481 540 mg/L of glyphosate for 96 h, respectively. A Neotropical hybrid fry fish, surubim, showed
482 increased swimming activity and ventilation frequency 96 h after exposure to 7.5 and 15 mg/L
483 of Roundup® (a.i.) (Sinhorin et al., 2014). In the other hand, Bridi et al., (2017) observed that
484 zebrafish larvae and adults exhibited significant reduction of distance travelled and mobility
485 when exposed to glyphosate and Roundup® formulations (0.01 to 0.5 mg/L a.i.) for 96 h. The
486 behavioural study of this work was performed in larvae after 21 days of glyphosate exposure.

487 **The absence of behavioural changes at the dark period could mean an adaptation of response**
488 **to stress.** It was shown that **sub**-chronic exposure to low concentrations of glyphosate
489 (0.1 mg/L a.i.) induced **an increase in swimming behaviour** in exposed rainbow trout larvae but
490 no effect on swimming activity was observed at 1 mg/L. **This apparent hyperactivity decreased**
491 **4 min later of light exposure. Same patrons were observed by Zhang et al. (2017) where**
492 **locomotive activities in day time of zebrafish larvae, exposed to low concentrations of**
493 **glyphosate (0.01 and 0.5 mg/L a.i.) were increased; however, at stronger concentrations**
494 **(5 mg/L a.i.) this increase was no longer observed when compared to non-exposed larvae.**

495 These alterations may have a consequence in the response face to predators or other danger
496 (Zhang et al., 2017).

497 In this study, the use Fpg-modified comet assay improved detection threshold for DNA
498 damage. The standard comet assay can detect single or double strand breaks and alkali-labile
499 sites, while the addition of Fpg enzyme can also detect lesions related to alkylation damage,
500 abasic sites (apuric or apyrimidic) and oxidative damage (8-oxoGua) induced by ROS
501 (Reactive Oxygen Species) production (Kienzler et al., 2012). In our exposure conditions,
502 glyphosate did not induce any DNA strand breaks on blood cells of rainbow trout larvae after
503 21 days of exposure. However, some studies have demonstrated the genotoxic potential of
504 Roundup® in different fish species like *Anguilla anguilla* (Guilherme et al., 2012, 2010),
505 *Corydoras paleatus* (De Castilhos Ghisi and Cestari, 2013), *Prochilodus lineatus* (Moreno et
506 al, 2014) and *Carassius auratus* (Çavaş and Könen, 2007). Guilherme et al., (2010) showed
507 Roundup®'s capacity to induce DNA single strand breaks and cytogenetic effects on blood
508 cells of European eel using low concentrations (58 and 116 µg/L a.i.) after 1 and 3 days of
509 exposure. Cavalcante et al., (2010) observed genotoxic potential of Roundup® on blood and
510 gill cells after 6 h of exposure to 10 mg/L (a.i.) on fish (*Prochilodus lineatus*), but DNA damage
511 returned to the baseline level after 24 and 96 h of exposure for erythrocytes and gill cells
512 respectively. The activation of the antioxidant and DNA repair systems after glyphosate
513 exposure have already been demonstrated by Cavalcante et al., (2010) and Marques et al.,
514 (2014). In our case, we may assume ROS were produced but larvae were able to activate
515 protective mechanisms such as DNA repair enzymes to prevent DNA damage on blood cells,
516 as reported in several articles (Marques et al., 2014; Ching et al., 2001; Kienzler et al., 2013).

517 On the other hand, Fpg-modified comet assay in RTL-W1 cell line indicated that both technical
518 grade glyphosate and Roundup® induced abasic sites and oxidative DNA damage at
519 concentrations of 0.1 and 1 mg/L (a.i.), but no significant increase in DNA damage was
520 observed with the classical comet assay. Observing a genotoxic on RTL-W1 (short exposure),
521 and not on larvae (longer exposure) favours the hypothesis of the activation of *in vivo* repair

522 systems. However, we must be cautious with this comparison because the studied cells are
523 not the same *in vivo* and *in vitro*. No genotoxicity studies of glyphosate have been performed
524 on RTL-W1 cell line. Using the human hepatoma cell line, HepG2, no DNA damage was
525 observed when glyphosate was tested as a pure form after an exposure of 4 h (Kašuba et al.,
526 2017). In human buccal epithelial cells, TR146, glyphosate and Roundup induced DNA
527 damage from 20 mg/L and DNA damage increased as a function of the exposure concentration
528 (Koller et al., 2012). Differences in genotoxicity activity were observed between *in vitro* and *in*
529 *vivo* exposure in tilapia erythrocytes after exposure to glyphosate (a.i.) (0.0007 - 0.7 mM)
530 (Alvarez-Moya et al., 2014). *In vitro*, DNA damage was proportional to glyphosate
531 concentration; however, *in vivo*, glyphosate was genotoxic to fish erythrocytes but not in a
532 concentration-dependent manner.

533 Malondialdehyde (MDA) is one of the secondary products that can be formed during lipid
534 peroxidation of uncontrolled oxidative stress in cells (Ayala et al., 2014). It is considered as the
535 most mutagenic product of lipid peroxidation, and once formed, MDA can react with proteins
536 or DNA to form adducts resulting in biomolecular damage (Ayala et al., 2014). Because of its
537 easy reaction with thiobarbituric acid (TBA), MDA has been used as a convenient biomarker
538 of lipid peroxidation using the thiobarbituric acid reacting substances test (TBARS) (Ayala et
539 al., 2014). Lipid peroxidation (LPO) has already been studied in fish exposed to glyphosate
540 based herbicides and results might be very variable according to fish species, exposure
541 duration (Gluszczak et al., 2007; Modesto and Martinez, 2010; Sinhorin et al., 2014), gender
542 (Harayashiki et al., 2013) and tissues (Gluszczak et al., 2007; Sinhorin et al., 2014). Juveniles
543 of *Prochilodus lineatus* have significantly increased LPO levels in liver after 6 h of exposure to
544 both 1 and 5 mg/L of Roundup Transorb. However, these alterations returned to control levels
545 after 24 h of exposure (Modesto and Martinez, 2010). On the other hand, Gluszczak et al. (2007)
546 did not observed TBARS alterations in liver of silver catfish (*Rhamdia quelen*) when exposed
547 to 0.2 and 0.4 mg/L, but they did in muscle tissue at both concentrations. Ferreira et al. (2010)
548 also studied the oxidative stress of different pesticides in silver catfish finding that methyl

549 ~~parathion and tebuconazole but glyphosate enhanced TBARS levels in liver of fish.~~ The hybrid
550 amazon fish surubim had significantly increased TBARS levels in both liver and muscle, but
551 not in the brain after exposure to 2.25 to 15 mg/L of Roundup (Sinhorin et al., 2014). Even
552 though several authors have studied TBARS levels in fish exposed to glyphosate, only few
553 analyses have been done on whole larvae. Our results show that TBARS levels were reduced
554 in whole larvae exposed to 0.1 mg/L of glyphosate when compared to control group. Fish have
555 a natural anti-oxidative defense system against free radicals, and are able to reduce oxidative
556 damage to below control levels (Marques et al., 2014). As hypothesized by Marques et al.
557 (2014), a development of antioxidant systems may occur as a response to ROS, reducing the
558 vulnerability of cells and their constituents. Reduced levels of lipid peroxidation have already
559 been observed in the livers of male guppy exposed to 700 µg/L of Roundup (a.i.) (Harayashiki
560 et al., 2013), in brain of piava fish (*Leporinus obtusidens*) exposed from 3 to 20 mg/L of
561 glyphosate commercial formulation (a.i.) (Gluszczak et al., 2011). Lipid peroxidation may not
562 only depend on ROS production, but may be also be affected by physiological transitions that
563 occur at different developmental stages (Cao et al., 2010; Mourente et al., 1999). The presence
564 of carbonyl groups in proteins induced by glyphosate was also studied in several reports (de
565 Moura et al., 2017; Gluszczak et al., 2011; Sinhorin et al., 2014) generally in liver since it is
566 consider as the main site of protein carbonyl production (Sinhorin et al., 2014). In contrast, the
567 absence of protein carbonyl changes in our results could also indicate, once again, that the
568 antioxidant system of rainbow trout larvae functions efficiently to defend against oxidative
569 stress. As for TBARS, only a few analyses have been done using whole fish larvae to analyze
570 carbonyl groups in proteins. Considering that protein carbonyl formation is non-reversible
571 (Zhang et al., 2008), it can be suggested that at this developmental stage of larvae, ROS
572 formation in rainbow trout larvae exposed to low or moderate concentrations of glyphosate
573 was weak or low enough to be detoxified by the antioxidant systems causing no changes in
574 TBARS and protein carbonyls groups.

575 Among the enzymes involved in ROS detoxification are SOD (superoxide dismutase), CAT
576 (catalase) and GST (glutathione-S-transferase). Inhibition of CAT and SOD activities in liver
577 were observed following exposure to glyphosate by Ferreira (2010) in silver catfish, Modesto
578 and Martinez (2010) in *Prochilodus lineatus* and by Sinhorin (2014) in surubim
579 (*Pseudoplatystoma sp.*). In contrast, CAT activity was induced in liver of *L. obtusidens* exposed
580 up to 6 mg/L of Roundup® (a.i.). We observed that *cat* gene was significantly repressed in
581 larvae exposed to 1 mg/L of glyphosate. Topal et al. (2015) studied both gene expression and
582 enzymatic activity in liver of juvenile rainbow trout exposed to different concentrations of
583 glyphosate (from 2.5 to 10 mg/L) from 6 to 96 h, observing that the expressions of *cat* and *sod*
584 were induced the first 6 h and then significantly decreased after 24 h of exposure. In the same
585 study, Topal et al. (2015), observed that the trend of the antioxidant enzymes activity of
586 catalase was opposed to the level of gene expression.

587 Interestingly, Webster and Santos (2015) studied the transcriptional profile, using RNA-seq, of
588 brown trout females exposed to glyphosate and Roundup (0.01, 0.5 and 10 mg/L) for 14 days.
589 They identified differentially expressed genes that encode antioxidant system proteins (up-
590 regulation of glutathione reductase, *gsr*) stress-responses proteins (heat shock proteins, *ddit*,
591 *ddit4l* and *gadd4l*) and pro-apoptotic signalling proteins (transcription factor tumour suppressor
592 protein *p53*). The nature of the response of the cell depends on the amount and the duration
593 of the stress, since cells respond in a variety of signalling pathways (Fulda et al., 2010; Webster
594 and Santos, 2015). According to Webster and Santos (2015), low concentrations of ROS may
595 help to induce pro-survival signalling, while higher levels of oxidative stress and cellular
596 damage might activate cell death signalling pathways as a protective mechanism. In addition,
597 in this same study (Webster and Santos, 2015), few changes in pro-apoptotic factors were
598 observed suggesting a pro-survival stress response at lower concentrations of glyphosate
599 producing low levels of oxidative stress.

600 The *cox1* gene code the cytochrome c oxidase subunit 1, which is one of the enzymes
601 involved in the respiratory electron chain transport in mitochondrial membrane. The

602 mitochondrial electron-transport chain is the main source of ROS during normal metabolism
603 (Chen et al., 2003). While cytochrome oxidase is not a source of ROS, its inhibition may
604 promote ROS production (Chen et al., 2003). Our results revealed a significant induction of
605 *cox1* (x2) gene expression on whole larvae exposed to 1 mg/L of glyphosate. An induction of
606 *cox1* could be a cell response to maintain respiratory chain function (Arini et al., 2015).
607 Induction of *cox1* gene could be viewed as a mechanism by which to restore mitochondria
608 activity and to efficiently consume O₂ and thus to limit ROS production. Induction of *cox1* gene
609 expression could be considered as a mechanism to avoid ROS production (Achard-Joris et al.,
610 2006).

611

612 **Conclusions**

613 This study provides an extensive evaluation of the toxicological effects of glyphosate using an
614 *in vivo* and *in vitro* approach. Results revealed that relatively low concentrations of glyphosate
615 induced hyperactive swimming behavior and morphological cranio-facial alterations on larvae.
616 In parallel, the studied cell line, RTL-W1, exhibited a DNA damage, which were not observed
617 in blood cells from exposed larvae using the same concentrations of glyphosate. This
618 difference may be explained by the duration of exposure, which was longer, and could have
619 led to an activation of the antioxidant and DNA repair system on blood cells. Decreased TBARS
620 levels and the differential regulation of *cat* and *cox1* gene expression observed on whole
621 exposed larvae could also confirm this hypothesis. It is important to consider the adjuvants in
622 commercial formulations, which can increase the toxicity of glyphosate for vertebrates, and not
623 only the active compound. Regarding the toxicity of glyphosate highlighted in rainbow trout
624 ELS at concentrations that can be found in aquatic ecosystems, we can conclude that
625 glyphosate can pose a potential risk for the most sensitive stage of fish.

626

627 **Acknowledgements**

628 This work was supported by grants from SENACYT (Secretaría Nacional de Ciencia y
629 Tecnología e Innovación) as a part of a PhD program. We want to thank Laurent Labbé and
630 Lionel Goardon (PEIMA, Sizun) for supplying rainbow trout embryos and for all the
631 recommendations that they have given us. Brigitte Delest (Irstea) is also thanked for her help
632 in preparing the pesticide samples prior to analysis. This work was part of the LABEX COTE
633 cluster of excellence “Continental to Coastal Ecosystems: evolution, adaptability and
634 governance”. The authors are grateful to James Emery for providing English proofreading
635 correction.

636

637 **References**

- 638 Achard-Joris, M., Gonzalez, P., Marie, V., Baudrimont, M., Bourdineaud, J.P., 2006.
639 Cytochrome c oxydase subunit I gene is up-regulated by cadmium in freshwater and
640 marine bivalves. *BioMetals* 19, 237–244. <https://doi.org/10.1007/s10534-005-5671-9>
- 641 Alvarez-Moya, C., Silva, M.R., Valdez Ramírez, C., Gallardo, D.G., León Sánchez, R.,
642 Aguirre, A.C., Velasco, A.F., 2014. Comparison of the in vivo and in vitro genotoxicity of
643 glyphosate isopropylamine salt in three different organisms. *Genet. Mol. Biol.* 37, 105–
644 110. <https://doi.org/10.1590/S1415-47572014000100016>
- 645 Anton, F., Laborda, E., de Ariz, M., 1994. Acute toxicity of the herbicide glyphosate to fish.
646 *Chemosphere* 28, 745–753.
- 647 Arini, A., Gourves, P.Y., Gonzalez, P., Baudrimont, M., 2015. Metal detoxification and gene
648 expression regulation after a Cd and Zn contamination: An experimental study on *Danio*
649 *rerio*. *Chemosphere* 128, 125–133. <https://doi.org/10.1016/j.chemosphere.2015.01.022>
- 650 Augustyniak, E., Adam, A., Wojdyla, K., Rogowska-Wrzesinska, A., Willetts, R., Korkmaz, A.,
651 Atalay, M., Weber, D., Grune, T., Borsa, C., Gradinaru, D., Chand Bollineni, R.,
652 Fedorova, M., Griffiths, H.R., 2015. Validation of protein carbonyl measurement: A multi-
653 centre study. *Redox Biol.* 4, 149–157. <https://doi.org/10.1016/j.redox.2014.12.014>
- 654 Ayala, A., Muñoz, M.F., Argüelles, S., 2014. Lipid Peroxidation: Production, Metabolism, and
655 Signaling Mechanisms of Malondialdehyde and 4-Hydroxy-2-Nonenal. *Oxid. Med. Cell.*
656 *Longev.* 2014, 1–31. <https://doi.org/10.1155/2014/360438>

- 657 Ayoola, S.O., 2008. Toxicity of glyphosate herbicide on Nile tilapia (*Oreochromis niloticus*)
658 juvenile. *African J. Agric. Res.* 3, 825–834.
- 659 Belpaeme, K., Cooreman, K., Kirsch-Volders, M., 1998. Development and validation of the in
660 vivo alkaline comet assay for detecting genomic damage in marine flatfish. *Mutat. Res. -*
661 *Genet. Toxicol. Environ. Mutagen.* 415, 167–184. [https://doi.org/10.1016/S1383-](https://doi.org/10.1016/S1383-5718(98)00062-X)
662 [5718\(98\)00062-X](https://doi.org/10.1016/S1383-5718(98)00062-X)
- 663 Bernier, N.J., Bedard, N., Peter, R.E., 2004. Effects of cortisol on food intake, growth, and
664 forebrain neuropeptide Y and corticotropin-releasing factor gene expression in goldfish.
665 *Gen. Comp. Endocrinol.* 135, 230–240. <https://doi.org/10.1016/j.ygcen.2003.09.016>
- 666 Bols, N., Dayeh, V.R., Lee, L.E.J., Schirmer, K., 2005. Use of fish cell lines in the toxicology
667 and ecotoxicology of fish. *Biochem. Mol. Biol. Fishes* 6, 43–85.
- 668 Bridi, D., Altenhofen, S., Gonzalez, J.B., Reolon, G.K., Bonan, C.D., 2017. Glyphosate and
669 Roundup® alter morphology and behavior in zebrafish. *Toxicology* 392, 32–39.
670 <https://doi.org/10.1016/j.tox.2017.10.007>
- 671 Cao, L., Huang, W., Liu, J., Yin, X., Dou, S., 2010. Accumulation and oxidative stress
672 biomarkers in Japanese flounder larvae and juveniles under chronic cadmium exposure.
673 *Comp. Biochem. Physiol. - C Toxicol. Pharmacol.* 151, 386–392.
674 <https://doi.org/10.1016/j.cbpc.2010.01.004>
- 675 Cardoso Giaquinto, P., Bordes de Sa, M., Seiko Sugihara, V., Bastos Gonçalves, B.B.,
676 Delicio, H.C., Barki, A., 2017. Effects of Glyphosate-Based Herbicide Sub-Lethal
677 Concentrations on Fish Feeding Behavior. *Bull. Environ. Contam. Toxicol.* 98, 460–464.
678 <https://doi.org/10.1007/s00128-017-2037-2>
- 679 Castaño, A., Bols, N., Braunbeck, T., Dierickx, P., Halder, M., Isomaa, B., Kawahara, K.,
680 Lee, L.E.J., Mothersill, C., Pärt, P., Repetto, G., Sintes, J.R., Rufli, H., Smith, R., Wood,
681 C., Segner, H., 2003. The Use of Fish Cells in Ecotoxicology. *Rep. Recomm. ECVAM*
682 *Work.* 47 ATLA 31, 317–351.
- 683 Castro Berman, M., Marino, D.J.G., Quiroga, M.V., Zagarese, H., 2018. Occurrence and
684 levels of glyphosate and AMPA in shallow lakes from the Pampean and Patagonian
685 regions of Argentina. *Chemosphere* 200, 513–522.
686 <https://doi.org/10.1016/j.chemosphere.2018.02.103>
- 687 Cavalcante, D.G.S.M., Martinez, C.B.R., Sofia, S.H., 2010. Genotoxic effects of Roundup®
688 on the fish *Prochilodus lineatus*. *Mutat. Res. - Genet. Toxicol. Environ. Mutagen.* 695,
689 41–46. <https://doi.org/10.1016/j.mrgentox.2008.06.010>

690 Çavaş, T., Könen, S., 2007. Detection of cytogenetic and DNA damage in peripheral
691 erythrocytes of goldfish (*Carassius auratus*) exposed to a glyphosate formulation using
692 the micronucleus test and the comet assay. *Mutagenesis* 22, 263–268.
693 <https://doi.org/10.1093/mutage/gem012>

694 Cericato, L., Neto, J.G.M., Fagundes, M., Kreutz, L.C., Quevedo, R.M., Finco, J., da Rosa,
695 J.G.S., Koakoski, G., Centenaro, L., Pottker, E., Anziliero, D., Barcellos, L.J.G., 2008.
696 Cortisol response to acute stress in jundiá *Rhamdia quelen* acutely exposed to sub-
697 lethal concentrations of agrichemicals. *Comp. Biochem. Physiol. - C Toxicol.*
698 *Pharmacol.* 148, 281–286. <https://doi.org/10.1016/j.cbpc.2008.06.008>

699 Chen, Q., Vazquez, E.J., Moghaddas, S., Hoppel, C.L., Lesnefsky, E.J., 2003. Production of
700 reactive oxygen species by mitochondria: Central role of complex III. *J. Biol. Chem.* 278,
701 36027–36031. <https://doi.org/10.1074/jbc.M304854200>

702 Ching, E.W.K., Siu, W.H.L., Lam, P.K.S., Xuà, L., Zhang, Y., Richardson, B.J., Wu, R.S.S.,
703 2001. DNA Adduct Formation and DNA Strand Breaks in Green-lipped Mussels (*Perna*
704 *viridis*) Exposed to Benzo [a] pyrene : Dose- and Time-Dependent Relationships. *Mar.*
705 *Pollut. Bull.* 42, 603–610.

706 Coupe, R.H., Kalkhoff, S.J., Capel, P.D., Gregoire, C., 2012. Fate and transport of
707 glyphosate and aminomethylphosphonic acid in surface waters of agricultural basins.
708 *Pest Manag. Sci.* 68, 16–30. <https://doi.org/10.1002/ps.2212>

709 De Boeck, G., Alsop, D., Wood, C., 2001. Cortisol Effects on Aerobic and Anaerobic
710 Metabolism , Nitrogen Excretion , and Whole-Body Composition in Juvenile Rainbow
711 Trout. *Physiol. Biochem. Zool.* 858–868.

712 De Castilhos Ghisi, N., Cestari, M.M., 2013. Genotoxic effects of the herbicide Roundup®in
713 the fish *Corydoras paleatus* (Jenyns 1842) after short-term, environmentally low
714 concentration exposure. *Environ. Monit. Assess.* 185, 3201–3207.
715 <https://doi.org/10.1007/s10661-012-2783-x>

716 de Moura, F.R., Brentegani, K.R., Gemelli, A., Sinhorin, A.P., Sinhorin, V.D.G., 2017.
717 Oxidative stress in the hybrid fish jundiara (*Leiarius marmoratus* × *Pseudoplatystoma*
718 *reticulatum*) exposed to Roundup Original®. *Chemosphere* 185, 445–451.
719 <https://doi.org/10.1016/j.chemosphere.2017.07.030>

720 El-Shebly, A.A., El-kady, M.A.H., 2008. Effects of glyphosate herbicide on serum growth
721 hormone (GH) levels and muscle protein content in Nile Tilapia (*Oreochromis niloticus*
722 L.). *Res. J. Fish. Hydrobiol.* 3, 84–88.

- 723 Fauvelle, V., Nhu-Trang, T.T., Feret, T., Madarassou, K., Randon, J., Mazzella, N., 2015.
724 Evaluation of Titanium Dioxide as a Binding Phase for the Passive Sampling of
725 Glyphosate and Aminomethyl Phosphonic Acid in an Aquatic Environment. *Anal. Chem.*
726 87, 6004–6009. <https://doi.org/10.1021/acs.analchem.5b00194>
- 727 Ferreira, D., Motta, A.C. da, Kreutz, L.C., Toni, C., Loro, V.L., Barcellos, L.J.G., 2010.
728 Assessment of oxidative stress in *Rhamdia quelen* exposed to agrichemicals.
729 *Chemosphere* 79, 914–921. <https://doi.org/10.1016/j.chemosphere.2010.03.024>
- 730 Folmar, L., Sanders, H., Julin, A., 1979. Toxicity of the herbicide glyphosate and several of
731 its formulations to fish and aquatic invertebrates, *Field Studies*.
- 732 Fulda, S., Gorman, A.M., Hori, O., Samali, A., 2010. Cellular stress responses: Cell survival
733 and cell death. *Int. J. Cell Biol.* 2010. <https://doi.org/10.1155/2010/214074>
- 734 Gasnier, C., Dumont, C., Benachour, N., Clair, E., Chagnon, M.C., Séralini, G.E., 2009.
735 Glyphosate-based herbicides are toxic and endocrine disruptors in human cell lines.
736 *Toxicology* 262, 184–191. <https://doi.org/10.1016/j.tox.2009.06.006>
- 737 Giesy, J.P., Dobson, S., Solomon, K.R., 2000. Ecotoxicological Risk Assessment for
738 Roundup Herbicide. *Rev. Environmetal Contam. Toxicol.* 167, 35–120.
- 739 Gluszczak, L., Dos Santos Miron, D., Moraes Silveira, B., Rodrigues Simões, R., Chitolina
740 Schetinger, M.R., Morsch, V.M., Loro, V.L., 2007. Acute effects of glyphosate herbicide
741 on metabolic and enzymatic parameters of silver catfish (*Rhamdia quelen*). *Comp.*
742 *Biochem. Physiol. - C Toxicol. Pharmacol.* 146, 519–524.
743 <https://doi.org/10.1016/j.cbpc.2007.06.004>
- 744 Gluszczak, L., Loro, V.L., Pretto, A., Moraes, B.S., Raabe, A., Duarte, M.F., Da Fonseca,
745 M.B., De Menezes, C.C., De Sousa Valladão, D.M., 2011. Acute exposure to
746 glyphosate herbicide affects oxidative parameters in piava (*Leporinus obtusidens*). *Arch.*
747 *Environ. Contam. Toxicol.* 61, 624–630. <https://doi.org/10.1007/s00244-011-9652-4>
- 748 Goulart, T.L.S., Boyle, R.T., Souza, M.M., 2015. Cytotoxicity of the association of pesticides
749 Roundup Transorb Ò and Furadan 350 SC Ò on the zebrafish cell line , ZF-L. *Toxicol.*
750 *Vitr.* 29, 1377–1384. <https://doi.org/10.1016/j.tiv.2015.06.007>
- 751 Grandcoin, A., Piel, S., Baurès, E., 2017. AminoMethylPhosphonic acid (AMPA) in natural
752 waters: Its sources, behavior and environmental fate. *Water Res.* 117, 187–197.
753 <https://doi.org/10.1016/j.watres.2017.03.055>
- 754 Guilherme, S., Gaivão, I., Santos, M.A., Pacheco, M., 2010. European eel (*Anguilla anguilla*)
755 genotoxic and pro-oxidant responses following short-term exposure to Roundup® - A

756 glyphosate-based herbicide. *Mutagenesis* 25, 523–530.
757 <https://doi.org/10.1093/mutage/geq038>

758 Guilherme, S., Santos, M.A., Barroso, C., Gaivão, I., Pacheco, M., 2012. Differential
759 genotoxicity of Roundup® formulation and its constituents in blood cells of fish (*Anguilla*
760 *anguilla*): Considerations on chemical interactions and DNA damaging mechanisms.
761 *Ecotoxicology* 21, 1381–1390. <https://doi.org/10.1007/s10646-012-0892-5>

762 Harayashiki, C.A.Y., Junior, A.S.V., Machado, A.A. de S., Cabrera, L. da C., Primel, E.G.,
763 Bianchini, A., Corcini, C.D., 2013. Toxic effects of the herbicide Roundup in the guppy
764 *Poecilia vivipara* acclimated to fresh water. *Aquat. Toxicol.* 142–143, 176–184.
765 <https://doi.org/10.1016/j.aquatox.2013.08.006>

766 Hildebrand, L.D., Sullivan, D.S., Sullivan, T.P., 1982. Experimental studies of rainbow trout
767 populations exposed to field applications of Roundup® herbicide. *Arch. Environ.*
768 *Contam. Toxicol.* 11, 93–98. <https://doi.org/10.1007/BF01055192>

769 Kašuba, V., Milić, M., Rozgaj, R., Kopjar, N., Mladinić, M., Žunec, S., Vrdoljak, A.L., Pavičić,
770 I., Marjanović Čermak, A.M., Pizent, A., Tariba Lovaković, B., Davor, Ž., 2017. Effects of
771 low doses of glyphosate on DNA damage , cell proliferation and oxidative stress in the
772 HepG2 cell line 19267–19281. <https://doi.org/10.1007/s11356-017-9438-y>

773 Kienzler, A., Bony, S., Devaux, A., 2013. DNA repair activity in fish and interest in
774 ecotoxicology: A review. *Aquat. Toxicol.* 134–135, 47–56.
775 <https://doi.org/10.1016/j.aquatox.2013.03.005>

776 Kienzler, A., Tronchère, X., Devaux, A., Bony, S., 2012. Assessment of RTG-W1, RTL-W1,
777 and PLHC-1 fish cell lines for genotoxicity testing of environmental pollutants by means
778 of a Fpg-modified comet assay. *Toxicol. Vitro.* 26, 500–510.
779 <https://doi.org/10.1016/j.tiv.2012.01.001>

780 Koakoski, G., Quevedo, R.M., Ferreira, D., Oliveira, T.A., da Rosa, J.G.S., de Abreu, M.S.,
781 Gusso, D., Marqueze, A., Kreutz, L.C., Giacomini, A.C.V., Fagundes, M., Barcellos,
782 L.J.G., 2014. Agrichemicals chronically inhibit the cortisol response to stress in fish.
783 *Chemosphere* 112, 85–91. <https://doi.org/10.1016/j.chemosphere.2014.02.083>

784 Koller, V.J., Fürhacker, M., Nersesyan, A., Mišák, M., Eisenbauer, M., Knasmueller, S., 2012.
785 Cytotoxic and DNA-damaging properties of glyphosate and Roundup in human-derived
786 buccal epithelial cells. *Arch. Toxicol.* 86, 805–813. [https://doi.org/10.1007/s00204-012-](https://doi.org/10.1007/s00204-012-0804-8)
787 [0804-8](https://doi.org/10.1007/s00204-012-0804-8)

788 Le Bihanic, F., Morin, B., Cousin, X., Le Menach, K., Budzinski, H., Cachot, J., 2014.

789 Developmental toxicity of PAH mixtures in fish early life stages. Part I: adverse effects in
790 rainbow trout. *Environ. Sci. Pollut. Res.* 21, 13720–13731.
791 <https://doi.org/10.1007/s11356-014-2804-0>

792 Lee, L.E.J., Clemons, J.H., Bechtel, D.G., Caldwell, S.J., Han, K.-B., Pasitschniak-Arts, M.,
793 Mosser, D.D., Bols, N.C., 1993. Development and characterization of a rainbow trout
794 liver cell line expressing cytochrome P450-dependent monooxygenase activity. *Cell*
795 *Biol. Toxicol.* 9, 279–294. <https://doi.org/10.1007/BF00755606>

796 Livak, K.J., Schmittgen, T.D., 2001. Analysis of relative gene expression data using real-time
797 quantitative PCR and the 2- $\Delta\Delta$ CT method. *Methods* 25, 402–408.
798 <https://doi.org/10.1006/meth.2001.1262>

799 Lopes, F.M., Sandrini, J.Z., Souza, M.M., 2018. Toxicity induced by glyphosate and
800 glyphosate-based herbicides in the zebrafish hepatocyte cell line (ZF-L). *Ecotoxicol.*
801 *Environ. Saf.* 162, 201–207. <https://doi.org/10.1016/j.ecoenv.2018.07.005>

802 Marques, A., Guilherme, S., Gaivão, I., Santos, M.A., Pacheco, M., 2014. Progression of
803 DNA damage induced by a glyphosate-based herbicide in fish (*Anguilla anguilla*) upon
804 exposure and post-exposure periods - Insights into the mechanisms of genotoxicity and
805 DNA repair. *Comp. Biochem. Physiol. Part - C Toxicol. Pharmacol.* 166, 126–133.
806 <https://doi.org/10.1016/j.cbpc.2014.07.009>

807 Martínez, A., Reyes, I., Reyes, N., 2007. Citotoxicidad del glifosato en células
808 mononucleares de sangre periférica humana. *Biomédica* 27, 594–604.

809 Mesnage, R., Bernay, B., Séralini, G.E., 2013. Ethoxylated adjuvants of glyphosate-based
810 herbicides are active principles of human cell toxicity. *Toxicology* 314, 122–128.
811 <https://doi.org/10.1016/j.tox.2012.09.006>

812 Modesto, K.A., Martinez, C.B.R., 2010. Effects of Roundup Transorb on fish: Hematology,
813 antioxidant defenses and acetylcholinesterase activity. *Chemosphere* 81, 781–787.
814 <https://doi.org/10.1016/j.chemosphere.2010.07.005>

815 Moreno, N.C., Sofia, S.H., Martinez, C.B.R., 2014. Genotoxic effects of the herbicide
816 Roundup Transorb® and its active ingredient glyphosate on the fish *Prochilodus*
817 *lineatus*. *Environ. Toxicol. Pharmacol.* 37, 448–454.
818 <https://doi.org/10.1016/j.etap.2013.12.012>

819 Morgan, J.D., Vigers, G.A., Farrell, A.P., Janz, D.M., Manville, J.F., 1991. Acute Avoidance
820 Reactions and Behavioral Responses of Juvenile Rainbow Trout (*Oncorhynchus*
821 *mykiss*) to Garlon, Garlon 3A and Vision Herbicides. *Environ. Toxicol. Chem.* 10, 73–79.

822 Morgan, M.J., Kiceniuk, J.W., 1992. Response of Rainbow-Trout To a 2 Month Exposure To
823 Vision(R), a Glyphosate Herbicide. *Bull. Environ. Contam. Toxicol.* 48, 772–780.

824 Mourente, G., Tocher, D.R., Díaz, E., Grau, A., Pastor, E., 1999. Relationships between
825 antioxidant enzyme activities and lipid peroxidation products during early development
826 in *Dentex dentex* eggs and larvae. *Aquaculture* 179, 309–324.

827 Myers, J.P., Antoniou, M.N., Blumberg, B., Carroll, L., Colborn, T., Everett, L.G., Hansen, M.,
828 Landrigan, P.J., Lanphear, B.P., Mesnage, R., Vandenberg, L.N., Vom Saal, F.S.,
829 Welshons, W. V., Benbrook, C.M., 2016. Concerns over use of glyphosate-based
830 herbicides and risks associated with exposures: A consensus statement. *Environ. Heal.*
831 *A Glob. Access Sci. Source* 15, 1–13. <https://doi.org/10.1186/s12940-016-0117-0>

832 Navarro, C.D.C., Martinez, C.B.R., 2014. Effects of the surfactant polyoxyethylene amine
833 (POEA) on genotoxic, biochemical and physiological parameters of the freshwater
834 teleost *Prochilodus lineatus*. *Comp. Biochem. Physiol. Part - C Toxicol. Pharmacol.* 165,
835 83–90. <https://doi.org/10.1016/j.cbpc.2014.06.003>

836 Nwani, C.D., Ibiam, U.A., Ibiam, O.U., Nworie, O., Onyishi, G., Atama, C., 2013. Investigation
837 on Acute Toxicity and Behavioral Changes in *Tilapia Zillii* due to glyphosate-based
838 herbicide, Forceup. *J. Anim. plan Sci.* 23, 888–892.

839 Pannetier, P., Fuster, L., Clérandeau, C., Lacroix, C., Gourves, P.Y., Cachot, J., Morin, B.,
840 2018. Usefulness of RTL-W1 and OLCAB-e3 fish cell lines and multiple endpoint
841 measurements for toxicity evaluation of unknown or complex mixture of chemicals.
842 *Ecotoxicol. Environ. Saf.* 150, 40–48. <https://doi.org/10.1016/j.ecoenv.2017.12.027>

843 Peruzzo, P.J., Porta, A.A., Ronco, A.E., 2008. Levels of glyphosate in surface waters,
844 sediments and soils associated with direct sowing soybean cultivation in north pampasic
845 region of Argentina. *Environ. Pollut.* 156, 61–66.
846 <https://doi.org/10.1016/j.envpol.2008.01.015>

847 Qin, Y., Li, X., Xiang, Y., Wu, D., Bai, L., Li, Z., Liang, Y., 2017. Toxic effects of glyphosate
848 on diploid and triploid fin cell lines from *Misgurnus anguillicaudatus*. *Chemosphere* 180,
849 356–364. <https://doi.org/10.1016/j.chemosphere.2017.03.098>

850 Salbego, J., Pretto, A., Gioda, C.R., De Menezes, C.C., Lazzari, R., Radünz Neto, J.,
851 Baldisserotto, B., Loro, V.L., 2010. Herbicide formulation with glyphosate affects growth,
852 acetylcholinesterase activity, and metabolic and hematological parameters in *Piava*
853 (*Ieporinus obtusidens*). *Arch. Environ. Contam. Toxicol.* 58, 740–745.
854 <https://doi.org/10.1007/s00244-009-9464-y>

- 855 Sinhorin, V.D.G., Sinhorin, A.P., Teixeira, J.M. dos S., Miléski, K.M.L., Hansen, P.C.,
856 Moreira, P.S.A., Kawashita, N.H., Baviera, A.M., Loro, V.L., 2014. Effects of the acute
857 exposition to glyphosate-based herbicide on oxidative stress parameters and
858 antioxidant responses in a hybrid Amazon fish surubim (*Pseudoplatystoma* sp).
859 *Ecotoxicol. Environ. Saf.* 106, 181–187. <https://doi.org/10.1016/j.ecoenv.2014.04.040>
- 860 Sinhorin, V.D.G., Sinhorin, A.P., Teixeira, J.M.S., Miléski, K.M.L., Hansen, P.C., Moeller,
861 P.R., Moreira, P.S.A., Baviera, A.M., Loro, V.L., 2014. Metabolic and Behavior Changes
862 in Surubim Acutely Exposed to a Glyphosate-Based Herbicide. *Arch Env. Contam*
863 *Toxicol* 659–667. <https://doi.org/10.1007/s00244-014-0073-z>
- 864 Sulukan, E., Mine, K., Ceylan, H., Beydemir, Ş., Işık, M., Atamanalp, M., Ceyhun, S.B., 2017.
865 An approach to clarify the effect mechanism of glyphosate on body malformations
866 during embryonic development of zebrafish (*Daino rerio*). *Chemosphere* 180, 77–85.
867 <https://doi.org/10.1016/j.chemosphere.2017.04.018>
- 868 Tierney, K.B., Singh, C.R., Ross, P.S., Kennedy, C.J., 2007. Relating olfactory neurotoxicity
869 to altered olfactory-mediated behaviors in rainbow trout exposed to three currently-used
870 pesticides. *Aquat. Toxicol.* 81, 55–64. <https://doi.org/10.1016/j.aquatox.2006.11.006>
- 871 Topal, A., Atamanalp, M., Uçar, A., Oruç, E., Kocaman, E.M., Sulukan, E., Akdemir, F.,
872 Beydemir, Ş., Kiliç, N., Erdoğan, O., Ceyhun, S.B., 2015. Effects of glyphosate on
873 juvenile rainbow trout (*Oncorhynchus mykiss*): Transcriptional and enzymatic analyses
874 of antioxidant defence system, histopathological liver damage and swimming
875 performance. *Ecotoxicol. Environ. Saf.* 111, 206–214.
876 <https://doi.org/10.1016/j.ecoenv.2014.09.027>
- 877 Tsui, M.T.K., Chu, L.M., 2003. Aquatic toxicity of glyphosate-based formulations:
878 Comparison between different organisms and the effects of environmental factors.
879 *Chemosphere* 52, 1189–1197. [https://doi.org/10.1016/S0045-6535\(03\)00306-0](https://doi.org/10.1016/S0045-6535(03)00306-0)
- 880 Üner, N., Oruç, E.Ö., Sevgiler, Y., Şahin, N., Durmaz, H., Usta, D., 2006. Effects of diazinon
881 on acetylcholinesterase activity and lipid peroxidation in the brain of *Oreochromis*
882 *niloticus*. *Environ. Toxicol. Pharmacol.* 21, 241–245.
883 <https://doi.org/10.1016/j.etap.2005.08.007>
- 884 Uren Webster, T.M., Laing, L.V., Florance, H., Santos, E.M., 2014. Effects of Glyphosate and
885 its Formulation, Roundup, on Reproduction in Zebra fish (*Danio rerio*). *Environ. Sci.*
886 *Technol.* 48, 1271–1279. <https://doi.org/10.1021/es404258h>
- 887 Van Bruggen, A.H.C., He, M.M., Shin, K., Mai, V., Jeong, K.C., Finckh, M.R., Morris, J.G.,

888 2018. Environmental and health effects of the herbicide glyphosate. *Sci. Total Environ.*
889 616–617, 255–268. <https://doi.org/10.1016/j.scitotenv.2017.10.309>

890 Vanlaeys, A., Dubuisson, F., Seralini, G.E., Travert, C., 2018. Formulants of glyphosate-
891 based herbicides have more deleterious impact than glyphosate on TM4 Sertoli cells.
892 *Toxicol. Vitro.* 52, 14–22. <https://doi.org/10.1016/j.tiv.2018.01.002>

893 Vera, M.S., Lagomarsino, L., Sylvester, M., Pérez, G.L., Rodríguez, P., Mugni, H., Sinistro,
894 R., Ferraro, M., Bonetto, C., Zagarese, H., Pizarro, H., 2010. New evidences of
895 Roundup®(glyphosate formulation) impact on the periphyton community and the water
896 quality of freshwater ecosystems. *Ecotoxicology* 19, 710–721.
897 <https://doi.org/10.1007/s10646-009-0446-7>

898 Webster, T.M.U., Santos, E.M., 2015. Global transcriptomic profiling demonstrates induction
899 of oxidative stress and of compensatory cellular stress responses in brown trout
900 exposed to glyphosate and Roundup. *BMC Genomics* 16, 1–14.
901 <https://doi.org/10.1186/s12864-015-1254-5>

902 Weeks Santos, S., Cachot, J., Gourves, P.Y., Clérandeau, C., Morin, B., Gonzalez, P., 2019.
903 Sub-lethal effects of waterborne copper in early developmental stages of rainbow trout
904 (*Oncorhynchus mykiss*). *Ecotoxicol. Environ. Saf.* 170, 778–788.
905 <https://doi.org/10.1016/j.ecoenv.2018.12.045>

906 Yusof, S., Ismail, A., Alias, M.S., 2014. Effect of glyphosate-based herbicide on early life
907 stages of Java medaka (*Oryzias javanicus*): A potential tropical test fish. *Mar. Pollut.*
908 *Bull.* 85, 494–498. <https://doi.org/10.1016/j.marpolbul.2014.03.022>

909 Zebral, Y.D., Costa, P.G., de Castro Knopp, B., Lansini, L.R., Zafalon-Silva, B., Bianchini, A.,
910 Robaldo, R.B., 2017. Effects of a glyphosate-based herbicide in pejerrey *Odontesthes*
911 *humensis* embryonic development. *Chemosphere* 185, 860–867.
912 <https://doi.org/10.1016/j.chemosphere.2017.07.069>

913 Zhang, S., Xu, J., Kuang, X., Li, S., Li, X., Chen, D., Zhao, X., Feng, X., 2017. Biological
914 impacts of glyphosate on morphology, embryo biomechanics and larval behavior in
915 zebrafish (*Danio rerio*). *Chemosphere* 181, 270–280.
916 <https://doi.org/10.1016/j.chemosphere.2017.04.094>

917 Zhang, X., Yang, F., Zhang, X., Xu, Y., Liao, T., Song, S., Wang, J., 2008. Induction of
918 hepatic enzymes and oxidative stress in Chinese rare minnow (*Gobiocypris rarus*)
919 exposed to waterborne hexabromocyclododecane (HBCDD). *Aquat. Toxicol.* 86, 4–11.
920 <https://doi.org/10.1016/j.aquatox.2007.07.002>

921

922

923 **FIGURE CAPTIONS**

924 **Figure 1.** Biometric analyzes of larvae after exposure to 0.1 and 1 mg/L of glyphosate. (A) total
925 body length (mm), (B) head length of larvae (mm) and (C) ratio of head size to total length of
926 larvae (%) are showed. Different letters indicate significant differences between conditions
927 (Mean \pm SD N = 3, ANOVA, $p < 0.05$).

928 **Figure 2.** Percentage of malformed rainbow trout larvae after 21 days of exposure to
929 glyphosate. Different letters indicate significant differences (Mean \pm SD, N = 3, ANOVA,
930 $p < 0.05$).

931 **Figure 3.** Mean velocity (cm/s) of larvae exposed to glyphosate after a light stimulation.
932 Velocity was recorded after 30 min video tracked analysis. Data was average over each 1 min
933 interval (A) and over each 10 min (B). Different letters indicate significant differences for each
934 period of illumination (Mean \pm SD N = 3, ANOVA, $p < 0.05$).

935 **Figure 4.** Cumulative time of high mobility (a) ; mobility (b); and immobility (c) on larvae
936 exposed to glyphosate. Different letters indicate significant differences between each period
937 of time (Mean \pm SD, N = 3, ANOVA, $p < 0.05$).

938 **Figure 5.** DNA damage in blood cells from rainbow trout larvae after exposure to 0.1 and 1
939 mg/L of glyphosate, with- and without addition of enzymatic Fpg treatment. Different letters
940 indicate significant differences between treatments (Mean \pm SD, N = 3, ANOVA, $p < 0.05$).

941 **Figure 6.** Lipid peroxidation (A) expressed as nanomoles of TBARS/mg of protein and protein
942 carbonyls (B) expressed as nanomoles of carbonyl/mg of protein in rainbow trout exposed to
943 glyphosate. Different letters represent significant differences. All values are expressed as
944 Mean \pm SD, N=3, ANOVA.


945 **Figure 7.** Comparative cytotoxicity of glyphosate (A) and Roundup (B) on the RTL-W1 cell line
946 after 24 h of exposure. Asterisks represent significant differences compared to control. Values
947 represent Mean \pm SD. (N=3, Kruskal-Wallis, $p < 0.05$).

948 **Figure 8.** DNA damage in RTL-W1 cell line induced by glyphosate (A) and Roundup (B)
949 measured by the comet assay with and without Fpg treatment. Values represent Mean \pm SD.
950 Different letters indicate significant differences. N=3, Kruskal-Wallis ($p < 0.05$).

951

952 Figure 1.

953



954

955


956

957

958

959 Figure 2.

960




961

962

963

964 **Figure 3.**

965




966

967

968 Figure 4.

969




970

971

972 Figure 5.

973




974

975

976

977 Figure 6.

978




979

980

981 Figure 7.


982



983


984

985



986 Figure 8.

987



988

989

990

991 **Table 1:** Accession number and specific primer pairs for the *Oncorhynchus mykiss* used in
 992 our study.

Gene	Accession number	Primer (5' – 3')
rpl7	NM_001160672.2	GGTCGCTCTCACAGACAACA ^a TTATGTCCGTCCTCTGGGT ^b
ef1α	NM_001124339.1	ATGGGCTGGTTCAAGGGATG ^a GATCATACCGGCCTTCAGGG ^b
cat	FJ226382.1	CAGGTGTCTTTCTTGTTTCAG ^a GTCCAGGATGGGAAGTTGC ^b
sod	NM_001124329.1	TGATTGGGGAGATCTCGGGT ^a CGGGTCCAGTGAGAGTCAAC ^b
gst	BT073173.1	ATTTTGGGACGGGCTGACA ^a CCTGGTGCTCTGCTCCAGT ^b
er-b	AJ242741	AGCCCTCTCCTCCACCCTACCA ^a ACAGCTGGCTGAGGAGGAGTT ^b
cox1	KP013084.1	TCGTTTGAGCCGTGCTAGTT ^a CTTCTGGGTGGCCGAAGAAT ^b
12s	KY798500.1	GCGCCAGCTTAAAACCCAAA ^a GCCCATTTCTTCCCACCTCA ^b
ogg1	XR_002474791.1	CTGATGGACAAGGCCAGTGT ^a GTAAGGACCCCATGGCTGTC ^b
rad51	XM_021612309.1	AGGCTGGAGGAGGACATCAT ^a GTATTTGAGGGTGGCAGCCT ^b
bax	BT074328.1	CAGAAAACCCAGGGAGGCAT ^a AGAACACATCCTGGGCACAG ^b
cyp19a1	XM_021598638	CTCTCCTCTCATACCTCAGGTT ^a AGAGGAACTGCTGAGTATGAAT ^b

^aForward primer

^bReverse primer

993

994

995

996

997

998 **Table 2:** Measured concentration of glyphosate in the exposure water for each studied
999 condition.

Nominal concentration (mg/L)	Measured concentration (mg/L)	
0.0	T0	0.0 ± 0.0
	T48	0.0 ± 0,0
0.1	T0	0.12 ± 0.0
	T48	0.12 ± 0.01
1.0	T0	1.22 ± 0.01
	T48	1.22 ± 0.01

1000

1001

1002

1003 **Table 3:** Effects on viability and development of rainbow trout during glyphosate exposure.

1004 Values represent Mean \pm SD (N = 3). The results show no significant difference.

	Control	Glyphosate 0.1 mg/L	Glyphosate 1 mg/L
Embryonic viability (%)	96.3 \pm 2.1	95.3 \pm 3.8	95.3 \pm 3.5
Larval viability (%)	91.9 \pm 3.4	93.2 \pm 3.6	92.2 \pm 6.2
Cumulative viability (%)	88.6 \pm 5.2	88.8 \pm 3.6	88.0 \pm 8.6
Hatching rate (%)	99.0 \pm 0.02	97.6 \pm 2.4	95.8 \pm 1.8
Development time (DD)	307.9 \pm 4.4	311.4 \pm 3.1	314.0 \pm 6.8

1005

1006