In vitro probiotic properties of selected lactobacilli and multi-strain consortium on immune function, gut barrier strengthening and gut hormone secretion

Yanath Belguesmia, Jeanne Alard, Rezak Mendil, Rozenn Ravallec, Corinne Grangette, Djamel Drider, Benoit Cudennec

► To cite this version:

Yanath Belguesmia, Jeanne Alard, Rezak Mendil, Rozenn Ravallec, Corinne Grangette, et al.. In vitro probiotic properties of selected lactobacilli and multi-strain consortium on immune function, gut barrier strengthening and gut hormone secretion. Journal of Functional Foods, 2019, 57, pp.382-391. 10.1016/j.jff.2019.04.028 . hal-02617752

HAL Id: hal-02617752

<https://hal.inrae.fr/hal-02617752>

Submitted on 22 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution - NonCommercial| 4.0 International License

In vitro probiotic properties of selected *lactobacilli* and multi-strain consortium on immune function, gut barrier strengthening and gut hormone secretion

Yanath Belguesmia^{1#}, Jeanne Alard^{2#}, Rezak Mendil¹, Rozenn Ravallec¹,
Corinne Grangette², Djamel Drider¹, Benoit Cudennec^{1*}

¹EA 7394, ICV–Institut Charles Viollette, UniversitéLille, INRA, ISA, UniversitéArtois, Université Littoral Côte d’Opale, F-59000 Lille, France

²Univ. Lille, CNRS, Inserm, CHU Lille, Institut Pasteur de Lille, U1019 – UMR 8204 – CIIL– Centre d’Infection et d’Immunité de Lille, F-59000 Lille, France

[#]These authors contributed equally to this work

*Corresponding author.

Email: benoit.cudennec@univ-lille.fr

Keywords: probiotics; inflammation; *lactobacilli*; multi-strain consortium; gut hormones; gut barrier

Abstract

Lactobacillus reuteri ICVB395, *L. gasseri* ICVB392 and *L. gasseri* ICVB396 strains, isolated from vaginal microbiota, were investigated for their probiotic traits. *L. reuteri* ICVB395 strain and the 3RG consortium, associating these three *Lactobacillus* strains, showed the best anti-inflammatory profile on peripheral blood mononuclear cell (PBMC) while *L. gasseri* ICVB392 was the most potent together with the 3RG consortium to strengthen a Caco-2-derived epithelial barrier. The three studied strains induced various secretion levels of glucagon-like peptide 1 (GLP-1) and cholecystokinin (CCK) by STC-1 enteroendocrine cells, whereas the 3RG consortium was globally less performing. Specific antagonists of protein G receptors, CaSR (Calcium-Sensing Receptor) and GPRC6A (G protein-coupled receptor family C group 6 member A), and inhibitor of the peptide transporter Pept-1 provoked differential modulation of the GLP-1 and CCK secretion by STC-1 cells, indicating that different mechanisms are involved in the capacity of *lactobacilli* and the 3RG to modulate gut hormones secretion.

1. Introduction

The gastro-intestinal tract houses billions of microorganisms, namely the gut microbiota, dominated by bacteria, and characterized by their extreme diversity and their key role in the gut homeostasis (Arulampalam, Greicius, & Pettersson, 2006; Sanders, 2016; Stubbendieck, Vargas-Bautista, & Straight, 2016). Nowadays this rich and opulent gut microbiota is generally considered as a full microbial “endocrine organ” (Clarke et al., 2014; Jayasinghe, Chiavaroli, Holland, Cutfield, & O’Sullivan, 2016). Amongst multitude roles of this microbiota we can quote the protection of the host against the invasion of unwanted microorganisms, the contribution to the proper functioning of the immune system, its influence on glucose and lipid homeostasis, and its role on the degradation of insoluble dietary fibres in short chain fatty acids known to exhibit a beneficial impact on the anti-inflammatory and metabolic responses (Delzenne & Cani, 2011; Drissi, Raoult, & Merhej, 2017; Sánchez et al., 2017). Nonetheless the gut microbiota has also a function of “dialogue” with the digestive tract, interacting with the intestinal cells (Cani & Knauf, 2016). The diversity of the gut microbiota can be overthrown by drastic changes related to clinical practice and external pressure, notably the mode of delivery and new-borns feeding, medical treatments such as antibiotic, nutrition and health care behaviours and environmental exposure. This can disrupt temporarily the balance of this complex ecosystem leading to digestive discomfort, and the development of chronic diseases (Beaugerie & Petit, 2004; Quigley, 2013). These perturbations are associated with a dramatic increase in incidence of immune-mediated diseases including allergic and inflammatory bowel diseases but also metabolic diseases including obesity and diabetes and most likely neurodegenerative and psychiatric diseases (Cani & Knauf, 2016; Doré, Multon, Béhier, & participants of Giens XXXII, Round Table No.2, 2017; Fernandez, Lasa, & Man, 2014). Therefore, the development of approaches targeting the key features of this altered host-microbes

67 interactions are highly relevant. Probiotics precisely aimed at preserving and/or restoring the
68 balance of this delicate ecosystem and therefore respond to the precise definition emitted in
69 2002 by the Food and Agriculture Organisation (FAO) and World Health Organization
70 (WHO), establishing that probiotics are "live microorganisms which, when ingested in
71 adequate amounts, exert positive effects on health, beyond traditional nutritional effects"
72 (FAO/WHO, 2002). Probiotic bacteria, mainly belonging to the *Lactobacillus* or
73 *Bifidobacterium* genera, are natural inhabitants of the gastrointestinal (GI) tract (Vaughan &
74 Mollet, 1999). Probiotics have to survive to the passage through the stomach and the upper
75 part of the small intestine before reaching their site of action to provide their benefits (Fooks
76 & Gibson, 2002). To overcome this challenge, the selection of potential probiotics bacteria
77 mainly focus on their ability to survive to the harsh conditions of the GI tract (Morelli, 2000).
78 Probiotics could be provided as a unique individual strain or as combination of multiple
79 strains. Multi-strain cocktails could present many advantages providing more benefits
80 compared to those of microorganisms taken alone (Timmerman, Koning, Mulder, Rombouts,
81 & Beynen, 2004). Indeed, combining multiple strains belonging to different species able to
82 colonize different parts of the digestive tract could be more effective to restore gut
83 homeostasis and to express their positive effects, as these strains can work simultaneously on
84 the different potential causes of the observed disorders (Collado, Meriluoto, & Salminen,
85 2007; Timmerman et al., 2004). Use of consortium strains could be justified by the individual
86 specific response of the host, as natural probiotic predominant species can vary from one
87 individual to another. Each strain also helps to generate specific enzyme activities and can
88 stimulate the immune system by different pathways (Sánchez et al., 2017). Bacteria belonging
89 to the two main families of known probiotics are found in various locations in the intestine.
90 *Lactobacilli* are natural residents of the small intestine while the bifidobacteria are dominant
91 in the colon, their natural habitat (Collado et al., 2007; Gionchetti, Lammers, Rizzello, &

92 Campieri, 2005; Perdigon, Galdeano, Valdez, & Medici, 2003; Timmerman et al., 2004).
93 Most *lactobacilli* and *bifidobacteria* species are considered as GRAS “Generally Recognized
94 As Safe” (Hugas & Monfort, 1997). The benefits of these bacteria are well known for decades
95 and many traditional and commercial *probiotic preparations contain them* (Di Cerbo, Palmieri,
96 Aponte, Morales-Medina, & Iannitti, 2015; Drissi et al., 2017). Probiotic strains express their
97 *benefits by strengthening of the intestinal barrier, regulating sugar and lipid metabolism,*
98 *reducing inflammation or preventing pathogens invasion* (Botta, Langerholc, Cencič, &
99 Cocolin, 2014; Di Cerbo et al., 2015; Park, Oh, & Cha, 2014; Takemura, Okubo, &
100 Sonoyama, 2010). *Moreover probiotics, especially lactobacilli, synthesize a wide variety of*
101 *proteases, implied in the food maturation process such as in fermented cheeses* (Liu, Bayjanov,
102 Renckens, Nauta, & Siezen, 2010). *Amongst which, serine proteases are produced by many*
103 **lactobacilli* species, notably by *L. gasseri*, *L. plantarum* and some *L. acidophilus* strains* (Law
104 & Haandrikman, 1997; Margono, Sumaryono, Malik, & Sadikin, 2014). Moreover they
105 produce, lactic acid promoting the balance of intestinal pH, and sometimes antimicrobial
106 substances that inhibit the growth of pathogens (Drissi et al., 2017).
107 Furthermore, recent works established that probiotic *lactobacilli* are able to interact with
108 intestinal cells and induce modulation of gut hormones; as glucagon-like peptide-1 (GLP-1);
109 cholecystokinin (CCK) and PYY peptide (Panwar et al., 2016; Yadav, Lee, Lloyd, Walter, &
110 Rane, 2013). These hormones, because of their influence in food intake regulation and
111 glucose homeostasis, represent promising lever to manage and reduce chronic metabolic
112 diseases like obesity and associated type 2-diabetes (Caron, Domenger, Dhulster, Ravallec, &
113 Cudennec, 2017).
114 During a previous work, some *Lactobacillus* strains, isolated from the vaginal microbiota,
115 exhibited many traits of probiotics microorganisms, including good *survivability in the harsh*
116 *conditions* of the GI tract and adhesion to epithelial cells without toxicity (Belguesmia et al.,

2016). During these researches some strains, including *L. gasseri* ICVB392, *L. reuteri* ICVB395 and *L. gasseri* ICVB396, initially designated CMUL34, CMUL67 and CMUL80 respectively, and renamed after re-identification by 16S rDNA sequencing, appeared good potential probiotics and were able, *in vitro*, to modulate gut hormone expression and secretion in murine enteroendocrine STC-1 cells.

The aim of this study is to investigate further probiotic traits of these strains, alone and associated in a consortium. We first evaluated their immunomodulation abilities and their capacity to strengthen the intestinal barrier. Using the enteroendocrine STC-1 cell line, we also studied deeper the impact of the consortium on the secretion of gut hormones, and we unravelled how the bacteria interacted with the cells. Finally, we studied their capacity to limit lipid accumulation in adipocytes.

2. Material and methods

2.1. Bacterial strains

The three *Lactobacillus* strains used in this research work were previously isolated from Lebanese vaginal microbiota (Al Kassaa, Hamze, Hober, Chihib, & Drider, 2014), and recently selected for their probiotic traits (Belguesmia et al., 2016). Previously named CMUL34, CMUL67 and CMUL80, they were re-identified by 16S rDNA sequencing and registered in the Institut Charles Viollette Laboratory Collection, under *L. gasseri* ICVB392, *L. reuteri* ICVB395 and *L. gasseri* ICVB396 strains respectively. The strains were also associated in equal amount within the multi-strain cocktail designated during this study as 3RG consortium. Before each experiment, the strains were grown for 18-24 h at 37°C in de Man-Rogosa-Sharpe (MRS) medium (De Man, Rogosa, & Sharpe, 1960). Two additional strains were used as control strains for immune cells stimulation: *Bifidobacterium longum* IPL

A7.5 was grown at 37°C in anaerobic condition (GENbag anaer, Biomérieux, France) in MRS (Difco, Detroit, USA) supplemented with 0.1% (w/v) L-cysteine hydrochloride (Sigma) and *Lactococcus lactis* MG1363 was cultured at 30°C, in M17 Broth supplemented with 0.5% glucose (Difco, Detroit, USA). For *in vitro* studies (stimulation of PBMCs, Caco-2 epithelial barrier and adipocytes), bacteria were grown overnight, washed twice in sterile phosphate buffered saline (PBS) buffer pH 7.2 and resuspended at a final concentration of 2×10^9 cfu.mL⁻¹ in PBS.

2.2. *In vitro* immunomodulation assays

Blood samples from five different healthy adult donors were obtained at the Etablissement Français du Sang (French National Blood Service), in accordance with our institution committees (INSERM, CNRS and Institut Pasteur de Lille, agreement N° DC 2013-2022). Peripheral Blood Mononuclear Cells (PBMCs) were isolated from the blood as already described (Foligne et al., 2007). Briefly, after Ficoll gradient centrifugation (GE Healthcare Bio-Sciences, Uppsala, Sweden), mononuclear cells were collected, washed in RPMI-1640 medium (Gibco, Life Technologies, Ghent, Belgium), and adjusted to 2×10^6 cells per mL in RPMI supplemented with gentamicin (150 µg.mL⁻¹), L-glutamine (2 mM), and 10% heat-inactivated FCS (Gibco, Life Technologies, Ghent, Belgium). PBMCs were stimulated with phosphate-buffered saline (PBS, Gibco, Life Technologies, Ghent, Belgium) or bacteria at a bacteria-to-cell ratio of 10:1 for 24 h at 37 °C with 5% CO₂. The supernatants were collected and stored at -20°C until cytokines (IL-10 IL-12 and IFN-γ) measurements performed using R&D Duoset ELISA kits (R&D, Minneapolis, MN, USA). *B. longum* and *Lactococcus lactis* were used as positive control for the induction of anti-inflammatory (IL-10) and Th-1/pro-inflammatory (IL-12 and IFN-γ) cytokine secretion by PBMCs, respectively.

2.3. *Epithelial barrier model*

The human colon epithelial cell line Caco-2 clone TC7 (Chantret et al., 1994) was used to study the impact of the *lactobacilli* on trans-epithelial electric resistance (TEER). The Caco-2 cells were grown at 37°C with 10% CO₂ in DMEM supplemented with 5% heat-inactivated foetal calf serum (FCS, Gibco, Life technologies, Ghent, Belgium), 1% Non-Essential Amino Acids (Gibco, Life Technologies, Ghent, Belgium), 100 U.mL⁻¹ penicillin and 100 µg.mL⁻¹ streptomycin (Gibco, Life Technologies, Ghent, Belgium) and 2mM L-glutamine (Gibco, Life Technologies, Ghent, Belgium).

For the permeability test, polarized Caco-2 monolayers were prepared by growing the epithelial cells on 12-wells Transwell® insert filters (polycarbonate membrane with 3 µm pore size, 12 mm diameters, Costar, Corning Life Science, Kennebunk, ME, USA) at a density of 10⁵ cells per cm². The medium was changed every two days until 14 days when optimal trans-epithelial resistance (TEER ≥ 1800 Ω/cm²) was reached (which was measured every 2 days 1 h after changing medium using a millicell-ERS (Electrical Resistance System; Millipore, Billerica, MA, USA). At day 14, fresh medium without FCS was added and cells were treated, in the apical compartment with bacteria (or not) at a bacteria-to-cell ratio of 10:1, 30 min before the addition of hydrogen peroxide (H₂O₂) in both basal and apical compartment (at 100µM final concentration). TEER was measured before H₂O₂ addition (T0) and every 30 min until 120 min. The results were compared to non-treated cells. Three different experiments were performed including duplicates of each condition and results were expressed in % TEER compared to T0 ± SEM.

2.4. *Lipid accumulation in adipocytes*

The effect of *lactobacilli* on lipid accumulation in adipocytes was studied using the pre-adipocyte 3T3-L1 murine cell line. The 3T3-L1 cells were grown at 37°C with 5% CO₂ in

DMEM supplemented with 10% heat-inactivated FCS, 100 U.mL⁻¹ penicillin and 100 µg.mL⁻¹ streptomycin and 2 mM L-glutamine. Cells were used between the 10th and the 16th passage. The 3T3-L1 cells were distributed in 12-wells plates at a concentration of 3500 cells per wells and were differentiated in adipocytes according to the protocol described by (Zebisch, Voigt, Wabitsch, & Brandsch, 2012). Briefly, cells were grown in medium supplemented with 0,5 mM 3-isobutyl-1-methylxanthine, 1 µg.mL⁻¹ insulin and 0.25 µM dexamethasone. After 48h, fresh medium supplemented with only insulin (1 µg.mL⁻¹) was replaced. Basal medium was changed every two days for 10 days, until cells were differentiated in mature adipocytes. Cells were then stimulated for 24h with bacteria (or not) at a bacteria-to-cell ratio of 10:1, in fresh medium in the presence of 150 µg.mL⁻¹ gentamicin. Lipid accumulation was quantified by Oil- Red-O staining. Briefly, cells were stained using 1 ml ready-to-use Oil-red O solution (DiaPath, Martinengo, Italy) for 15 min., washed 3 times with PBS (Gibco, Life Technologies, Ghent, Belgium). Oil-Red-O was eluting with isopropanol for 30 min incubation. Optic density was measured at 490 nm by a spectrophotometer (ELx808, Biotech instruments). The percentage of Oil-red-O stained cells relative to control cells without bacteria was calculated as (A490nm [probiotic sample]/A490nm[control])*100.

2.5. Gut hormones secretion study

The STC-1 murine cell line, derived from the intestinal tumour of double transgenic mice, gratefully received from Dr. C. Roche (INSERM U865, Lyon, France), was used for gut hormones study. The cells were grown in Dulbecco's Modified Eagle's Medium (DMEM, Invitrogen, France), supplemented with 10% foetal bovine serum, 5 mM of L-glutamine and 100 U.mL⁻¹ of penicillin and streptomycin, at 37°C in 5% CO₂-95% air atmosphere. The STC-1 cells were passed twice a week, and were used between the 60th and the 65th passage for the different assays. STC-1 cells were seeded in 24 wells plate at 40,000 cells/well and

217 were grown in DMEM culture medium for 48-72h. Cells were washed twice with Hepes
218 buffer without glucose (NaCl 140 mM, Hepes 20 mM, KCl 4.5 mM, CaCl₂ 1.2 mM, MgCl₂
219 1.2 mM, adjusted to pH 7.4 with NaOH 3M) and then co-incubated with 10⁸ CFU.mL⁻¹ of the
220 selected *Lactobacillus* strains, alone or in combination, for 8 hours at 37°C in 5% CO₂-95%
221 air atmosphere. Purified peptidoglycan from *L. acidophilus* (Macho-Fernandez et al., 2011),
222 and purified flagellin from *Salmonella enterica* serovar Typhimurium (gratefully supplied by
223 Dr Jean Claude Sirard from Institut Pasteur de Lille) were tested at 50 and 10 µg.mL⁻¹
224 respectively, to evaluate their impact on gut hormones secretion. The resulting supernatants
225 were centrifuged (8000 g for 10 min) and were kept at -20°C. GLP-1 and CCK quantifications
226 were realized by Radio-Immuno Assay (RIA) using EMD Milipore (USA) and Cisbio
227 International (France) kits for each hormone, respectively.

228 To examine the capacity of bacteria to degrade the gut hormones, pure active GLP-1 1-26
229 (EMD Milipore, USA) and CCK8S (Sigma Aldrich Merk, Germany) peptide hormones were
230 used at initial concentration of 800 pM and 600 pM, respectively. *Lactobacilli* (at 10⁷
231 CFU.mL⁻¹) were incubated in Hepes buffer solution containing active GLP-1 or CCK8S, in
232 presence or absence of DPP-IV enzyme inhibitor (Ile-Pro-Ile, Sigma-Aldrich Germany) at 1
233 mg.mL⁻¹, during 8 h at 37°C. The supernatants were recovered by centrifugation at 8000 g,
234 4°C for 10 min, and quantification of remaining active GLP- 1 or CCK8S in the supernatants
235 was performed by RIA as previously described. Negative control without bacteria was
236 incubated in the same conditions for both hormones. The results were expressed in percentage
237 of initial concentration of active GLP-1 or CCK8S.

238 The effects of CaSR and GPRC6A antagonists and Pept-1 inhibitor on the secretion of GLP-1
239 and CCK hormones by STC-1 after contact with selected *Lactobacillus* strains were studied.
240 The two antagonists and the inhibitor were prepared according to the suppliers
241 recommendations. NPS 2143 (Sigma Aldrich, Merck Germany), CpD (Enamine, Ukraine),

and 4-aminomethylbenzoic acid (AMBA-4, Sigma Aldrich, Merck Germany) were used at a final concentration of 25 μ M, 50 μ M and 10 mM in Hepes Buffer, respectively. The STC-1 cells were washed after reaching 80% confluence culture and were incubated for 15 minutes at 37°C in 5% CO₂-95% air atmosphere with 100 μ L of described above solutions. Then the treated STC-1 cells were incubated with *lactobacilli* in the same conditions as described above.

2.6. RT-PCR analysis

The relative transcript levels of PepT1 (Peptide transporter) and β -actin were analysed by quantitative real-time PCR. Briefly, total RNA was extracted from cultured cells using the NucleoSpin® RNA XS (Macherey-Nagel, Germany) according to the manufacturer's instructions. Concentration and purity of each sample were evaluated on a NanoDrop Lite (Thermo Scientific, USA). cDNA was obtained by reverse transcription on a Mastercycler gradient (Eppendorf, Germany) using the RevertAid H Minus First Strand cDNA Synthesis Kit (Thermo Scientific, USA). Finally, reverse transcribed cDNAs were quantified by comparative Ct experiment on a StepOne™ Plus system (Applied BioSystems, Life Technologies, USA) using the Power SYBR Green PCR Master Mix (Applied BioSystems, Life Technologies, USA) and specific oligonucleotides: forward (F) 5'-ACACCCTTAACGAGATGGTCAC-3' and reverse (R) 5'-CCGCCGTGGTGTTTATTGTG-3' for PepT1 and (F) 5'-TGCCCTGAGGCTCTTTTCCA-3' and (R) 5'-GGCATAGAGGTCTTTACGGATGTC-3' for β -actin, all purchased from Eurogentec (France). The cycling conditions were 10 min at 95 °C, 40 cycles of 15 sec at 95 °C, 30 sec at 60 or 61 °C and 30 sec at 72 °C, followed by a melting curve step.

2.7. Statistical analysis

Data were expressed as mean $\pm$ standard deviation (SD) calculated over three independent experiments performed in triplicate. SigmaPlot 11.0 software (Germany) was used to carry out statistical analysis. One-Way ANOVA followed by a pairwise comparison with Tukey's test was used for comparison of data with normal distribution. p values < 0.05 were regarded as significant.

3. Results

3.1 Immunomodulatory capacities of the *Lactobacilli*

The three *Lactobacillus* strains, incubated alone and in 3RG multi-strain consortium with the peripheral blood mononuclear cells induced different cytokine secretion profiles. The two strains of *L. gasseri* ICV396 and ICV392 were not able to induce significant release of IL-10 conversely to *L. reuteri* ICVB395 strain which induced significant levels in comparison to untreated cells, reaching similar level (900 pg.mL⁻¹) observed with the control anti-inflammatory strain *B. longum*. The consortium 3RG was also able to induce IL-10 secretion after PBMC stimulation, but at a lower level, reaching approximately 400 pg.mL⁻¹ despite no significant difference when compared to the control (Figure 1A). The three individual *Lactobacillus* strains and the 3RG consortium were not able to induce detectable IL-12 and IFN γ secretion by PBMC (Figure 1B, 1C), while the control *L. lactis* strain induced significant amount of these two Th1/pro-inflammatory cytokines.

3.2 Capacity of the strains to strengthen the epithelial barrier

We evaluated the capacity of the strains to restore the gut barrier function using an *in vitro* model of H₂O₂-sensitized Caco-2 cells monolayers, as previously reported (J. Alard et al., 2018). As expected, H₂O₂ sensitization induced permeability as shown by a significant and time-dependent drop in the trans-epithelial electric resistance (TEER). All the strains were

able to restore the epithelial barrier, as shown by an increase in the TEER as compared to H₂O₂-sensitized control cells (Figure 2), which was however not significant for *L. gasseri* ICVB396 and *L. reuteri* ICVB395 which was able to maintain the TEER at the level of untreated control only until 60 min. Interestingly, the 3RG consortium and the *L. gasseri* ICVB392 strain not only restored but even reinforced the epithelial barrier, the TEER being higher than the H₂O₂ non-sensitized control monolayer.

3.3 Impact on lipid accumulation in adipocytes

Lipid accumulation observed in 3T3-L1 derived mature adipocytes treated with the *lactobacilli* were generally lower than the level found in non-treated adipocytes. The most important reduction was observed with the *L. reuteri* ICVB395 and *L. gasseri* ICVB396 strains, which induced significant drop of lipid level comparatively to the control, reaching respectively 30% and 20% decrease in comparison to the control level (Figure 3). The third strain, *L. gasseri* ICVB392, as well as the 3RG consortium showed also a similar tendency which was however not significant (Figure 3).

3.4 Capacity of the *lactobacilli* strains to modulate gut hormones secretion

3.4.1 Effects on active-GLP-1 secretion

We evaluated the capacity of the strains to induce the release of GLP-1 using the STC-1 enteroendocrine cell line. After 8 hours bacterial stimulation of STC-1 cells, the highest GLP-1 secretion was obtained with *L. gasseri* ICVB392 and *L. reuteri* ICVB395, which induced hormone secretion estimated to 6 fold higher than the control level. *L. gasseri* ICVB396 and the 3RG consortium induced also significant GLP-1 release, which was, however lower, reaching 4 fold and 3 fold of the control GLP-1 secretion level, respectively (Figure 4A).

3.5.2 Effects on CCK secretion

The *Lactobacillus* strains tested in this study exhibited moderate impact on the CCK secretion by stimulated STC-1 cells (Figure 4B). *L. reuteri* ICVB395 and *L. gasseri* ICVB396 showed the most important effect inducing significant CCK secretion, reaching level more than one fold and half of the control level. The association of these strain with *L. gasseri* ICVB392, in the 3RG consortium led to a decrease of the quantified CCK level compared to those obtained with ICVB395 and ICVB396 tested alone (Figure 4B). Purified peptidoglycan from *L. acidophilus* and flagellin from *S. enteritidis* serovar Thyphimurium didn't stimulate the secretion of both CCK and GLP-1 by STC-1 cells (data not shown).

3.5.3 Investigation of the different pathways involved on gut hormone secretion modulation

Since GLP-1 and CCK secretion are under the influence of three major signalling pathways implying protein G receptors, notably the calcium-sensing receptor (CaSR) and the GPR family C group 6 member A (GPRC6A), and the di/tripeptide transporter 1 (Pept1), we evaluated their respective role using CaSR and GPRC6A antagonists and Pept-1 inhibitor.

We first studied *Pept-1* gene expression in the enteroendocrine STC-1 cells. RT-PCR analysis unveiled a significant gene expression of Pept-1 with CT value obtained of 26.47 ± 0.44 while reference β -actin gene expression showed CT of 14.02 ± 0.22 (data not shown)

The addition of CaSR and GPRC6A antagonists induced a significant decrease of GLP-1 secretion induced by the stimulation of the STC-1 cells with the *lactobacilli* alone or with the 3RG consortium, except for GPRC6A inhibitor with ICVB396 strain (Figure 4A). Analysis of effect of CaSR antagonist indicated that the most important decrease is observed for the ICVB392 and ICVB395 strains for which the GLP-1 level decreased from 6 to around 4 fold the control level. The GLP-1 secretion level reduction observed for the ICVB396 strain and the 3RG consortium was moderate decreasing from 4 to 2.5 fold and from 3 to less than 2 fold

of the control level, respectively. Regarding the effect of the GPRC6A antagonist, we observed more or less the same effect with major decrease for the ICVB392 and ICV395 strains, and moderate for the 3RG consortium. However the decrease of GLP-1 level measured for the ICV396 pre-treated STC-1 was not significant (Figure 4A). Interestingly Pept-1 inhibitor induced a reduction of GLP-1 secretion for all strains and the 3RG consortium, nearly to the same levels obtained with GPRC6A inhibitor, except for the strain ICVB395 for which GLP-1 secretion was not impacted by AMBA-4 Pept-1 inhibitor. The results obtained for CCK were slightly different, as we observed a statistically significant effect just for the ICVB396 strains and 3RG consortium after pre-treatment with CaSR antagonist. GPRC6A and Pept-1 antagonist and inhibitor didn't appear to affect the CCK secretion level in a significant manner in all cases studied in this work, excepted for a slight effect which was observed with Pept-1 inhibitor and GPRC6A antagonists on the effect of ICVB392 strain and the 3RG consortium respectively (Figure 4B).

3.5.4 Capacity of selected *Lactobacilli* strains to degrade GLP-1 1-26 and CCK8S

When the GLP-1 1-26 (800 pM) and CCK8S (600 pM) peptide solutions (Figure 5, black bars) were incubated with the selected strains and the consortium for 8h, significant decreases of peptides were observed in the presence of *L. gasseri* ICVB396 and the 3RG multi-strains cocktail. Resulting concentrations of around 550 pM of active GLP-1 1-26, and of around 180 and 340 pM of CCK8S, were measured after incubation with *L. gasseri* ICVB396 strain and 3RG consortium, respectively. A significant impact was also observed for *L. gasseri* ICVB392 on the measured amount of CCK with a decrease from 600 to 400 pM after incubation.

In the presence of the tripeptide (Ile-Pro-Ile), a DPP-IV inhibitor (grey bars), the consumption of GLP-1 1-26 and CCK8S by *L. gasseri* ICVB392 and ICVB396 was reduced to nearly the

control level. However, the DPP-IV inhibitor did not significantly affect the consumption of GLP-1 and CCK by the 3RG consortium (Figure 5 A, B).

4. Discussion

In the present study, we investigated additional probiotic traits of *Lactobacillus* strains selected during previous works (Al Kassaa et al., 2014; Belguesmia et al., 2016). We previously established, using Caco-2 cells stimulation, that the three strains, selected in the present study, exhibited promising anti-inflammatory abilities, limiting the level of IL1- β -induced IL-8 release and inducing the secretion of IL-10 (Belguesmia et al., 2016). It has been pointed out that in vitro immunomodulation assays have to be considered carefully regarding the type and the physiological state of the eukaryotic cell models. Indeed probiotic strains could display different immunomodulatory profiles using epithelial cell model (i.e. HT-29 cell line) and PBMC stimulation used (Kechaou et al., 2013). Taking in account this postulate and in order to consolidate the results obtained on epithelial intestinal cells, we evaluated the immunomodulation capacities of the selected *Lactobacillus* strains, considered alone or in combination in the 3RG consortium, using in vitro PBMCs stimulation. We notably unravel their ability to induce the secretion of the anti-inflammatory IL-10 versus the Th1/pro-inflammatory IL-12 and IFN γ cytokines, We previously observed that in vitro immunomodulation abilities of lactobacilli are strain-specific and linked to their in vivo protective effects in murine models of colitis (Foligne et al., 2007).

We revealed during PBMC study that the *L. reuteri* ICVB395 strain was the most potent strain to induce the secretion of the anti-inflammatory IL-10 cytokine, while the two other *L. gasseri* strains, were not able to induce significant IL-10 response comparatively to the negative control. The 3RG consortium was able to induce moderate but significant secretion of this interleukin.

Nonetheless in the present study, none of the strains alone or in the 3RG consortium was able to induce IL-12 or interferon γ (IFN- γ) which overall remained undetectable (Figure 1C). IL-12 cytokine plays an important role in activating the Th1 immune response by promoting the differentiation of naïve T cell to Th1 cells to produce interferon γ (IFN- γ) (Meijerink et al., 2012; Watson, Sargianou, & Panos, 2012). In contrast, IL-10 is known to inhibit natural killer (NK) and Th1 cells, by down-regulating the IL-12 production and facilitating Th2 immune response (Tripp, Wolf, & Unanue, 1993; Uyemura et al., 1996). The balance between these two interleukins defines the immune response and plays a major role in the anti-inflammatory/pro-inflammatory state which is dysregulated during colitis, allergy, irritable bowel syndrome and other inflammatory diseases. Indeed, a high IL-10/IL-12 ratio could predict favourable anti-inflammatory abilities of probiotic strains, especially for *lactobacilli* (Foligne et al., 2007; Meijerink et al., 2012; Watson et al., 2012). Hence the bacterial strains tested showed differential abilities to influence the inflammatory state.

Chronic inflammatory diseases are often associated with an increased intestinal permeability known as leaky gut which facilitates the translocation of commensal bacteria, thus contributing to the development of a chronic inflammatory state. We then evaluated the capacity of the strains to strengthen the epithelial barrier using an in vitro model of epithelial barrier. All the strains were able to attenuate the H₂O₂-induced permeability. However, the best ability to strengthen the epithelial barrier was observed with *L. gasseri* ICVB392 and the 3RG consortium which were not only able to restore the H₂O₂-sensitized monolayer but were also able to reinforce the trans-epithelial resistance. Among studied probiotic properties of microorganisms, TEER assay appeared as a reliable method to anticipate and establish effect of potential probiotics strains on epithelium (J. Alard et al., 2018; Klingberg, Pedersen, Cencic, & Budde, 2005; Messaoudi et al., 2012). Most studies showed protective effect of probiotic strains, improving permeability of sensitized epithelial monolayer, with some strains

416 able also to reinforce the epithelium barrier with increased TEER values over the basal level
417 observed for untreated control epithelium (Anderson et al., 2013; Botta et al., 2014; Ramos,
418 Thorsen, Schwan, & Jespersen, 2013). Mechanisms implied in this phenomenon are still not
419 completely elucidated and seems to be strain-specific (Anderson et al., 2013; Ramos et al.,
420 2013). Some strains seem to display a negative effect (i.e. *L. fermentum* RGR1487) whereas
421 other (i.e. *L. fermentum* RGR1485) have neutral/positive effect on the TEER of Caco-2 cells
422 epithelium (Anderson et al., 2013). Ramos et al. (2013) isolated a number of probiotic
423 *lactobacilli*, belonging to *L. plantarum*, *L. brevis* and *L. fermentum* species, from different
424 Brazilian food products. Most of these strains showed enhanced TEER, but not at the same
425 level and independently from species consideration. In a previous study, we were also able to
426 select strains (*L. acidophilus* PI11, *L. helveticus* PI5 and *L. gasseri* LA806) able to restore and
427 reinforce the epithelial barrier using the same in vitro model (J. Alard et al., 2018). Recently,
428 Kawano et al., stated that the probiotic *L. gasseri* SBT2055 (LG2055) exhibits anti-obesity
429 effects by improving the intestinal integrity and thus reducing the entry of inflammatory
430 substances like endotoxin from the gut lumen, which may improve the inflammation state
431 within metabolic organs (Kawano, Miyoshi, Ogawa, Sakai, & Kadooka, 2016).

432 Another aspect of probiotic property investigated during our study is the ability of the strains
433 to influence the accumulation of lipid in adipocytes. We showed that two of the tested strains
434 (*L. reuteri* ICV39 and *L. gasseri* ICV396) induced significant reduction of lipid accumulation
435 in adipocytes. The third strain, *L. gasseri* ICVB392, doesn't reach such reduction but achieve
436 a lowering tendency. Similar behaviour was also observed for the 3RG consortium containing
437 the three strains with an intermediate effect but a lower efficiency than the *L. reuteri*
438 ICVB395 and *L. gasseri* ICVB396 strains considered alone. In previous work, Park et al.,
439 showed that a probiotic strain of *L. brevis*, designated KLEB, inhibited lipid accumulation in
440 the differentiated 3T3-L1 adipocytes by downregulating the expression of adipogenic

transcription factors and other specific genes involved in lipid metabolism, leading to the inhibition of adipocyte differentiation, intracellular triglyceride accumulation and a decrease of glycerol-3-phosphate dehydrogenase (GPDH) activity (Park et al., 2014). In an *in vivo* study using high fat diet-fed C57BL/6 mice, a strain of *L. bulgaricus* N°14 was clearly able to reduce adipocytes size, the weight of white adipose tissue and the serum leptin and cholesterol levels (Takemura et al., 2010).

In a recent review, Drissi et al. analysed the results of experimental and clinical studies which evaluated the impact of *Lactobacilli* on animal and human body weight and reported a strain specific effect. The administration of *L. reuteri*, *L. sakei*, *L. acidophilus* and *L. casei* was associated with weight gain in human, while the consumption of specific strains of *L. gasseri*, *L. amylovorus*, *L. plantarum* and some *L. acidophilus* strains was associated with weight loss in obese humans and body fat loss in overweight healthy individuals (Drissi et al., 2017). Stenman et al. also identified promising probiotics for preclinical studies including several *Lactobacillus* strains, notably *L. acidophilus* NCFM, *L. gasseri* 2055, *L. reuteri* GMNL-263 and the multistrain consortium LGG/Bb12 and VSL#3, which showed proven benefits on insulin insensitivity, fat accumulation and weight loss during *in vivo* assays in human and animal trials (Stenman, Burcelin, & Lahtinen, 2016).

Gut hormones, notably PYY and GLP-1, released from enteroendocrine cells within the gastrointestinal tract are known to play crucial role not only in the control of satiety and energy balance, but also, notably for GLP-1, numerous effects as incretin hormone on glucose homeostasis (Holst, 2007). GLP-1 was also recently shown to exhibit anti-inflammatory effects and to promote gut barrier integrity (Lebrun et al., 2017). In the present study, using the STC-1 cell line, we confirmed that the selected strains, *L. gasseri* ICVB392, *L. reuteri* ICVB395 and *L. gasseri* ICVB396, were able to modulate the secretion of active GLP-1 and CCK gut hormones. GLP-1 and CCK were demonstrated to be potential targets of probiotic

466 preparations to control food intake and regulate the appetite on mouse model (Yadav et al.,
467 2013). Interestingly, the two strains, *L. gasseri* ICVB392 and *L. reuteri* ICVB395, induced
468 strong GLP-1 secretion, reaching around 6 fold the control level, while the *L. gasseri*
469 ICVB396 induced lower amount however reaching 4 fold the basal level. Surprisingly,
470 mixing the 3 strains in the 3RG consortium seemed to provoke a reduction of the secreted
471 GLP-1, as compared to the levels obtained with individual strains, notably with the two
472 highest ones. Interestingly, we demonstrated that the *L. gasseri* ICVB396 was able, in contrast
473 to the two other strains, to degrade in vitro the active GLP-1 and this was also observed with
474 the mixture. We can thus hypothesize that the GLP-1 released upon STC-1 stimulation with
475 the *L. gasseri* strain, or by the 3RG consortium could be under-evaluated following its
476 degradation by proteases. This result was not surprising regarding the diversity of the
477 proteases present in **lactobacilli** (Law & Haandrikman, 1997; Liu et al., 2010). Dipeptidyl
478 peptidase IV (DPP-IV) is a prolyl oligopeptidase, member of the serine proteases, able to
479 cleave the incretin hormone GLP-1, playing thus a determinant role on its metabolic and
480 immune functions. Two forms of DPP-IV have been described, a soluble circulating form and
481 a transmembrane serine exopeptidase (Aso et al., 2012). DPP-IV hydrolyses the GLP-1(7–36)
482 amide to generate GLP-1(9–36) amide and the N-terminal histidine–alanine dipeptide
483 (Nadkarni, Chepurny, & Holz, 2014). We therefore use a competitive inhibitor of the DPP-IV,
484 the Ile-Pro-Ile tripeptide called diprotin A, to ensure that observed consumption, or
485 degradation, of GLP-1 in the STC-1 culture supernatants was not linked to the presence of
486 serine protease DPP-IV-like enzymes produced by the *Lactobacillus* strains. We showed that
487 the GLP-1 degradation was inhibited by the addition of the competitive inhibitor, while the
488 effect of the 3RG consortium on GLP-1 release seemed to be insensitive to the treatment with
489 the Ile-Pro-Ile tripeptide, maintaining the same decreased level of GLP-1, in the presence or
490 absence of the inhibitor. The peptidic nature of the DPP-IV inhibitor made it susceptible to be

degraded by enzymes secreted by the other *lactobacilli*, explaining the absence of inhibitory effects on serine proteases implied in the degradation of GLP-1 1-26 and CCK-8S hormones used in this study. However it is difficult to elucidate the potential effect of enzymes interaction when the three strains consortium partners are mixed in the 3RG multi-strains cocktail. Indeed this interesting observation pointed out the limits of the in vitro models used. However this observation need to be relativized in in vivo conditions, in which GLP-1 is secreted at the basal side of the intestinal epithelium, making the degradation of this hormone by the bacteria not conceivable, since the microorganisms interact with these cells on their apical side (Bohórquez & Liddle, 2011).

Similarly to the results observed on GLP-1, CCK secretion level varied significantly when the three strains were mixed in the 3RG consortium. The three *Lactobacillus* strains appeared to be able to increase the level of the CCK secreted by the STC-1 cells. Although the CCK level obtained after incubation with the 3RG multi-strains cocktail was lower (by 20%) than the level obtained with *L. reuteri* ICVB395 or *L. gasseri* ICVB396 tested alone. As for GLP-1 experiment, we estimate the ability of the *selected Lactobacilli* strains selected for this study to consume or degrade the CCK8S hormone. We also observed a strong reduction of the CCK8S level when co-incubated with the ICVB392 and ICVB396 strains, or the 3RG consortium. Moreover as for the GLP-1 1-26, the CCK consumption by the two strains was dramatically reduced by the addition of diprotin A. Again, the impact of the 3RG consortium seemed to be insensitive to the addition of the serine protease inhibitor, as the level of the CCK remained the same observed with and without the addition of the diprotin A (Figure 5). This could be explained also in this case by the complex interactions occurring in a multi-strains mixture between strains and STC-1 cells. Rose et al. established that CCK-8 can be cleaved by serine peptidases, an isoform of tripeptidyl peptidase II notably present in the rat brain, and producing CCK-5 and GWM as major fragments. Interestingly CCK-8 degradation

516 was limited by serine-alkylating reagents, permitting recovery of complete initial amount of
517 CCK-8 (C. Rose, Camus, & Schwartz, 1988; Christiane Rose et al., 1996). Interaction of
518 probiotic preparations with intestinal cells, in the control of gut hormone secretion, is the
519 subject of many researches. Most of them, notably using experimental murine models of
520 obesity, pointed the importance of short chain fatty acids (SCFA), which are bacterial
521 fermentation products of soluble fibers by commensal bacteria, on the food intake and weight
522 regulation (Jeanne Alard et al., 2016; Wang et al., 2015; Yadav et al., 2013). However none of
523 the selected strains used in this study are able to produce the main SCFA involved in energy
524 balance, namely butyrate, propionate or acetate, since no detectable level were measured
525 using gas chromatography analysis in the supernatant of the STC-1 cells culture after 8h of
526 incubation with the chosen *Lactobacillus* strains (data not shown). We then investigated the
527 mechanism involved in the interaction between the *lactobacilli* strains and the STC-1 cells.
528 Enteroendocrine mammalian cells (EECs) can interact with bacteria, or derived components,
529 through different receptors. The apical portion of enteroendocrine cells, in contact the gut
530 lumen, may sense bacterial inputs through pattern recognition receptors (PRR) such as toll-
531 like receptors (e.g., TLRs 4, 5, and 9), which are expressed in EECs. Stimulation with
532 bacterial ligands (e.g., LPS or flagellin) following direct contact, have been shown to promote
533 the secretion of gut hormones (Bohórquez & Liddle, 2011). However many other receptors
534 could be involved in interaction with bacteria, we therefore evaluated other possible pathways
535 which could affect the gut hormones secretion.

536 GLP-1 and CCK secretion are under the influence of three major signalling pathways
537 implying protein G receptors, notably the calcium-sensing receptor (CaSR) and the GPR
538 family C group 6 member A (GPRC6A), and the di/tripeptide transporter 1 (Pept1), involving
539 ERK 1/2 phosphorylation or proton uptake coupled to peptide transport (Caron et al., 2017).
540 These receptors were demonstrated to be sensitive to the action of peptide fragments and

541 amino-acids, and their activation led to the release of GLP-1 and CCK gut hormones in the
542 circulatory system (Caron et al., 2017).

543 Previous research established the expression of the gene encoding Pept-1 in the STC-1 cell
544 line (Liou et al., 2011), while other published work indicated low expression levels
545 (Diakogiannaki et al., 2013). We were able, using quantitative PCR, to highlight *pept1* gene
546 expression in the STC-1 cells we used. We therefore evaluate the possible implication of this
547 transporter in the interaction between the *Lactobacillus* strains and the STC-1 cells, as well as
548 the two GRPs.

549 We demonstrated that the inhibitory action of specific antagonists and inhibitors of those
550 receptors and transporter led to different effects on the impact of the tested strains ICVB392,
551 ICVB395 and ICVB396, and the consortium 3RG. The response to the two strains of *L.*
552 *gasseri* ICVB392 and the ICVB396 was modified by the antagonists of CaSR and GPRC6A,
553 and the inhibitor of the Pept1 transporter, leading globally to the decrease GLP-1 secretion in
554 the medium. Although the effect obtained with the *L. reuteri* ICVB395 strain appeared to be
555 insensitive to the action of the Pept1 inhibitor AMBA-4, interestingly the effects of the
556 inhibitor and antagonists were significant for the 3RG consortium. These observations
557 indicated differential interaction of the *Lactobacillus* strains with the STC-1 cells involved in
558 GLP-1 secretion. Conversely the effect of Pept1 inhibitors and antagonists of CasR and
559 GPRC6A on CCK secretion were more limited. However the CaSR antagonist and the Pept1
560 inhibitor affected significantly the interaction of STC-1 with *L. gasseri* ICVB392 and
561 ICVB396, involved in CCK secretion. However the ability of these strains to degrade the
562 GLP-1 and the CCK could minimise the observed induction effects on hormone secretion
563 measured in the supernatant of STC-1 cells culture. The calcium-sensing receptor (CaSR) is a
564 Class C G-protein coupled receptor which senses extracellular levels of calcium ion (Vezzoli,
565 Soldati, & Gambaro, 2009). Various metabolites could activate CaSR response leading to

566 GLP-1 and CCK secretion, like Ca^{++} ions and basic L-amino acids like L-phenylalanine
567 (Caron et al., 2017). Previous studies on *L. gasseri* strains, isolated from vaginal environment,
568 established that those lactic acid bacteria, with good aggregation abilities, possess specific
569 surface protein designated aggregation promoting factor (*afp*) (Boris, Suarez, & Barbes, 1997).
570 This protein appeared to contain basic amino acids and phenylalanine residues which can
571 interact with the surface of host cells (Boris et al., 1997; Jankovic et al., 2003).

572 The GPRC6A receptor is a newly orphanized class C GPCR, for which the first allosteric
573 antagonist, based on the 2-arylindole privileged structure scaffold, was recently reported
574 (Johansson et al., 2015). GPRC6A activation led to initiating signalling cascade,
575 intermediated by $\text{G}\alpha_q$ subunit, which induce intracellular Ca^{++} increase and GLP-1 hormone
576 secretion (Caron et al., 2017). Wellendorph et al. established that the GPRC6A was activated
577 by basic α -amino acids, L-Argenin, L-Lysine and L-Ornithine being the most active agonists
578 for this receptor. The role of these amino acids in activation of GPRC6A receptor was
579 confirmed by Oya et al. using GLUTag and STC-1 enteroendocrine cell lines. Furthermore
580 GPRC6A receptor antagonists, a phospholipase C inhibitor or an IP3 receptor antagonist,
581 suppress significantly the L-Ornithine-induced intracellular Ca^{++} concentration and GLP-1
582 secretion, identifying the involvement of this pathway in the hormone secretion (Oya et al.,
583 2013; Wellendorph et al., 2005).

584 Interestingly, peptides and amino acids are generally sensed by EEC via the proton-coupled
585 peptide-transporter PepT1, this nutrient-sensing being linked to hormone secretion (Zietek &
586 Rath, 2016). We previously demonstrated that the muropeptide M-tri-Lys, derived from the
587 peptidoglycan of a selected *Lactobacillus* strain, was able to protect mice from colitis in a
588 NOD2-dependant manner, while the M-tri-Lys-N was not, suggesting that only the M-tri-Lys
589 could be internalized by PepT1 or PepT2 transporters into the cell to interact with the
590 cytosolic receptor NOD2 (Macho-Fernandez et al., 2011). The importance of the main TLR

591 adaptator MyD88 was also linked to the increase of GLP-1 and glucose-dependent
592 insulintropic polypeptide (GIP) secretion when STC-1 cells were co-cultured with
593 *Lactobacilli* strains (Panwar et al., 2016). Moreover it was established that the modulation of
594 the microbiota/SCFA/bile-acid-signalling, impacted the metabolism and the inflammatory
595 processes via GLP-1 and other markers, representing an interesting target for the treatment of
596 chronic diseases (Zietek & Rath, 2016).

597 In vitro tests realized during this study showed that the 3RG consortium exhibited different
598 response profiles on GLP-1 and CCK hormone secretion, in comparison to the strains
599 considered alone. However the association of these strains could retain our interest as it could
600 combine the potential of each strain maximizing their combined effect on the metabolism,
601 even if this was not observed in our experimental in vitro conditions. It remains thus
602 important to evaluate such potential in in vivo animal models to definitively establish the
603 beneficial impact of the proposed 3RG consortium.

604 In conclusion our study established that the selected *Lactobacillus* strains we evaluated in this
605 work, presented interesting probiotic potential regarding in vitro experiment results. Positive
606 impacts on anti-inflammatory cytokine secretion, decrease in lipid accumulation in adipocytes
607 and restoration of the epithelial barrier, were highlighted for ICVB395 strain and in a more
608 limited tendency for 3RG consortium. ICV 392 was the best potent strain to strengthen the
609 epithelial barrier together with the 3RG consortium. The impact of the *lactobacilli* on gut
610 hormone GLP-1 and CCK secretion appeared more difficult to analyse due to the limit of the
611 in vitro enteroendocrine STC-1 cell model used. We indeed established that some of the
612 *Lactobacillus* strains have the ability to degrade the GLP-1 1-26 and CCK-8S, probably,
613 through the action of serine peptidases. To resolve this question a relevant alternative would
614 be to use a polarized EEC in vitro model using insert (transwell) system, in which gut
615 hormones would be secreted at the basolateral compartment while bacteria would be added at

the apical surface. A recent study published by Zhang et al., described such approach using secreting GLP-1 Caco-2 cells grown in vitro on inserts (Zhang, Liu, Chen, & Luo, 2018). However, since all these in vitro models present certain limits, we plan to unravel the probiotic potential of these strains in appropriate in vivo models to confirm their beneficial capacities in the context of chronic diseases such as IBD and obesity.

Acknowledgments

This work was supported by a grant of the Nord-Pas-de-Calais Region: “2nd appel à projet, Programme projets émergents.” It has also been carried out in the framework of Alibiotech project which is financed by the European Union, the French State, and the French Region of Hauts-de-France and was also supported by the Institut Pasteur de Lille and the Centre National de la Recherche Scientifique (CNRS). The experiments were partly performed at IUT A which is gratefully acknowledged.

Bibliography

- Al Kassaa, I., Hamze, M., Hober, D., Chihib, N.-E., & Drider, D. (2014). Identification of Vaginal Lactobacilli with Potential Probiotic Properties Isolated from Women in North Lebanon. *Microbial Ecology*, 67(3), 722-734. <https://doi.org/10.1007/s00248-014-0384-7>
- Alard, J., Peucelle, V., Boutillier, D., Breton, J., Kuylle, S., Pot, B., Holowacz, S., Grangette, C. (2018). New probiotic strains for inflammatory bowel disease management identified by combining in vitro and in vivo approaches. *Beneficial Microbes*, 9(2), 317-331. <https://doi.org/10.3920/BM2017.0097>
- Alard, J., Lehrter, V., Rhimi, M., Mangin, I., Peucelle, V., Abraham, A.L., Mariadassou, M., Maguin, E., Waligora-Dupriet, A.J., Pot, B., Wolowczuk, I., & Grangette C. (2016).

641 Beneficial metabolic effects of selected probiotics on diet-induced obesity and insulin
642 resistance in mice are associated with improvement of dysbiotic gut microbiota.
643 *Environmental Microbiology*. 18(5):1484-97. doi: 10.1111/1462-2920.13181.

644 Anderson, R. C., Young, W., Clerens, S., Cookson, A. L., McCann, M. J., Armstrong, K. M.,
645 & Roy, N. C. (2013). Human oral isolate *Lactobacillus fermentum* AGR1487 reduces
646 intestinal barrier integrity by increasing the turnover of microtubules in Caco-2 cells.
647 *PLoS ONE*, 8(11), e78774.

648 Arulampalam, V., Greicius, G., & Pettersson, S. (2006). The long and winding road to gut
649 homeostasis. *Current opinion in gastroenterology*, 22(4), 349–353.

650 Aso, Y., Ozeki, N., Terasawa, T., Naruse, R., Hara, K., Suetsugu, M., Takebayashi, K.,
651 Shibazaki, M., Haruki, K., Morita, K., & Inukai, T.. (2012). Serum level of soluble
652 CD26/dipeptidyl peptidase-4 (DPP-4) predicts the response to sitagliptin, a DPP-4
653 inhibitor, in patients with type 2 diabetes controlled inadequately by metformin and/or
654 sulfonylurea. *Translational Research*, 159(1), 25–31.

655 Beaugerie, L., & Petit, J.-C. (2004). Microbial-gut interactions in health and disease.
656 Antibiotic-associated diarrhoea. *Best Practice & Research. Clinical Gastroenterology*,
657 18(2), 337-352. <https://doi.org/10.1016/j.bpg.2003.10.002>

658 Belguesmia, Y., Domenger, D., Caron, J., Dhulster, P., Ravallec, R., Drider, D., & Cudennec,
659 B. (2016). Novel probiotic evidence of *Lactobacilli* on immunomodulation and
660 regulation of satiety hormones release in intestinal cells. *Journal of Functional Foods*,
661 24, 276-286.

662 Bohórquez, D. V., & Liddle, R. A. (2011). Axon-like basal processes in enteroendocrine cells:
663 characteristics and potential targets. *Clinical and translational science*, 4(5), 387–391.

664 Boris, S., Suarez, J. E., & Barbes, C. (1997). Characterization of the aggregation promoting
665 factor from *Lactobacillus gasseri*, avaginal isolate. *Journal of applied microbiology*,
666 83(4), 413–420.

667 Botta, C., Langerholc, T., Cencič, A., & Cocolin, L. (2014). In vitro selection and
668 characterization of new probiotic candidates from table olive microbiota. *PLoS One*,
669 9(4), e94457.

670 Cani, P. D., & Knauf, C. (2016). How gut microbes talk to organs: the role of endocrine and
671 nervous routes. *Molecular metabolism*, 5(9), 743–752.

672 Caron, J., Domenger, D., Dhulster, P., Ravallec, R., & Cudennec, B. (2017). Protein
673 digestion-derived peptides and the peripheral regulation of food intake. *Frontiers in*
674 *endocrinology*, 8, 85.

675 Chantret, I., Rodolosse, A., Barbat, A., Dussaulx, E., Brot-Laroche, E., Zweibaum, A., &
676 Rousset, M. (1994). Differential expression of sucrase-isomaltase in clones isolated
677 from early and late passages of the cell line Caco-2: evidence for glucose-dependent
678 negative regulation. *Journal of Cell Science*, 107(1), 213–225.

679 Clarke, G., Stilling, R. M., Kennedy, P. J., Stanton, C., Cryan, J. F., & Dinan, T. G. (2014).
680 Minireview: Gut microbiota: the neglected endocrine organ. *Molecular Endocrinology*
681 *(Baltimore, Md.)*, 28(8), 1221–1238. <https://doi.org/10.1210/me.2014-1108>

682 Collado, M. C., Meriluoto, J., & Salminen, S. (2007). In vitro analysis of probiotic strain
683 combinations to inhibit pathogen adhesion to human intestinal mucus. *Food Research*
684 *International*, 40(5), 629–636.

685 De Man, J. C., Rogosa, deM, & Sharpe, M. E. (1960). A medium for the cultivation of
686 *Lactobacilli*. *Journal of applied Bacteriology*, 23(1), 130–135.

687 Delzenne, N. M., & Cani, P. D. (2011). Gut microbiota and the pathogenesis of insulin
688 resistance. *Current diabetes reports*, 11(3), 154.

689 Di Cerbo, A., Palmieri, B., Aponte, M., Morales-Medina, J. C., & Iannitti, T. (2015).
690 Mechanisms and therapeutic effectiveness of Lactobacilli. *Journal of clinical*
691 *pathology*, jclinpath–2015.

692 Diakogiannaki, E., Pais, R., Tolhurst, G., Parker, H. E., Horscroft, J., Rauscher, B., ...
693 Reimann, F. (2013). Oligopeptides stimulate glucagon-like peptide-1 secretion in mice
694 through proton-coupled uptake and the calcium-sensing receptor. *Diabetologia*, 56(12),
695 2688-2696.

696 Doré, J., Multon, M.-C., Béhier, J.-M., & participants of Giens XXXII, Round Table No. 2.
697 (2017). The human gut microbiome as source of innovation for health: Which
698 physiological and therapeutic outcomes could we expect? *Thérapie*, 72(1), 21-38.
699 <https://doi.org/10.1016/j.therap.2016.12.007>

700 Drissi, F., Raoult, D., & Merhej, V. (2017). Metabolic role of Lactobacilli in weight
701 modification in humans and animals. *Microbial pathogenesis*, 106, 182–194.

702 FAO/WHO. (2002). *Joint FAO/WHO working group report on drafting guidelines for the*
703 *evaluation of probiotics in food*. FAO/WHO London, ON.

704 Fernandez, L. M. B., Lasa, J. S., & Man, F. (2014). Intestinal microbiota: its role in digestive
705 diseases. *Journal of clinical gastroenterology*, 48(8), 657–666.

706 Foligne, B., Nutten, S., Grangette, C., Dennin, V., Goudercourt, D., Poiret, S., Dewulf, J.,
707 Brassart, D., Mercenier, A., & Pot, B. (2007). Correlation between in vitro and in vivo
708 immunomodulatory properties of lactic acid bacteria. *World journal of*
709 *gastroenterology: WJG*, 13(2), 236.

710 Fooks, L. J., & Gibson, G. R. (2002). Probiotics as modulators of the gut flora. *British*
711 *Journal of Nutrition*, 88(S1), s39–s49.

712 Gionchetti, P., Lammers, K. M., Rizzello, F., & Campieri, M. (2005). VSL# 3: an analysis of
 713 basic and clinical contributions in probiotic therapeutics. *Gastroenterology Clinics*,
 714 34(3), 499–513.

715 Holst, J. J. (2007). The physiology of glucagon-like peptide 1. *Physiological reviews*, 87(4),
 716 1409–1439.

717 Hugas, M., & Monfort, J. M. (1997). Bacterial starter cultures for meat fermentation. *Food*
 718 *chemistry*, 59(4), 547–554.

719 Jankovic, I., Ventura, M., Meylan, V., Rouvet, M., Elli, M., & Zink, R. (2003). Contribution
 720 of aggregation-promoting factor to maintenance of cell shape in *Lactobacillus gasseri*
 721 4B2. *Journal of bacteriology*, 185(11), 3288–3296.

722 Jayasinghe, T. N., Chiavaroli, V., Holland, D. J., Cutfield, W. S., & O’Sullivan, J. M. (2016).
 723 The new era of treatment for obesity and metabolic disorders: evidence and
 724 expectations for gut microbiome transplantation. *Frontiers in cellular and infection*
 725 *microbiology*, 6, 15.

726 Johansson, H., Boesgaard, M. W., Nørskov-Lauritsen, L., Larsen, I., Kuhne, S., Gloriam, D.
 727 E., Bräuner-Osborne, H., & Sejer Pedersen, D. (2015). Selective Allosteric
 728 Antagonists for the G Protein-Coupled Receptor GPRC6A Based on the 2-
 729 Phenylindole Privileged Structure Scaffold. *Journal of Medicinal Chemistry*, 58(22),
 730 8938-8951. <https://doi.org/10.1021/acs.jmedchem.5b01254>

731 Kawano, M., Miyoshi, M., Ogawa, A., Sakai, F., & Kadooka, Y. (2016). *Lactobacillus gasseri*
 732 SBT2055 inhibits adipose tissue inflammation and intestinal permeability in mice fed
 733 a high-fat diet. *Journal of nutritional science*, 5.

734 Kechaou, N., Chain, F., Gratadoux, J.-J., Blugeon, S., Bertho, N., Chevalier, C., Le Goffic, R.,
 735 Courau, S., Molimard, P., Chatel, J.M., Langella, P., Bermúdez-Humarán, L. G.
 736 (2013). Identification of One Novel Candidate Probiotic *Lactobacillus plantarum*

737 Strain Active against Influenza Virus Infection in Mice by a Large-Scale Screening.
738 *Applied and Environmental Microbiology*, 79(5), 1491-1499.
739 <https://doi.org/10.1128/AEM.03075-12>

740 Klingberg, T. D., Pedersen, M. H., Cencic, A., & Budde, B. B. (2005). Application of
741 measurements of transepithelial electrical resistance of intestinal epithelial cell
742 monolayers to evaluate probiotic activity. *Applied and environmental microbiology*,
743 71(11), 7528–7530.

744 Law, J., & Haandrikman, A. (1997). Proteolytic enzymes of lactic acid bacteria. *International*
745 *Dairy Journal*, 7(1), 1–11.

746 Lebrun, L. J., Lenaerts, K., Kiers, D., de Barros, J.-P. P., Le Guern, N., Plesnik, J., Thomas,
747 C., Bourgeois, T., Dejong, C.H.C., Kox, M., Hundscheid, I.H.R., Khan, N.A.,
748 Mandard, S., Deckert, V., Pickkers, P., Drucker, D.J., Lagrost, L., & Grober, J.. (2017).
749 Enteroendocrine L cells sense LPS after gut barrier injury to enhance GLP-1 secretion.
750 *Cell reports*, 21(5), 1160–1168.

751 Liou, A. P., Chavez, D. I., Espero, E., Hao, S., Wank, S. A., & Raybould, H. E. (2011).
752 Protein hydrolysate-induced cholecystokinin secretion from enteroendocrine cells is
753 indirectly mediated by the intestinal oligopeptide transporter PepT1. *American*
754 *Journal of Physiology - Gastrointestinal and Liver Physiology*, 300(5), G895-G902.
755 <https://doi.org/10.1152/ajpgi.00151.2011>

756 Liu, M., Bayjanov, J. R., Renckens, B., Nauta, A., & Siezen, R. J. (2010). The proteolytic
757 system of lactic acid bacteria revisited: a genomic comparison. *BMC genomics*, 11(1),
758 36.

759 Macho-Fernandez, E., Valenti, V., Rockel, C., Hermann, C., Pot, B., Boneca, I. G., &
760 Grangette, C. (2011). Anti-inflammatory capacity of selected Lactobacilli in
761 experimental colitis is driven by NOD2-mediated recognition of a specific

762 peptidoglycan-derived muropeptide. *Gut*, gut.2010.232918.
 763 <https://doi.org/10.1136/gut.2010.232918>

764 Margono, T., Sumaryono, W., Malik, A., & Sadikin, M. (2014). Characterization of Trypsin-
 765 Like Protease of *Lactobacillus plantarum* FNCC 0270. *HAYATI Journal of*
 766 *Biosciences*, 21(2), 87–94.

767 Meijerink, M., Wells, J. M., Taverne, N., de Zeeuw Brouwer, M.-L., Hilhorst, B., Venema, K.,
 768 & van Bilsen, J. (2012). Immunomodulatory effects of potential probiotics in a mouse
 769 peanut sensitization model. *FEMS Immunology & Medical Microbiology*, 65(3), 488–
 770 496.

771 Messaoudi, S., Madi, A., Prévost, H., Feuilleley, M., Manai, M., Dousset, X., & Connil, N.
 772 (2012). In vitro evaluation of the probiotic potential of *Lactobacillus salivarius*
 773 SMXD51. *Anaerobe*, 18(6), 584–589.

774 Morelli, L. (2000). In vitro selection of probiotic Lactobacilli: a critical appraisal. *Current*
 775 *Issues in Intestinal Microbiology*, 1(2), 59–67.

776 Nadkarni, P., Chepurny, O. G., & Holz, G. G. (2014). Regulation of glucose homeostasis by
 777 GLP-1. In *Progress in molecular biology and translational science* (Vol. 121, p. 23–
 778 65). Elsevier.

779 Oya, M., Kitaguchi, T., Pais, R., Reimann, F., Gribble, F., & Tsuboi, T. (2013). The G
 780 protein-coupled receptor family C group 6 subtype A (GPRC6A) receptor is involved
 781 in amino acid-induced glucagon-like peptide-1 secretion from GLUTag cells. *Journal*
 782 *of Biological Chemistry*, 288(7), 4513–4521.

783 Panwar, H., Calderwood, D., Gillespie, A. L., Wylie, A. R., Graham, S. F., Grant, I. R.,
 784 Grover, S. & Green, B. D. (2016). Identification of lactic acid bacteria strains
 785 modulating incretin hormone secretion and gene expression in enteroendocrine cells.
 786 *Journal of Functional Foods*, 23, 348–358.

787 Park, J.-E., Oh, S.-H., & Cha, Y.-S. (2014). *Lactobacillus brevis* OPK-3 isolated from kimchi
788 inhibits adipogenesis and exerts anti-inflammation in 3T3-L1 adipocyte. *Journal of the*
789 *Science of Food and Agriculture*, 94(12), 2514–2520.

790 Perdigon, G., Galdeano, C. M., Valdez, J. C., & Medici, M. (2003). Interaction of lactic acid
791 bacteria with the gut immune system. *European journal of clinical nutrition*, 56(S4),
792 S21.

793 Quigley, E. M. M. (2013). Bugs on the brain; brain in the gut—seeking explanations for
794 common gastrointestinal symptoms. *Irish journal of medical science*, 182(1), 1–6.

795 Ramos, C. L., Thorsen, L., Schwan, R. F., & Jespersen, L. (2013). Strain-specific probiotics
796 properties of *Lactobacillus fermentum*, *Lactobacillus plantarum* and *Lactobacillus*
797 *brevis* isolates from Brazilian food products. *Food microbiology*, 36(1), 22–29.

798 Rose, C., Camus, A., & Schwartz, J. C. (1988). A serine peptidase responsible for the
799 inactivation of endogenous cholecystokinin in brain. *Proceedings of the National*
800 *Academy of Sciences*, 85(21), 8326–8330.

801 Rose, Christiane, Vargas, F., Facchinetti, P., Bourgeat, P., Bambal, R. B., Bishop, P. B., Chan,
802 S.M., Moore, A.N., Ganellin, C.R., & Schwartz, J.-C. (1996). Characterization and
803 inhibition of a cholecystokinin-inactivating serine peptidase. *Nature*, 380(6573), 403.

804 Sánchez, B., Delgado, S., Blanco-Míguez, A., Lourenço, A., Gueimonde, M., & Margolles, A.
805 (2017). Probiotics, gut microbiota, and their influence on host health and disease.
806 *Molecular nutrition & food research*, 61(1), 1600240.

807 Sanders, M. E. (2016). Probiotics and microbiota composition. *BMC medicine*, 14(1), 82.

808 Stenman, L. K., Burcelin, R., & Lahtinen, S. (2016). Establishing a causal link between gut
809 microbes, body weight gain and glucose metabolism in humans—towards treatment
810 with probiotics. *Beneficial microbes*, 7(1), 11–22.

811 Stubbendieck, R. M., Vargas-Bautista, C., & Straight, P. D. (2016). Bacterial communities:
812 interactions to scale. *Frontiers in microbiology*, 7, 1234.

813 Takemura, N., Okubo, T., & Sonoyama, K. (2010). *Lactobacillus plantarum* strain No. 14
814 reduces adipocyte size in mice fed high-fat diet. *Experimental biology and medicine*,
815 235(7), 849–856.

816 Timmerman, H. M., Koning, C. J. M., Mulder, L., Rombouts, F. M., & Beynen, A. C. (2004).
817 Monostrain, multistrain and multispecies probiotics—a comparison of functionality
818 and efficacy. *International journal of food microbiology*, 96(3), 219–233.

819 Tripp, C. S., Wolf, S. F., & Unanue, E. R. (1993). Interleukin 12 and tumor necrosis factor
820 alpha are costimulators of interferon gamma production by natural killer cells in
821 severe combined immunodeficiency mice with listeriosis, and interleukin 10 is a
822 physiologic antagonist. *Proceedings of the National Academy of Sciences*, 90(8),
823 3725–3729.

824 Uyemura, K., Demer, L. L., Castle, S. C., Jullien, D., Berliner, J. A., Gately, M. K., Warrier,
825 R.R., Pham, N., Fogelman, A.M., & Modlin, R. L. (1996). Cross-regulatory roles of
826 interleukin (IL)-12 and IL-10 in atherosclerosis. *The Journal of clinical investigation*,
827 97(9), 2130–2138.

828 Vaughan, E. E., & Mollet, B. (1999). Probiotics in the new millennium. *Food/Nahrung*, 43(3),
829 148–153.

830 Vezzoli, G., Soldati, L., & Gambaro, G. (2009). Roles of calcium-sensing receptor (CaSR) in
831 renal mineral ion transport. *Current pharmaceutical biotechnology*, 10(3), 302–310.

832 Wang, J., Tang, H., Zhang, C., Zhao, Y., Derrien, M., Rocher, E., van-Hylckama Vlieg, J.E.,
833 Strissel, K., Zhao, L., Obin, M., & Shen, J. (2015). Modulation of gut microbiota
834 during probiotic-mediated attenuation of metabolic syndrome in high fat diet-fed mice.
835 *The ISME journal*, 9(1), 1.

- Watson, D. C., Sargianou, M., & Panos, G. (2012). Interleukin-12 (IL-12)/IL-10 Ratio as a Marker of Disease Severity in Crimean-Congo Hemorrhagic Fever. *Clinical and Vaccine Immunology*, 19(5), 823-824. <https://doi.org/10.1128/CVI.00030-12>
- Wellendorph, P., Hansen, K. B., Balsgaard, A., Greenwood, J. R., Egebjerg, J., & Bräuner-Osborne, H. (2005). Deorphanization of GPRC6A: a promiscuous L-alpha-amino acid receptor with preference for basic amino acids. *Molecular Pharmacology*, 67(3), 589-597. <https://doi.org/10.1124/mol.104.007559>
- Yadav, H., Lee, J.-H., Lloyd, J., Walter, P., & Rane, S. G. (2013). Beneficial metabolic effects of a probiotic via butyrate-induced GLP-1 hormone secretion. *The Journal of Biological Chemistry*, 288(35), 25088-25097. <https://doi.org/10.1074/jbc.M113.452516>
- Zebisch, K., Voigt, V., Wabitsch, M., & Brandsch, M. (2012). Protocol for effective differentiation of 3T3-L1 cells to adipocytes. *Analytical Biochemistry*, 425(1), 88-90. <https://doi.org/10.1016/j.ab.2012.03.005>
- Zhang, C., Liu, H., Chen, S., & Luo, Y. (2018). Evaluating the effects of IADHFL on inhibiting DPP-IV activity and expression in Caco-2 cells and contributing to the amount of insulin released from INS-1 cells in vitro. *Food & Function*, 9(4), 2240-2250. <https://doi.org/10.1039/C7FO01950E>
- Zietek, T., & Rath, E. (2016). Inflammation Meets Metabolic Disease: Gut Feeling Mediated by GLP-1. *Frontiers in Immunology*, 7, 154. <https://doi.org/10.3389/fimmu.2016.00154>

Figures captions:

Figure. 1 Immunomodulation capacities of the selected strains

Peripheral blood mononuclear cells (PBMCs) derived from human blood were stimulated *in vitro* with freshly cultured bacteria at a ratio of 10:1 (bacteria/cells) for 24h. *B. longum* strain was added as anti-inflammatory control and *L. lactis* MG1363 strain as pro-inflammatory control. Control PBMCs were only treated with PBS buffer. Levels of IL-10 (A), IL-12 (B) and IFN γ (C) were measured by ELISA in the supernatant after 24 h stimulation with the selected bacteria. GraphPad Prism was employed for graph preparation and statistical evaluation. Differences between groups were assessed using ANOVA, followed by nonparametric Mann-Whitney test. Data with p value ≤ 0.05 were considered to be significant.

Figure. 2 Capacity of the selected **lactobacilli** to restore or strengthen epithelial barrier in polarized Caco-2 monolayers sensitized with hydrogen peroxide (H₂O₂).

Caco-2 monolayers were treated at the apical side with the bacteria (at 10:1 bacteria/cell ratio) 30 mn before the addition of H₂O₂ (100 μ M). Changes in trans-epithelial electrical resistance (TEER) across Caco-2 cell monolayers were measured before the addition of H₂O₂ (T₀) and every 30 mn until 120 mn. Results were expressed as % TEER compared to T₀. Values represent the mean of 3 repeated experiments. Means without a common letter are different (p<0.05) using one way ANOVA with Tukey *post hoc* test for pairwise comparisons.

Figure. 3 Capacity of the strains to limit lipid accumulation.

Mature 3T3-L1 derived adipocytes were treated for 24h with the bacteria (10:1 bacteria/ cell ratio). **Lipid content was compared** using red-oil staining **and reported** in percentage of level obtained with **lactobacilli**-treated cells versus control untreated cells **(normalized at 100%)**. Values are the mean of four repeated experiments. Means without a common letter are different (p<0.05) using one way ANOVA with Tukey *post hoc* test for pairwise comparisons.

Figure. 4 Capacity of the strains to induce the release of gut peptides and impact of specific antagonists of CasR and GPRC6A and Pept-1 inhibitor.

CasR antagonist NPS 2143 (25 μ M, dark grey bars), GPRC6A antagonist CpD (50 μ M, grey bars) and Pept-1 inhibitor AMBA-4 (10 mM, black bars) were added (or not, white bars) 15 mn prior the addition of the bacteria and active Glucagon-Like Peptide 1 “GLP-1” (A) and

cholecystokinin “CCK” (B), secreted by STC-1 cells were measured by RIA. Values are expressed in fold of untreated control cells without inhibitors treatment and are means $\pm$ SD of three repeated experiments. Means without a common letter are different ($p < 0.05$) using one way ANOVA with Tukey *post hoc* test for pairwise comparisons in each inhibitor condition. * $p < 0.05$ vs. control.

Figure. 5 Capacity of selected strains to degrade GLP-1 (A) and CCK (B).

(A) GLP-1 1-26 (800 pM) and (B) CCK8S (600 pM) were treated with bacteria (at 10^7 CFU.mL⁻¹) in the presence (grey bars) or absence (black bars) of DDP-IV inhibitor for 8h. Remaining quantities of each hormone were measured by RIA and expressed in percentage of control without treatment. Values are expressed in percentage of control and are means $\pm$ SD of three repeated measurements. Means without a common letter are different ($p < 0.05$) using one way ANOVA with Tukey *post hoc* test for pairwise comparisons.

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.