



HAL
open science

Effect of mono or co-culture of EPS-producing *Streptococcus thermophilus* strains on the formation of acid milk gel and the appearance of texture defects

Thi Binh An Nguyen, Laetitia Picart-Palmade, Michaël Nigen, Luciana Jimenez, Hassina Ait-Abderahim, Sylvie Marchesseau

► To cite this version:

Thi Binh An Nguyen, Laetitia Picart-Palmade, Michaël Nigen, Luciana Jimenez, Hassina Ait-Abderahim, et al.. Effect of mono or co-culture of EPS-producing *Streptococcus thermophilus* strains on the formation of acid milk gel and the appearance of texture defects. *International Dairy Journal*, 2019, 98, pp.17-24. 10.1016/j.idairyj.2019.06.007 . hal-02618041

HAL Id: hal-02618041

<https://hal.inrae.fr/hal-02618041>

Submitted on 25 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Effect of mono or co-culture of EPS-producing *Streptococcus thermophilus* strains on**
2 **the formation of acid milk gel and the appearance of texture defects**

3

4

5

6

7 An Thi-Binh Nguyen ^a, Laetitia Picart-Palmade ^a, Michaël Nigen ^a, Luciana Jimenez ^b,
8 Hassina Ait-Abderahim ^b, Sylvie Marchesseau ^{a*}

9

10

11

12

13 ^a *IATE, Université de Montpellier, CIRAD, INRA, Montpellier SupAgro, Montpellier, France*

14 ^b *Chr Hansen France A/S, Rue d'Aulnay, F-91180 Saint-Germain-lès-Arpajon, France*

15

16

17

18

19 * Corresponding author. Tel.:

20 *E-mail address:* sylvie.marchesseau@umontpellier.fr (S. Marchesseau)

21

22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37

ABSTRACT

One acidifying (ST1) and two texturing strains (ST2 and ST3) of *Streptococcus thermophilus* were used as pure or co-cultures to identify and understand their effects on the structuring of acid milk gels and on the appearance of texture defects, i.e., syneresis and graininess. Symbiosis between specific texturing and acidifying strains reduced acidification time and increased exocellular polysaccharide (EPS) content. The texturing strain could simultaneously produce low and high molar mass EPS and their distribution in mass and/or in number were influenced by the proportion of acidifying to texturing strain used. The results of this study suggest that the high molar mass EPS contributes to acid gel firmness, but less so compared with the acidification rate. The ability of strain ST3 to prevent texture defects, specifically graininess, did not depend on the acidification kinetics or final EPS content, but rather on the structural properties of EPS and/or the bacterial chain morphology.

38 **1. Introduction**

39

40 Fermented milk such as yoghurt is one of the most popular dairy products in many
41 countries. In the literature, set-type fermented milk is defined as a viscoelastic gel formed
42 from the aggregation of casein micelles due to milk acidification (Tamime, Robinson, &
43 Latrill, 2001). For manufacturers, a set-type acid milk gel must have a firm body, smooth
44 texture, thick mouthfeel and structural stability. The most common texture defects in milk
45 gels are spontaneous syneresis and graininess. Spontaneous syneresis, or wheying-off,
46 indicates the separation of serum on the yoghurt surface, and graininess usually refers to
47 lumpiness or granular texture due to the presence of large protein aggregates that range in
48 size from 1 to 5 mm (Folkenberg, Dejmek, Skriver, & Ipsen, 2006; Lucey & Singh, 1997).
49 The three main factors determining gel texture and the appearance of defects are the milk
50 composition, the manufacturing process and the starter cultures (Lucey & Singh, 1997;
51 Sodini, Remeuf, Haddad, & Corrieu, 2004). A positive correlation between graininess and
52 high content of denatured whey proteins was reported in the literature (Lucey, 2004; Nguyen
53 et al., 2018a; Remeuf, Mohammed, Sodini, & Tissier, 2003). However, the effect of the
54 starter culture on the graininess has not been clarified.

55 Starter cultures can influence milk gel texture by their acidification rate and/or by
56 their exocellular polysaccharide (EPS) production. EPS from “texturing” or “ropy” cultures
57 are well known to improve the gel viscosity and reduce syneresis (Folkenberg et al., 2006;
58 Marshall & Rawson, 1999; Mende, Mentner, Thomas, Rohm, & Jaros, 2012). Some authors
59 have also observed a reduction in graininess when using a ropy strain or a high-level EPS-
60 producing culture, but no further details on the mechanism of this phenomenon have been
61 reported (Hassan, Ipsen, Janzen, & Qvist, 2003; Küçükçetin, Weidendorfer, & Hinrichs,
62 2009). In a previous study (Nguyen et al., 2018a), the effects of heat treatment, milk

63 composition and two EPS-producing starter cultures on the microstructure and the texture
64 defects of acid milk gel have suggested that the presence of denatured whey proteins is
65 critical for grain formation. Gel viscosity and texture defects appeared to be culture-
66 dependent and were correlated with the amount, molar mass and intrinsic viscosity of the
67 EPS produced. Furthermore, the gel microstructure obtained from the two cultures showed
68 major differences in the location of bacterial cells in the gels, suggesting that simultaneously
69 with the EPS effect, the length of bacterial cell chains could also interfere with the formation
70 of the protein network during acidification due to steric hindrance. However, the different
71 acidification kinetics of the two cultures used might influence gel structuring.

72 In industrial applications, commercial starter cultures for yoghurt fermentation usually
73 consist of several strains of *Streptococcus thermophilus*, which are chosen for their specific
74 properties such as fast initial acidification or texture promoting capacity. The interaction
75 between different *S. thermophilus* strains could affect the total EPS production, but different
76 trends were observed and no details on the acidification kinetics or EPS characterisation have
77 been reported (Folkenberg et al., 2006). In addition, the influence of co-cultures of *S.*
78 *thermophilus* strains on milk gelation has not yet been studied. The aim of this study was to
79 identify the effects of mono or co-culture of EPS-producing *S. thermophilus* on the formation
80 of acid milk gels and the appearance of texture defects. For this purpose, three *S.*
81 *thermophilus* strains were used alone or in combination: one for its fast acidification
82 characteristics and the other two for their texturing properties. Hence, seven starter cultures
83 were formed with different ratios between acidifying and texturing strains to study the
84 correlations between the gel texture characteristics (e.g., firmness, viscosity, syneresis,
85 graininess), the kinetics of acidification and gelation, but also the quantity and structural
86 features of EPS produced.

87

88 2. Materials and methods

89

90 2.1. Solvents and reagents

91

92 Low-heat skim milk powder (Spray 0) was purchased from Ingredia (France). The
93 three *S. thermophilus* strains were provided by Chr. Hansen A/S, Arpajon (France). Lactose
94 monohydrate and trichloroacetic acid (TCA) were from GPR Rectapur, VWR chemicals
95 (Belgium). D(+)-Glucose monohydrate and 1 N sodium hydroxide (NaOH) were from Merck
96 (Germany). Phenol, LiNO₃ and NaN₃ were from Sigma (USA). Ethanol (95%, v/v) was from
97 TechniSolv, VWR chemicals (France). Concentrated sulphuric acid (95%, w/w) was from
98 Normapur, VWR chemicals (France).

99

100 2.2. Starter cultures

101

102 Three different frozen lyophilised strains of *S. thermophilus* ST1, ST2, ST3 from the
103 Chr Hansen culture collection (Chr Hansen A/S) were used alone (pure culture) or in
104 combination (co-culture) in seven different starter cultures. From manufacturer information,
105 strain ST1 possess a cell wall protease (PrtS+) and could play a role as acidifying strain. ST2
106 or ST3 were texturing strains without cell wall protease (PrtS-) and were chosen to provide
107 different textural types of milk gels: firm or viscous, respectively.

108

109 Cultures ST1, ST2 and ST3 were composed of a single *S. thermophilus* strain
110 corresponding to the name of the culture. The four co-cultures ST1+2 (1/1.7), ST1+2 (5/1),
111 ST1+3 (1/1.7)-and ST1+3 (5/1) consisted of the combination of acidifying strain ST1 with a
texturing-strain (ST2 or ST3) with the cfu ratios of acidifying strain/texturing strain

112 corresponding to the name of the culture. All strains were stored at $-80\text{ }^{\circ}\text{C}$. Strains were
113 thawed and mixed then 2 g of the final culture were added to 100 mL of UHT skim milk. Ten
114 millilitres of this pre-inoculated milk were immediately added to 1 L of pasteurised milk to
115 start the lactic fermentation. The final inoculation rate for the seven starter cultures was 10^6
116 cfu mL⁻¹ pasteurised milk. Sampling, thawing and formulation were performed in the
117 laboratory under sterile conditions.

118

119 2.3. *Acid milk gel manufacture*

120

121 The milk-based preparation was carried out as described by Nguyen et al. (2018b).
122 The milk formulation contained 11.8% (w/w) dry matter, 4.0% (w/w) proteins and 6.7%
123 (w/w) lactose. The skim milk powder was dissolved in ultra-pure water during 1h at room
124 temperature and then kept for 12 h at $4\text{ }^{\circ}\text{C}$ before the pasteurisation at $95\text{ }^{\circ}\text{C}$ for 6 min. After
125 the heating step, milk suspensions were cooled to $43\text{ }^{\circ}\text{C}$ and immediately inoculated with the
126 starter cultures. Fermentation procedures, pH monitoring and gel structuring have previously
127 been reported by Nguyen et al. (2018b). In brief, the fermentation was carried out at $43\text{ }^{\circ}\text{C}$
128 with pH control until a final pH of 4.65 was reached. In parallel, gel formation was monitored
129 using a Physica MCR 300 rheometer (Anton Paar, St Albans, UK) equipped with concentric
130 cylinder geometry (CC27) that operated in continuous oscillation mode at a frequency of 1
131 Hz and 1% strain. The gel point was defined as the moment when the storage modulus (G')
132 first exceeds the loss modulus (G'') (Winter, 1987). After acidification, the gels were stored
133 undisturbed in plastic cups at $4\text{ }^{\circ}\text{C}$ for 7 days before determination of texture properties.

134

135 2.4. *Gel texture properties*

136

137

138 The textural parameters of set-type acid milk gel at day 7 including the firmness, the
139 viscosity, the spontaneous syneresis and the grain defect were characterised as described by
140 Nguyen et al. (2018a).

141 The firmness of acid milk gel was determined immediately after removing out of the
142 cold room (4 °C) using a texture analyser equipped with an acrylic cylinder compression
143 probe (diameter = 40 mm, thickness = 5 mm). The compression test was carried out with a
144 speed of 5 mm s⁻¹, the trigger force of 5 g and a penetration depth of 15 mm. The gel firmness
145 (N) corresponding to the maximum compression force, i.e., the rupture point of the gel, was
146 calculated with the Exponent software (Stable Micro Systems, Godalming, UK).

147 For the viscosity measurement, the set-type acid milk gels were stirred 20 times with a
148 spoon and 20 g of sample was placed in the rheometer cup. The viscosity of the stirred gel (η ,
149 Pa s) at a shear rate of 300 s⁻¹ (13 °C) was determined using Physica MCR 300 rheometer
150 equipped with concentric cylinder geometry and the shear rates $\dot{\gamma}$ (s⁻¹) increased with a linear
151 ramp from 0.271 s⁻¹ to 300 s⁻¹ in 210 s.

152 The spontaneous syneresis (% w/w) was the percentage of whey on the surface of the
153 undisturbed gel to the total gel weight and was measured according to the siphon method
154 (Amatayakul, Sherkat, & Shah, 2006).

155 The grain defect was determined by image analysis. After stirring 5 times with a
156 spoon, 1 g of acid gel was dispersed in 100 mL of distilled water by stirring at 400 rpm for
157 2 min. 10 mL of diluted sample was poured into a plastic Petri dish and visualised with a
158 digital camera in black and white mode. The image analysis was performed using ImageJ
159 1.49v software (National Institutes of Health, USA) with Fiji plug-in. Grains with a diameter
160 > 1 mm were enumerated (Remeuf et al., 2003). The grains were classified into different size

161 groups by their diameter: 1–1.5 mm, 1.5–2.0 mm, 2.0–2.5 mm, 2.5–3 mm or > 3 mm. The
162 results were expressed as the number of grains of each size per gram of gel.

163

164 2.5. *EPS quantification*

165

166 EPS quantification was performed using the protocol 1 described by Nguyen et al.
167 (2018b). The EPS were extracted in hot-acidic medium and dialysed to obtain a high
168 extraction yield. The total sugar content was determined by the phenol-sulfuric acid method
169 using glucose as the standard (range of concentration from 0 to 200 mg L⁻¹). The acidified
170 milk with glucono- δ -lactone (GDL) was used as blank and the residual milk sugars after
171 dialysis were subtracted from all data.

172

173 2.6. *EPS characterisation*

174

175 EPS were extracted without a heating step using the protocol 3 described by Nguyen
176 et al. (2018b) to avoid EPS denaturation. The extraction process included a step of pH
177 correction to 7, the protein precipitations in an acidic medium, followed by the precipitations
178 of EPS by ethanol before dialysis and lyophilisation. The structural properties of extracted
179 EPS were characterised using high-performance size exclusion chromatography coupled with
180 multi-angle laser light scattering detector (HPSEC-MALS). The material and operating
181 conditions were reported in Nguyen et al. (2018b). The weight-average molar mass (M_w) and
182 the intrinsic viscosity ($[\eta]$) were determined using the software ASTRA 6.1.2 (Wyatt
183 Technologies, Santa Barbara, CA). The M_w was calculated using the Berry's model (2nd
184 order). The data were analysed using a refractive index increment (dn/dc) of 0.145 mL g⁻¹
185 according to the literature (Lambo-Fodje et al., 2007).

186

187 2.7. *Statistical analysis*

188

189 All analyses were conducted at least in triplicate and the values were expressed as the
190 mean \pm standard deviation. The experimental data were analysed statistically using the
191 Fischer's and Student's tests with 95% confidence level and using Pearson's correlation
192 coefficient functions for the correlation between means.

193

194 3. **Results and discussion**

195

196 3.1. *Characterisation of starter cultures*

197

198 3.1.1. *Acidification kinetics*

199 The kinetics of milk acidification induced by lactic fermentation were monitored by
200 pH evolution between pH 6.5 and 4.6 (Fig. 1). The parameters evaluated were the total
201 acidification time and the maximum acidification rate (V_{\max}) (Table 1). As an acidifying
202 strain, ST1 presented the highest V_{\max} value of all pure cultures and the shortest acidification
203 time (~300 min) compared with ST2 and ST3. The acidification curve profiles of strains ST2
204 and ST3 (Fig. 1) were very different from that of strain ST1. ST2 and ST3 had significantly
205 slower acidification rates during the first 160 min and also a stronger slowdown in
206 acidification after the shoulder observed around pH 6.25. This specific profile corresponded
207 to the strains' urease activity, which produced ammonia from milk urea and induced a slight
208 alkalisation of the medium. Furthermore, the final count of bacterial cells at the end of the
209 fermentation was $\sim 10^9$, 10^7 or 10^6 cfu g⁻¹ for ST1, ST2 or ST3, respectively, and was
210 consistent with the kinetics observed. These differences in acidification kinetics and cell

211 growth were related to the proteolytic activity of these 3 strains. Indeed, strain ST1 has a cell
212 envelope-associated proteinase (PrtS+) that hydrolyses caseins into peptides and essential
213 amino acids to ensure high cell density growth and thereby generating high acidification
214 activity (Galia, Perrin, Genay, & Dary, 2009). When this enzymatic activity is lacking, such
215 as for ST2 and ST3 (PrtS-), the growth of *S. thermophilus* strains in milk is less optimal
216 depending on the specific intracellular peptidases and amino acids requirements of the strain
217 (Galia et al., 2009).

218 The addition of ST1 to ST2 or ST3 in four co-cultures resulted in a considerable
219 decrease in acidification time and a significant increase in V_{\max} (Fig. 1, Table 1). For
220 example, the total acidification time for cultures of ST2 was reduced from 1351 min (pure
221 culture) to 528 min or 304 min when ST1 was added with a ST1/ST2 ratio of 1/1.7 or 5/1,
222 respectively. Furthermore, while pure culture of ST3 had the slowest acidification kinetics of
223 all cultures tested, the ST1+3 co-cultures had better acidification performance as compared
224 with ST1+2 co-cultures for the same ST1 ratio. These results suggested that the proteinase
225 activity of ST1 produced peptides and essential amino acids that stimulated growth of both
226 texturing strains. However, the amino acid requirements seemed to be different between ST2
227 and ST3, as indicated by their differing rates of acidification. ST1 metabolites may be more
228 adapted to the needs of ST3 than to those of ST2, which allowed for a better symbiotic
229 relation between ST1 and ST3 than between ST1 and ST2.

230

231 3.1.2. *EPS production and characterisation*

232 The EPS produced by the seven starter cultures were extracted for quantification (Fig.
233 2) and for macromolecular characterisation (molar mass distribution (M_w) and intrinsic
234 viscosity ($[\eta]$) (Fig. 3). Although the strain ST1 was not considered as textural strain, it was
235 able to produce a substantial amount of EPS of $\sim 55 \pm 4$ mg glucose kg^{-1} of gel in a short time

236 of acidification (~300 min) (Fig. 2). However, the EPS produced have low M_w of $\sim 9.2 \times 10^4$ g
237 mol^{-1} and low $[\eta]$ of ~ 42 mL g^{-1} , which could explain the poor texturing property of this
238 strain. These results also confirmed that macromolecular properties are a more critical
239 characteristic for the functional properties of EPS than the total amount of polymers (Mende,
240 Rohm, & Jaros, 2016). Even after an acidification time that was 4–5 times longer than for
241 culture ST1, cultures ST2 and ST3 did not produce a greater quantity of EPS (~ 30 mg
242 glucose kg^{-1} and ~ 52 mg glucose kg^{-1} , respectively), particularly for ST2 (Fig. 2). However,
243 the EPS produced by these two texturing strains had very different macromolecular properties
244 from each other and from those of the ST1 culture. Based on elution profiles of HPSEC-
245 MALS analysis, EPS produced by ST2 or ST3, could be divided into two groups of low and
246 high M_w as also observed in the literature (Petry et al., 2003; Vaningelgem et al., 2004). The
247 EPS produced by the strain ST3 has a M_w and intrinsic viscosity significantly higher than
248 those produced by cultures ST2 or ST1 (Fig. 3). Furthermore, EPS from ST3 consisted of
249 80% high M_w polymers in number, in contrast to EPS from ST2, which contained only 35%
250 high M_w molecules.

251 Regarding the strain ST2 and its co-cultures with ST1, the increase in the proportion
252 of ST1 from 0 to 37% [culture ST1+2 (1/1.7)] and to 83 % [culture ST1+2 (5/1)] induced a
253 significant increase in the amount of EPS produced from 30 to ~ 52 and to ~ 67 mg glucose kg^{-1}
254 1 , respectively (Fig. 2), while the time of acidification was significantly decreased (Table 1).
255 The same trend was also observed when ST1 was added to strain ST3. Furthermore, the
256 macromolecular characteristics of the EPS produced by ST2 and its co-cultures were broadly
257 similar (Fig. 3A,D). For the strain ST3 and its co-cultures ST1+3, a significant decrease of
258 EPS M_w was observed when the acidification time decreased (Pearson's $r = 0.951$) due to the
259 increase in ST1/ST3 ratio. These results suggested a chain length extension of EPS from ST3
260 during longer acidification time. Laws, Leivers, Chacon-Romero, and Chadha (2009)

261 reported a similar trend in which EPS M_w produced by *Lactobacillus acidophilus* increased
262 3.7 times from 6 h to 24 h of fermentation in skim milk supplemented with glucose at 42 °C
263 and at a controlled pH of 5.8. In association with cell counting, the authors observed that EPS
264 synthesis and chain length extension continued after the end of the exponential phase, using
265 sugar nucleotides already produced during in the latter phase.

266 When looking in more details at the results of EPS M_w distribution (in mass or in
267 number, Fig. 3B,C), different trends could be observed. Firstly, the presence of the acidifying
268 strain ST1 at a level of 37% in culture ST1+2 (1/1.7) and ST1+3 (1/1.7) did not change the
269 M_w distribution characteristics as compared with pure strains (Fig. 3B,C), suggesting that in
270 these conditions ST1 produced a negligible quantity of EPS, but had an enhancing effect on
271 the EPS production of texture strains. In contrast, a significant change in the M_w distribution
272 of EPS was observed with 83% ST1 in cultures ST1+2 (5/1) and ST1+3 (5/1), in that the low
273 M_w EPS then became predominant. These findings could be related to the short acidification
274 time observed with these co-cultures and/or to the prevalence of the acidifying strain over the
275 texturing strain. Furthermore, we observed that the high M_w EPS group from culture ST1+3
276 (5/1) presented a lower average M_w , but a higher $[\eta]$ than for ST1+2 (5/1). These findings
277 suggested that the structure of EPS produced by ST2 could be more flexible and/or branched
278 than those produced by ST3. It has been shown that for two EPS with identical structure (i.e.
279 repeating units branching, stiffness of side chain), a higher M_w could lead to a higher $[\eta]$ and
280 therefore to a higher gel viscosity (Faber, Zoon, Kamerling, & Vliegthart, 1998;
281 Looijesteijn, van Casteren, Tuinier, Doeswijk-Voragen, & Hugenholtz, 2000; Mende et al.,
282 2016). However, other structural factors such as side chain stiffness and branching also made
283 a considerable contribution to the $[\eta]$ of EPS. Generally, for a given M_w , a rigid linear
284 polysaccharide will tend to extend under shearing, as opposed to a flexible branched

285 molecules which will tend to compact, and will then contribute to increase the viscosity of the
286 solution (Whistler & BeMiller, 1997).

287 Overall, the results presented in this section showed that the acidification kinetics, the
288 production and the macromolecular characteristics of EPS depended not only on the strains
289 but also on the interaction between textural and acidifying strains. The combination between
290 ST1 and ST3 seemed to be more efficient than the combination between ST1 and ST2.
291 Furthermore, all pure cultures could produce EPS, but these EPS differed in terms of
292 quantity, distribution in mass or in number and intrinsic viscosity. In the same way, the
293 overall distribution of low and high M_w EPS is strongly correlated with the interaction
294 between acidifying and texturing strains. It is quite difficult to compare these findings with
295 other studies because very few data are reported on the EPS production of mixtures of *S.*
296 *thermophilus*. Folkenberg et al. (2006) observed different trends of EPS production in
297 fermented milk by 5 *S. thermophilus* strains in pure or in mix cultures of two strains. Their
298 results suggested that the interaction between two *S. thermophilus* strains could be
299 compatible or not, depending on the strain used. However, they did not report acidification
300 time or molecular characteristics of EPS, which made difficult the comparison with our
301 results.

302

303 3.2. *Influence of starter cultures on milk gelation and textural properties of gels*

304

305 3.2.1. *Gelation kinetics*

306 The kinetics of gelation were monitored by the evolution of the storage modulus (G')
307 and the loss modulus (G''), characterised by time and pH of the gel point. As presented in
308 Table 1, the time and pH of gelation depended on the culture used. The pH of the gel point
309 was approximately 5.2 for the cultures ST1, ST2, ST1+2 (1/1.7) and ST1+2 (5/1), and was

310 not influenced by the rate of acidification. The same pH value was reported for the gel point
311 of heated milk acidified chemically using glucono-delta-lactone (Lucey, Tamehana, Singh, &
312 Munro, 1998; Tamime et al., 2001). In the case of ST3, the pH of the gel point was found to
313 be significantly higher at 5.57 and decreased to 5.4 and 5.2 when strain ST1 was added at 37
314 and 83%, respectively (Table 1). The occurrence of the gel point at a higher pH could be
315 related to the presence of EPS and their interactions with milk proteins (Mende et al., 2013b).

316 In the case of strain ST3 and its co-cultures, a positive correlation (Pearson's $r =$
317 0.992) was observed between the pH of gelation and EPS macromolecular properties (M_w ,
318 $[\eta]$), but not with the EPS quantity. The decrease in the pH of the gel point could be related
319 to a decrease of the average M_w of EPS and/or of the proportion of high M_w EPS (Fig. 3). On
320 the other hand, for all cultures, a predominant proportion in number of low M_w EPS
321 corresponded to a gel point occurring at a lower pH of ~ 5.2 (Fig. 3C). Mende et al. (2013a,b)
322 also reported a similar trend. The authors observed early gelation due to the depletion effect
323 between casein micelles and added bacterial EPS or commercial dextran (Tuinier,
324 Grotenhuis, Holt, Timmins, & de Kruif, 1999). These two neutral polysaccharides, bacterial
325 EPS and dextran had different M_w of 2.6×10^6 Da and 5×10^5 Da, respectively. Compared with
326 bacterial EPS, addition of 15 – 80 times more dextran was required to achieve a similar effect
327 on gelation point.

328

329 3.2.2. *Gel texture*

330 Gel texture was characterised by four parameters: (i) firmness, (ii) viscosity, (iii)
331 spontaneous syneresis (Table 2) and (iv) graininess (Fig. 4). Gels from the ST3 culture are
332 not mentioned in this section because after 24 h of acidification, the gels obtained at a final
333 pH of ~ 4.8 were heterogeneous with an upper gel-like part and a lower liquid-like part.

334

335 *Firmness.* Gel firmness is the main texture property of set-type acid milk gel and this
336 parameter is mainly governed by the protein network and not by bacterial EPS (Folkenberg et
337 al., 2006; van Marle & Zoon, 1995). For the monocultures used in the present study, ST1
338 provided the lowest gel firmness while ST2 produced the highest gel firmness, which could
339 be related to its long acidification time of nearly 24 h. Moreover, it was observed for strain
340 ST2 and its co-cultures ST1+2 that the higher the proportion of ST1, the faster the
341 acidification and the lower the gel firmness (Pearson's $r = 0.998$). According to the literature,
342 slow protein aggregation could result in continuous network formation with more connected
343 proteins, reducing the rearrangement of particles during gel formation and contributing to an
344 increase in gel rigidity during fermentation at low temperatures (Lucey & Singh, 1997;
345 Sodini et al., 2004).

346 In the present study, it is interesting to note that although the cultures ST1, ST1+2
347 (5/1) and ST1+3 (5/1) induced similar acidification times (Table 1), the resulting gel firmness
348 was significantly different in the order: ST1+2 (5/1) > ST1+3 (5/1) > ST1, consistent with the
349 order of the average M_w of EPS produced. In other words, the M_w of EPS from ST1+2 (5/1)
350 was higher than from those of ST1+3 (5/1) and ST1 (Pearson's $r = 0.987$). These findings
351 suggested that the presence of high M_w EPS was also an important determining parameter for
352 gel firmness, though not as much as the acidification rate.

353 When comparing the gels of co-cultures with the same ST1 ratio, the gels containing
354 ST2 had a higher firmness than those containing ST3 (Table 2). Other studies have also
355 reported a decreased gel firmness when using a ropy culture which is defined as a high EPS
356 producing culture with high $[\eta]$ such as the culture ST3 (Hassan et al., 2003; Hess, Roberts,
357 & Ziegler, 1997; Mende et al., 2012). Using microscopic analyses, Hassan et al. (2003)
358 observed that the EPS of ropy cultures were mainly located and segregated in the pores of the
359 gel rather than attached to the protein network. Hence, the gel structure from a ropy culture

360 was less dense than that of non-ropy culture. Mende et al. (2012) and Rohm and Kovac
361 (1994) also suggested that the large amount of uncharged EPS produced by ropy culture
362 could have partially prevented the development of protein-protein bonds during gel formation
363 and thus reduced the rigidity of acid milk gel.

364

365 *Stirred gel viscosity.* Stirred gel viscosity is correlated with a thick mouthfeel and with
366 the smoothness of fermented milk products (Folkenberg et al., 2006). In this study, the stirred
367 gel viscosity varied depending on the culture and the macromolecular properties of the
368 produced EPS. ST1 produced only low M_w EPS with low $[\eta]$ and formed gels presenting the
369 lowest viscosity. When ST2 was used, even if a low quantity of EPS was produced, the
370 viscosity of the stirred gels formed was higher than for strain ST1. This result could be
371 explained by the high percentage (80%) in mass of EPS with high M_w produced by culture
372 ST2.

373 Concerning the co-cultures of ST2 or ST3, increasing the ratio of the acidifying strain
374 decreased the stirred gel viscosity from 193 to 164 mPa s for ST1+2 (1/1.7) and (5/1),
375 respectively, or from 301 to 239 mPa s for ST1+3 (1/1.7) and (5/1), respectively (Table 2).
376 These results could be directly linked to the macromolecular properties of the EPS produced
377 (Fig. 3) and suggested a positive correlation between the quantity of high M_w EPS and the
378 stirred gel viscosity (Pearson's $r = 0.890$). Petry et al. (2003) also reported a striking
379 correlation between an elevated proportion of the high M_w EPS fraction and viscosity of acid
380 milk gel. In the present study, we also observed that even though the amount and molar mass
381 distribution of EPS were similar, gels from culture ST1+3 (1/1.7) were more viscous than
382 those from culture ST1+2 (1/1.7). This result could be partially explained by the higher $[\eta]$ of
383 the EPS produced by ST3 and probably by a different molecular rigidity as already
384 mentioned in section 3.1.2. Also when comparing ST1+2 (5/1) and ST1+3 (5/1), despite both

385 co-cultures exhibiting a low proportion of high M_w EPS and similar $[\eta]$ of EPS, the viscosity
386 of stirred gels was significantly different, suggesting that EPS were not the only factor
387 contributing to this textural property. In a previous study based on gel microstructure
388 observations, Nguyen et al. (2018a) suggested that in parallel to EPS production, the
389 morphology of bacterial cell chain length should also be taken into account to understand the
390 texture results.

391

392 *Spontaneous syneresis defect.* Spontaneous syneresis is the contraction of a gel
393 without exerting external forces and is related to an instability of the gel network that results
394 in the loss of ability to entrap the entire serum phase (Lucey & Singh, 1997). Gels obtained
395 with pure culture of ST1 presented a medium level of syneresis while those from ST2 had the
396 highest syneresis, possibly related to the long acidification time. For co-cultures of ST2, this
397 texture defect decreased from 2.7 to 1.4% when the ratio of ST1/ST2 changed from 1/1.7 to
398 5/1. In this case, a positive correlation was observed between syneresis, gel firmness and time
399 of acidification (Pearson's $r = 0.999$) suggesting that gel contraction could occur during the
400 slow fermentation, thereby hardening the gel and expelling the serum. In contrast, low
401 syneresis levels were observed for co-culture of ST1+3. Decreased gel syneresis has been
402 reported with the use of ropy strains that produce a high quantity of high M_w EPS
403 (Amatayakul et al., 2006; Kristo, Miao, & Corredig, 2011). It can be assumed that the ropy
404 EPS located in the gel pores could bind the serum and thus reduce the syneresis.

405

406 *Graininess defect.* Graininess or lumpiness is considered an undesirable defect
407 because consumers usually expect a smooth product (Lucey, 2004). Remeuf et al. (2003)
408 measured graininess by the number of grains with a diameter above 1 mm in 1 g of acid milk
409 gel. In the present study, the graininess defect was determined by the number of grains, but

410 also by their diameter from 1 mm to above 3 mm. Larger grains were more damaging for
411 texture perception even in low number (Fig. 4). The graininess was high when using pure
412 cultures ST1 and ST2. In the mixtures, regardless of the strain ratio, co-cultures of ST3
413 presented a low graininess while the co-cultures of ST2 showed a medium granular texture
414 (Fig. 4).

415 According to the literature, grains are large aggregates of proteins that appear under
416 certain conditions such as a high fermentation temperature, excessive whey protein to casein
417 ratio or with the use of certain types of starter culture (Lucey, 2004). However, only few
418 studies have focused on the impact of starter culture on the appearance of grain. In agreement
419 to our result, Küçükçetin et al. (2009) reported that the number of grains and their size were
420 significantly reduced when using a high-level EPS-producing starter culture, rather than a
421 medium- or low-level EPS-producing culture. Results from the present study provided
422 additional data on this phenomenon. Indeed, it is worth noting that cultures ST1+2 (5/1) and
423 ST1+3 (5/1) had a similar kinetics of acidification and gelation (Table 1) and similar
424 production of EPS (Fig. 2) that presented identical intrinsic viscosity and EPS molar mass
425 distribution (Fig. 3 B,C,D). However, despite these similarities, the texture of the resulting
426 gels from ST1+2 (5/1) presented more major defects (syneresis and graininess) as compared
427 with gels from culture ST1+3 (5/1). These new data support the inhibitory effect of strain
428 ST3 against appearance of texture defects, specifically graininess, which has been observed
429 in a previous study (Nguyen et al., 2018a). Therefore, we suggested that this interesting
430 technological property of culture ST3 could be related to the bacterial chains morphology
431 and/or the structural properties of the EPS produced and that it did not depend on the
432 acidification kinetics or final EPS content.

433

434 **4. Conclusion**

435

436 It is known that the symbiosis between texturing and acidifying strains of *S.*
437 *thermophilus* reduces the acidification time and has an impact on the final acid milk gel
438 structure. In this work, we selected one acidifying strain (ST1) and two texturing strains
439 (ST2, ST3) strains, and used them singly or in co-cultures (ST1+2 or ST1+3) at two different
440 ratios (1/1.7 or 5/1) in pasteurised skim milk to assess and understand the influence of their
441 interaction on the acidification kinetics, EPS production, gel structure and appearance of
442 texture defects.

443 In pure culture, due to the presence of proteinases in the cell wall, the acidifying strain
444 provided rapid acidification and produced a medium amount of EPS. However, these EPS
445 had a low M_w and intrinsic viscosity $[\eta]$ hence the gel obtained had the lowest firmness and
446 viscosity, and the most graininess defect. Both texturing strains did not have the proteinase
447 activity and showed a slow acidification. They both produced EPS with high molar mass but
448 with different macromolecular properties.

449 In co-culture, the strains ST2 and ST3 had positive symbiotic relations with ST1, as
450 showed by the reduction in acidification time and the increase in EPS production.
451 Furthermore, the viscosity of gels increased significantly, probably due to the production of
452 high M_w EPS by the texturing strains. Texture defects such as syneresis and graininess
453 disappeared with use of the co-cultures of ST1+3. Moreover, co-cultures ST1+3 (5/1) and
454 ST1+2 (5/1) had some similarities in terms of (i) acidification, (ii) gelation kinetics, (iii) EPS
455 production and molar mass distribution and (iv) gel firmness. However, despite these
456 similarities, gels from co-culture ST1+2 (5/1) showed worse texture defects as compared with
457 ST1+3 (5/1).

458 These results highlight the fact that the inhibitory effect of a strain against the
459 appearance of texture defects is not correlated to the acidification rate or EPS content, but
460 that it could instead be explained by differences in the structural properties of EPS and/or in
461 the morphology of bacterial chains.

462

463 **Acknowledgements**

464

465 The authors acknowledge the financial support of the French National Association for
466 Research and Technology (ANRT) and Chr. Hansen France SAS for the PhD fellowship. We
467 are grateful to Charles Cunault (UMR IATE, Montpellier, France) for his instruction in the
468 analyse image by ImageJ and plug-in Figi, and to Ophelia Pedersen for improving the
469 manuscript.

470

471 **References**

472

473 Amatayakul, T., Sherkat, F., & Shah, N. P. (2006). Syneresis in set yogurt as affected by EPS
474 starter cultures and levels of solids. *International Journal of Dairy Technology*, 59, 216–
475 221.

476 Faber, E. J., Zoon, P., Kamerling, J. P., & Vliegenthart, J. F. G. (1998). The
477 exopolysaccharides produced by *Streptococcus thermophilus* Rs and Sts have the same
478 repeating unit but differ in viscosity of their milk cultures. *Carbohydrate Research*, 310,
479 269–276.

480 Folkenberg, D. M., Dejmek, P., Skriver, A., & Ipsen, R. (2006). Interactions between EPS-
481 producing *Streptococcus thermophilus* strains in mixed yoghurt cultures. *Journal of*
482 *Dairy Research*, 73, 385–393.

483 Galia, W., Perrin, C., Genay, M., & Dary, A. (2009). Variability and molecular typing of
484 *Streptococcus thermophilus* strains displaying different proteolytic and acidifying
485 properties. *International Dairy Journal*, *19*, 89–95.

486 Hassan, A. N., Ipsen, R., Janzen, T., & Qvist, K. B. (2003). Microstructure and rheology of
487 yogurt made with cultures differing only in their ability to produce exopolysaccharides.
488 *Journal of Dairy Science*, *86*, 1632–1638.

489 Hess, S. J., Roberts, R. F., & Ziegler, G. R. (1997). Rheological properties of nonfat yogurt
490 stabilized using *Lactobacillus delbrueckii* ssp. *bulgaricus* producing exopolysaccharide
491 or using commercial stabilizer systems. *Journal of Dairy Science*, *80*, 252–263.

492 Kristo, E., Miao, Z., & Corredig, M. (2011). The role of exopolysaccharide produced by
493 *Lactococcus lactis* subsp. *cremoris* in structure formation and recovery of acid milk gels.
494 *International Dairy Journal*, *21*, 656–662.

495 Küçükçetin, A., Weidendorfer, K., & Hinrichs, J. (2009). Graininess and roughness of stirred
496 yoghurt as influenced by processing. *International Dairy Journal*, *19*, 50–55.

497 Lambo-Fodje, A. M., Leeman, M., Wahlund, K. G., Nyman, M., Öste, R., & Larsson, H.
498 (2007). Molar mass and rheological characterisation of an exopolysaccharide from
499 *Pediococcus damnosus* 2.6. *Carbohydrate Polymers*, *68*, 577–586.

500 Laws, A. P., Leivers, S., Chacon-Romero, M., & Chadha, M. J. (2009). Variation in the
501 molecular mass of exopolysaccharides during the time course of extended fermentations
502 of skimmed milk by lactic acid bacteria. *International Dairy Journal*, *19*, 768–771.

503 Looijesteijn, P. J., van Casteren, W. H., Tuinier, R., Doeswijk-Voragen, C. H., &
504 Hugenholtz, J. (2000). Influence of different substrate limitations on the yield,
505 composition and molecular mass of exopolysaccharides produced by *Lactococcus lactis*
506 subsp. *cremoris* in continuous cultures. *Journal of Applied Microbiology*, *89*, 116–122.

507 Lucey, J. A. (2004). Cultured dairy products: an overview of their gelation and texture

508 properties. *International Journal of Dairy Technology*, 57, 77–84.

509 Lucey, J. A., & Singh, H. (1997). Formation and physical properties of acid milk gels: A
510 review. *Food Research International*, 30, 529–542.

511 Lucey, J. A., Tamehana, M., Singh, H., & Munro, P. A. (1998). Effect of interactions
512 between denatured whey proteins and casein micelles on the formation and rheological
513 properties of acid skim milk gels. *Journal of Dairy Research*, 65, 555–567.

514 Marshall, V. M., & Rawson, H. L. (1999). Effects of exopolysaccharide-producing strains of
515 thermophilic lactic acid bacteria on the texture of stirred yoghurt. *International Journal
516 of Food Science & Technology*, 34, 137–143.

517 Mende, S., Mentner, C., Thomas, S., Rohm, H., & Jaros, D. (2012). Exopolysaccharide
518 production by three different strains of *Streptococcus thermophilus* and its effect on
519 physical properties of acidified milk. *Engineering in Life Sciences*, 12, 466–474.

520 Mende, S., Peter, M., Bartels, K., Dong, T., Rohm, H., & Jaros, D. (2013a). Concentration
521 dependent effects of dextran on the physical properties of acid milk gels. *Carbohydrate
522 Polymers*, 98, 1389–1396.

523 Mende, S., Peter, M., Bartels, K., Rohm, H., & Jaros, D. (2013b). Addition of purified
524 exopolysaccharide isolates from *Streptococcus thermophilus* to milk and their impact on
525 the rheology of acid gels. *Food Hydrocolloids*, 32, 178–185.

526 Mende, S., Rohm, H., & Jaros, D. (2016). Influence of exopolysaccharides on the structure,
527 texture, stability and sensory properties of yoghurt and related products. *International
528 Dairy Journal*, 52, 57–71.

529 Nguyen, A. T.-B., Nigen, M., Jimenez, L., Ait-Abderahim, H., Cunault, C., Marchesseau, S.,
530 et al. (2018a). A multi-scale approach to identify the role of heat treatment, milk protein
531 composition and starter culture on the gel formation and the texture defects of acid milk
532 gel. *Food Hydrocolloids*, 85, 299–310.

533 Nguyen, A. T.-B., Nigen, M., Jimenez, L., Ait-Abderrahim, H., Marchesseau, S., & Picart-
534 Palmade, L. (2018b). Performances of different protocols for exocellular
535 polysaccharides extraction from milk acid gels: Application to yogurt. *Food Chemistry*,
536 239, 742–750.

537 Petry, S., Furlan, S., Waghorne, E., Saulnier, L., Cerning, J., & Maguin, E. (2003).
538 Comparison of the thickening properties of four *Lactobacillus delbrueckii* subsp.
539 *bulgaricus* strains and physicochemical characterization of their exopolysaccharides.
540 *FEMS Microbiology Letters*, 221, 285–291.

541 Remeuf, F., Mohammed, S., Sodini, I., & Tissier, J. P. (2003). Preliminary observations on
542 the effects of milk fortification and heating on microstructure and physical properties of
543 stirred yogurt. *International Dairy Journal*, 13, 773–782.

544 Rohm, H., & Kovac, A. (1994). Effects of starter cultures on linear viscoelastic and physical
545 properties of yogurt gels. *Journal of Texture Studies*, 25, 311–329.

546 Sodini, I., Remeuf, F., Haddad, S., & Corrieu, G. (2004). The relative effect of milk base,
547 starter, and process on yogurt texture: A Review. *Critical Reviews in Food Science and*
548 *Nutrition*, 44, 113–137.

549 Tamime, A. Y., Robinson, R. K., & Latrielle, E. (2001). Yoghurt and other fermented milks.
550 In A. Y. Tamime & B. A. Law (Eds.), *Mechanisation and automation in dairy*
551 *technology* (pp. 152–203). Sheffield, UK: Sheffield Academic Press.

552 Tuinier, R., Grotenhuis, E. T., Holt, C., Timmins, P. A., & de Kruif, C. G. (1999). Depletion
553 interaction of casein micelles and an exocellular polysaccharide. *Physical Review E*, 60,
554 848–856.

555 van Marle, M. E., & Zoon, P. (1995). Permeability and rheological properties of microbially
556 and chemically acidified skim-milk gels. *Netherlands Milk and Dairy Journal*, 49, 47–
557 65

- 558 Vaningelgem, F., Zamfir, M., Mozzi, F., Adriany, T., Vancanneyt, M., Swings, J., et al.
559 (2004). Biodiversity of exopolysaccharides produced by *Streptococcus thermophilus*
560 strains is reflected in their production and their molecular and functional characteristics.
561 *Applied and Environmental Microbiology*, 70, 900–912.
- 562 Whistler, R. L., & BeMiller, J. N. (1997). *Carbohydrate chemistry for food scientists*. St.
563 Paul, MI, USA: Eagan Press.
- 564 Winter, H. H. (1987). Can the gel point of a cross-linking polymer be detected by the $G' - G''$
565 crossover? *Polymer Engineering & Science*, 27, 1698–1702.

Figure legends

Fig. 1. Kinetics of acidification monitored during skim milk fermentation by cultures ST1 (— ■ —), ST2 (——), ST1+2 (1/1.7) (●●●●●), ST1+2 (5/1) (— —), ST3 (——), ST1+3 (1/1.7) (●●●●●), ST1+3 (5/1) (— —). Skim milk suspensions were pasteurised at 95 °C for 6 min.

Fig. 2. Quantity of EPS (mg glucose kg⁻¹) produced by the 7 cultures/co-cultures (ST1, ST2, ST1+2 (1/1.7), ST1+2 (5/1), ST3, ST1+3 (1/1.7), ST1+3 (5/1)) during fermentation of skim milk pasteurised at 95 °C for 6 min. Mean values ± standard deviation (n = 3). Values affected with different letters were significantly different for $p = 0.05$.

Fig. 3. Macromolecular properties of extracted EPS from the 7 cultures/co-cultures (ST1, ST2, ST1+2 (1/1.7), ST1+2 (5/1), ST3, ST1+3 (1/1.7), ST1+3 (5/1): A, average molar mass; B, molar mass distribution in mass; C, molar mass distribution in number of particles; D, intrinsic viscosity. The EPS were classified in high molar mass (□) or low molar mass (■).

Fig. 4. Graininess defect of acid milk gels made by 6 different cultures [ST1, ST2, ST1+2 (1/1.7), ST1+2 (5/1), ST1+3 (1/1.7), ST1+3 (5/1)]. Grains were classified by their diameters: 1–1.5 mm (□), 1.5–2 mm (□), 2–2.5 mm (▣), 2.5–3 mm (▣) and > 3 mm (■). Mean values ± standard deviation (n = 9).




Figure 1




Figure 2




Figure 3




Figure 4

Table 1

Acidification parameters of skim acid milk gel obtained using 3 pure strain cultures (ST1, ST2, ST3) and 4 co-cultures [ST1+2 (1/1.7), ST1+2 (5/1), ST1+3 (1/1.7), ST1+3 (5/1)]. ^a

Parameters	ST1	ST2	ST1+ST2		ST3	ST1+ST3	
			1/1.7	5/1		1/1.7	5/1
Acidification							
Acidification time (min)	348 ± 49 ^{ad}	1351 ± 109 ^b	528 ± 33 ^c	304 ± 24 ^a	ND	395 ± 27 ^d	287 ± 11 ^a
V_{\max} (10 ⁻³ pH unit min ⁻¹)	17.1 ± 1.7 ^a	14.0 ± 0.3 ^b	16.9 ± 1.3 ^a	20.0 ± 0.3 ^c	6.5 ± 0.1 ^d	18.1 ± 1.7 ^{ac}	20.0 ± 0.1 ^c
Gelation							
Time of gel point (min)	206 ± 13 ^a	388 ± 36 ^c	208 ± 10 ^a	157 ± 11 ^b	371 ± 18 ^c	179 ± 11 ^b	164 ± 10 ^b
pH of gel point	5.16 ± 0.05 ^a	5.15 ± 0.05 ^a	5.23 ± 0.01 ^a	5.21 ± 0.05 ^a	5.57 ± 0.01 ^c	5.39 ± 0.06 ^b	5.22 ± 0.01 ^a

^a Reconstituted skim milk was pasteurised at 95 °C, 6 min. V_{\max} is the maximum acidification rate. Data were obtained by monitoring the pH evolution (acidification) and rheological parameters (gelation) during fermentation at 43 °C; values of each characteristic affected with different letters were significantly different for $p = 0.05$ (ND, not determined).

Table 2Textural parameters of pasteurised acid milk gel using 6 different cultures ^a

Cultures	ST1	ST2	ST1+ST2		ST1+ST3	
			1/1.7	5/1	1/1.7	5/1
Firmness (N)	3.36 ± 0.18 ^a	6.44 ± 0.15 ^d	4.78 ± 0.08 ^c	4.14 ± 0.13 ^b	4.43 ± 0.25 ^b	3.85 ± 0.11 ^e
Viscosity (mPa s) at $\gamma = 300 \text{ s}^{-1}$	138 ± 12 ^a	186 ± 4 ^c	193 ± 3 ^c	164 ± 2 ^b	301 ± 10 ^e	239 ± 15 ^d
Spontaneous syneresis (% w/w)	1.8 ± 0.3 ^a	6.3 ± 0.3 ^c	2.7 ± 0.4 ^b	1.4 ± 0.4 ^a	0.4 ± 0.2 ^e	0 ^d

^a Reconstituted skim milk was pasteurised at 95 °C for 6 min; values of each characteristic affected with different letters were significantly different at $p = 0.05$.