Acoustic detection of **radiotracked foraging** bats in temperate lowland forests

Laurent TILLON^a, Michel BARATAUD^b, Sylvie GIOSA^b & Stéphane AULAGNIER^c

a- ONF Réseau mammifères, Direction des Forêts et des Risques Naturels, 2 avenue de Saint-Mandé, 75570 Paris cedex 12, France, laurent.tillon@onf.fr

b- SFEPM, Colombeix, 23400 Saint-Amand-Jartoudeix, France, michel.barataud1@orange.fr

c- Comportement et Ecologie de la Faune Sauvage, INRA, Université de Toulouse, CS

52627, 31326 Castanet Tolosan cedex, France, Stephane.Aulagnier@inra.fr (corresponding author)

Abstract

During the recent years studies of bat activity are predominantly based on ultrasound detection. However this method suffers from several biases such as different species call ranges, temporal and habitat-related variability. In order to test the bias linked to the detection of whispering gleanings bats in temperate lowland forests, we equipped several individuals of *Myotis bechsteinii* and *Plecotus auritus* with transmitters and followed them on their foraging grounds where we simultaneously recorded echolocation calls. Our results highlight the very low detectability of these species whose presence was ascertained at the recording station. On the other hand, we detected the presence of many other species. We suggest methodological recommendations for ultrasound detection whenever gleanings bat species are concerned.

Keywords: Acoustic; Detection; Habitat; *Myotis bechsteinii*; *Plecotus auritus*.

Most inventories or monitoring to identify habitat selection by species are based on presence / absence data. Yet it is often difficult to associate the absence of data with the absence of the target taxon (Brotons *et al.*, 2004; Mackenzie, 2005; Royle, Nichols & Kery, 2005). A gap in a set of presence-absence data may have important implications for the production of high-quality analysis, and thus for understanding ecological phenomena, or even for implementing conservation strategies. Such analysis can have various biases that depend on the method used, the observers, the environment in which the observations take place, the viewing conditions (such as the weather), the detectability of species, the behaviour of animals at specific and/or individual levels, or the modelling approach (Nichols *et al.*, 2000; Kéry & Schmid, 2004; Pellet & Schmidt, 2005; Gooch *et al.*, 2006). The identification of the most appropriate method to answer questions relating to the distribution of a taxon, or the selection of habitats by that taxon, is a priority before beginning a study (Tyre *et al.*, 2003; Mackenzie, 2005; Mackenzie & Royle, 2005; Lobo *et al.*, 2010; Tanadini & Schmidt, 2011).

For bats most recent studies use ultrasonic detection, a method which overcomes the ethics of capturing protected species and has the advantage of recording all bat species emitting a signal likely to be caught by a suitable device (Parsons & Szewczak, 2009; Britzke, Gillam, & Murray, 2013; Fenton, 2013). In addition, this method can be implemented by an observer or by passive recording (Adams *et al.*, 2012, Skalak, Sherwin & Brigham, 2012; Stahlschmidt & Brühl, 2012; Froidevaux *et al.*, 2014; Kubista and Bruckner, 2017). However, studies based on ultrasonic detection have many methodological biases due to the equipment, the sampling effort, the acoustic identification and the measurement of activity, which amplify observer effects (Archaux *et al.*, 2013), making it difficult to compare results (Britzke *et al.*, 2013). In addition, the probability of detection may be relatively low and very different from one species to another (Barataud, 2012). For example, specific detection probabilities obtained over large areas by combining capture and ultrasonic detection, from the historical data of

presence and pseudo-absence varied from 0.239 for *Myotis evotis* to 0.532 for *M. californicus* (Weller, 2008). Detectability is even more different among groups of species (e.g. *Eptesicus/Nyctalus*, *Pipistrellus* and *Myotis*, Barataud, 2012; Archaux *et al.*, 2013; Froidevaux *et al.*, 2016), especially in forests where the signals are generally low due to attenuation by the foliage, so detection is better in open stands than in dense stands (Patriquin *et al.*, 2003; Barataud, 2012). Given the different ecology of taxa and their foraging behaviour which are closely linked to their echolocation strategies (e.g. Aldridge and Rautenbach, 1987; Barataud, 2012; Müller *et al.*, 2012), we can assume that the ultrasound detection of each bat will depend on the species, the habitat where the study takes place and the animals' activity (related to the season and the weather). Moreover, the information is limited to the number of contacts per species at each recording station, and does not provide any estimate of the actual number of individuals (Barataud, 2012). Therefore ultrasonic detection requires an evaluation of the probability of species detection to assess its usefulness as a survey method for investigating habitat selection (Gorresen *et al.*, 2008; Weller, 2008; Hayes, Ober & Sherwin, 2009; Archaux *et al.*, 2013; Pauli *et al.*, 2017).

Therefore, we investigated the probability of detecting echolocation calls of two whispering bats (emitting weak echolocation calls detected at 5-20 m according to Barataud, 2012 and Skiba, 2009) in forest whilst being sure of their presence **by radio telemetry**. **We detected** simultaneously other species or other individuals of the same species for **checking** the efficiency of the bat detector.

Recording stations were located in the forest of Tronçais (Auvergne, France; 46°39'N, 02°41'E). This forest is dominated by sessile oak (*Quercus petraea*), which is regularly accompanied by beech (*Fagus sylvatica*). The understory is dominated by oak (*Quercus petraea*), beech (*Fagus sylvatica*) and holly (*Ilex aquifolium*), more rarely by hornbeam

(*Carpinus betulus*) and hazel (*Corylus avellana*). Forest stands can reach more than 40m high, a height which can therefore limit the detection of bats from the ground when they are hunting in the canopy. The canopy is closed, and scattered bushes cover between 20% and 50% of the understory.

We were able to capture three adult females of *M. bechsteinii* (8.5, 9.1 and 11.5 g), and six adult females of *P. auritus* (7.9 ± 0.3 g) using mist nets across pathways and streams in July 2008 (good weather conditions, temperatures more than 10°C). Following Aldridge & Brigham (1988), we equipped them with BD2N transmitters (0.43 g, manufactured by Holohil Systems Ltd., Carp, Ontario, Canada for a total weight, including surgical glue, less than 6% of each bat weight). This study is part of a broader project licensed by the Ministry of Environment. These bats were then monitored by radio telemetry (using a Regall 2000 receiver and a three element Yagi antenna manufactured by Titley Inc., Australia), and located using the “homing in” technique, a close approach method (see Amelon *et al.*, 2009). When the animal’s position was ascertained, a second grounded operator stood underneath the foraging animal recorded all bat activity at this station for ten minutes using a Pettersson D1000x bat detector. Bats were identified using both heterodyne and time expansion (with Batsound 3.3 software) proposed by Barataud (2012). For *M. bechsteinii* identification was ascertained by the context of sequences following Barataud (2005). One contact was counted every five seconds for all bat species (maximum 120 contacts). Two individuals of one species recorded at the same time were therefore counted as two contacts.

Using a binomial distribution $[B(N, 0.5)]$, where N is the number of radiotracked individuals] we calculated the probability of acoustic detection of both species $p(x \leq n)$, where n is the number of times the species should be detected given that at least one individual with a

transmitter was present at the recording station (number of possible contacts) and x the number of times it was detected (number of recorded contacts).

M. bechsteinii was recorded during 10 minutes at 25 different stations (8.3 ± 4.7 per individual), four in young plots (trees under 12m), 17 in old-growth plots and four at a border of old-growth plots. *P. auritus* was recorded at 35 stations (5.8 ± 1.6 per individual), all in old-growth plots.

During the twenty-five recording periods conducted in presence of radiotracked *Myotis bechsteinii*, this species was contacted four times for a total of 108 contacts (Poisson distribution mean: 0.036), with 97 contacts in one old-growth plot (Table 1); for the latter we could not discriminate the presence of one or several individuals. During the thirty-five recording periods of *Plecotus auritus* this species was contacted two times with only one contact (Poisson distribution mean: 4.76^{E-4}). These detection scores are very low. The probability that *Myotis bechsteinii* was detected four times is $p(x \leq 4) = 4.55^{E-4}$. The probability that *Plecotus auritus* was detected two times is $p(x \leq 2) = 1.84^{E-8}$. Moreover, we detected bats of these two species but not necessarily the radiotracked individuals and always one animal per contact.

At the sixty recording stations we also recorded at least 12 additional bat species and some *Myotis* sp. that were not identified (16 contacts at 8 stations). The most detected species was *Pipistrellus pipistrellus* (595 contacts at 47 stations), followed by *Myotis myotis* (32 contacts at seven stations), and *Nyctalus* sp. (18 contacts at only two stations). *M. nattereri* and *M. mystacinus* were detected in ten and nine periods respectively for 14 contacts each. Other taxa were occasionally detected (Table 1).

Our results show that gleaning bats were only slightly detected in forest whilst they were present. In addition, the number of contacts remained very low, except for one recording station of *Myotis bechsteinii* with ninety-seven contacts in ten minutes, which means almost a permanent detection of the likely radiotracked individual. The ultrasound detection of these gleaner bats is reported to be difficult. Over 38,371 call sequences obtained with ground automatic detectors, Müller *et al.* (2012) recorded *M. bechsteinii* sixty-nine times (for 11530 call sequences only identified as medium-sized *Myotis*) and *Plecotus* only three times. Other tests provided similar results on North American gleaning bats (Skalak *et al.*, 2012; Kennedy, Sillett & Szewczak, 2014; Luszcz *et al.*, 2016). This possibly explains why Kubista and Bruckner (2017) found a perplexingly high variability with batcorders placed ca. 10m apart. Most often we did not detect the target species, even if we detected other taxa, sometimes with a high number of contacts for *P. pipistrellus*, but most of the time with only a small number of contacts, including loud calling species such as *E. serotinus*. Thus, an absence of ultrasonic detection or a very small number of contacts does not necessarily mean the absence or the scarcity of the target species. It is indeed common to confuse the lack of contact with the absence of species (Mackenzie, 2005; Albert & Thuiller, 2008). We know for example that populations are often poorly estimated and are even considered absent when the detectability of species is low, as with some amphibians and birds (Mackenzie & Royle, 2005; Tanadini & Schmidt, 2011). Integrating detection bias is nevertheless an essential component for any ecological study, and generally requires methodological adjustments during field observations (Kéry & Schmid, 2004; Pellet & Schmidt, 2005), in particular for bats (Hayes, 2000; Weller, 2008; Barataud, 2012; Archaux *et al.*, 2013).

Our results are mainly explained by the echolocation parameters and hunting behaviour of the two studied species. These gleaning bats emit weak echolocation calls and forage in cluttered habitats. *Myotis bechsteinii* and *Plecotus auritus* are two species of forest bats which forage in

the foliage (Plank *et al.*, 2012). The range of their echolocation calls is low: ten meters for *Myotis bechsteinii* and five meters for *Plecotus auritus* in forest understory (Barataud, 2012). Moreover, *M. bechsteinii* can hunt insects directly on foliage without emitting ultrasounds (Siemers & Swift, 2006), and could prey on insects in the canopy (Plank *et al.*, 2012), becoming inaudible to a ground observer. *P. auritus* catch moths flying or resting in the foliage (Robinson, 1990; Anderson & Racey, 1991; Shiel *et al.*, 1991; Swift, 1998; Ashrafi *et al.*, 2011), emitting then ultrasonic signals with very low range (Swift, 1998; Barataud, 2012) or hunt by passive listening (Anderson & Racey, 1993). In most situations, the two studied bats are poorly audible by a ground observer unless one animal flies near the bat detector. Our results demonstrate that it is not possible to use acoustic detection from the ground only to survey and monitor forest bats. The use of automatic detectors could be particularly relevant as they afford the installation of microphones in the canopy, as suggested by Rieger & Nagel (2007), Collins & Jones (2009) and Müller *et al.* (2012). However, while automatic recorders offer very long listening periods (Skalak *et al.*, 2012; Stahlschmidt & Brühl, 2012), active detection by an observer can maximize the number of recording stations (Tyre *et al.*, 2003). Moreover, the analysis of recorded calls requires differentiating groups of species, between those which emit their signals over long distances and therefore are easily detectable, and those with a very limited signal range (a few meters) (Hayes *et al.*, 2009; Barataud, 2012). Some studies only provide generic results due to the difficulty in obtaining correct species identification of several groups of taxa, including medium-sized *Myotis* species (Rieger & Nagel, 2007; Collins & Jones, 2009, Barataud 2012). This has little consequence regarding *Plecotus* whose species share quite similar ecology (Ashrafi *et al.*, 2011; Swift, 1998). This is detrimental for *Myotis* species which have very different niches and population trends (Meschede & Heller, 2003; Müller *et al.*, 2012).

In addition, we can issue two types of recommendations, according to the purpose of the study. For inventories, as detection rate is locally low for several bat species we recommend increasing the number of ground recording stations per night in the same forest. For studies of habitat selection, we support using several automatic detectors recording simultaneously bats from the ground to the canopy overnight or, depending on target species, ground observers with microphones also from the ground to the canopy for a long time (one hour or more). One limit then could be the huge amount of files to analyze and timetable and financial trade-offs will drive survey protocols depending on the objectives.

Acknowledgements.

We thank the members of the mammal network of the Office National des Forêts for their help in the field, and Dorine Pasqualini for checking the English language. We also thank reviewers and associate-editor for useful comments on our manuscript.

References

- 1 Adams, A.M., Jantzen, M.K., Hamilton, R.M., Fenton, M.B., 2012. Do you hear what I hear? Implications of detector selection for acoustic monitoring of bats. *Meth. Ecol. Evol.* 3, 992–998.
- 2 Albert, C.H., Thuiller, W., 2008. Favourability functions versus probability of presence: Advantages and misuses. *Ecography* 31, 417–422.
- 3 Aldridge, H.D.J.N., Rautenbach, I.L., 1987. Morphology, echolocation and resource partitioning in insectivorous bats. *J. Anim. Ecol.*, 56, 763–778.
- 4 Aldridge, H.D.J.N., Brigham, R.M., 1988. Load carrying and maneuverability in an insectivorous bat: a test of the 5% “rule” of radio-telemetry. *J. Mammal.*, 69, 379–382

- 197 5 Amelon, S.K., Dalton, D.C., Millsbaugh, J.J., Wolf, S.A., 2009. Radiotelemetry.
198 Techniques and analysis. In: Kunz, T.H., Parsons, S. (Eds), Ecological and behavioral
199 methods for the study of bats (second edition). The Johns Hopkins University Press,
200 Baltimore, pp. 57–77.
- 201 6 Archaux, F., Tillon, L., Fauvel, B., Martin, H., 2013. Foraging habitat use by bats in a large
202 temperate oak forest: Importance of mature and regeneration stands. *Rhinolophe* 19:
203 47–58.
- 204 7 Ashrafi, S., Beck, A., Rutishauser, M., Arlettaz, R., Bontadina, F., 2011. Trophic niche
205 partitioning of cryptic species of Long-eared bats in Switzerland: Implications for
206 Conservation. *Eur. J. Wildl. Res.* 57, 843–849.
- 207 8 Barataud, M., 2005. Acoustic variability, and identification possibilities for seven European
208 bats of the genus *Myotis*. *Rhinolophe*, 17, 43–62.
- 209 9 Barataud, M., 2012. Ecologie acoustique des Chiroptères d'Europe. Identification des
210 espèces, étude de leurs habitats et comportements de chasse. Biotope - Muséum
211 National d'Histoire Naturelle, Mèze - Paris.
- 212 10 Britzke, E.R., Gillam, E.H., Murray, K.L., 2013. Current state of understanding of
213 ultrasonic detectors for the study of bat ecology. *Acta Theriol.* 58, 109–117.
- 214 11 Brotons, L., Thuiller, W., Araujo, M.B., Hirzel, A.H., 2004. Presence-absence versus
215 presence-only modelling methods for predicting bird habitat suitability. *Ecography* 27,
216 437–448.
- 217 12 Collins, J., Jones, G., 2009. Differences in bat activity in relation to bat detector height:
218 Implications for bat surveys at proposed windfarm sites. *Acta Chiropt.* 11, 343–350.
- 219 13 Fenton, M.B. 2013. Chapter 3 - Evolution of echolocation. In: Adams, R.A., Pedersen,
220 S.C. (Eds), Bat evolution, ecology, and conservation. Springer Science and Business
221 Media, New York, pp. 47-70.

- 222 14 Froidevaux, J.S.P., Zellweger, F., Bollmann, K., Obrist, M.K., 2014. Optimizing passive
223 acoustic sampling of bats in forests. *Ecol. Evol.* 4, 4690–4700.
- 224 15 Froidevaux, J.S.P., Zellweger, F., Bollmann, K., Jones, G., Obrist, M.K., 2016. From field
225 surveys to LIDAR: Shining a light on how bats respond to forest structure. *Remote*
226 *Sensing Environ.* 175, 242–250.
- 227 16 Gannon, W.L., Sherwin, R.E., 2004. Are acoustic detectors a ‘silver bullet’ for assessing
228 habitat use by bats? in: Brigham, R.M., Kalko, E.K.V., Jones, G., Parsons, S.,
229 Limpens, H.J.G.A. (Eds), *Bat echolocation research: Tools, techniques and analysis*.
230 *Bat Conservation International*, Austin, pp. 38-45.
- 231 17 Gooch, M.M., Heupel, A.M., Price, S.J., Dorcas, M.E., 2006. The effects of survey
232 protocol on detection probabilities and site occupancy estimates of summer breeding
233 anurans. *Appl. Herpetol.* 3, 129–142.
- 234 18 Gorresen, P.M., Miles, A.C., Todd, C.M., Bonaccorso, F.J., Weller, T.J., 2008. Assessing
235 bat detectability and occupancy with multiple automated echolocation detectors. *J.*
236 *Mammal.* 89, 11–17.
- 237 19 Hayes, J.P., 2000. Assumptions and practical considerations in the design and
238 interpretation of echolocation-monitoring studies. *Acta Chiropt.* 2, 225–236.
- 239 20 Hayes, J.P., Ober, H.K., Sherwin, R.E., 2009. Survey and monitoring of bats. In: Kunz,
240 T.H., Parsons, S. (Eds), *Ecological and behavioral methods for the study of bats*.
241 Second edition. The Johns Hopkins University Press, Baltimore, pp. 112-129.
- 242 21 Kennedy, J.-P., Sillett, S.C., Szewczak, J.M., 2014. Bat activity across the vertical gradient
243 of an old-growth *Sequoia sempervirens* forest. *Acta Chiropt.* 16, 53–63.
- 244 22 Kéry, M., Schmid, H., 2004. Monitoring programs need to take into account imperfect
245 species detectability. *Basic appl. Ecol.* 5, 65–73.

23 Kubista, C.E., Bruckner, A., 2017. Within-site variability of field recordings from stationary, passively working detectors. *Acta Chiropt.* 19, 189–197.

24 Limpens, H.J.G.A., McCracken, G.F., 2004. Choosing a bat detector: Theoretical and practical aspects. In: Brigham, R.M., Kalko, E.K.V., Jones, G., Parsons, S., Limpens, H.J.G.A. (Eds), *Bat echolocation research: Tools, techniques and analysis*. Bat Conservation International, Austin, pp. 28-37.

25 Lobo, J.M., Jiménez-Valverde, A., Hortal, J., 2010. The uncertain nature of absences and their importance in species distribution modelling. *Ecography* 33, 103–114.

26 Luszcz, T.M.J., Barclay, R.M.R., 2016. Influence of forest composition and age on habitat use by bats in southwestern British Columbia. *Can. J. Zool.* 94, 145–153.

27 Mackenzie, D.I., 2005. Was it there? Dealing with imperfect detection for species presence/absence data. *Austr. N. Zeal. J. Stat.* 47, 65–74.

28 Mackenzie, D.I., Royle, J.A., 2005. Designing occupancy studies: general advice and allocating survey effort. *J. appl. Ecol.* 42, 1105–1114.

29 Meschede, A., Heller, K.-G., 2003. Ecologie et protection des chauves-souris en milieu forestier. *Rhinolophe* 16, 1–248.

30 Müller, J., Mehr, M., Bässler, C., Fenton, M.B., Hothorn, T., Pretzsch, H., Klemmt, H.-J., Brandl, R., 2012. Aggregative response in bats: Prey abundance versus habitat. *Oecologia* 169, 673–684.

31 Nichols, J.D., Hines, J.E., Sauer, J.R., Fallon, F.W., Fallon, J.E., Heglund, P.J., 2000. A double-observer approach for estimating detection probability and abundance from point counts. *Auk* 117, 393–408.

32 Parsons, S., Szewczak, J.M., 2009. Detecting, Recording, and analysis the vocalizations of bats. In: Kunz, T.H., Parsons, S. (Eds), *Ecological and behavioral methods for the*

270 study of bats. Second edition. The Johns Hopkins University Press, Baltimore, pp. 91-
 271 111.

272 33 Patriquin, K.J., Hogberg, L.K., Chruszcz, B.J., Barclay, R.M.R., 2003. The influence of
 273 habitat structure on the ability to detect ultrasound using bat detectors. Wildl. Soc.
 274 Bull. 31, 475–481.

275 34 Pauli, B.P., Zollner, P.A., Haulton, G.S., 2017. Nocturnal habitat selection of bats using
 276 occupancy models. J. Wildl. Manage. 81, 878–891.

277 35 Pellet, J., Schmidt, B.R., 2005. Monitoring distributions using call surveys: estimating site
 278 occupancy, detection probabilities and inferring absence. Biol. Conserv. 123, 27–35.

279 36 Plank, M., Fiedler, K., Reiter, G., 2012. Use of forest strata by bats in temperate forests. J.
 280 Zool. 286, 154–162.

281 37 Rieger, I., Nagel, P., 2007. Vertical stratification of bat activity in a deciduous forest. In:
 282 Unterseher, M., Morawetz, W., Klotz, S., Arndt, E. (Eds), The canopy of a temperate
 283 floodplain forest - Results from five years of research at the Leipzig Canopy Crane.
 284 Universität Leipzig, The Leipzig Canopy Crane Project, Leipzig, pp. 141-149.

285 38 Robinson, M.F., 1990. Prey selection by the brown long-eared bat (*Plecotus auritus*).
 286 Myotis 28, 5–18.

287 39 Royle, J.A., Nichols, J.D., Kery, M., 2005. Modelling occurrence and abundance of
 288 species when detection is imperfect. Oikos 110, 353–359.

289 40 Shiel, C.B., McAney, C.M., Fairley, J.S., 1991. Analysis of the diet of Natterer's bat
 290 *Myotis nattereri* and the common long-eared bat *Plecotus auritus* in the West of
 291 Ireland. J. Zool. 223, 299–305.

292 41 Siemers, B.M., Swift, S.M., 2006. Differences in sensory ecology contribute to resource
 293 partitioning in the bats *Myotis bechsteinii* and *Myotis nattereri* (Chiroptera:
 294 Vespertilionidae). Behav. ecol. Sociobiol. 59, 373–380.

- 295 42 Skalak, S.L., Sherwin, R.E., Brigham, R.M., 2012. Sampling period, size and duration
296 influence measures of bat species richness from acoustic surveys. *Meth. Ecol. Evol.* 3,
297 490–502.
- 298 43 Stahlschmidt, P., Brühl, C.A., 2012. Bats as bioindicators – the need of a standardized
299 method for acoustic bat activity surveys. *Meth. Ecol. Evol.* 3, 503–508.
- 300 44 Swift, S.M., 1998. Long-eared bats. Poyser Natural History, T and AD Poyser Ltd,
301 London.
- 302 45 Tanadini, L.G., Schmidt, B.R., 2011. Population size influences amphibian detection
303 probability: Implications for biodiversity monitoring programs. *PLoS ONE* 6, e28244.
- 304 46 Tyre, A.J., Tenhumberg, B., Field, S.A., Niejalke, D., Parris, K., Possingham, H.P., 2003.
305 Improving precision and reducing bias in biological surveys: estimating false-negative
306 error rates. *Ecol. Applic.* 13, 1790–1801.
- 307 47 Weller, T.J., 2008. Using occupancy estimation to assess the effectiveness of a regional
308 multiple-species conservation plan: Bats in the Pacific Northwest. *Biol. Conserv.* 141,
309 2279–2289.

310

311 Table 1: Number of contacts of bat species at recording stations performed in the presence of
312 *Myotis bechsteinii* (3 adult females, 25 stations, left – 1-4: young plots, 5-21: old-
313 growth plots, 22-25: border of old-growth plots) and *Plecotus auritus* (6 adult females,
314 35 stations in old-growth plots, right) in temperate lowland forests.

Station	<i>P. pipistrellus</i>	<i>P. kuhlii</i>	<i>E. serotinus</i>	<i>Plecotus</i> sp.	<i>B. barbastellus</i>	<i>M. bechsteinii</i>	<i>M. emarginatus</i>	<i>M. nattereri</i>	<i>M. mystacinus</i>	<i>M. myotis</i>	<i>Myotis</i> sp.
1	1							1			
2	3					5	1			4	
3	3					1		1	23		
4		1									
5	72							1			
6	4								1	1	
7	5										
8	21										
9	35	3									
10		1				97					
11					1					4	
12		1						1			
13	1							1			
14	1						1			1	
15	4	1						1	3		
16	37			3							
17											
18	4	3			1						
19	4										
20	9	1				5					
21	7								1		
22	6	1			1						
23	8			2				3			
24	31	3	4								1
25	7										
a	263	8	11	5	3	108	2	9	1	28	6
b	20	4	6	2	3	4	2	7	1	4	3

Station	<i>R. ferrumequinum</i>	<i>P. pipistrellus</i>	<i>P. kuhlii</i>	<i>E. serotinus</i>	<i>Nyctalus</i> sp.	<i>Plecotus</i> sp.	<i>B. barbastellus</i>	<i>M. daubentonii</i>	<i>M. nattereri</i>	<i>M. mystacinus</i>	<i>M. myotis</i>	<i>Myotis</i> sp.
1		5						21				
2									2			
3		1								4		4
4							2					
5		1					5		1			3
6												
7												1
8				1						1	2	
9		9								1		
10		6										
11		3	2									
12		6										
13		26				1						
14						1						
15		1								1	1	1
16	1	1										
17		23										
18		18										
19										2		
20		3							2			
21		19								2		
22		8										
23		3										
24		5										
25		17	4									
26		11	1								1	
27		23			6							
28		3			12							1
29		5										
30		42								1		
31		72						2				
32		16										
33		2										
34		3		2								
35											1	
a	1	332	7	3	18	2	7	23	5	13	4	10
b	1	27	3	2	2	2	2	2	3	8	3	5

315