

The Escherichia coli colibactin resistance protein ClbS is a novel DNA binding protein that protects DNA from nucleolytic degradation

Katja Molan, Zdravko Podlesek, Vesna Hodnik, Matej Butala, Eric Oswald

▶ To cite this version:

Katja Molan, Zdravko Podlesek, Vesna Hodnik, Matej Butala, Eric Oswald. The Escherichia coli colibactin resistance protein ClbS is a novel DNA binding protein that protects DNA from nucleolytic degradation. DNA Repair, 2019, 79, pp.50-54. 10.1016/j.dnarep.2019.05.003 . hal-02618217

HAL Id: hal-02618217 https://hal.inrae.fr/hal-02618217

Submitted on 25 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Contents lists available at ScienceDirect

DNA Repair

journal homepage: www.elsevier.com/locate/dnarepair

The *Escherichia coli* colibactin resistance protein ClbS is a novel DNA binding protein that protects DNA from nucleolytic degradation

Katja Molan^a, Zdravko Podlesek^a, Vesna Hodnik^{a,1}, Matej Butala^a, Eric Oswald^b, Darja Žgur Bertok^{a,*}

^a Department of Biology, Biotechnical Faculty, University of Ljubljana, Večna pot 111, Ljubljana, Slovenia
^b IRSD, Université de Toulouse, INSERM, INRA, ENVT, UPS, Toulouse, France

ARTICLE INFO

Keywords: ClbS DNA-binding protein DNA damage protection Pks genomic island Escherichia coli

ABSTRACT

Cells employ specific and nonspecific mechanisms to protect their genome integrity against exogenous and endogenous factors. The *clbS* gene is part of the polyketide synthase machinery (*pks* genomic island) encoding colibactin, a genotoxin implicated in promoting colorectal cancer. The *pks* is found among the *Enterobacteriaceae*, in particular *Escherichia coli* strains of the B2 phylogenetic group. Several resistance mechanisms protect toxin producers against toxicity of their products. ClbS, a cyclopropane hydrolase, was shown to confer colibactin resistance by opening its electrophilic cyclopropane ring. Here we report that ClbS sustained viability and enabled growth also of *E. coli* expressing another genotoxin, the Usp nuclease. The *recA::gfp* reporter system showed that ClbS protects against Usp induced DNA damage. To elucidate the mechanism of ClbS mediated protection, we studied the DNA binding ability of the ClbS protein. We show that ClbS directly interacts with single-stranded DNA (ssDNA) and double-stranded DNA (dsDNA), whereas ssDNA seems to be the preferred substrate. Thus, the ClbS DNA-binding characteristics may serve bacteria to protect their genomes against DNA degradation.

1. Introduction

To protect the integrity of their genomes against exogenous and endogenous assault, cells employ specific and as well as nonspecific mechanisms. Exogenous factors are radiation, chemical agents and others, while endogenous factors are reactive oxygen species, metabolic products [1] and in a number of bacterial species potentially also genotoxins. Genotoxins such as colibactin are significant bacterial virulence factors that provoke DNA damage in eukaryotic cells. However, the genotoxin producing cell must be protected against its cognate toxin.

Colibactin is encoded by the polyketide synthase (*pks*) genomic island present among the *Enterobacteriaceae*, particularly pathogenic and commensal *Escherichia coli* strains of the B2 phylogenetic group [2]. The prevalence of these strains has in developed countries increased significantly [3,4]. Colibactin generates DNA cross-links that are, during DNA damage repair, converted to double strand breaks [5]. Recently, alkylation of DNA by colibactin has been demonstrated [6]. It has also been shown to induce colorectal cancer [7–11].

Colibactin is initially produced as an inactive prodrug that is

activated by ClbP-mediated cleavage upon export to the periplasm [12,13]. Further, it was postulated that ClbS, also encoded by the *pks* island, is a colibactin resistance protein [14] that cleaves the colibactin cyclopropane ring, essential for genotoxicity, via its cyclopropane hydrolase activity [15].

As we previously demonstrated that immunity to the Usp genotoxin is assured by the Imu3 protein via nonspecific DNA binding [16], we postulated that ClbS might also provide direct genome protection. We therefore investigated whether ClbS exhibits DNA binding ability. Our results show that ClbS is a multifunctional protein, that besides directly inactivating colibactin, as shown previously, also associates with DNA to presumably protect the genome against certain genotoxic agents.

2. Material and methods

2.1. Bacterial strains and plasmid construction

Laboratory *E. coli* strains and plasmids used in this study are presented in Table 1. The *clbS* gene was PCR amplified from BAC pks^+ plasmid [2] with ClbS-F (5'- TGGATCCGCTGTTCCATCATCAAAAGAA

* Corresponding author at: Department of Biology, Biotechnical Faculty, University of Ljubljana, Večna pot 111, 1000 Ljubljana, Slovenia. *E-mail address:* darja.zgur.bertok@bf.uni-lj.si (D. Žgur Bertok).

https://doi.org/10.1016/j.dnarep.2019.05.003

Received 2 December 2018; Received in revised form 29 March 2019; Accepted 18 May 2019 Available online 19 May 2019

1568-7864/ © 2019 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (http://creativecommons.org/licenses/BY/4.0/).





¹ Present address: Sandoz, Kolodvorska 27, SI-1234 Mengeš, Slovenia.

Table 1

Bacterial strains and plasmids used in this study.

DH5a (DE3)Expression host, harboring the lambda DE3.Thermo Fisher S $E. coli BL21(DE3)$ Expression host, harboring the lambda DE3; Cm^R .Thermo Fisher S $BAC pks^+$ The source of the $clbS$ gene.(2) $pUSP4$ The source of the usp gene; Ap^R .(17) $pET8c$ $E. coli T7$ expression vector, containing an N-terminal hexahistidine tag (His ₆) and thrombin cleavage site.Novagen $pCLB-his$ $pET8c$ with $clbS$ gene.Thermo Fisher S $pLVSS$ $pLys$ with removed lysozyme gene; Cm^R .Thermo Fisher S $pLCUBI$ $pLysS$ with $clbS$ and usp genes; Cm^R and Ap^R .This study $pLUBI$ $pLysS$ with usp gene; Cm^R and Ap^R .This study $pCO201$ Description of W^R .This study	Scientific Scientific

GAG-3`) and ClbS-R (5`-TTACGCGTTATTCTGCAAGACATTTCTGCAG-3`) primers (Sigma-Aldrich) at 55 °C annealing temperature. The PCR product, digested with *Bam*HI and *Mlu*I, was cloned into the expression

vector pET8c, in a fusion with a thrombin-cleavable N-terminal hexahistidine tag (His₆-tag). This plasmid, designated pCLB-his, was subsequently used for ClbS protein isolation and for the construction of vectors, used for in vivo studies. Sequencing of the inserted clbS fragment was performed to check for potential PCR introduced mutations. For the construction of vectors, the BamHI and EcoRV digested and blunted fragment of the low copy number pLysS plasmid was employed. This backbone fragment contained only the replication region and the chloramphenicol resistance gene, with the T7 lysozyme gene removed. The source of the usp gene for subcloning into the pLysS fragment was plasmid pUSP4 [17]. Two plasmids were thus constructed, pLUBI, carrying the usp gene but without its immunity genes, and pLCUBI harbouring usp and clbS, and were transformed into E. coli BL21 (DE3). Both genes were controlled by independent T7 polymerase promoters. To follow the SOS response, plasmid pSC201 with a recA::gfp fusion was used [18], transformed into either E. coli BL21 (DE3) with pLCUBI, or E. coli BL21 (DE3) with pLUBI.

2.2. Expression, Isolation, purification and detection of the ClbS protein

ClbS was expressed in E. coli strain BL21 (DE3) pCLB-his. An overnight culture was used to inoculate liquid Luria Bertani medium supplemented with 25 µg/mL chloramphenicol and grown at 37 °C to an OD₆₀₀ 0.5 when ClbS production was induced by the addition of isopropyl β-D-1-thiogalactopyranoside (IPTG) at a final concentration of 0.8 mM. The culture was subsequently grown for an additional 4 h when biomass was collected by 10 min centrifugation at 4.000 \times g. His₆-tagged ClbS was purified by Ni-NTA metal affinity chromatography (Qiagen) and stored in elution buffer (50 mM NaH₂PO₄, 300 mM NaCl, 250 mM imidazole, pH 8) at 4 °C. Further, overnight dialysis was performed against a final buffer (50 mM NaH₂PO₄·H₂O, 200 mM NaCl, 20 mM $\beta\text{-mercaptoethanol},$ pH 8) at 4 °C. The concentration of the purified ClbS protein was determined using a NanoDrop 1000 (Thermo Fisher Scientific) with the extinction coefficient at 280 nm of 37530 M⁻¹ cm⁻¹ and by Pierce BCA protein assay kit, using Bovine Serum Albumin (BSA) (Sigma-Aldrich) as the standard. Additionally, the His6-tag was removed from His6-ClbS with the Thrombin cleavage capture kit (Novagen) and the buffer exchanged by overnight dialysis as described above. To assay the removal of the affinity tag, the thrombin-digested or undigested His6-ClbS samples were resolved on a 12% acrylamide gel (Fig. S1).

2.3. Electrophoretic mobility shift assays (EMSA)

Various concentrations of ClbS carrying or lacking the affinity His₆tag (0.15 μ M – 2.4 μ M) and 60 ng/ μ L of DNA (Table 2) were used to establish the nucleic acid-binding ability of ClbS. ClbS was incubated with DNA in 1 X FastDigest Green Buffer (Fermentas) at 37 °C for 15 min, prior to the analysis of the protein-DNA complexes by EMSA analysis on 1.5–2 % agarose gels. As a negative control BSA was included at a final concentration of 1.6 μ M. Additionally, mixtures of pUC19/*Eco*RI (60 ng/ μ L) and ClbS (0.15 μ M – 2.4 μ M) were pre-incubated at 37 °C for 5–15 minutes when colicin E7 (ColE7), isolated from *E. coli* [19] or commercial DNase I (Sigma-Aldrich) were added in final concentrations 0.42 mg/ml and 0.5 mg/mL, respectively, followed by an additional 30 min incubation. As a control, DNA without ClbS was incubated with ColE7 or commercial DNase I. Results were analyzed employing 2% agarose gels.

2.4. Surface plasmon resonance assays (SPR)

Measurements were performed at 25 °C on a Biacore T100 (GE Healthcare, UK) at the Infrastructural Centre for Analysis of Molecular Interactions (University of Ljubljana, Slovenia). To prepare biotinylated DNA for immobilization on the streptavidin sensor chip, plasmid pJBPL-A was used as a matrix DNA to amplify the 403 bp long DNA sequence (from Salmonella bongori). Annealing temperature of 55 °C was used to amplify with primers Bong-biotin (5' -CATAAAACCTCGTGTA TGGTGCGC-3 and Bong-R (5'-GAATTCCATTTTACGACATCCGAATAT AAATTAAACGAA-3`) (Integrated DNA Technology). Approximately 50 response units (RU) of the 3'-biotinylated 403 bp long PCR product were immobilized on the second flow cell of the streptavidin (SA) sensor chip (GE Healthcare) in SPR buffer (20 mM Tris (pH 7.4), 140 mM NaCl, 2 mM DTT, 0.005% surfactant P20 and 0.1 mg/ml BSA). The first flow cell was left empty and used to control any non-specific binding to the sensor surface. The interaction between ClbS and chipimmobilized DNA was studied by injecting various concentrations of ClbS in SPR buffer. Injections of ClbS over the immobilized DNA fragment were performed at a flow rate of $10\,\mu l \, min^{-1}$ for $2 \, min$ with dsDNA and 4 min with ssDNA. The sensor chip with bound DNA was regenerated by injection of SPR buffer containing 0.03% SDS for 15 s at a flow rate of 30 µl min⁻¹. Further, chip-immobilized dsDNA denaturation was applied with 2 pulses of 50 mM NaOH for 30 s to obtain ssDNA. The response after NaOH treatment dropped for approximately 20 RU, indicating that one DNA chain was removed. Injections of various concentrations of ClbS in SPR buffer was applied at a flow rate of 10 µl min⁻¹ for 4 min. SPR experiments were performed in triplicate and representative experiments are shown. The data were analyzed with the Biacore T100 Evaluation Software and equilibrium dissociation constants (K_D) were determined by fitting the data to the steady state affinity model.

2.5. Quenching of intrinsic ClbS tryptophan fluorescence by ss- and dsDNA

Intrinsic tryptophan fluorescence of ClbS ($40 \mu g/mL$ in 50 mM phosphate buffer, 200 mM NaCl, β -mercaptoethanol, pH 8.0) was measured. DNA used for titrations are presented in Table 2. After adding DNA, a 10 s incubation followed when measurements were made in a constantly stirred 1 cm path length quartz cuvette with a scan

Table 2

DNA	Length (bp or bases)	Application	Reference
pUC19	2686 3377	EMSA Matrix DNA for PCB reaction	Fermentas This laboratory
PCR amplicon	403	SPR (ds- and ssDNA), EMSA,	This study
ds- and ssDNA phage genome M13	6407	Trp fluorescence quenching EMSA, Trp fluorescence quenching	[20]

rate of 250 nm/min (37 $^{\circ}$ C, excitation and emission slits at 5 nm each). The fluorescence emission spectra were recorded in the wavelength range of 300–450 nm after excitation at 295 nm, specifically monitoring the tryptophan fluorescence emission. Fluorescent emissions of the buffer titrated with DNA (background intensities) were subtracted.

2.6. Growth of E. coli $clbS^+$ usp^+ and E. coli usp^+ strains and cell morphology analysis

The two *E. coli* BL21 (DE3) strains were cultivated at 37 °C in liquid BHI medium with ampicillin and chloramphenicol at final concentrations 100 and 20 μ g/ml, respectively. To suppress basal level expression of the T7 polymerase promoter, glucose was added at a final concentration of 0.4%. When cultures reached OD₆₀₀ 0.5, protein synthesis was induced by addition of IPTG at final concentration 0.8 mM. One ml of both cultures was sampled before induction and each hour following induction (at least three hours). The OD₆₀₀ was measured and plotted in time.

Prior to microscopy, the *E. coli* BL21 (DE3) $clbS^+$ usp^+ and *E. coli* usp^+ strains were transformed with a plasmid carrying a *recA::gfp* gene fusion and cultured as described above in media supplemented with 30 µg/ml kanamycin. Both cultures were sampled before and every hour following induction. Cells were washed twice with NaCl and attached to glass slides coated with 0.1% (w/v) poly-L-lysine (Sigma). Cell morphology was observed under light and *recA::gfp* expression under fluorescence microscopy, using a Zeiss AxioImager Z1 microscope equipped with a HRc digital camera.

3. Results and discussion

3.1. ClbS binds ss- and dsDNA in a non-specific manner

Initially, EMSA was applied and showed that ClbS causes circular and linear DNA retardation (Fig. 1A) and, that it interacts with ss- or dsDNAs of various lengths and source (Table 2, Fig. S2). Thus, EMSA experiments implied that ClbS binds DNA in a non-specific manner. It is of note that ClbS binding activity was not affected by the His₆ tag (Fig. S3). SPR spectrometry measurements of ClbS binding to ssDNA (Fig. 1B) or dsDNA (Fig. 1C) revealed an approximately 40-times higher affinity of ClbS for ssDNA exhibiting an apparent equilibrium dissociation constant (K_D) of 24.5 ± 1.2 nM compared to dsDNA exhibiting a K_D of 936.8 \pm 12.3 nM (Fig. 1 D). Additionally, to confirm that ClbS interacts with DNA we assayed the tryptophan fluorescence spectra of ClbS, which carries five tryptophan residues, in the presence or in the absence of DNA. We observed quenching of intrinsic ClbS tryptophan fluorescence by various ss- and dsDNAs. The most pronounced decrease in intensity of emission maximum was observed upon titration with ssDNA (Fig. 1E). Our results implied that the five ClbS tryptophan residues behave as two populations of fluorophores, one accessible to quenchers and the other buried in the protein structure [21], presented as downward curvature in the Stern-Volmer plot (Fig. 1F).

3.2. ClbS protects DNA from deoxyribonucleases

DNA protection by ClbS was examined by measuring growth of the E. coli strain carrying a plasmid encoded usp gene and its cognate immunity genes replaced with *clbS*, in comparison to a strain harbouring only the usp gene (Table 1). Expression of usp and clbS was controlled by the T7 polymerase promoter. Following IPTG induction of usp and *clbS* the strain encoding only *usp* (lacking its immunity genes or *clbS*), exhibited dramatically reduced growth in comparison to the strain with both usp and clbS (Fig. 2A). DNA damage induces the SOS response involved in DNA repair. The principal regulators of the SOS response are the LexA repressor and RecA activator. Upon DNA damage, RecA is activated by binding to exposed ssDNA which stimulates LexA autocleavage, derepressing SOS genes. SOS induction inhibits cell division, resulting in cell filamentation [22]. Light microscopy of IPTG-induced cells expressing usp revealed pronounced filamentation (Fig. 2B), while typical cell shape and size were detected among E. coli expressing both usp and clbS genes (Fig. 2C). The "two examined strains also carried a plasmid encoded recA::gfp, a DNA damage SOS reporter gene fusion. Fluorescent microscopy revealed that usp+ cells without clbS, additionally exhibited elevated expression of recA::gfp (Fig. 2D) when compared to the $usp^+ clbS^+$ cells (Fig. 2E). Filamentation and elevated expression of recA are hallmarks of induction of the DNA damage inducible SOS response. Thus, typical cell shape and size, as well as basal level expression of *recA* in $usp^+ clbS^+$ cells indicates, that ClbS protects cells from Usp nuclease activity.

We subsequently hypothesized that binding of ClbS could provide DNA protection against other deoxyribonucleases. Co-incubation of DNA and ClbS protected plasmid DNA against DNase I (Fig. 2F) and the ColE7 nuclease (Fig. 2G) in a concentration dependent manner, confirming that ClbS provides protection against various nucleases by presumably interacting non-specifically with DNA. ClbS binding to DNA is evident as DNA retardation (Fig. 2G, and F). Other examples of protective bacterial proteins have been characterized previously namely, two *E. coli* nucleoid associated proteins, the multifunctional Dps and CbpA, that non-specifically bind DNA, to drive nucleoid compactation and DNA protection [23–26]. Further, the α/β type small acid-soluble spore proteins that are synthesized during sporulation, bind DNA changing its conformation to an A-like helix, that is more resilient against damage [27,28].

In conclusion, our results imply that ClbS, besides directly interacting with colibactin is also a novel DNA binding protein. Thus, beyond colibactin, ClbS also protects DNA against degradation/genotoxic activity presumably providing a selective advantage for bacterial strains encountering stressful conditions also within the host.

Conflict of interest

The authors have no conflict of interest.

Footnotes

Data accessibility available via Mendeley Data: https://data.mendeley.com/datasets/4y3wpkrfzc/draft?a =



Fig. 1. ClbS is a DNA binding protein. (A) EMSA of ClbS interacting with the DNA. M designates marker (bp). Lane 1 contains pUC19 plasmid DNA, lane 2 pUC19 with BSA and lane 3 pUC19 with ClbS. Lane 4 contains pUC19 linearized with *Eco*RI, lane 5 pUC19/*Eco*RI with BSA and lane 6 pUC19/*Eco*RI with ClbS. (B) SPR sensorgrams showing the interaction of ClbS with 403 bp long chip-immobilized ssDNA or (C) dsDNA. (D) The apparent equilibrium dissociation constant (K_D) values of ClbS interacting with ssDNA, red or with dsDNA, black. The average K_Ds and standard deviations were determined from three titrations of each protein. (E) Quenching of intrinsic tryptophan fluorescence of ClbS titrated with ssDNA. The legend denotes ssDNA concentrations preincubated with ClbS. (F) Stern-Volmer plot of tryptophan quenching within ClbS due to interaction with ssDNA or dsDNA.



Fig. 2. ClbS protects DNA from deoxyribonucleases. (A) Growth curves of E. coli usp⁺clbS⁺ (black squares) and E. coli usp⁺ strains (white squares). Induction of proteins at time 0 h. The experiment was performed seven times, and the means \pm standard errors of the means (error bars) are shown. (B) Light and (D) fluorescent microscopy of the E. coli usp⁺ strain following 3 h - induction. (C) Light and (E) fluorescent microscopy of the E. coli $usp^+ clbS^+$ strain following 3 h - induction. (F and G) ClbS protect DNA against DNase degradation. (F) M designates marker (bp). Lane 1 contains pUC19/EcoRI plasmid DNA, lane 2 DNA with DNase I (0.5 mg/mL), lanes 3-7 DNA with ClbS (2.4-0.15µM) and DNase I. (G) M designates marker (bp). Lane 1 contains pUC19/EcoRI DNA, lane 2 DNA with ColE7 (0.42 mg/ml) and lanes 3-6 DNA with ClbS (2.4-0.15µM) and ColE7.



Acknowledgements

dr. Kristina Sepčič with interpretation of results obtained from intrinsic ClbS tryptophan fluorescence quenching. This work was supported by grant P1-0198 and J1-8150 from the Slovene Research Agency. Katja Molan is a recipient of a PhD grant from the Slovene Research Agency.

We thank dr. Miloš Vittori for assistance with microscopy and prof.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:https://doi.org/10.1016/j.dnarep.2019.05.003.

References

- N. Chatterjee, G.C. Walker, Mechanisms of DNA damage, repair and mutagenesis, Environ. Mol. Mutagen. 58 (2017) 235–263, https://doi.org/10.1002/em.22087.
- [2] J.P. Nougayrède, S. Homburg, F. Taieb, M. Boury, E. Brzuszkiewicz, G. Gottschalk, C. Buchrieser, J. Hacker, U. Dobrindt, E. Oswald, *Escherichia coli* induces DNA double-strand breaks in eukaryotic cells, Science 313 (2006) 848–851, https://doi. org/10.1126/science.1127059.
- [3] P. Escobar-Páramo, K. Grenet, A. Le Menac'h, L. Rode, E. Salgado, C. Amorin, S. Gouriou, B. Picard, M.C. Rahimy, A. Andremont, E. Denamur, R. Ruimy, Largescale population structure of human commensal *Escherichia coli* isolates, Appl. Environ. Microbiol. 70 (2004) 5698–5700, https://doi.org/10.1128/AEM.70.9. 5698-5700.2004.
- [4] F.L. Nowrouzian, A.E. Wold, I. Adlerberth, *Escherichia coli* strains belonging to phylogenetic group B2 have superior capacity to persist in the intestinal microflora of infants, J. Infect. Dis. 191 (2005) 1078–1083, https://doi.org/10.1086/427996.
- [5] N. Bossuet-Greif, J. Vignard, F. Taieb, G. Mirey, D. Dubois, C. Petit, E. Oswald, J.P. Nougayrède, The colibactin genotoxin generates DNA interstrand cross-links in infected cells, MBio 9 (2018) e02393–17, https://doi.org/10.1128/mBio.02393-17.
- [6] M.R. Wilson, Y. Jiang, P.W. Villalta, A. Stornetta, P.D. Boudreau, A. Carrá, C.A. Brennan, E. Chun, L. Ngo, L.D. Samson, B.P. Engelward, W.S. Garrett, S. Balbo, E.P. Balskus, The human gut bacterial genotoxin colibactin alkylates DNA, Science 363 (2019), https://doi.org/10.1126/science.aar/785.
- [7] T. Secher, A. Samba-Louaka, E. Oswald, J.P. Nougayrède, *Escherichia coli* producing colibactin triggers premature and transmissible senescence in mammalian cells, PLoS One 8 (2013) e77157, https://doi.org/10.1371/journal.pone.0077157.
- [8] J.C. Arthur, E. Perez-Chanona, M. Mühlbauer, S. Tomkovich, J.M. Uronis, T.-J. Fan, B.J. Campbell, T. Abujamel, B. Dogan, A.B. Rogers, J.M. Rhodes, A. Stintzi, K.W. Simpson, J.J. Hansen, T.O. Keku, A.A. Fodor, C. Jobin, Intestinal inflammation targets cancer-inducing activity of the microbiota, Science 338 (2012) 120–123, https://doi.org/10.1126/science.1224820.
- [9] G. Dalmasso, A. Cougnoux, J. Delmas, A. Darfeuille-Michaud, R. Bonnet, The bacterial genotoxin colibactin promotes colon tumor growth by modifying the tumor microenvironment, Gut Microbes 5 (2014) 675–680, https://doi.org/10.4161/ 19490976.2014.969989.
- [10] A. Cougnoux, G. Dalmasso, R. Martinez, E. Buc, J. Delmas, L. Gibold, P. Sauvanet, C. Darcha, P. Déchelotte, M. Bonnet, D. Pezet, H. Wodrich, A. Darfeuille-Michaud, R, Bonnet. Bacterial genotoxin colibactin promotes colon tumour growth by inducing a senescence-associated secretory phenotype, Gut 63 (2014) 1932–1942, https://doi.org/10.1136/gutjnl-2013-305257.
- [11] C.M. Dejea, P. Fathi, J.M. Craig, A. Boleij, R. Taddese, A.L. Geis, X. Wu, C.E. DeStefano Shields, E.M. Hechenbleikner, D.L. Huso, R.A. Anders, F.M. Giardiello, E.C. Wick, H. Wang, S. Wu, D.M. Pardoll, F. Housseau, C.L. Sears, Patients with familial adenomatous polyposis harbor colonic biofilms containing tumorigenic bacteria, Science 359 (2018) 592–597, https://doi.org/10.1126/ science.aah3648.
- [12] D. Dubois, O. Baron, A. Cougnoux, J. Delmas, N. Pradel, M. Boury, B. Bouchon, M.A. Bringer, J.P. Nougayrède, E. Oswald, R. Bonnet, ClbP is a prototype of a

peptidase subgroup involved in biosynthesis of nonribosomal peptides, J. Biol. Chem. 286 (2011) 35562–35570, https://doi.org/10.1074/jbc.M111.221960.

- [13] C.A. Brotherton, E.P. Balskus, A prodrug resistance mechanism is involved in colibactin biosynthesis and cytotoxicity, J. Am. Chem. Soc. 135 (2011) 3359–3362, https://doi.org/10.1021/ja312154m.
- [14] N. Bossuet-Greif, D. Dubois, C. Petit, S. Tronnet, P. Martin, R. Bonnet, E. Oswald, J.P. Nougayrède, *Escherichia coli* ClbS is a colibactin resistance protein, Mol. Microbiol. 99 (2016) 897–908, https://doi.org/10.1111/mmi.13272.
- [15] P. Tripathi, E.E. Shine, A.R. Healy, C.S. Kim, S.B. Herzon, S.D. Bruner, J.M. Crawford, ClbS is a cyclopropane hydrolase that confers colibactin resistance, J. Am. Chem. Soc. 139 (2017) 17719–17722, https://doi.org/10.1021/jacs. 7b09971.
- [16] M. Črnigoj, Z. Podlesek, M. Budič, D. Žgur Bertok, The *Escherichia coli* uropathogenic-specific-protein-associated immunity protein 3 (Imu3) has nucleic acid binding activity, BMC Microbiol. 14 (2014) 16, https://doi.org/10.1186/1471-2180-14-16.
- [17] D. Nipič, Z. Podlesek, M. Budič, M. Črnigoj, D. Žgur Bertok, *Escherichia coli* uropathogenic-specific protein, Usp, is a bacteriocin-like genotoxin, J. Infect. Dis. 208 (2013) 1545–1552, https://doi.org/10.1093/infdis/jit480.
- [18] M. Ronen, R. Rosenberg, B.I. Shraiman, U. Alon, Assigning numbers to the arrows: parameterizing a gene regulation network by using accurate expression kinetics, Proc. Natl. Acad. Sci. 16 (2002) 10555–10560, https://doi.org/10.1073/pnas. 152046799.
- [19] M. Budič, M. Rijavec, Ž. Petkovšek, D. Žgur Bertok, *Escherichia coli* bacteriocins: antimicrobial efficacy and prevalence among isolates from patients with bacteraemia, PLoS One 6 (2011) e28769, https://doi.org/10.1371/journal.pone. 0028769.
- [20] P.M. van Wezenbeek, T.J. Hulsebos, J.G. Schoenmakers, Nucleotide sequence of the filamentous bacteriophage M13 DNA genome: comparison with phage fd, Gene 11 (1980) 129–148, https://doi.org/10.1016/0378-1119(80)90093-1.
- [21] J.R. Lakowicz, Quenching of fluorescence, Principles of Fluorescence Spectroscopy, Springer, Boston, MA, 2006, pp. 277–330, https://doi.org/10.1007/978-0-387-46312-4_8.
- [22] R. D'Ari, O. Huisman, Novel mechanism of cell division inhibition associated with the SOS response in *Escherichia coli*, J. Bacteriol. 156 (1983) 243–250 https:// www.ncbi.nlm.nih.gov/pmc/articles/PMC215076/#reference-sec.
- [23] J.G. Bird, S. Sharma, S.C. Roshwalb, J.R. Hoskins, S. Wickner, Functional analysis of CbpA, a DnaJ homolog and nucleoid-associated DNA-binding protein, J. Biol. Chem. 281 (2006) 34349–34356, https://doi.org/10.1074/jbc.M603365200.
- S. Cosgriff, K. Chintakayala, Y.T. Chim, X. Chen, S. Allen, A.L. Lovering, D.C. Grainger, Dimerization and DNA-dependent aggregation of the *Escherichia coli* nucleoid protein and chaperone CbpA, Mol. Microbiol. 77 (2010) 1289–1300, https://doi.org/10.1111/j.1365-2958.2010.07292.x.
 M. Almirón, A.J. Link, D. Furlong, R. Kolter, A novel DNA-binding protein with
- [25] M. Almirón, A.J. Link, D. Furlong, R. Kolter, A novel DNA-binding protein with regulatory and protective roles in starved *Escherichia coli*, Genes Dev. 6 (1992) 2646–2654, https://doi.org/10.1101/gad.6.12b.2646.
- [26] A. Martinez, R. Kolter, Protection of DNA during oxidative stress by the nonspecific DNA-binding protein Dps, J. Bacteriol. 179 (1997) 5188–5194, https://doi.org/10. 1128/jb.179.16.5188-5194.1997.
- [27] P. Setlow, Mechanisms which contribute to the long-term survival of spores of Bacillus species, J. Appl. Bacteriol. 76 (1994) 49S–60S.
- [28] W.L. Nicholson, N. Munakata, G. Horneck, H.J. Melosh, P. Setlow, Resistance of Bacillus endospores to extreme terrestrial and extraterrestrial environments, Microbiol. Mol. Biol. Rev. 64 (2000) 548–572.