High-protein diets for weight management: Interactions with the intestinal microbiota and consequences for gut health. A position paper by the my new gut study group

Francois F. Blachier, Martin Beaumont, Kevin Joseph Portune, Nils Steuer, Annaig Lan, Marc Audebert, Nadezda N. Khodorova, Mireille M. Andriamihaja, Gheorghe Airinei, Robert Benamouzig, et al.

► To cite this version:

Francois F. Blachier, Martin Beaumont, Kevin Joseph Portune, Nils Steuer, Annaig Lan, et al.. High-protein diets for weight management: Interactions with the intestinal microbiota and consequences for gut health. A position paper by the my new gut study group. *Clinical Nutrition*, 2019, 38 (3), pp.1012-1022. 10.1016/j.clnu.2018.09.016 . hal-02618730

HAL Id: hal-02618730

<https://hal.inrae.fr/hal-02618730>

Submitted on 22 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution - NonCommercial| 4.0 International License

High-protein diets for weight management: Interactions with the intestinal microbiota and consequences for gut health. A position paper by the My New Gut study group

François Blachier^{a*}, Martin Beaumont^a, Kevin Joseph Portune^b, Nils Steuer^c, Annaïg Lan^a, Marc Audebert^d, Nadezda Khodorova^a, Mireille Andriamihaja^a, Gheorghe Airinei^c, Robert Benamouzig^c, Anne-Marie Davila^a, Lucie Armand^a, Simone Rampelli^e, Patrizia Brigidi^e, Daniel Tomé^a, Sandrine Paule Claus^f, Yolanda Sanz^b

^aUMR PNCA, AgroParisTech, INRA, Université Paris-Saclay, Paris, France

^bMicrobial Ecology, Nutrition and Health Research Unit, Institute of Agronomy and Food Technology, Spanish National Research Council, Valencia, Spain

^cDepartment of Gastroenterology, Avicenne Hospital, Assistance Publique-Hôpitaux de Paris, Bobigny, France

^dToxalim, Research Centre in Food Toxicology, INRA, Toulouse, France

^eDepartment of Pharmacy and Biotechnology, University of Bologna, Bologna, Italy

^fDepartment of Food Nutritional Sciences, University of Reading, Reading, United Kingdom

*Corresponding author. UMR PNCA, 16 rue Claude Bernard, 75005 Paris, France.

E-mail address: francois.blachier@agroparistech.fr

Summary

Background and aim: This review examines to what extent high-protein diets (HPD), which may favor body weight loss and improve metabolic outcomes in overweight and obese individuals, may also impact the gut environment, shaping the microbiota and the host-microbe (co)metabolic pathways and products, possibly affecting large intestine mucosa homeostasis.

Methods: PubMed-referenced publications were analyzed with an emphasis on dietary intervention studies involving human volunteers in order to clarify the beneficial vs. deleterious effects of HPD in terms of both metabolic and gut-related health parameters; taking into account the interactions with the gut microbiota.

Results: HPD generally decrease body weight and improve blood metabolic parameters, but also modify the fecal and urinary contents in various bacterial metabolites and co-metabolites. The effects of HPD on the intestinal microbiota composition appear rather heterogeneous depending on the type of dietary intervention. Recently, HPD consumption was shown to modify the expression of genes playing key roles in homeostatic processes in the rectal mucosa, without evidence of intestinal inflammation. Importantly, the effects of HPD on the gut were dependent on the protein source (i.e. from plant or animal sources), a result which should be considered for further investigations.

Conclusion: Although HPD appear to be efficient for weight loss, the effects of HPD on microbiota-derived metabolites and gene expression in the gut raise new questions on the impact of HPD on the large intestine mucosa homeostasis leading the authors to recommend some caution regarding the utilization of HPD, notably in a recurrent and/or long-term ways.

Keywords: High-protein diet, microbiota, bacterial metabolites and co-metabolites, large intestine mucosa

1. Introduction

In a context of a high proportion of overweight and obese individuals, notably in populations from Europe and the USA (1), numerous different types of weight-loss diets are currently proposed and consumed (2). Among them, high-protein diets (HPD), which represent a heterogeneous group of diets with different composition (3), are all characterized by a higher proportion of protein (25 – 30 % of total energy intake) among the two other dietary macronutrients (i.e. carbohydrates and fat) when compared with the usual macronutrient proportion. These HPD are used by millions of individuals around the world for weight-loss (4). One of the main rationales for the consumption of HPD is that it is generally recognized that, on a basis of equal energy content, protein is more satiating than carbohydrates and fats (5). Considering that weight gain is primarily observed when energy recovered from food is superior to energy expenditure, notably in relationship with physical exercise (6), HPD, by reducing dietary energy intake, is likely to help, at least transiently, in the process of body weight reduction (7).

However, there is presently no definition of the maximal amount of dietary protein that can be consumed without short- and/or long-term metabolic and physiopathological side effects. Indeed, if the benefits of decreased body weight in overweight and obese individuals in terms of metabolic and general health outcomes appear obvious based on numerous studies (8), then the interest of HPD consumption for such outcomes must be confronted with possible undesirable effects upon different tissues and organs in a beneficial over deleterious ratio perspective. For instance, it is well known that HPD are contraindicated in individuals with chronic kidney diseases or at risk for such diseases, as HPD may accelerate kidney dysfunctions (9, 10). Regarding the impact of HPD on gut health, this remains an emerging but important topic.

The aim of the present review is to present the available evidence, including recent data obtained in the MyNewGut European research project, in order to balance the advantages of HPD for weight loss and metabolic health against the potential risks of such unbalanced diets focusing on the gut ecosystem homeostasis. As a matter of fact, there are indications from clinical and experimental studies that dietary changes may modify the large intestine luminal environment with a potential impact on the colonic mucosa (11).

PubMed-referenced publications were analyzed using the following terms in combination: [high-protein diet OR dietary protein OR protein source] / [intestinal microbiota OR bacterial metabolites OR co-metabolites] / [large intestine OR colon OR rectum] / [weight-loss OR overweight OR obesity]. Among the numerous papers found, priority was given for references related to dietary intervention studies with human volunteers, notably those reporting consequences in terms of intestinal physiology and physiopathology.

This review is part of a series of position paper of the MyNewGut project aiming at providing recommendations for dietary guidelines based on project results and the latest advantages in the field regarding insights gained in the role of the gut microbiome, as described in the introductory paper (12).

2. High-protein diet, weight loss, and metabolic effects

2.1 High-protein diet and weight loss

HPD can be defined in regards to the absolute amount of dietary protein (in grams) consumed per day, or to the proportion of dietary protein in the total energy intake; or to the amount of dietary protein per unit of body weight. A useful reference can be found on the recommended daily amount of dietary protein which has been determined to be equal to 0.83 g of protein per kg body weight per day (13), thus representing 58.1 g dietary protein per day for an individual weighting 70 kg. As a matter of fact, mean dietary protein consumption is largely above these recommended value for instance in France since it averages 87.3 g/day (average value for men and women) (14), and in the USA where it averages 82.8 g/day taking into account men and women dietary protein consumption (15), thus representing approximately 1.5 fold the recommended daily amount of protein. HPD can represent as much as 5 fold higher than the recommended daily amount (4), but it is generally considered that diets containing at least 25-30% of energy in the form of protein are HPD (16). As a matter of comparison, in France, 16.8% of the dietary energy comes from protein in typical diets (14). Incidentally, HPD are also largely consumed by athletes who wish to increase their muscle mass and performance, but this

aspect is out of the scope of the present review and will not be described here, although the readers are referred to excellent reviews on that topic (17, 18).

Two main types of controlled clinical intervention studies with HPD have been performed. The first one is the “*ad libitum*” studies in which volunteers consume the amount of HPD or control normoproteic diet (NPD) until they naturally stop their food consumption. In these studies, due to the satiating effects of HPD, volunteers on HPD generally eat less food than the control NPD subjects, and consequently significantly decrease their body weight compared to the body weight measured at the onset of the dietary intervention. In the study of Weigle et al. (19), HPD given *ad libitum* for 2 weeks resulted in a decrease of body weight. Johnstone et al. (20) also reported reduction of food intake and body weight following 4-week-consumption of HPD. *Ad libitum* consumption of HPD for 6 months resulted in a marked decrease of body fat when compared with individuals receiving a NPD (21). In a study on weight loss maintenance after dietary energy restriction, it has been shown that HPD, when given for 12 weeks (22) or 12 months (23), is efficient for weight control. However, in the “real life” condition, a vast majority of individuals, after initial body weight reduction, recover their initial body weight in the long term (24), leading possibly to recurrent episodes of weight-loss HPD consumption. A study using *ad libitum* HPD has shown that meat-based HPD is not more efficient for body weight decrease than protein from plant origin (25).

The second type of HPD intervention studies consists of increasing the proportion of protein in the diet compared to the control, but in that case, the amount of energy consumed between groups is maintained constant. This is generally done by decreasing the relative proportion of another macronutrient in the diet, namely carbohydrates or fats. In that kind of isocaloric clinical protocol, the studies generally found no or little effect of such diets for body weight reduction (16, 26) corresponding to the view that the amount of dietary energy intake, at a constant level of physical exercise, is a major parameter for fixing the evolution of body weight for one given individual.

A third type of studies related to the use of HPD in obese patients are those related to the use of such diet for maintaining the lean mass in malnourished obese patients. Since we will not develop this aspect in our review, the readers are referred to a recent review paper on that topic (27).

2.2 High-protein diet and metabolic parameters

The interpretation of the effect of HPD on metabolic parameters can be somewhat complicated. For instance, if a HPD is given to overweight individuals in an “*ad libitum*” protocol, it will be difficult to determine what part the increased proportion of protein in the diet plays in the normalization of metabolic parameters in comparison with the part played by the decrease of energy intake due to the satiating effect of HPD and the resultant decrease of body weight. In overweight and obese individuals, marked decrease of body weight, whatever the cause, allows the normalization of metabolic parameters (8, 28).

In protocols in which the experimental diets are isocaloric, the HPD, as said above, are necessarily decreased in another macronutrient, thus rendering it difficult to attribute the effects of HPD solely to the increased content of protein and/or to the reduced amount of the other macronutrient. In a recent randomized, parallel, double-blind controlled study in which the HPD (using milk casein or soy protein as supplements) were given to volunteers for 3 weeks, no significant changes on any of the biochemical and anthropometric parameters were measured in blood in fasting conditions when compared with control subjects receiving a normoproteic isocaloric diet. A notable exception to this lack of change in parameters was observed for systolic blood pressure, which was decreased in the group of volunteers receiving the soy protein supplementation; an effect that was likely due to the presence of protein-associated isoflavones in the protein extract (26). Thus, under condition of equal energy consumption, HPD appear to exert no short-term sizeable effect on the metabolic and anthropometric parameters.

3. High-protein diet and changes in the gut ecosystem

The process of protein digestion in the small intestine is a very efficient process with digestibility usually ranging from 89 to 95%, depending on the nature of the protein (29, 30). Generally speaking, proteins from animal sources are overall more digestible than proteins from plant sources (31). Some sources of protein, for instance rapeseed protein, are known for their lower digestibility (32). In addition, food cooking (33, 34) and food matrix structure (35) can impact protein digestibility. Importantly, and as a result of incomplete digestion in the small intestine, a residual amount of

undigested protein and peptides, together with individual amino acids are transferred through the ileo-caecal junction in the large intestine (36). Based on a regular western diet, it has been determined that approximately 12 g of protein and peptide from both dietary and endogenous origin escape digestion in the small intestine, thus reaching the colonic lumen (37). This amount of nitrogenous material is increased nearly proportionally when the amount of dietary protein increases (29). From studies evaluating the proportion of dietary and endogenous protein which escape digestion and move from the ileum to the large intestine, it has been determined that the majority of the ileal nitrogen is originating from endogenous losses (1-2 g/day), while the nitrogen from dietary origin represents 0.7-1.2 g/day (36). The results obtained in animal models suggest that the part ascribed to endogenous protein is not vastly different according to the amount of protein consumed (38). Since the large intestine luminal content is characterized by a much more abundant microbiota than what is measured in the small intestine (39), and also by a much slower transit time (40), the proteins and peptides which enter the large intestinal luminal content undergo the catalytic action of bacterial proteases and peptidases which release sequentially shorter peptides and amino acids (41). The large intestinal epithelium, in contrast with the small intestinal epithelium which is very efficient for oligopeptide and amino acid absorption, is not believed to transfer any significant amount of amino acids from the lumen to the bloodstream, except in the neonatal period (42, 43). Therefore, protein and peptide-derived amino acids are metabolized by the large intestinal microbiota which use them for protein synthesis and catabolic pathways with the production of numerous intermediates and final metabolites (44); a net amount of these latter being able to accumulate within the luminal content (Figure 1). This process of protein degradation is more active in the distal part than in the proximal part of the large intestine (45). In the case of HPD consumption, the increased transfer of nitrogenous compounds in the large intestine is liable to modify the microbiota composition, and/or to change the microbiota diversity, and/or its metabolic activity, and finally to change the production of bacterial metabolites with possible consequences for the large intestinal mucosa metabolism, physiology and health (46-50) as described below.

3.1 High-protein diets and intestinal microbiota composition

Relatively few human intervention studies have examined the short-term (less than 4 weeks) effects of HPD on the gut microbiota composition (Table 1). Two main factors preclude direct comparison between the studies presented in Table 1: (i) differences in energy intake (e.g. calorie restriction) and (ii) differences in fiber intake. These two parameters are known to have a profound influence on the gut microbiota composition and should therefore be considered as important potential confounding factors with the effects of dietary protein intake. Moreover, there are large variations between the studies in terms of methods used to analyze the composition of the gut microbiota. With these limitations in mind, it is still possible to propose some general conclusion regarding the effects of dietary protein intake on the gut microbiota.

Two of the studies in Table 1 used HPD without modification of dietary fiber and energy intake. (26, 45). Using 16S rDNA sequencing for fecal or rectal biopsy samples, and denaturing gradient gel electrophoresis (DGGE) for fecal samples, respectively, these two studies did not detect changes in the gut microbiota composition after the HPD (Table 1). In a study by David et al. (51) a diet containing dietary protein from animal origin containing almost no fibers was given *ad libitum* for 5 days. This dietary intervention resulted in almost doubling the protein intake (i.e. 30.1% of energy intake) as compared to the protein consumption at the onset of intervention, and was found to impact the microbiota composition by increasing the abundance of bile-tolerant microorganisms (*Alistipes*, *Bilophila*, and *Bacteroides*), and by decreasing the levels of Firmicutes that metabolize plant polysaccharides (*Roseburia*, *Eubacterium rectale*, and *Ruminococcus*). Such a HPD was found to change the microbiota β -diversity within 2 days. However, this latter effect appeared to be transient, as the β -diversity returned to the initial configuration within 2 days after the end of the intervention (51). However, these changes could not be attributed solely to the level of protein intake since there was considerable concomitant modification of fat intake (in addition to fiber intake) in this study.

The other studies presented in Table 1 used HPD with caloric restriction that resulted in weight-loss. Two of them (different analysis of the same samples), showed that the HPD induced an alteration of the gut microbiota composition with a decreased abundance of presumed beneficial bacteria such as *Bifidobacterium* or *Rosburia/Eubacterium rectale* (52, 53) However, both resistant starch and total

carbohydrates were also lower in the high-protein/weight loss diet compared to the maintenance diet (52). This is an important point to consider as resistant starch has been positively associated with the abundance of *Bifidobacterium* and *Eubacterium spp* (54, 55).; and a reduction in carbohydrates led to decreases in both genera (56). In another study, a weight-loss HPD combined with an increase in fiber intake also induced a decrease in *Eubacterium rectale* but increased bacterial gene richness in individuals with low gene counts together with an increase abundance of bacteria considered protective such as *Faecalibacterium prausnitzii* and *Roseburi* (57). Lastly, two other studies using weight-loss HPD combined with a low fiber intake observed a decrease in the total bacterial biomass and in the abundance of *Bifidobacterium* and *Rosburia/Eubacterium rectale* (56, 58). Overall, the studies presented in Table 1 show that HPD have a limited effect on the gut microbiota composition when they are not associated with calorie restriction or with a modification of fiber intake. This conclusion may also be connected to the observed relatively little changes in the microbiota composition according to the diet when compared with the inter-individual variations (< 10%) (53).

3.2 High-protein diets and impact on gut mucosa: potential role of bacterial metabolites

The mixture of bacterial metabolites in the intestinal content is complex (59) and far from being fully characterized. Among these compounds, numerous metabolites are produced by the intestinal microbiota from amino acid substrates (60). The concentrations of these metabolites are usually measured in the feces, which are related to the concentrations of the luminal content within the most distal part of the large intestine, namely in the rectum. These metabolite concentrations depend on the bacterial production from the available substrates, on the bacterial composition and overall metabolic activity, on the absorption through the large intestinal epithelium, and on the transit time (61). (Figure 1). Other parameters may influence the concentrations of the different forms of the bacterial metabolites within the large intestine content. For instance, the luminal pH, which will result from the overall acid/base balance in this compartment, will in turn determine the ratio of the different non-ionized and ionized forms of ionic bacterial metabolites (62), which will affect their uptake from the luminal content to the colonocyte intracellular content. In addition, the situation is complicated by the

fact that some bacterial metabolites (for instance hydrogen sulfide) can bind to fecal components, thus reducing the concentration of free (unbound) metabolites presumed to act on the epithelial cells (Figure 2) (63). We present below the effects of HPD on bacterial metabolites and their main effects observed in Humans and experimental animal models but the reader is referred to another recent review for more exhaustive description of the metabolites produced by the microbiota from amino acids (41).

3.2.1 Effects of high-protein diets on the fecal composition and effects of individual bacterial metabolites on colonic epithelial cells

Several intervention studies in humans have shown that HPD with different sources of dietary protein induce a shift from carbohydrate to protein degradation by the gut microbiota (26, 58, 59, 64), with an alteration of numerous bacterial metabolite concentrations in feces, thus indicating changes in the luminal environment of the colonic epithelial cells. In contrast with the high variability described above between human intervention studies regarding the effects of HPD on microbiota composition (Table 1), the effects of HPD on bacterial metabolites are more homogeneous despite differences in experimental design (Table 2). This observation emphasizes the importance of substrate availability, namely amino acids in our case, rather than taxonomic composition of the microbiota for determining the metabolic output in the large intestine. This could also be due to redundancy of functions and metabolic pathways in the microbiome, the collective genome of the microbiota (65).

Most of the studies in Table 2 reported that HPD consumption induced an increase in amino acid-derived short-chain fatty acids (SCFA) such as isobutyrate, isovalerate, and 2-methylbutyrate (26, 49, 58). In contrast, a decrease in the SCFA butyrate was consistently found after HPD consumption (26, 51, 56, 58) albeit several of these studies included decreases in fiber content among the HPD. However, in a recent study by Beaumont et al. (26) volunteers from the HPD and control groups consumed a similar amount of dietary fibers and energy than the NPD group thus suggesting that the reduction of fecal butyrate concentration in HPD can be attributed primarily to the amount of protein in the diet. As butyrate is well-known as a major oxidative substrate and a regulator of histone

acetylation, and thus of gene transcription in human colonocytes (66, 67), the measured decrease in its fecal concentration after HPD is presumably detrimental for the rectal mucosa homeostasis.

Two studies in volunteers receiving a HPD found a marked increase in fecal ammonia concentrations (59, 64), while two others did not (26, 58), likely due to the different experimental protocols. Also, HPD were found to increase the concentrations of several S-containing metabolites (59, 68). For most of these metabolites, there is surprisingly no indication on the impact of such changes on the colonic/rectal epithelium renewal and functions. However, from *in vitro* studies with human or rodent colonocytes, there are indications that several amino acid-derived bacterial metabolites including hydrogen sulfide (H₂S), ammonia and *p*-cresol act as metabolic troublemakers towards colonocyte mitochondrial energy metabolism within the range of concentrations that are measured in the colonic content or in feces (69, 70).

In contrast, some bacterial metabolites derived from amino acids were found to exert beneficial effects on the intestinal epithelial barrier (reviewed in 11). For instance, indole which is produced from L-tryptophan has been shown to increase epithelial cell tight-junction resistance as will be detailed in the part 3.3. Another bacterial metabolites derived from tryptophan, namely indole propionic acid, has been shown recently to be efficient for decreasing the intestinal permeability in rodents (71). Thus, in order to document the beneficial versus deleterious effects of the mixture of bacterial metabolites contained within the intestinal content, it is clearly necessary to take into account the fact that these contents contain compounds with both positive and negative effects on the intestinal mucosa.

3.2.2 Genotoxic and cytotoxic potential of fecal water recovered after high-protein diet consumption

In order to get information on the possible overall cytotoxic and genotoxic potential of fecal water-soluble components after controlled dietary intervention, it is feasible to prepare the so-called “fecal water” samples by diluting and homogenizing fecal samples in aqueous medium, and test the supernatant on human colonocytes. Although fecal water samples do not contain all the luminal compounds and dilute the bacterial metabolites, fecal water toxicity has been proposed to represent a potential biomarker for intestinal disease risk (72). When an isocaloric HPD was given for 2 weeks to healthy human subjects in a crossover design, the mixture of water-soluble components recovered

from the feces shown no increased genotoxicity or cytotoxicity potential towards human colonocytes when compared to the NPD (45). Similarly, in a study by Benassi-Evans et al. (73), the authors performed a nutritional intervention with HPD during 52 weeks using a parallel design with overweight and obese volunteers. They found that the fecal water recovered from individuals consuming HPD was not more genotoxic than ones recovered from control volunteers consuming isocaloric NPD. In accordance with the results presented above, in a study by Beaumont et al. (26), supplementation of the diet with either casein or soy protein for 3 weeks, did not result in higher cytotoxic potential of the fecal water when compared with the results obtained from isocaloric NPD volunteers. Thus, collectively, the available data indicate that the fecal water samples recovered from volunteers consuming HPD in short- and medium terms show no increased genotoxic and cytotoxic potential *in vitro* towards colonic epithelial cells than samples recovered from control NPD.

3.3 High-protein diet and urinary metabolome

Urinary metabolomic analysis is useful in order to identify the bacterial metabolites and cometabolites (produced by the microbiota and metabolized by the host) which have been produced by the gut microbiota, absorbed from the lumen to the bloodstream through the intestinal epithelium (with or without metabolism in colonocytes), possibly further metabolized by the host in the liver or other organs outside the splanchnic area, and finally excreted in the urine where they accumulate (Figure 3). For instance, HPD ingestion results in the increased urinary excretion of the bacterial metabolite phenol (64). This is of interest as phenol has been shown to act as a cytotoxic compound towards colonocytes (74); and as impaired phenol detoxification has been associated with ulcerative colitis (75).

In addition, the cometabolite *p*-cresyl sulfate is produced in the colon mucosa and the liver from the bacterial metabolite *p*-cresol, which itself is produced by the microbiota from the amino acid L-tyrosine (76). Urinary concentration of *p*-cresyl sulfate has been repetitively found to be increased after HPD consumption (26, 59, 77) when compared with control NPD (Table 2). Since *p*-cresol has been shown to inhibit colonocyte oxygen consumption, and to be genotoxic towards colonocytes (70), *p*-cresyl sulfate synthesis has been hypothesized to correspond in colonic epithelial cells to a

detoxifying metabolic pathway for this bacterial metabolite. This possibility has been challenged by the fact that *p*-cresyl sulfate displayed pro-inflammatory and cytotoxic effects on renal tubular epithelial cells (78, 79), and that serum *p*-cresyl sulfate level may help in predicting progression of chronic kidney disease (80, 81).

In a study by Beaumont et al. (26), the relative concentration of another urinary cometabolite, namely indoxyl sulfate, increased after HPD (Table 2). Since indole, the precursor for the synthesis of indoxyl sulfate in the liver, has been shown to contribute to the maintenance of the colonic barrier function (82, 83) and to alleviate hepatic inflammation (84), this bacterial metabolite can be considered as beneficial for the host. However, in order to establish the beneficial vs. deleterious effects of indole on the colon epithelium, it is important to consider that this bacterial metabolite activates the aryl hydrocarbon receptor (AhR)-mediated transcription of Cyp 1a1 and Cyp 1b1 in human colonocytes (85, 86). These two enzymes belongs to the cytochrome P450 family which, apart from their role in the deactivation of deleterious compounds and xenobiotics, can catalyse the bioactivation of procarcinogen compounds into carcinogens (87-89). In addition, indoxyl sulfate is suspected to act as a uremic toxin contributing to renal disease progression (90-92).

Thus, the analysis of the urinary metabolome gives important information regarding the exposure of the intestinal mucosa to bacterial metabolites (Figure 3), even if the results obtained emphasizes the difficulty to predict how changes of a complex mixture of bacterial metabolites will impact the colonic/rectal mucosa according to the time of exposition and respective concentrations.

3.4 High-protein diets and gut mucosa inflammation

Although the results of epidemiological studies regarding the association between HPD consumption and risk of inflammatory bowel diseases (IBD) are heterogeneous (93), two studies have shown that a high amount of animal protein intake is associated with increased inflammatory bowel disease incidence and relapse (94, 95). However, short-term supplementation (3 weeks) with casein or soy protein, did not show any sign of rectal mucosal inflammation based on the measurement of pro-

inflammatory cytokines in rectal biopsies, and on the fecal concentrations of calprotectin and secreted IgA, when compared with an isocaloric NPD (26).

Participation of some bacterial metabolites on the process of mucosal inflammation in pre-disposed subjects may be related to a reduced capacity of the mucosa for deleterious metabolite detoxification. For instance, it has been reported that impaired H₂S detoxification in intestinal mucosa is associated with Crohn's disease (96) and ulcerative colitis (97). These results are important to be taken into account, knowing that increased protein consumption is correlated with increased H₂S fecal excretion in volunteers (68), and that excessive luminal H₂S decreases colonocyte respiration and increases the expression of several genes involved in IBD in a rodent model (98). It can therefore be predicted that there might be differences between individuals in terms of mucosal response to HPD according to individual detoxification capacities.

3.5 High-protein diets and gene expression in gut mucosa

The first experimental evidence using transcriptomic analysis which has shown that casein-containing HPD can modify gene expression in the colonic mucosa were obtained in the rat model by Mu et al. (99) using a 6-week- dietary intervention protocol with isocaloric experimental (HPD) and control (NPD) diets. Beaumont et al. (100) used a 2 week-intervention protocol with whole milk protein-containing HPD in the rat model to demonstrate that HPD down-regulates colonic epithelial cell gene expression notably in relationship with cell metabolism, NF-κB signaling, DNA repair, glutathione metabolism and cellular adhesion, when compared with gene expression in colonocytes recovered from isocaloric NPD. In this latter study, the HPD was found to up-regulate the expression of genes related to cell proliferation and chemical barrier function. These animal studies allow to establish the new proof of concept according to which increasing the amount of protein in the diet will result in a modification of gene expression in the colonic mucosa, and more specifically in the colonic epithelial cells. Further, a randomized controlled study with overweight volunteers reported that 3 week-dietary supplementation with either casein or soy protein resulted in small amplitude changes in the

expression of numerous genes in the rectal mucosa, notably for genes involved in homeostatic processes such as cell cycle or cell death (26).

3.6 The effects of high-protein diets on the fecal and urinary metabolome and on the large intestine mucosa according to different protein sources

It can be hypothesized that the source of protein used in the HPD studies may represent an important parameter for modulating the colonic epithelium luminal environment and gene expression in the rectal mucosa. First, as presented above, different dietary proteins displayed different digestibility characteristics. Second, the differences in the amino acid composition between proteins provide the intestinal microbiota with different amounts of individual amino acids as substrates for the microbiota metabolic activity, thus potentially resulting in different fecal bacterial metabolite compositions and urinary bacterial/host cometabolites in the urine. Up to now, this hypothesis has been little explored but one recent study reported that when the habitual diet is supplemented with either milk casein or soy protein, differences are observed in the fecal and urinary metabolome, with such differences coinciding with changes in gene expression in the rectal mucosa (26). Indeed, in the case of supplementation with casein, when compared with the isocaloric NPD group, the feces were characterized by increased relative concentration of 2-methylbutyrate; while in the case of supplementation with soy protein, an increase of this bacterial metabolite was also measured but together with an increase of valerate, tyramine, and phenylacetate. Regarding the urinary metabolome, casein supplementation resulted in increased urea, isobutyrate, 3-hydroxybutyrate, 3-hydroxyisovalerate, *p*-cresyl sulfate, phenylacetylglutamine and indoxylsulfate relative concentration; while supplementation with soy protein resulted in an increased of the same metabolites but not of the uremic toxin *p*-cresyl sulfate, the co-metabolite produced from *p*-cresol (79).

More importantly, casein and soy protein HPD were found to differentially modify the expression of genes playing key roles in the maintenance of the rectal mucosa homeostasis maintenance in general, and in colonic health (gastrointestinal diseases and cancer) in particular. At the cellular level, the casein diet was specifically associated with increased expression of genes related to extracellular matrix, cell adhesion, and mucus production; while the soy protein diet was specifically associated

with modification of the expression of genes associated with oxidative stress and detoxification processes. Expression of other genes associated with cellular processes like apoptosis, cell cycle and proliferation, and cytoskeleton formation were modified by both casein and soy protein (26). To determine if such changes in gene expression impact the rectal epithelium renewal and functions, and/or if it corresponds to an adaptation towards a changing luminal environment, new experiments are required. Regarding this latter aspect, the fact that the expression of genes related to mucus production was solely increased in the rectal mucosa of volunteers after casein supplementation but not after soy protein supplementation, may indicate an adaptation of the rectal mucosa towards a more aggressive luminal environment following casein-based HPD consumption.

4. Conclusion and perspectives

Although it appears that HPD can help in diminishing the dietary intake, and thus favor weight loss, there are some results which raise new questions on the safety of their utilization. It must be recognized that, according to the available literature, there is no definitive evidence that such diets are deleterious for gut health in short- and medium- term intervention studies conducted so far. Indeed, as presented above, short-term consumption of HPD by itself neither increases the inflammation of the large intestinal mucosa, nor increases the *in vitro* genotoxicity and cytotoxicity of the mixture of compounds contained in the fecal water extracts in healthy subjects. However, HPD have been shown in a repetitive manner to decrease fecal butyrate concentrations. Since butyrate is generally considered as a fuel substrate and a regulator of gene expression in the rapidly renewing colonic epithelial cells, this decrease must be seen as potentially deleterious for the colonic mucosa homeostasis. The same remark can be made regarding the finding that HPD consumption results in increased exposition of the intestinal mucosa to *p*-cresol, a bacterial metabolite with genotoxic and metabolic troublemaker characteristics towards colonocytes (70). In addition, *p*-cresol is the precursor of *p*-cresyl sulfate, a cometabolite with reported cytotoxic activity towards renal cells (78, 79) (Figure 3). Conversely, there is evidence that HPD increases the exposure of the large intestine mucosa to indole, a bacterial metabolite considered as an important player in the maintenance of the epithelial barrier function. However, this positive effect of indole on the intestinal epithelium must be

counterbalanced by the suspicion that indoxyl sulfate, a cometabolite of indole produced in the liver, is also acting as a uremic toxin (90, 101) (Figure 3). Then, the different effects of bacterial metabolites and cometabolites on different cell types, either within the intestinal mucosa as detailed in the present paper, or at the periphery, makes it difficult to predict if one given compound in a mixture should be considered as overall beneficial or deleterious. The finding that an increased consumption of dietary protein modifies within 3 weeks the normal expression of genes known to be involved in processes related to the maintenance of the rectal mucosal homeostasis (26), represents an important new finding which should be taken into consideration before formulating any recommendation on HPD consumption.

Regarding the effects of amino acid-derived bacterial metabolites on metabolic parameters, recent data suggest that some of these metabolites might contribute to an improvement of some of these parameters. For instance, indole has been shown *in vitro* to modulate the secretion of the incretin glucagon-like peptide 1 (GLP-1) (102). Moreover, hydrogen sulfide produced by the gut microbiota has been shown to lower blood pressure in rats (103), to improve glucose metabolism, and to increase GLP-1 secretion in mice (104). Lastly, several neurotransmitters can be produced by the gut microbiota from amino acids (41), and it can be speculated that this may contribute to the dietary protein- induced satiety. Further studies, notably with larger groups of human volunteers, and of longer duration are needed to determine whether the potential effects of amino acid-derived bacterial metabolites, depending on the protein sources, could participate in the beneficial metabolic effects of HPD associated with body weight reduction.

5. Implications for dietary recommendation regarding high-protein diet consumption

Although body weight reduction associated with *ad libitum* HPD consumption in overweight and obese individuals is obviously associated with favorable outcomes, the data obtained principally from clinical trials with human volunteers, dietary intervention in animal models, and *in vitro* experiments with human colonic epithelial cells have shown that HPD modifies the luminal environment of the rectal epithelium and impacts gene expression in the mucosa. We therefore recommend caution in the utilization of HPD diets for body weight loss, taking into account the possible regain of body weight

after HPD consumption, which may lead to redundant and long-term utilization of HPD. Considering the most recent evidence showing that the effects of HPD on the gut depend on the protein source (i.e. from plant and animal sources), not only the quantity, but also the quality of dietary protein should be considered for further investigations and possibly for future dietary recommendations.

Acknowledgement

The MyNewGut project is financially supported by a grant from the EU 7th Framework Programme under Grant Agreement 613979. The EU is not liable for the content presented in this publication.

Conflict of interest

FB, MB, KJP, NS, AL, MA, NK, MA, GA, RB, AMD, LA, SR, PB, DT, SPC, and YS declare no competing interest in relation to this paper.

Legends of figures:

Figure 1. Schematic view of the fate of undigested proteins in case of High-Protein Diet (HPD) consumption.

HPD diet consumption increases the transfer of dietary proteins from the ileum to the large intestine. The proteases and peptidases of the microbiota release amino acids which can be incorporated in the bacterial proteins or lead to a multitude of metabolic end products, notably in the distal parts of the large intestine. Some of these metabolites are known to be transferred by the colonic epithelial cells from the luminal content to the portal bloodstream with or without prior metabolism in the colonocytes. The concentrations of bacterial metabolites in the lumen are the net result of production/utilization by the microbiota, and absorption through the colonic epithelium. The metabolites measured in the feces is a reflection of the metabolites present in the rectum.

Figure 2. Schematic view of the entry and metabolism of bacterial metabolites in the colonic epithelial cells.

Several bacterial metabolites in the luminal content can enter colonocytes by processes of diffusion or transport. Although some of them can be released as such in the bloodstream, several bacterial metabolites are known to undergo intracellular metabolism leading to the production of co-metabolites. Bacterial metabolites and co-metabolites can be released in the portal bloodstream and reach the liver and peripheral organs outside the splanchnic area. Finally, these compounds can accumulate in urine after glomerular filtration and/or tubular secretion by kidneys.

Figure 3. Schematic view of the impact of high-protein diet (HPD) consumption on the bacterial metabolite and co-metabolite concentrations in feces and urine.

Undigested proteins and peptides enter the large intestine and are metabolized by the microbiota which produce various metabolites from amino acids. Some of these metabolites are partly absorbed through the large intestine epithelium, while the residual amount of metabolites are excreted in the feces. Absorbed metabolites reach the liver where some of them undergo further metabolism. Co-metabolites and metabolites are finally excreted in the urine. HPD consumption results in measurable modifications of the concentration of bacterial metabolites in feces and urine. As indicated in the text, some compounds originating from the microbial metabolic activity (like butyrate and H₂S) are known to impact energy metabolism and gene expression in colonocytes, while some of them (like indole) are implicated in the maintenance of the epithelial barrier function. Some co-metabolites measured in urine (like indoxylsulfate and *p*-cresylsulfate) are suspected to act as uremic toxins.

Table 1: Effects of high-protein diet on intestinal microbiota composition. The main characteristics and findings from human intervention studies using high-protein diets are summarized. BMI: body mass index, DGGE: denaturing gradient gel electrophoresis, FISH: fluorescence in situ hybridization, % E: % of energy intake, g/d: grams/day. For carbohydrates and fat intake, the readers are referred to the original publications.

Study design	BMI	Duration	Protein intake	Protein source	Fiber intake	Calorie restricted	Method	Intestinal microbiota composition	Reference
n=12-13 Parallel	25 - 30	3 weeks	14 % E	Mixed	17.0 g/d	No	16S rDNA sequencing (feces and rectal biopsies)	Control diet	(26)
			34 % E	Mixed + casein	14.4 g/d	No		No detectable differences	
			31 % E	Mixed + soy protein	17.9 g/d	No		No detectable differences	
n=20 Cross-over	19 - 26	2 weeks	12 % E	Mixed	17.4 g/d	No	DGGE (feces)	No detectable differences	(45)
			15 % E	Mixed	16.3 g/d	No		Control diet	
			27 % E	Mixed	15.4 g/d	No		No detectable differences	
n = 10 Cross-over	19 - 32	5 days	10 % E	Plant protein	41.2 g/d	No	16S rDNA sequencing (feces)	↓ <i>Bilophila wadsworthia</i>	(51)
			16 % E	Mixed	21.1 g/d	No		Control diet	
			30 % E	Animal protein	0 g/d	No		↑ <i>Bilophila wadsworthia</i> , <i>Alistipes putredinis</i> ; ↓ <i>Bifidobacterium adolescentis</i> , <i>Roseburia faecis</i> , <i>Ruminococcus bromii</i>	
n=14 Parallel	28 - 51	3 weeks	103.3 g/d	Mixed	27.7 g/d	No	Phylogenetic (HITchip) microarray (feces)	Control diet	(53)
			144.1 g/d	Mixed	25.1 g/d	Yes		↓ <i>Bifidobacterium</i> , <i>Aerococcus</i> , <i>Granulicatella</i> , <i>Dialister</i> , <i>Papillibacter cinnamivorans</i> ; ↑ <i>Lactococcus</i> , <i>Bacteroides vulgatus</i> , <i>Anaerotruncus colihominis</i>	
n=14 Parallel	28 - 51	3 weeks	103.3 g/d	Mixed	27.7 g/d	No	16S rDNA sequencing, DGGE, qPCR (feces)	Control diet	(52)
			144.1 g/d	Mixed	25.1 g/d	Yes		↓ <i>Collinsella aerofaciens</i> , <i>Roseburia/Eubacterium rectale</i> ; ↑ <i>Oscillibacter valericigenes</i>	
n=49 Non-randomized	33 (mean)	6 weeks	19 % E	Mixed	14.5 g/d	No	Metagenomic sequencing (feces)	Control diet	(57)
			37 % E	Mixed	19.0 g/d	Yes		↓ <i>Eubacterium rectale</i> ; ↑ <i>Parabacteroides distasonis</i> , <i>Faecalibacterium prausnitzii</i> , <i>Bacteroides dorei</i> , <i>Parabacteroides merdae</i> , <i>Eubacterium eligens</i> , <i>Ruminococcus sp.</i> , <i>Roseburia hominis</i> , <i>Odoribacter splanchnicus</i> , <i>Subdoligranulum sp.</i> , Gene richness in low gene counts individuals	
n=17 Cross-over	30 - 49	28 days	13 % E	Mixed	21.9 g/d	No	FISH (feces)	Control diet	(58)
			28 % E	Mixed	12.8 g/d	Yes		↓ total bacteria	
			29 % E	Mixed	8.8 g/d	Yes		↓ total bacteria, <i>Bacteroides</i> , <i>Roseburia/Eubacterium rectale</i>	
n=20 Cross-over	30 - 42	28 days	94.4 g/d	Mixed	27.9 g/d	No	FISH (feces)	Control diet	(56)
			127.2 g/d	Mixed	11.7 g/d	Yes		↓ total bacteria, <i>Roseburia/Eubacterium rectale</i> , <i>Bifidobacterium</i>	
			119.5 g/d	Mixed	6.1 g/d	Yes		↓ total bacteria, <i>Roseburia/Eubacterium rectale</i> , <i>Bifidobacterium</i>	

Table 2: Effects of high-protein diet on the metabolic activity of the gut microbiota. The main characteristics and findings from human intervention studies using high-protein diets are summarized. BMI: body mass index, % E: % of energy intake, g/d: grams/day, NMR: nuclear magnetic resonance, GC: gas chromatography, MS: mass spectrometry, LC: liquid chromatography. For carbohydrates and fat intake, the readers are referred to the original publications.

Study design	BMI	Duration	Protein intake	Protein source	Fiber intake	Calorie restricted	Method	Intestinal microbiota metabolites	Reference
n=12-13 Parallel	25 - 30	3 weeks	14 % E	Mixed	17.0 g/d	No	NMR metabolomics, GC (feces)	Control diet	(26)
			34 % E	Mixed + casein	14.4 g/d	No		↓ butyrate; ↑ branched-chain amino acids, 2-methylbutyrate	
			31 % E	Mixed + soy protein	17.9 g/d	No		↓ butyrate; ↑ 2-methylbutyrate, isovalerate, valerate, phenylacetate, tyramine,	
n=12-13 Parallel	25 - 30	3 weeks	14 % E	Mixed	17.0 g/d	No	NMR metabolomics (urines)	Control diet	(26)
			34 % E	Mixed + casein	14.4 g/d	No		↑ isobutyrate, indoxylsulfate, phenylacetylglutamine, <i>p</i> -cresylsulfate	
			31 % E	Mixed + soy protein	17.9 g/d	No		↑ isobutyrate, indoxylsulfate, phenylacetylglutamine	
n = 10 Cross-over	19 - 32	5 days	10 % E	Plant protein	41.2 g/d	No	GC (feces)	No detectable differences	(51)
			16 % E	Mixed	21.1 g/d	No		Control diet	
			30 % E	Animal protein	0 g/d	No		↑ isobutyrate, isovalerate; ↓ acetate, butyrate	
n=14 Parallel	28 - 51	3 weeks	103.3 g/d	Mixed	27.7 g/d	No	GC (feces)	Control diet	(53)
			144.1 g/d	Mixed	25.1 g/d	Yes		↑ isobutyrate, isovalerate, lactate; ↓ Acetate, butyrate	
n=20 Cross-over	19 - 26	2 weeks	12 % E	Mixed	17.4 g/d	No	GC-MS metabolomics (feces)	No detectable differences	(45)
			15 % E	Mixed	16.3 g/d	No		Control diet	
			27 % E	Mixed	15.4 g/d	No		↑ isobutyrate	
n=20 Cross-over	19 - 26	2 weeks	12 % E	Mixed	17.4 g/d	No	GC-MS (urine)	No detectable differences	(45)
			15 % E	Mixed	16.3 g/d	No		Control diet	
			27 % E	Mixed	15.4 g/d	No		↑ <i>p</i> -cresol	
n=17 Cross-over	30 - 49	28 days	13 % E	Mixed	21.9 g/d	No	GC, LC-MS (Feces)	Control diet	(58)
			28 % E	Mixed	12.8 g/d	Yes		↑ isobutyrate, isovalerate, valerate, phenylacetate	
			29 % E	Mixed	8.8 g/d	Yes		↓ butyrate; ↑ isobutyrate, isovalerate, valerate, phenylacetate, phenylpropionate	
n=20 Cross-over	30 - 42	28 days	94.4 g/d	Mixed	27.9 g/d	No	GC (feces)	Control diet	(56)
			127.2 g/d	Mixed	11.7 g/d	Yes		↓ acetate, propionate, butyrate, valerate, lactate	
			119.5 g/d	Mixed	6.1 g/d	Yes		↓ acetate, propionate, butyrate, isovalerate, valerate, lactate; ↑ ammonia	

References

1. Hruby, A., and Hu, F. B. (2015) The Epidemiology of Obesity: A Big Picture. *Pharmacoeconomics* **33**, 673-689
2. Thom, G., and Lean, M. (2017) Is There an Optimal Diet for Weight Management and Metabolic Health? *Gastroenterology* **152**, 1739-1751
3. Santesso, N., Akl, E. A., Bianchi, M., Mente, A., Mustafa, R., Heels-Ansdell, D., and Schunemann, H. J. (2012) Effects of higher- versus lower-protein diets on health outcomes: a systematic review and meta-analysis. *European journal of clinical nutrition* **66**, 780-788
4. Pesta, D. H., and Samuel, V. T. (2014) A high-protein diet for reducing body fat: mechanisms and possible caveats. *Nutr Metab (Lond)* **11**, 53
5. Leidy, H. J., Clifton, P. M., Astrup, A., Wycherley, T. P., Westerterp-Plantenga, M. S., Luscombe-Marsh, N. D., Woods, S. C., and Mattes, R. D. (2015) The role of protein in weight loss and maintenance. *Am J Clin Nutr*
6. Blundell, J. E., Gibbons, C., Caudwell, P., Finlayson, G., and Hopkins, M. (2015) Appetite control and energy balance: impact of exercise. *Obes Rev* **16 Suppl 1**, 67-76
7. Clifton, P. M., Condo, D., and Keogh, J. B. (2014) Long term weight maintenance after advice to consume low carbohydrate, higher protein diets--a systematic review and meta analysis. *Nutr Metab Cardiovasc Dis* **24**, 224-235
8. Rueda-Clausen, C. F., Ogunleye, A. A., and Sharma, A. M. (2015) Health Benefits of Long-Term Weight-Loss Maintenance. *Annual review of nutrition* **35**, 475-516
9. Friedman, A. N. (2004) High-protein diets: potential effects on the kidney in renal health and disease. *Am J Kidney Dis* **44**, 950-962
10. Ko, G. J., Obi, Y., Tortorici, A. R., and Kalantar-Zadeh, K. (2017) Dietary protein intake and chronic kidney disease. *Curr Opin Clin Nutr Metab Care* **20**, 77-85
11. Blachier, F., Beaumont, M., Andriamihaja, M., Davila, A. M., Lan, A., Grauso, M., Armand, L., Benamouzig, R., and Tome, D. (2017) Changes in the Luminal Environment of the Colonic Epithelial Cells and Physiopathological Consequences. *The American journal of pathology* **187**, 476-486
12. Sanz, Y., Romani-Perez, M., Benitez-Paez, A., Portune, K. J., Brigidi, P., Rampelli, S., Dinan, T., Stanton, C., Delzenne, N., Blachier, F., Neyrinck, A. M., Beaumont, M., Olivares, M., Holzer, P., Gunther, K., Wolters, M., Ahrens, W., Claus, S. P., Campoy, C., Murphy, R., Sadler, C., Fernandez, L., and Kamp, J. V. (2018) Towards microbiome-informed dietary recommendations for promoting metabolic and mental health: Opinion papers of the MyNewGut project. *Clin Nutr*
13. Rand, W. M., Pellett, P. L., and Young, V. R. (2003) Meta-analysis of nitrogen balance studies for estimating protein requirements in healthy adults. *Am J Clin Nutr* **77**, 109-127
14. Dubuisson, C., Lioret, S., Touvier, M., Dufour, A., Calamassi-Tran, G., Volatier, J. L., and Lafay, L. (2010) Trends in food and nutritional intakes of French adults from 1999 to 2007: results from the INCA surveys. *Br J Nutr* **103**, 1035-1048
15. Pasiakos, S. M., Agarwal, S., Lieberman, H. R., and Fulgoni, V. L., 3rd. (2015) Sources and Amounts of Animal, Dairy, and Plant Protein Intake of US Adults in 2007-2010. *Nutrients* **7**, 7058-7069
16. Westerterp-Plantenga, M. S., Lemmens, S. G., and Westerterp, K. R. (2012) Dietary protein - its role in satiety, energetics, weight loss and health. *Br J Nutr* **108 Suppl 2**, S105-112
17. Phillips, S. M. (2014) A brief review of higher dietary protein diets in weight loss: a focus on athletes. *Sports Med* **44 Suppl 2**, S149-153
18. Tipton, K. D., and Wolfe, R. R. (2004) Protein and amino acids for athletes. *J Sports Sci* **22**, 65-79

19. Weigle, D. S., Breen, P. A., Matthys, C. C., Callahan, H. S., Meeuws, K. E., Burden, V. R., and Purnell, J. Q. (2005) A high-protein diet induces sustained reductions in appetite, ad libitum caloric intake, and body weight despite compensatory changes in diurnal plasma leptin and ghrelin concentrations. *Am J Clin Nutr* **82**, 41-48
20. Johnstone, A. M. (2012) Safety and efficacy of high-protein diets for weight loss. *The Proceedings of the Nutrition Society* **71**, 339-349
21. Skov, A. R., Toubro, S., Ronn, B., Holm, L., and Astrup, A. (1999) Randomized trial on protein vs carbohydrate in ad libitum fat reduced diet for the treatment of obesity. *Int J Obes Relat Metab Disord* **23**, 528-536
22. Claessens, M., van Baak, M. A., Monsheimer, S., and Saris, W. H. (2009) The effect of a low-fat, high-protein or high-carbohydrate ad libitum diet on weight loss maintenance and metabolic risk factors. *International journal of obesity (2005)* **33**, 296-304
23. Aller, E. E., Larsen, T. M., Claus, H., Lindroos, A. K., Kafatos, A., Pfeiffer, A., Martinez, J. A., Handjieva-Darlenska, T., Kunesova, M., Stender, S., Saris, W. H., Astrup, A., and van Baak, M. A. (2014) Weight loss maintenance in overweight subjects on ad libitum diets with high or low protein content and glycemic index: the DIOGENES trial 12-month results. *International journal of obesity (2005)* **38**, 1511-1517
24. Greenway, F. L. (2015) Physiological adaptations to weight loss and factors favouring weight regain. *International journal of obesity (2005)* **39**, 1188-1196
25. Neacsu, M., Fyfe, C., Horgan, G., and Johnstone, A. M. (2014) Appetite control and biomarkers of satiety with vegetarian (soy) and meat-based high-protein diets for weight loss in obese men: a randomized crossover trial. *Am J Clin Nutr* **100**, 548-558
26. Beaumont, M., Portune, K. J., Steuer, N., Lan, A., Cerrudo, V., Audebert, M., Dumont, F., Mancano, G., Khodorova, N., Andriamihaja, M., Airinei, G., Tome, D., Benamouzig, R., Davila, A. M., Claus, S. P., Sanz, Y., and Blachier, F. (2017) Quantity and source of dietary protein influence metabolite production by gut microbiota and rectal mucosa gene expression: a randomized, parallel, double-blind trial in overweight humans. *Am J Clin Nutr* **106**, 1005-1019
27. Theodorakopoulos, C., Jones, J., Bannerman, E., and Greig, C. A. (2017) Effectiveness of nutritional and exercise interventions to improve body composition and muscle strength or function in sarcopenic obese older adults: A systematic review. *Nutr Res* **43**, 3-15
28. Dattilo, A. M., and Kris-Etherton, P. M. (1992) Effects of weight reduction on blood lipids and lipoproteins: a meta-analysis. *Am J Clin Nutr* **56**, 320-328
29. Silvester, K. R., and Cummings, J. H. (1995) Does digestibility of meat protein help explain large bowel cancer risk? *Nutr Cancer* **24**, 279-288
30. Bos, C., Juillet, B., Fouillet, H., Turlan, L., Dare, S., Luengo, C., N'Tounda, R., Benamouzig, R., Gausseres, N., Tome, D., and Gaudichon, C. (2005) Postprandial metabolic utilization of wheat protein in humans. *Am J Clin Nutr* **81**, 87-94
31. Tome, D. (2012) Criteria and markers for protein quality assessment - a review. *Br J Nutr* **108 Suppl 2**, S222-229
32. Bos, C., Airinei, G., Mariotti, F., Benamouzig, R., Berot, S., Evrard, J., Fenart, E., Tome, D., and Gaudichon, C. (2007) The poor digestibility of rapeseed protein is balanced by its very high metabolic utilization in humans. *J Nutr* **137**, 594-600
33. Evenepoel, P., Claus, D., Geypens, B., Hiele, M., Geboes, K., Rutgeerts, P., and Ghoo, Y. (1999) Amount and fate of egg protein escaping assimilation in the small intestine of humans. *Am J Physiol* **277**, G935-943
34. Oberli, M., Marsset-Baglieri, A., Airinei, G., Sante-Lhoutellier, V., Khodorova, N., Remond, D., Foucault-Simonin, A., Piedcoq, J., Tome, D., Fromentin, G., Benamouzig, R., and Gaudichon, C. (2015) High True Ileal Digestibility but Not Postprandial Utilization of Nitrogen from Bovine Meat Protein in Humans Is Moderately Decreased by High-Temperature, Long-Duration Cooking. *J Nutr* **145**, 2221-2228
35. Chacko, A., and Cummings, J. H. (1988) Nitrogen losses from the human small bowel: obligatory losses and the effect of physical form of food. *Gut* **29**, 809-815

36. Gaudichon, C., Bos, C., Morens, C., Petzke, K. J., Mariotti, F., Everwand, J., Benamouzig, R., Dare, S., Tome, D., and Metges, C. C. (2002) Ileal losses of nitrogen and amino acids in humans and their importance to the assessment of amino acid requirements. *Gastroenterology* **123**, 50-59
37. Gibson, J. A., Sladen, G. E., and Dawson, A. M. (1976) Protein absorption and ammonia production: the effects of dietary protein and removal of the colon. *Br J Nutr* **35**, 61-65
38. Bax, M. L., Buffiere, C., Hafnaoui, N., Gaudichon, C., Savary-Auzeloux, I., Dardevet, D., Sante-Lhoutellier, V., and Remond, D. (2013) Effects of meat cooking, and of ingested amount, on protein digestion speed and entry of residual proteins into the colon: a study in minipigs. *PLoS one* **8**, e61252
39. Schippa, S., and Conte, M. P. (2014) Dysbiotic events in gut microbiota: impact on human health. *Nutrients* **6**, 5786-5805
40. Dinning, P. G. (2016) Recording In Vivo Human Colonic Motility: What Have We Learnt Over the Past 100 Years? *Advances in experimental medicine and biology* **891**, 213-222
41. Portune K J, B. M., Davila AM, Tome D, Blachier F, Sanz Y. (2016) Gut microbiota role in dietary protein metabolism and health-related outcomes: The two sides of the coin. *Trends in Food Science and Technology* **57**, 213-232
42. Fuller, M. (2012) Determination of protein and amino acid digestibility in foods including implications of gut microbial amino acid synthesis. *Br J Nutr* **108 Suppl 2**, S238-246
43. van der Wielen, N., Moughan, P. J., and Mensink, M. (2017) Amino Acid Absorption in the Large Intestine of Humans and Porcine Models. *J Nutr* **147**, 1493-1498
44. Davila, A. M., Blachier, F., Gotteland, M., Andriamihaja, M., Benetti, P. H., Sanz, Y., and Tome, D. (2013) Intestinal luminal nitrogen metabolism: Role of the gut microbiota and consequences for the host. *Pharmacological research : the official journal of the Italian Pharmacological Society* **68**, 95-107
45. Windey, K., De Preter, V., Louat, T., Schuit, F., Herman, J., Vansant, G., and Verbeke, K. (2012) Modulation of protein fermentation does not affect fecal water toxicity: a randomized cross-over study in healthy subjects. *PLoS one* **7**, e52387
46. Mottawea, W., Chiang, C. K., Muhlbauer, M., Starr, A. E., Butcher, J., Abujamel, T., Deeke, S. A., Brandel, A., Zhou, H., Shokralla, S., Hajibabaei, M., Singleton, R., Benchimol, E. I., Jobin, C., Mack, D. R., Figeys, D., and Stintzi, A. (2016) Altered intestinal microbiota-host mitochondria crosstalk in new onset Crohn's disease. *Nat Commun* **7**, 13419
47. Nyangale, E. P., Mottram, D. S., and Gibson, G. R. (2012) Gut microbial activity, implications for health and disease: the potential role of metabolite analysis. *Journal of proteome research* **11**, 5573-5585
48. Sharon, G., Garg, N., Debelius, J., Knight, R., Dorrestein, P. C., and Mazmanian, S. K. (2014) Specialized metabolites from the microbiome in health and disease. *Cell metabolism* **20**, 719-730
49. Louis, P., Hold, G. L., and Flint, H. J. (2014) The gut microbiota, bacterial metabolites and colorectal cancer. *Nature reviews. Microbiology* **12**, 661-672
50. Boleij, A., and Tjalsma, H. (2012) Gut bacteria in health and disease: a survey on the interface between intestinal microbiology and colorectal cancer. *Biol Rev Camb Philos Soc* **87**, 701-730
51. David, L. A., Maurice, C. F., Carmody, R. N., Gootenberg, D. B., Button, J. E., Wolfe, B. E., Ling, A. V., Devlin, A. S., Varma, Y., Fischbach, M. A., Biddinger, S. B., Dutton, R. J., and Turnbaugh, P. J. (2014) Diet rapidly and reproducibly alters the human gut microbiome. *Nature* **505**, 559-563
52. Walker, A. W., Ince, J., Duncan, S. H., Webster, L. M., Holtrop, G., Ze, X., Brown, D., Stares, M. D., Scott, P., Bergerat, A., Louis, P., McIntosh, F., Johnstone, A. M., Lobley, G. E., Parkhill, J., and Flint, H. J. (2011) Dominant and diet-responsive groups of bacteria within the human colonic microbiota. *The ISME journal* **5**, 220-230
53. Salonen, A., Lahti, L., Salojärvi, J., Holtrop, G., Korpela, K., Duncan, S. H., Date, P., Farquharson, F., Johnstone, A. M., Lobley, G. E., Louis, P., Flint, H. J., and de Vos, W. M.

- (2014) Impact of diet and individual variation on intestinal microbiota composition and fermentation products in obese men. *The ISME journal* **8**, 2218-2230
54. Martinez-Garcia, M., Diaz-Valdes, M., and Anton, J. (2010) Diversity of pufM genes, involved in aerobic anoxygenic photosynthesis, in the bacterial communities associated with colonial ascidians. *FEMS Microbiol Ecol* **71**, 387-398
 55. Martinez, I., Kim, J., Duffy, P. R., Schlegel, V. L., and Walter, J. (2010) Resistant starches types 2 and 4 have differential effects on the composition of the fecal microbiota in human subjects. *PloS one* **5**, e15046
 56. Duncan, S. H., Belenguer, A., Holtrop, G., Johnstone, A. M., Flint, H. J., and Lobley, G. E. (2007) Reduced dietary intake of carbohydrates by obese subjects results in decreased concentrations of butyrate and butyrate-producing bacteria in feces. *Appl Environ Microbiol* **73**, 1073-1078
 57. Cotillard, A., Kennedy, S. P., Kong, L. C., Prifti, E., Pons, N., Le Chatelier, E., Almeida, M., Quinquis, B., Levenez, F., Galleron, N., Gougis, S., Rizkalla, S., Batto, J. M., Renault, P., consortium, A. N. R. M., Dore, J., Zucker, J. D., Clement, K., Ehrlich, S. D., Blottiere, H., Leclerc, M., Juste, C., de Wouters, T., Lepage, P., Fouqueray, C., Basdevant, A., Henegar, C., Godard, C., Fondacci, M., Rohia, A., Hajdich, F., Weissenbach, J., Pelletier, E., Le Paslier, D., Gauchi, J. P., Gibrat, J. F., Loux, V., Carre, W., Maguin, E., van de Guchte, M., Jamet, A., Boumezbear, F., and Layec, S. (2013) Dietary intervention impact on gut microbial gene richness. *Nature* **500**, 585-588
 58. Russell, W. R., Gratz, S. W., Duncan, S. H., Holtrop, G., Ince, J., Scobbie, L., Duncan, G., Johnstone, A. M., Lobley, G. E., Wallace, R. J., Duthie, G. G., and Flint, H. J. (2011) High-protein, reduced-carbohydrate weight-loss diets promote metabolite profiles likely to be detrimental to colonic health. *Am J Clin Nutr* **93**, 1062-1072
 59. Geypens, B., Claus, D., Evenepoel, P., Hiele, M., Maes, B., Peeters, M., Rutgeerts, P., and Ghoo, Y. (1997) Influence of dietary protein supplements on the formation of bacterial metabolites in the colon. *Gut* **41**, 70-76
 60. Wikoff, W. R., Anfora, A. T., Liu, J., Schultz, P. G., Lesley, S. A., Peters, E. C., and Siuzdak, G. (2009) Metabolomics analysis reveals large effects of gut microflora on mammalian blood metabolites. *Proc Natl Acad Sci U S A* **106**, 3698-3703
 61. Roager, H. M., Hansen, L. B., Bahl, M. I., Frandsen, H. L., Carvalho, V., Gobel, R. J., Dalgaard, M. D., Plichta, D. R., Sparholt, M. H., Vestergaard, H., Hansen, T., Sicheritz-Ponten, T., Nielsen, H. B., Pedersen, O., Lauritzen, L., Kristensen, M., Gupta, R., and Licht, T. R. (2016) Colonic transit time is related to bacterial metabolism and mucosal turnover in the gut. *Nat Microbiol* **1**, 16093
 62. Nugent, S. G., Kumar, D., Rampton, D. S., and Evans, D. F. (2001) Intestinal luminal pH in inflammatory bowel disease: possible determinants and implications for therapy with aminosalicylates and other drugs. *Gut* **48**, 571-577
 63. Blachier, F., Davila, A. M., Mimoun, S., Benetti, P. H., Atanasiu, C., Andriamihaja, M., Benamouzig, R., Bouillaud, F., and Tome, D. (2010) Luminal sulfide and large intestine mucosa: friend or foe? *Amino Acids* **39**, 335-347
 64. Cummings, J. H., Hill, M. J., Bone, E. S., Branch, W. J., and Jenkins, D. J. (1979) The effect of meat protein and dietary fiber on colonic function and metabolism. II. Bacterial metabolites in feces and urine. *Am J Clin Nutr* **32**, 2094-2101
 65. Lozupone, C. A., Stombaugh, J. I., Gordon, J. I., Jansson, J. K., and Knight, R. (2012) Diversity, stability and resilience of the human gut microbiota. *Nature* **489**, 220-230
 66. Hamer, H. M., Jonkers, D., Venema, K., Vanhoutvin, S., Troost, F. J., and Brummer, R. J. (2008) Review article: the role of butyrate on colonic function. *Alimentary pharmacology & therapeutics* **27**, 104-119
 67. Thibault, R., Blachier, F., Darcy-Vrillon, B., de Coppet, P., Bourreille, A., and Segain, J. P. (2010) Butyrate utilization by the colonic mucosa in inflammatory bowel diseases: a transport deficiency. *Inflamm Bowel Dis* **16**, 684-695

68. Magee, E. A., Richardson, C. J., Hughes, R., and Cummings, J. H. (2000) Contribution of dietary protein to sulfide production in the large intestine: an in vitro and a controlled feeding study in humans. *Am J Clin Nutr* **72**, 1488-1494
69. Mimoun, S., Andriamihaja, M., Chaumontet, C., Atanasiu, C., Benamouzig, R., Blouin, J. M., Tome, D., Bouillaud, F., and Blachier, F. (2012) Detoxification of H₂S by differentiated colonic epithelial cells: implication of the sulfide oxidizing unit and of the cell respiratory capacity. *Antioxid Redox Signal* **17**, 1-10
70. Andriamihaja, M., Lan, A., Beaumont, M., Audebert, M., Wong, X., Yamada, K., Yin, Y., Tome, D., Carrasco-Pozo, C., Gotteland, M., Kong, X., and Blachier, F. (2015) The deleterious metabolic and genotoxic effects of the bacterial metabolite p-cresol on colonic epithelial cells. *Free Radic Biol Med* **85**, 219-227
71. Dodd, D., Spitzer, M. H., Van Treuren, W., Merrill, B. D., Hryckowian, A. J., Higginbottom, S. K., Le, A., Cowan, T. M., Nolan, G. P., Fischbach, M. A., and Sonnenburg, J. L. (2017) A gut bacterial pathway metabolizes aromatic amino acids into nine circulating metabolites. *Nature* **551**, 648-652
72. Pearson, J. R., Gill, C. I., and Rowland, I. R. (2009) Diet, fecal water, and colon cancer--development of a biomarker. *Nutrition reviews* **67**, 509-526
73. Benassi-Evans, B., Clifton, P., Noakes, M., and Fenech, M. (2010) High-protein/high red meat and high-carbohydrate weight-loss diets do not differ in their effect on faecal water genotoxicity tested by use of the WIL2-NS cell line and with other biomarkers of bowel health. *Mutation research* **703**, 130-136
74. Pedersen, G., Brynskov, J., and Saermark, T. (2002) Phenol toxicity and conjugation in human colonic epithelial cells. *Scand J Gastroenterol* **37**, 74-79
75. Ramakrishna, B. S., Roberts-Thomson, I. C., Pannall, P. R., and Roediger, W. E. (1991) Impaired sulphation of phenol by the colonic mucosa in quiescent and active ulcerative colitis. *Gut* **32**, 46-49
76. Gryp, T., Vanholder, R., Vaneechoutte, M., and Glorieux, G. (2017) p-Cresyl Sulfate. *Toxins (Basel)* **9**
77. Windey, K., De Preter, V., and Verbeke, K. (2012) Relevance of protein fermentation to gut health. *Mol Nutr Food Res* **56**, 184-196
78. Poveda, J., Sanchez-Nino, M. D., Glorieux, G., Sanz, A. B., Egido, J., Vanholder, R., and Ortiz, A. (2014) p-cresyl sulphate has pro-inflammatory and cytotoxic actions on human proximal tubular epithelial cells. *Nephrol Dial Transplant* **29**, 56-64
79. Watanabe, H., Miyamoto, Y., Honda, D., Tanaka, H., Wu, Q., Endo, M., Noguchi, T., Kadowaki, D., Ishima, Y., Kotani, S., Nakajima, M., Kataoka, K., Kim-Mitsuyama, S., Tanaka, M., Fukagawa, M., Otagiri, M., and Maruyama, T. (2013) p-Cresyl sulfate causes renal tubular cell damage by inducing oxidative stress by activation of NADPH oxidase. *Kidney international* **83**, 582-592
80. Wu, G. D., Chen, J., Hoffmann, C., Bittinger, K., Chen, Y. Y., Keilbaugh, S. A., Bewtra, M., Knights, D., Walters, W. A., Knight, R., Sinha, R., Gilroy, E., Gupta, K., Baldassano, R., Nessel, L., Li, H., Bushman, F. D., and Lewis, J. D. (2011) Linking long-term dietary patterns with gut microbial enterotypes. *Science* **334**, 105-108
81. Lin, C. J., Pan, C. F., Chuang, C. K., Sun, F. J., Wang, D. J., Chen, H. H., Liu, H. L., and Wu, C. J. (2014) P-cresyl sulfate is a valuable predictor of clinical outcomes in pre-ESRD patients. *Biomed Res Int* **2014**, 526932
82. Bansal, T., Alaniz, R. C., Wood, T. K., and Jayaraman, A. (2010) The bacterial signal indole increases epithelial-cell tight-junction resistance and attenuates indicators of inflammation. *Proc Natl Acad Sci U S A* **107**, 228-233
83. Shimada, Y., Kinoshita, M., Harada, K., Mizutani, M., Masahata, K., Kayama, H., and Takeda, K. (2013) Commensal bacteria-dependent indole production enhances epithelial barrier function in the colon. *PLoS one* **8**, e80604

84. Beaumont, M., Neyrinck, A. M., Olivares, M., Rodriguez, J., de Rocca Serra, A., Roumain, M., Bindels, L. B., Cani, P. D., Evenepoel, P., Muccioli, G. G., Demoulin, J. B., and Delzenne, N. M. (2018) The gut microbiota metabolite indole alleviates liver inflammation in mice. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*, fj201800544
85. Hubbard, T. D., Murray, I. A., Bisson, W. H., Lahoti, T. S., Gowda, K., Amin, S. G., Patterson, A. D., and Perdew, G. H. (2015) Adaptation of the human aryl hydrocarbon receptor to sense microbiota-derived indoles. *Sci Rep* **5**, 12689
86. Cheng, Y., Jin, U. H., Allred, C. D., Jayaraman, A., Chapkin, R. S., and Safe, S. (2015) Aryl Hydrocarbon Receptor Activity of Tryptophan Metabolites in Young Adult Mouse Colonocytes. *Drug Metab Dispos* **43**, 1536-1543
87. Santes-Palacios, R., Ornelas-Ayala, D., Cabanas, N., Marroquin-Perez, A., Hernandez-Magana, A., Del Rosario Olguin-Reyes, S., Camacho-Carranza, R., and Espinosa-Aguirre, J. J. (2016) Regulation of Human Cytochrome P4501A1 (hCYP1A1): A Plausible Target for Chemoprevention? *Biomed Res Int* **2016**, 5341081
88. He, X., and Feng, S. (2015) Role of Metabolic Enzymes P450 (CYP) on Activating Procarcinogen and their Polymorphisms on the Risk of Cancers. *Curr Drug Metab* **16**, 850-863
89. Badal, S., and Delgoda, R. (2014) Role of the modulation of CYP1A1 expression and activity in chemoprevention. *J Appl Toxicol* **34**, 743-753
90. Leong, S. C., and Sirich, T. L. (2016) Indoxyl Sulfate-Review of Toxicity and Therapeutic Strategies. *Toxins (Basel)* **8**
91. Tan, X., Cao, X., Zou, J., Shen, B., Zhang, X., Liu, Z., Lv, W., Teng, J., and Ding, X. (2017) Indoxyl sulfate, a valuable biomarker in chronic kidney disease and dialysis. *Hemodial Int* **21**, 161-167
92. Ramezani, A., and Raj, D. S. (2014) The gut microbiome, kidney disease, and targeted interventions. *Journal of the American Society of Nephrology : JASN* **25**, 657-670
93. Spooren, C. E., Pierik, M. J., Zeegers, M. P., Feskens, E. J., Masclee, A. A., and Jonkers, D. M. (2013) Review article: the association of diet with onset and relapse in patients with inflammatory bowel disease. *Alimentary pharmacology & therapeutics* **38**, 1172-1187
94. Jantchou, P., Morois, S., Clavel-Chapelon, F., Boutron-Ruault, M. C., and Carbonnel, F. (2010) Animal protein intake and risk of inflammatory bowel disease: The E3N prospective study. *The American journal of gastroenterology* **105**, 2195-2201
95. Jowett, S. L., Seal, C. J., Pearce, M. S., Phillips, E., Gregory, W., Barton, J. R., and Welfare, M. R. (2004) Influence of dietary factors on the clinical course of ulcerative colitis: a prospective cohort study. *Gut* **53**, 1479-1484
96. Arijs, I., Vanhove, W., Rutgeerts, P., Schuit, F., Verbeke, K., and De Preter, V. (2013) Decreased mucosal sulfide detoxification capacity in patients with Crohn's disease. *Inflamm Bowel Dis* **19**, E70-72
97. De Preter, V., Arijs, I., Windey, K., Vanhove, W., Vermeire, S., Schuit, F., Rutgeerts, P., and Verbeke, K. (2012) Decreased mucosal sulfide detoxification is related to an impaired butyrate oxidation in ulcerative colitis. *Inflamm Bowel Dis* **18**, 2371-2380
98. Beaumont, M., Andriamihaja, M., Lan, A., Khodorova, N., Audebert, M., Blouin, J. M., Grauso, M., Lancha, L., Benetti, P. H., Benamouzig, R., Tome, D., Bouillaud, F., Davila, A. M., and Blachier, F. (2016) Detrimental effects for colonocytes of an increased exposure to luminal hydrogen sulfide: The adaptive response. *Free Radic Biol Med* **93**, 155-164
99. Mu, C., Yang, Y., Luo, Z., Guan, L., and Zhu, W. (2016) The Colonic Microbiome and Epithelial Transcriptome Are Altered in Rats Fed a High-Protein Diet Compared with a Normal-Protein Diet. *J Nutr* **146**, 474-483
100. Beaumont, M., Andriamihaja, M., Armand, L., Grauso, M., Jaffrezic, F., Laloe, D., Moroldo, M., Davila, A. M., Tome, D., Blachier, F., and Lan, A. (2017) Epithelial response to a high-protein diet in rat colon. *BMC Genomics* **18**, 116

101. Vanholder, R., Schepers, E., Pletinck, A., Nagler, E. V., and Glorieux, G. (2014) The uremic toxicity of indoxyl sulfate and p-cresyl sulfate: a systematic review. *Journal of the American Society of Nephrology : JASN* **25**, 1897-1907
102. Chimerel, C., Emery, E., Summers, D. K., Keyser, U., Gribble, F. M., and Reimann, F. (2014) Bacterial metabolite indole modulates incretin secretion from intestinal enteroendocrine L cells. *Cell Rep* **9**, 1202-1208
103. Tomasova, L., Dobrowolski, L., Jurkowska, H., Wrobel, M., Huc, T., Ondrias, K., Ostaszewski, R., and Ufnal, M. (2016) Intracolonic hydrogen sulfide lowers blood pressure in rats. *Nitric Oxide* **60**, 50-58
104. Pichette, J., Fynn-Sackey, N., and Gagnon, J. (2017) Hydrogen Sulfide and Sulfate Prebiotic Stimulates the Secretion of GLP-1 and Improves Glycemia in Male Mice. *Endocrinology* **158**, 3416-3425

Figure 1

Figure 2

Figure 3