



# **The Arabidopsis NRT1.1 transceptor coordinately controls auxin biosynthesis and transport to regulate root branching in response to nitrate**

Amel Maghiaoui, Eléonore Bouguyon, Candela Cuesta, Francine Perrine-Walker, Carine Alcon, Gabriel Krouk, Eva Benkova, Philippe Nacry, Alain Gojon, Lien L. Bach

## **► To cite this version:**

Amel Maghiaoui, Eléonore Bouguyon, Candela Cuesta, Francine Perrine-Walker, Carine Alcon, et al.. The Arabidopsis NRT1.1 transceptor coordinately controls auxin biosynthesis and transport to regulate root branching in response to nitrate. *Journal of Experimental Botany*, 2020, 71 (15), pp.4480-4494. 10.1093/jxb/eraa242/5840670 . hal-02619371

**HAL Id: hal-02619371**

**<https://hal.inrae.fr/hal-02619371>**

Submitted on 25 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# **The *Arabidopsis* NRT1.1 transceptor coordinately controls auxin biosynthesis and transport to regulate root branching in response to nitrate**

Amel Maghiaoui<sup>\*1</sup>, Eléonore Bouguyon<sup>\*1,3</sup>, Candela Cuesta<sup>2,4</sup>, Francine Perrine-Walker<sup>1,5</sup>, Carine Alcon<sup>1</sup>, Gabriel Krouk<sup>1</sup>, Eva Benkova<sup>2</sup>, Philippe Nacry<sup>1</sup>, Alain Gojon<sup>1</sup> and Liên Bach<sup>1,\*\*</sup>

\* These authors contributed equally to this work.

<sup>1</sup>BPMP, Univ Montpellier, CNRS, INRAE, Institut Agro, Montpellier, France

<sup>2</sup>Institute of Science and Technology Austria, Klosterneuburg, 3400, Austria

<sup>3</sup>Present address: Site de Bellepierre, UFR Santé, Université de la Réunion, 1 Allée des Aigues Marines, 97487 Saint Denis Cedex, France

<sup>4</sup>Present address: Instituto Universitario de Biotecnología de Asturias, Departamento de Biología de Organismos y Sistemas, Universidad de Oviedo, Spain

<sup>5</sup>Present address: School of Life and Environmental Sciences, F22 Life, Earth and Environmental Sciences Building, University of Sydney NSW 2006, Australia

\*\* Address Correspondence to [lien.bach@supagro.fr](mailto:lien.bach@supagro.fr)

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors is: Liên BACH ([lien.bach@supagro.fr](mailto:lien.bach@supagro.fr)).

## **HIGHLIGHT**

The NRT1.1 transceptor controls both upstream auxin biosynthesis for lateral root primordia emergence and downstream action of auxin for lowering the mechanical resistance of the primary root tissues allowing LRP outgrowth.

Accepted Manuscript

## ABSTRACT

In agricultural systems, nitrate is the main source of nitrogen available for plants. Besides its role as a nutrient, nitrate has been shown to act as a signal molecule for plant growth, development and stress responses. In *Arabidopsis*, the NRT1.1 nitrate transceptor represses lateral root (LR) development at low nitrate availability by promoting auxin basipetal transport out of the LR primordia (LRPs). In addition, our present study shows that NRT1.1 acts as a negative regulator of the *TAR2* auxin biosynthetic gene expression in the root stele. This is expected to repress local auxin biosynthesis and thus to reduce acropetal auxin supply to the LRPs. Moreover, NRT1.1 also negatively affects expression of the LAX3 auxin influx carrier, thus preventing cell wall remodeling required for overlying tissues separation during LRP emergence. Both NRT1.1-mediated repression of *TAR2* and LAX3 are suppressed at high nitrate availability, resulting in the nitrate induction of *TAR2* and LAX3 expression that is required for optimal stimulation of LR development by nitrate. Altogether, our results indicate that the NRT1.1 transceptor coordinately controls several crucial auxin-associated processes required for LRP development, and as a consequence that NRT1.1 plays a much more integrated role than previously anticipated in regulating the nitrate response of root system architecture.

Key words: auxin signaling, development, lateral root primordia, nitrate, NRT1.1, transceptor

## INTRODUCTION

Nitrogen (N) is an essential element that plants can acquire from various inorganic or organic sources in the soil, but nitrate ( $\text{NO}_3^-$ ) is the main form taken up and assimilated by most species (Nacry *et al.*, 2013). Soil  $\text{NO}_3^-$  concentration is often a major factor limiting the growth and yield of crop plants as it fluctuates dramatically in both time and space (Crawford and Glass 1998; Miller *et al.*, 2007). To cope with this, plants are able to quickly modulate their  $\text{NO}_3^-$  acquisition efficiency according to changes both in their own internal N status and in the external  $\text{NO}_3^-$  availability (Forde 2002; Nacry *et al.*, 2013; Kiba and Krapp 2016; O'Brien *et al.*, 2016). In addition to the regulation of root  $\text{NO}_3^-$  transport systems (Kiba and Krapp 2016; Bellegarde *et al.*, 2017; Jacquot *et al.*, 2017), modification of root system architecture (RSA) is another key component of this response (Nacry *et al.*, 2013; Forde, 2014; Giehl and von Wirén, 2014; Kiba and Krapp, 2016; O'Brien *et al.*, 2016). In more detail,  $\text{NO}_3^-$  has a dual effect on RSA in many species. On the one hand, high  $\text{NO}_3^-$  provision to the plant has a systemic repressive effect (Zhang and Forde, 1998; Giehl and von Wirén, 2014), and as a consequence mild  $\text{NO}_3^-$  limitation results in a general stimulation of root growth for improved foraging of the soil (Forde, 2002; Hermans *et al.*, 2006; Giehl and von Wirén, 2014). On the other hand,  $\text{NO}_3^-$  supply locally stimulates growth of the roots in contact with a high  $\text{NO}_3^-$  concentration, therefore contributing to preferential colonization and exploitation of the  $\text{NO}_3^-$ -rich areas of the soil (Drew, 1975; Forde, 2002; Ruffel *et al.*, 2011, Nacry *et al.*, 2013). In dicotyledonous species where the root system is generally constituted of a primary root (PR) originating from the embryo, onto which lateral roots (LRs) develop post-embryonically,  $\text{NO}_3^-$  was generally reported to markedly affect LR development and growth, and to a much lesser extent PR growth (Forde, 2014). Furthermore, many responses of RSA to changes in  $\text{NO}_3^-$  availability were shown to be due to a signaling rather than a nutritional effect, pointing out the crucial role of specific  $\text{NO}_3^-$  sensing and signaling systems (O'Brien *et al.*, 2016; Guan, 2017; Sun *et al.*, 2017).

Compelling evidence indicates that the membrane  $\text{NO}_3^-$  transporter NRT1.1 (CHL1/NPF6.3), first identified as an influx carrier participating in the root uptake of  $\text{NO}_3^-$  (Tsay *et al.*, 1993; Liu and Tsay, 2003), is a central component of the  $\text{NO}_3^-$  sensing and signaling systems in *Arabidopsis thaliana* (Gojon *et al.*, 2009, 2011; Nacry *et al.*, 2013; Forde, 2014; Bouguyon *et al.*, 2015). This led to the proposal that it acts as a  $\text{NO}_3^-$  transceptor (transporter/receptor), with a dual transport/sensing function (Muños *et al.*, 2004; Walch-Liu and Forde, 2008; Ho *et*

*al.*, 2009; Wang *et al.*, 2009; Gojon *et al.*, 2011). In particular, NRT1.1 is a master player in the NO<sub>3</sub><sup>-</sup> regulation of RSA as it governs many responses of LR growth to NO<sub>3</sub><sup>-</sup> (Remans *et al.*, 2006; Krouk *et al.*, 2010; Bouguyon *et al.*, 2015; O'Brien *et al.*, 2016; Sun *et al.*, 2017). In *Arabidopsis*, LRs originate from pericycle founder cells that divide and differentiate in order to initiate a LRP (Stage I), which has to develop across overlying tissues to eventually emerge out of the parent root (Stage VIII) (Malamy and Benfey, 1997). Through this series of well-defined stages, auxin plays a central role in controlling many steps of LRP development (Dubrovsky *et al.*, 2008; Benkova *et al.*, 2003; Bhalerao, 2002; Fukaki *et al.*, 2007; Ljung *et al.*, 2001). Within the LRP, an auxin gradient is progressively established with a maximum at the apex that is required for proper growth of the LRP (Benkova *et al.*, 2003). This gradient is created by the coordinated activity of efflux proteins (encoded by the *PIN* and *ABCB* gene families) that facilitate movement out of cells and influx transporters (encoded mostly by the *AUX/LAX* gene family) that facilitate auxin entry into cells (Petrasek and Friml, 2009; Vieten *et al.*, 2007). A “fountain model” was proposed for auxin flow in LRP based on auxin accumulation and auxin transporters expression patterns (Benkova *et al.*, 2003). According to this model, an acropetal flow (involving PIN1/7 located in the inner cell layers) allows auxin to come from the PR stele and to reach the LRP apex where it accumulates, while a basipetal flow (mediated by PIN3/4 in the outer cell layer) redirects the hormone to the PR. In addition, AUX1, ABCB transporters and LAX3 have been shown to be involved in LRP development: AUX1 facilitates LRP initiation and emergence (Marchant *et al.*, 2002; Swarup and Péret, 2012) and ABCB1 and ABCB19 physically interact with PIN1 as well with AUX1 for normal LR initiation and development (Mravec *et al.*, 2008). The influx/efflux auxin transporter ABCB4 (Yang and Murphy, 2009; Wu *et al.*, 2010; Kubes *et al.*, 2012) was also identified as a negative regulator of LRP initiation (Santelia *et al.*, 2005) through a mechanism remaining to be elucidated. Finally, LAX3 plays an important role in LR emergence process by increasing the level of auxin in LRP overlying cells, thus leading to a remodeling of cortical/epidermal cell walls that facilitates their separation and allows the outgrowth of the LRP (Swarup *et al.*, 2008).

Within this general scheme, we recently proposed a model for the mechanism by which NRT1.1 regulates LR growth in response to NO<sub>3</sub><sup>-</sup> (Krouk *et al.*, 2010; Bouguyon *et al.*, 2015; Bouguyon *et al.*, 2016). Using heterologous expression systems, we found that NRT1.1 is not only able to transport NO<sub>3</sub><sup>-</sup> but also auxin. The auxin influx activity of NRT1.1 is however inhibited by NO<sub>3</sub><sup>-</sup> in a concentration-dependent manner (Krouk *et al.*, 2010). Furthermore, we

observed that NRT1.1 is expressed in LRPs only in the absence or at low  $\text{NO}_3^-$  availability (Bouguyon *et al.*, 2016). Taking into account the perfect match between the localization of the NRT1.1 protein with that of the basipetal transport route for auxin (outer cell layer of LRP), we postulated from the above observations that NRT1.1 mediates auxin influx into the outer cell layer of the LRPs, and thus facilitates the basipetal export of auxin out of these organs in the absence of  $\text{NO}_3^-$  or at low  $\text{NO}_3^-$  concentration (Krouk *et al.* 2010). Accordingly, in the absence of  $\text{NO}_3^-$  or at low  $\text{NO}_3^-$  concentration NRT1.1 lowers auxin accumulation/signaling (assessed by *DR5* activity) in the LRPs, which in turn leads to an inhibition of the emergence of the LRPs (Krouk *et al.*, 2010; Bouguyon *et al.*, 2015; Bouguyon *et al.*, 2016; Zhang *et al.*, 2019). At higher ( $>0.5$  mM)  $\text{NO}_3^-$  supply, both NRT1.1 expression in LRPs and NRT1.1-dependent auxin transport activity are inhibited. This results in an increased auxin accumulation/signaling in LRPs, which favors their emergence. Therefore, the local stimulation of LR growth by  $\text{NO}_3^-$  is explained by the relief of the repressive action of NRT1.1. Accordingly, null mutants for *NRT1.1* (*chl1* mutants) have a much higher LR density than wild-type plants in the absence, but not in the presence of  $\text{NO}_3^-$  (Krouk *et al.* 2010, Bouguyon *et al.* 2015, Bouguyon *et al.* 2016)

An important point that remains unclear in this model is to know if, in addition to its direct effect on auxin transport in the LRPs, NRT1.1 also acts indirectly through modified expression/activity of other molecular actors determining auxin distribution in these organs. As matter of fact, auxin transport in LRP also involves the action of up to ten other auxin carriers, *i.e.*, PIN1-4, PIN7, ABCB1, ABCB4, ABCB19, AUX1 and LAX3 (see Petrášek and Friml, 2009). Interestingly, earlier transcriptome studies at the whole root or whole plant level have revealed that several auxin transporter genes are responsive to  $\text{NO}_3^-$  (Gutierrez *et al.*, 2007) in an NRT1.1-dependent manner (c Furthermore, there are reports indicating that responses of RSA to nutrients, including N, can be accounted for by changes in expression of *AUX* and *PIN* genes (Li *et al.*, 2011; Giehl *et al.*, 2012; Sun *et al.*, 2017; Naz *et al.*, 2019). Nevertheless, these considerations do not allow any conclusion on a putative role of NRT1.1 in regulating other auxin carriers for modulating LR development because little is known on the regulation of PIN, ABCB and AUX/LAX expression by  $\text{NO}_3^-$  signaling in LRPs. In addition, recent evidence highlights the importance of not only auxin transport, but also local auxin biosynthesis in determining the changes in auxin accumulation in plant tissues in response to external factors (Brumos *et al.*, 2018; Zhao *et al.*, 2018). This suggests that the nutrient regulation of RSA may also rely on a control exerted on auxin biosynthesis pathways. Of particular interest is the observation that the *Arabidopsis* *TAR2* gene, encoding



a Tryptophane amino transferase catalyzing the first step of the Trp-dependent auxin biosynthetic pathway, is required for stimulation of LR growth in response to mild N limitation (Ma *et al.*, 2014). However, nothing is known on the role of this gene in the local NO<sub>3</sub><sup>-</sup> stimulation of LRP development in connection with NRT1.1-dependent signaling.

In the current work, we thus aimed at investigating how NO<sub>3</sub><sup>-</sup> regulates the expression of the major auxin transporters (PINs, ABCBs, AUX1/LAX3) and of the TAR2 auxin biosynthetic enzyme during LRP development, and at determining if this regulation may account for an indirect effect of NRT1.1 on LR growth, through its NO<sub>3</sub><sup>-</sup> signaling action on auxin transporter or auxin biosynthetic enzyme expression. As shown by the results detailed below, our work revealed that NRT1.1 has a much more integrated role in regulating LRP development than anticipated before. Indeed, our data indicate that NRT1.1 not only participates in auxin transport in LRPs, but also controls both the upstream biosynthesis of the hormone as a source of auxin for the LRPs, and the downstream action of auxin for lowering the mechanical resistance of the PR tissues to LRP outgrowth.

## MATERIALS & METHODS

### Plant material

The *Arabidopsis thaliana* accessions used in this study were Columbia (Col-0) and Wassilewskija (WS). The following GFP-tagged lines were used and crossed with the *chl1-5* mutant (Tsay *et al.*, 1993): *pDR5::VENUS* (Brunoud *et al.*, 2012), *pPIN1::PIN1-GFP* (Benkova, 2003), *pPIN2::PIN2-GFP* (Xu and Scheres, 2005), *pPIN3::PIN3-GFP* (Zadnikova *et al.*, 2010), *pPIN4::PIN4-GFP* (Vieten, 2005), *pPIN7::PIN7-GFP* (Blilou *et al.*, 2005; Mravec *et al.*, 2008), *pAUX1::AUX1-YFP* (Swarup, 2004), *pLAX3::LAX3-YFP* (Swarup *et al.*, 2008), *DR5::GFP* (Ulmasov *et al.*, 1997), *pTAR2::GUS* and *tar2-c* (Ma *et al.*, 2014). The following GFP-tagged lines were used and crossed with the *chl1-10* mutant (Munos *et al.*, 2004): *pABCB1::ABCB1-GFP* (Mravec *et al.*, 2008), *pABCB4::ABCB4-GFP* (Cho *et al.*, 2007), *pABCB19::ABCB19-GFP* (Bailly *et al.*, 2008). Homozygous plants for both *chl1-5* mutation and genes encoding tagged proteins were screened on F2 and F3 offsprings by PCR for *chl1-5* deletion as described in Mounier *et al.*, (2014) and fluorescence. Transgenic *Arabidopsis* line expressing *NRT1.1-GFPLoop* in WT background were obtained as described by Bouguyon *et al.*, 2016.



### Plant culture and root growth analysis

Seedlings were grown for 12 days in vertical petri dishes according to Krouk *et al.*, (2010). Basal culture medium without N was supplemented with the appropriate concentration of KNO<sub>3</sub>, or Gln for each experiment as described in figure's caption. For analysis of root system architecture, the petri dishes were scanned using a Epson Perfection V700 PHOTO scanner (24-bits-color; resolution 600 ppp). The root system parameters such as primary root length, lateral roots length and number as well as lateral root density were measured by the Neuron J plugin from Image J software. Root system pictures for all genotypes were taken using Nikon digital camera D90. To analyze root growth parameters, Student T Test's was used to show statistical difference between genotypes and treatments.

### RNA analysis

Root samples were frozen in liquid nitrogen and disrupted for 1 min at 30 s<sup>-1</sup> in a Retch mixer mill MM301 homogenizer. Total RNA was extracted using TRIzol reagent (Invitrogen), DNase treated (Qiagen) purified using an RNeasy MinElute Cleanup Kit (Qiagen) and reverse transcription was achieved with 4 µg of RNAs with ThermoScript™ RT-PCR System for First-Strand cDNA Synthesis (Invitrogen) using an anchored oligo-dT<sub>(20)</sub> primer. Accumulation of transcript was measured by quantitative real-time PCR (LightCycler 480; Roche Diagnostics) using the Light Cycler 480 SYBR Green 1 Master kit (Roche Diagnostics). All steps were performed according manufacturer's recommendations. The gene expression was normalized using *CLATHRIN* as an internal standard. The specific primers used are summarized in table 1. In order to examine whether gene expression was significantly different between treatments or genotypes, ANOVA followed by a Turkey test or Student T Test's were applied.

### Gus staining

Samples were prefixed for 45 minutes at room temperature in a solution containing 50 mM NaPO<sub>4</sub> pH7, 1,5% formaldehyde and 0,05% Triton X100. Then plants are washed 3 times in 50 mM NaPO<sub>4</sub> pH7 with 0.05% Triton X100. Thereafter, samples were stained in 50 mM NaPO<sub>4</sub> pH7, 0,5 mM Ferricyanide, 0,5 mM Ferrocyanide, 0,05% Triton X100 and 1 mM X-Gluc at 37°C for 48h in the dark. After GUS staining, plants were washed in 50 mM NaPO<sub>4</sub> pH7 for 5 minutes and chlorophyll was removed by EtOH increasing concentrations solutions.

The samples were mounted in 20% EtOH and were imaging by using x10 N.A. 0,3 or x40 N.A. 0,75 objectives on an Olympus BX61 microscope equipped with a camera Jenoptik Progres.

### Confocal microscopy

For confocal microscopy observation, plant samples were mounted in osmoted water and viewed on a Leica SP8 Confocal Microscope (Leica). GFP/YFP were excited at 488nm by an Argon source and re-emitted light was filtered by a passing band of 505-530nm. Laser intensity was adjusted for each line and pinhole was settled at 1 unit, resolution at 1024 x 1024 to compare expression and accumulation of reporter proteins. Quantification of fluorescence in roots was performed using Image J software. The data represent the mean pixel values in a ROI.

## **RESULTS**

### **Regulation of auxin carriers mRNA levels in response to NO<sub>3</sub><sup>-</sup>.**

To determine how auxin transporter genes respond to NO<sub>3</sub><sup>-</sup> and whether this response is dependent on NRT1.1, we measured by quantitative RT-PCR the transcript levels of ten auxin transporters in roots of both WT and *chl1-5* (*NRT1.1* null mutant) plants grown for 8 days on either N-free medium or on medium containing 1mM KNO<sub>3</sub>. The ten auxin transporter genes reported to contribute to auxin traffic in LR<sub>s</sub> (*PIN1-4*, *PIN7*, *ABCB1*, *ABCB4*, *ABCB19*, *AUX1* and *LAX3*) were sorted in three different groups according to their expression patterns (Figure 1). The first one is constituted by *PIN2*, *PIN3* and *ABCB1*, for which mRNA levels are not significantly influenced either by NO<sub>3</sub><sup>-</sup> supply or by *chl1-5* mutation (Figure 1A). The second group gathers *ABCB4*, *ABCB19*, *PIN1*, *PIN4*, and *PIN7*, for which mRNA levels are modulated by NO<sub>3</sub><sup>-</sup>. Among these genes, *ABCB4* is the only one to be repressed upon NO<sub>3</sub><sup>-</sup> supply, whereas the others are induced (Figure 1B). The NO<sub>3</sub><sup>-</sup>-responses of these genes are the same in both WT and *chl1-5* plants, suggesting that NRT1.1 plays no role in these responses. The last group contains *AUX1* and *LAX3* (Figure 1C). Both genes show higher expression in presence of NO<sub>3</sub><sup>-</sup> as compared to N-free medium in WT plants, but the difference is suppressed or strongly attenuated in *chl1-5* mutant. This is due to the up-regulation of these two genes in the *chl1-5* mutant as compared to the WT in N-free

condition, but not on 1 mM  $\text{NO}_3^-$ . Therefore, NRT1.1 appears to act as a negative regulator of the expression of both *AUX1* and *LAX3* in the absence, but not in the presence of  $\text{NO}_3^-$ . Altogether, these data show that although the majority of the auxin carrier genes investigated (7 out of 10) respond to  $\text{NO}_3^-$  provision in the roots, only two of them display a  $\text{NO}_3^-$  response that seems to involve NRT1.1 as a regulator.

### Protein level and localization of auxin carriers in response to $\text{NO}_3^-$ .

The response to  $\text{NO}_3^-$  of NRT1.1 and of the ten auxin transporters was evaluated at the protein level in LRPs using reporter-tagged transgenic plants (translational fusions with GFP or YFP) in WT or *chl1* mutant genetic background. Confocal sections through roots of transgenic plants grown in similar conditions to those used for the quantitative RT-PCR measurements were performed to compare protein accumulation and localization in LRPs in different conditions and at different developmental stages.

As reported previously (Bouguyon *et al.*, 2016), the protein expression of NRT1.1 in LRPs is strongly regulated by  $\text{NO}_3^-$ . Indeed, the GFP signal in roots of plants expressing a *NRT1.1::GFP* transgene under the control of the *NRT1.1* promoter in *chl1-5* background indicates that NRT1.1 is present in pre-emerged LRPs in the absence, but not in the presence of 1 mM  $\text{NO}_3^-$  in the medium (Supplemental Fig. 1). In  $\text{NO}_3^-$ -fed plants, NRT1.1 expression is only observed in LRPs after emergence. Most auxin transporters (8 out of 10) do not show any significant  $\text{NO}_3^-$ -dependent change either in protein accumulation or localization in the context of LRP development and emergence. In our conditions and regardless the presence of  $\text{NO}_3^-$ , ABCB4 and PIN4 are not detectable in LRPs and PIN7 appears to be only transiently expressed at stages I and II but then rapidly disappears as LRPs progress to next phase (data not shown). Moreover, as previously shown (Benkova *et al.*, 2003), PIN2 is not expressed in LRPs before emergence (Supplemental Fig. 2). Neither the localization nor the protein abundance of PIN1, ABCB19 and AUX1 are significantly affected by  $\text{NO}_3^-$  or *chl1* mutation (Supplemental Figs 3, 4 and 5). Finally, although the GFP signal associated with the ABCB1-GFP fusion protein seemed slightly increased in the presence of  $\text{NO}_3^-$ , this was not found to be statistically significant after quantification (Supplemental Fig. 6).

In contrast, PIN3, and LAX3 are responsive to  $\text{NO}_3^-$  treatments. In WT plants and in presence of  $\text{NO}_3^-$ , PIN3-GFP spatio-temporal expression pattern in LRPs is similar to that previously

shown (Benkova *et al.*, 2003). Indeed, PIN3 is expressed from the first developmental stage to emergence and its early widespread localization is progressively restricted to the meristematic zone (Figure 2A- a-c right panels). However, on N-free medium, PIN3-GFP signal becomes undetectable in LRPs from stage IV-V up to emergence (Figure 2A b-c left panel). To determine whether this is due to a  $\text{NO}_3^-$  specific effect, PIN3-GFP expression was investigated in plants grown on 0.5 mM glutamine as an alternative N source (Figure 2A, central panels). Supply of glutamine results in a PIN3-GFP expression similar to that recorded with  $\text{NO}_3^-$ , indicating that lack of expression on N-free medium is not due to the absence of  $\text{NO}_3^-$  *per se*, but most probably results from N-deficiency. In agreement with the hypothesis that PIN3-GFP is not specifically regulated by  $\text{NO}_3^-$ , the effect of N on this protein is NRT1.1-independent since the same pattern is observed in a WT or a *chl1* mutant background (Figure 2B, 2C).

As previously shown (Swarup *et al.*, 2008), LAX3-YFP signal in WT background is detected from stage II to stage VII in mature stele and in cortical or epidermal cells overlaying LRPs under  $\text{NO}_3^-$  supply (Figure 3A- a to c right panels). However, LAX3-YFP signal is strongly decreased in cells overlaying LRPs, regardless of the LRP developmental stage, in response to either N-starvation (Figure 3A- a to c left panel) or supply of glutamine as a N source (Figure 3A, middle panel). Interestingly, this down-regulation of LAX3-YFP expression is not recorded in *chl1-5* mutant background (Figure 3B, 3C). These data show that NRT1.1 acts as a negative regulator of LAX3 expression in the absence of  $\text{NO}_3^-$ . Interestingly, the *chl1-5* mutation restores LAX3-YFP signal in LRP overlying cells only after LRPs have reached stage IV (Figure 3A and B-compare a with b and c stages), indicating that NRT1.1 is responsible for LAX3 extinction at late, but not at early stages of LRP development. This is fully consistent with the observation that NRT1.1 starts repressing LRP development at stages IV/V (Krouk *et al.*, 2010; Bouguignon *et al.*, 2016). In conclusion, the pattern of LAX3-YFP expression around LRPs in WT and *chl1* mutant background is in agreement with that of LAX3 transcript accumulation in the roots (*i.e.*, both are up-regulated by  $\text{NO}_3^-$  in WT plants and by *chl1* mutation in the absence of  $\text{NO}_3^-$ ).

It has been reported that the upregulation of LAX3 expression that facilitates LRP outgrowth is triggered by the LBD29 transcription factor (Swarup *et al.*, 2008; Porco *et al.*, 2016). To test whether LBD29 responds to  $\text{NO}_3^-$  in a NRT1.1-dependent manner, its transcript accumulation in roots was investigated in roots of Col or *chl1-5* plants grown on N-free

medium or on 1 mM  $\text{NO}_3^-$ . The data indicate that expression of *LBD29* is strongly induced by  $\text{NO}_3^-$  supply as compared to N-free conditions in WT plants, and that it is markedly overexpressed in the *chl1-5* mutant as compared with the WT on N-free medium, but not in the presence of  $\text{NO}_3^-$  (Figure 3D). This is fully consistent with the expression patterns of both *LAX3* gene (Figure 1) and LAX3-GFP protein (Figure 3), which therefore suggests that LAX3 response to  $\text{NO}_3^-$  is mediated by the NRT1.1-dependent regulation of *LBD29* expression.

### **The TAR2 auxin biosynthesis enzyme is up-regulated by $\text{NO}_3^-$ in a NRT1.1-dependent manner.**

The *TAR2* gene was reported to be N-responsive and to contribute to the regulation of LR development by  $\text{NH}_4\text{NO}_3$  supply (Ma *et al.*, 2014). However, its putative role in the specific response to  $\text{NO}_3^-$  in connection with the NRT1.1-dependent pathway is not known. Therefore, we first investigated the effect of the three N regimes (N-free, 0.5 mM glutamine or 1 mM  $\text{NO}_3^-$ ) on *TAR2* mRNA accumulation in the roots of WT and *chl1-5* plants, as well as in those of the *tar2c* knock-out mutant of *TAR2*. Although *TAR2* mRNA accumulation was previously shown to be down-regulated by increasing  $\text{NH}_4\text{NO}_3$  concentrations (Ma *et al.*, 2014), it was in our conditions markedly induced in the WT by N supply as compared with N-free condition, with a much more pronounced response on  $\text{NO}_3^-$  as compared with glutamine (Figure 4A). This  $\text{NO}_3^-$  induction was fully lost in the *chl1-5* mutant because *TAR2* expression was already high in this line on N-free medium or on gln. As expected, no significant *TAR2* mRNA level was recorded in the *tar2c* knock-out mutant.

Both the induction of *TAR2* expression by  $\text{NO}_3^-$ , and its up-regulation in N-starved *chl1-5* plants as compared to WT were confirmed by histological staining of GUS activity in *pTAR2::GUS* plants (Figure 4B). As shown previously (Ma *et al.*, 2014), *TAR2* expression in roots is predominantly localized in the stele, and we did not observe *pTAR2* activity in endodermis, cortex or epidermis (Figure 4B and 4C). In presence of either 0.5 mM gln or 1 mM  $\text{NO}_3^-$ , strong GUS staining was also observed inside the vast majority (>90%) of young LRPs (stages I-III) in both WT and *chl1-5* plants (Figure 4C). On N-free medium, GUS staining in LRPs was only visible in ca~60% of the young LRPs of *chl1-5* plants, and at a lower intensity than that recorded in plants fed with either gln or  $\text{NO}_3^-$  (Figure 4C). In all cases, *pTAR2::GUS* expression in LRPs was no more recorded as soon as these LRPs

evolved to later stages of development (stages IV-V onwards). Altogether, the above results show that in the absence of  $\text{NO}_3^-$ , NRT1.1 acts as a strong negative regulator of *TAR2* expression in the stele and to a lower extent in the young LRPs. This indicates that NRT1.1 may control LRP development not only through its auxin transport facilitation within the LRPs (Krouk *et al.*, 2010), but also by modulating auxin biosynthesis in the stele. In addition, these results also show that *TAR2* expression may not only be responsive to NRT1.1-mediated  $\text{NO}_3^-$  signaling but also to another N stimulatory effect that can be activated by gln supply.

### **TAR2 is required for the stimulation of LR development by $\text{NO}_3^-$ and its regulation by NRT1.1**

The induction of *TAR2* expression in the roots by  $\text{NO}_3^-$  suggests that increased auxin biosynthesis is associated with  $\text{NO}_3^-$  regulation of RSA. To determine whether *TAR2* may contribute to the  $\text{NO}_3^-$  stimulation of LR development by locally increasing auxin provision to LRPs, expression of *DR5::GFP* was investigated in both WT and *tar2c* mutant backgrounds in the roots of plants grown on 1 mM  $\text{NO}_3^-$  (Figure 5). The GFP signal was predominantly found in the stele and LRPs of both genotypes, but with a strongly reduced intensity in *tar2c* as compared to WT (Figure 5). The negative effect of *TAR2* mutation on *DR5::GFP* expression in LRPs was found at almost all stages of development, with the exception of stage VII, just prior to emergence (Figures 5B and 5C). This indicates that *TAR2*-dependent auxin biosynthesis in the stele and/or the LRPs is required for increasing auxin accumulation/signaling in LRPs in response to  $\text{NO}_3^-$ . Note however that although important, the presence of a functional *TAR2* gene is not strictly required to generate *DR5::GFP* expression in the root. This expression is still significant in the stele and in LRPs at all developmental stages in the *tar2c* mutant (Figures 5A, 5B and 5C), suggesting that auxin partly comes from another origin than the *TAR2*-dependent biosynthesis pathway.

*TAR2* mutation alone (*tar2c* mutant) had only very limited effect, if any, on the growth of the primary root in the three conditions tested (Figures 6A and 6B). However, it very significantly decreased (by nearly 50%) the density of LR in the plants grown on  $\text{NO}_3^-$ , but not in those supplied with either glutamine or N-free medium (Figures 6A and 6C). This demonstrates that *TAR2* participates in the stimulation of LR development by  $\text{NO}_3^-$ , in agreement with the observations that *TAR2* expression is induced by  $\text{NO}_3^-$  and that its



mutation lowers *DR5* activity in LRPs (Figures 4 and 5). Combining *NRT1.1* and *TAR2* mutations (double *chl1-5tar2c* mutant) had no additional effect on LR development as compared to the single *TAR2* mutation, but strikingly affected it as compared to the single *NRT1.1* mutation (Figures 6A and 6C). Indeed, the dramatic increase of LR density classically observed in *chl1-5* as compared to the WT on  $\text{NO}_3^-$ -free medium (Krouk *et al.*, 2010; Bouguyon *et al.*, 2015) was totally abolished in the double *chl1-5tar2c* mutant. These data indicate that the up-regulation of *TAR2* expression in response to *NRT1.1* mutation (Figure 4) is most likely one key response required to trigger enhanced LR development in *chl1-5* plants.

## DISCUSSION

### Nitrate regulation of auxin carriers in lateral root primordia

One striking conclusion arising from our work is that, with the noticeable exception of LAX3, *NRT1.1* has little effect on the expression of the auxin carriers involved in LRP development. However, as suggested by previous reports on *Arabidopsis* (Li *et al.*, 2011) and rice (Naz *et al.*, 2019; Sun *et al.*, 2017), our data show that the root expression of key auxin carriers is responsive to N supply in WT plants. Although not dramatic, changes in mRNA levels in the whole root were found in response to  $\text{NO}_3^-$  for 7 out of the 10 auxin carriers investigated (Figure 1). When considering more precisely the expression at the protein level in or around the LRPs of WT background plants, only 2 of them (namely, PIN3 and LAX3) displayed a visible increase in the fluorescence from GFP or YFP fusion proteins on 1 mM  $\text{NO}_3^-$  as compared to N-free medium (Figures 2 and 3). Most importantly, for PIN3 this effect was not specific of  $\text{NO}_3^-$  since the same increase in fluorescence was observed upon provision of 0.5 mM gln as a N source (Figure 2). This suggests that the response to  $\text{NO}_3^-$  of PIN3 is not specifically due to  $\text{NO}_3^-$  signaling, but either to a signaling action of an N metabolite downstream of  $\text{NO}_3^-$  assimilation, or to more general effect of N nutrition. As a matter of fact, organic forms of N, and in particular amino acids, have been shown to specifically control several mechanisms involved in root growth and development in *Arabidopsis* (Walch-Liu *et al.*, 2006; Gifford *et al.*, 2008; Vidal *et al.*, 2010). Our data indicate that none of the auxin carriers expressed in LRPs (PIN1-4, PIN7, ABCB1, ABCB4, ABCB19 and AUX1) displays a regulation pattern correlated with that previously reported for *DR5* activity in LRPs (*e.g.*, repression in the absence of  $\text{NO}_3^-$  that is suppressed in the



*chl1-5* mutant, (see Krouk *et al.*, 2010). This supports the hypothesis that NRT1.1 modulates auxin accumulation in LRP directly through its auxin transport capacity, and not indirectly through modified expression of other auxin carriers. However, it cannot be ruled out that NRT1.1 may act on these carriers at a regulatory level that was not investigated in our study, by triggering posttranslational modifications affecting transport activity for instance. Furthermore, although not predominantly expressed in LRPs, ABCB21 has recently been reported to markedly affect *DR5* activity in these organs (Jenness *et al.*, 2019), but its regulation by  $\text{NO}_3^-$  is unknown. Therefore, further investigation is needed to conclude whether the above listed auxin carriers are regulated by NRT1.1, or not.. In contrast, our data on LAX3 mRNA accumulation and of LAX3-YFP fusion protein expression clearly indicate that the expression of this carrier in overlying tissues of the LRPs is under control by NRT1.1-dependent  $\text{NO}_3^-$  signaling. First, the stimulation by  $\text{NO}_3^-$  of LAX3-YFP expression in cortical cells cannot be mimicked by gln supply (Figure 3). Second, both the increases in LAX3 gene and LAX3-YFP protein expression in response to  $\text{NO}_3^-$  were suppressed in the *chl1-5* mutant (Figures 1 and 3). This is further supported by the fact that the gene expression of LBD29, the direct positive regulator of LAX3 (Porco *et al.*, 2016), displays the same pattern of regulation (Figure 3). In agreement with what was previously recorded for LR density and for auxin accumulation/signaling in LRP (*DR5* activity), the stimulatory effect of  $\text{NO}_3^-$  was suppressed because LAX3 and LBD29 expression were already high in the *chl1-5* mutant in the absence of  $\text{NO}_3^-$ , indicating a negative regulatory role of NRT1.1 under this condition (Figures 1 and 3). The mechanisms by which NRT1.1 may lead to repression of LBD29 and LAX3 expression are not known. However, the stimulation of both LBD29 and LAX3 expression in cells overlying LRPs was proposed to be due to the induction of LBD29 and LAX3 genes by the auxin diffusing from the LRP to the overlying tissues (Swarup *et al.*, 2008; Porco *et al.*, 2016). Therefore, it is conceivable that the NRT1.1-mediated repression of auxin accumulation in LRPs in the absence of  $\text{NO}_3^-$  (Krouk *et al.*, 2010; Bouguyon *et al.*, 2015; Bouguyon *et al.*, 2016; Zhang *et al.*, 2019), may lead to a lowered auxin diffusion out of the LRPs, which may in turn limit induction of both LBD29 and LAX3 genes.

## Auxin biosynthesis is a target of NRT1.1-dependent NO<sub>3</sub><sup>-</sup> signaling in roots

The importance of local auxin biosynthesis for controlling plant development has been recently reconsidered, and shown to be important for responses to internal or external signals (ethylene, light, temperature and pathogens, Ljung *et al.*, 2001; Brumos *et al.*, 2018; Zhao *et al.*, 2018). Data concerning RSA responses to nutrients are scarce on this point (Guan *et al.*, 2017; Olatunji *et al.*, 2017), with the exception of the report by Ma *et al.*, (2014), indicating that the *TAR2* auxin biosynthetic gene contributes to the root foraging response triggered by mild N-limitation in *Arabidopsis*. This response is considered to be due to a repression exerted by high N supply, which is not specific to NO<sub>3</sub><sup>-</sup>, and may involve systemic signaling (Forde, 2014; Giehl and von Wirén, 2014). Our data confirm a key role of *TAR2* in the N regulation of LR growth, but in a different context that is the local stimulation of LRP development by NO<sub>3</sub><sup>-</sup> signaling. Indeed, *TAR2* expression is induced by NO<sub>3</sub><sup>-</sup> (Figure 4), and *tar2* mutants display a markedly reduced increase in LR density in response to NO<sub>3</sub><sup>-</sup> supply as compared with WT plants (Figure 6). This is correlated with reduced *DR5* activity in both the stele and LRPs of plants with *tar2* as compared to WT background (Figure 5). This suggests that the *TAR2* biosynthetic pathway is an important source of auxin for ensuring proper accumulation of the hormone in LRPs and for yielding full stimulation of LR development by NO<sub>3</sub><sup>-</sup>. However, because neither *DR5* activity nor LR emergence is totally suppressed in the *tar2* mutant, it is likely that another auxin source, coming from acropetal transport from the shoot, other biosynthesis and/or conjugation pathway (Bhalerao *et al.*, 2002; Petrasek and Friml, 2009; Olatunji *et al.*, 2017), also contributes to supply auxin to the LRPs. As it is the case for LR density and key factors associated with it (*DR5* activity in LRPs, *LBD29/LAX 3* expression), the stimulation of *TAR2* expression by NO<sub>3</sub><sup>-</sup> in the stele is actually mostly due to the relief of the repression exerted by NRT1.1 in the absence of NO<sub>3</sub><sup>-</sup>. This is illustrated by the upregulation of *TAR2* in the *chl1-5* mutant on N-free medium (Figure 4). This *TAR2* upregulation appears to be crucial for the strongly increased LR growth phenotype of the *chl1-5* plants because the double *chl1-5tar2c* mutant did not differ in LR density from the WT in NO<sub>3</sub><sup>-</sup>-free media (Figure 4). The mechanism associated with the NRT1.1-dependent repression of *TAR2* expression in the stele in the absence of NO<sub>3</sub><sup>-</sup> is unclear. One hypothesis is that the stimulation of basipetal auxin transport out of the LRPs by NRT1.1 (Krouk *et al.*, 2010) may locally increase auxin levels in the stele. Because the *TAR2* auxin biosynthetic pathway has been recently reported to be feedback repressed by auxin (Suzuki *et al.*, 2015; Takato *et al.*, 2017), this may explain why *TAR2* expression is

reduced in the stele of WT plants deprived of  $\text{NO}_3^-$  (Figure 4). Alternatively, because NRT1.1 has been shown to be also expressed in the central cylinder of mature portions of the PR, where it contributes to xylem loading of  $\text{NO}_3^-$  (Léran *et al.*, 2013), it is also possible that NRT1.1 may act on *TAR2* expression directly in the stele when  $\text{NO}_3^-$  is absent from the medium.

Interestingly, *TAR2* was found to be expressed in a N-stimulated manner in young LRPs at stages I-III (Figure 4C), *i.e.*, at earlier developmental stages than those (IV-V) where NRT1.1 starts to act as a regulator of LRP emergence (Bouguyon *et al.*, 2016). Because *TAR2* expression in young LRPs is responsive to both gln and  $\text{NO}_3^-$ , this suggests that N can modulate early LRP development by controlling auxin biosynthesis *in situ* independently of NRT1.1. This is in line with the previous proposal that N can act on various developmental check points of LR development through different mechanisms (Forde, 2014; Giehl and von Wirén, 2014).

### **A more integrated model for NRT1.1-dependent regulation of LR development by $\text{NO}_3^-$ .**

Our previous working model for explaining the NRT1.1-dependent repression of LR growth in the absence or at low availability ( $<0.5$  mM) of  $\text{NO}_3^-$  almost entirely relied on the hypothesis that NRT1.1 inhibits LRP outgrowth (more specifically at developmental stage IV) because it facilitates the basipetal auxin transport out of the LRP, therefore preventing the hormone accumulation at the LRP tip (Krouk *et al.*, 2010; Bouguyon *et al.*, 2015). This hypothesis was supported by the auxin transport capacity of NRT1.1 recorded in heterologous systems (Krouk *et al.*, 2010; Bouguyon *et al.*, 2015; Wei *et al.*, 2018 ; Zhang *et al.*, 2019), and by its largely predominant localization in the outer cell layer of the LRP (Krouk *et al.*, 2010), which corresponds to the basipetal auxin transport route (Benkova *et al.*, 2003). Because increasing  $\text{NO}_3^-$  concentrations ( $>0.5$  mM) results both in an inhibition of the NRT1.1-mediated auxin transport facilitation (Krouk *et al.*, 2010) and in a repression of the NRT1.1 protein expression in LRPs (Bouguyon *et al.*, 2016), we proposed that the local stimulation of LR growth by  $\text{NO}_3^-$  is due to a relief of the repression exerted by NRT1.1. The data reported here strengthen our previous conclusions, but show that NRT1.1 affects other processes in addition to auxin distribution within the LRP, such as the biosynthesis of the hormone and the mechanical resistance of the PR tissues to LRP outgrowth. As summarized in Figure 7, the repressive role of NRT1.1 on LRP development can now be extended to three separate but synergistic actions. Indeed, NRT1.1 prevents auxin accumulation in the LRP not

only by favoring the basipetal transport of the hormone, but also by triggering the repression of *TAR2* expression in the stele. As a consequence of reduced auxin biosynthesis, it is likely that the acropetal flux channeling the hormone to the LRP tip is also lowered, therefore contributing to slowing down LRP development and emergence. In addition, NRT1.1 acts as a negative regulator of the expression of *LAX3* in the PR tissues overlying the LRPs. Given the proposed function of *LAX3* in LRP development (Swarup *et al.*, 2008), this is expected to prevent the cell wall relaxation in the LRP overlying tissues that is required to lower the mechanical resistance to LRP emergence (Lucas *et al.*, 2013).

Beyond the repression of LRP development at low  $\text{NO}_3^-$  availability summarized in Figure 7, NRT1.1 has also been reported to favor LR elongation in  $\text{NO}_3^-$ -rich patches of the external medium (Remans *et al.*, 2006), and to trigger the induction of the *AFB3* auxin receptor gene by  $\text{NO}_3^-$  that results in repressed PR growth and stimulated LR initiation (Vidal *et al.*, 2010; Vidal *et al.*, 2014). By interfering with auxin biosynthesis, transport, signaling, and thus by controlling various key checkpoints of LR development, our studies reveal that NRT1.1 has a much more integrated function in governing the  $\text{NO}_3^-$  regulation of RSA in *Arabidopsis* than previously anticipated (Vidal *et al.*, 2014; Bouguyon *et al.*, 2015). It may therefore be expected that all aspects and mechanisms of this integrated function are not fully unraveled yet.

## ACKNOWLEDGMENTS

P.N., G.K., E.Ba., L.B. and A.G. designed and supervised the experiments; A.M., E.Bo., C.C.M., F.P.W., P.N. and L.B. performed experiments; A.M. and E.Bo. generated plant material; A.M., E.Bo., C.C.M., G.K., P.N., E.Ba., L.B. and A.G. analyzed the data, A.M., L.B., and A.G. wrote the article with contribution of all the authors.

This work was supported by the Agence Nationale de la Recherche program “Programme PCRI ANR-16-CE20-0022”. We would like to thank members of the lab for discussion. We are grateful to Yiping Tong for the gift of tar2-c (SALK\_143208.56.00.x) and TAR2pro::GUS (CS16434) seeds from the Arabidopsis Bio-logical Resource Center (ABRC). We thank the Montpellier Rio Imaging platform for microscopy observations.

## BIBLIOGRAPHY

- Bailly A, Sovero V, Vincenzetti V, Santelia D, Bartnik D, Koenig BW, Mancuso S, Martinoia E, Geisler M.** 2008. Modulation of P-glycoproteins by auxin transport inhibitors is mediated by interaction with immunophilins. *Journal of Biological Chemistry* **283**, 21817-21826.
- Bellegarde F, Gojon A, Martin A.** 2017. Signals and players in the transcriptional regulation of root responses by local and systemic N signaling in *Arabidopsis thaliana*. *Journal of Experimental Botany* **68**, 2553-2565.
- Benkova E, Michniewicz M, Sauer M, Teichmann T, Seifertova D, Jurgens G, Friml J.** 2003. Local, efflux-dependent auxin gradients as a common module for plant organ formation. *Cell* **115**, 591-602.
- Bhalerao RP, Eklof J, Ljung K, Marchant A, Bennett M, Sandberg G.** 2002. Shoot-derived auxin is essential for early lateral root emergence in *Arabidopsis* seedlings. *The Plant Journal* **29**, 325-332.
- Blilou I, Xu J, Wildwater M, Willemsen V, Paponov I, Friml J, Heidstra R, Aida M, Palme K, Scheres B.** 2005. The PIN auxin efflux facilitator network controls growth and patterning in *Arabidopsis* roots. *Nature* **433**, 39-44.
- Bouguyon E, Brun F, Meynard D, et al** 2015. Multiple mechanisms of nitrate sensing by *Arabidopsis* nitrate transceptor NRT1.1. *Nature Plants* **1**, 15015.
- Bouguyon E, Perrine-Walker F, Pervent M, et al.** 2016. Nitrate Controls Root Development through Posttranscriptional Regulation of the NRT1.1/NPF6.3 Transporter/Sensor. *Plant Physiology* **172**, 1237-1248.
- Brumos J, Robles LM, Yun J, Vu TC, Jackson S, Alonso JM, Stepanova AN.** 2018. Local Auxin Biosynthesis Is a Key Regulator of Plant Development. *Developmental Cell* **47**, 306-318.e305.
- Brunoud G, Wells DM, Oliva M, et al.** 2012. A novel sensor to map auxin response and distribution at high spatio-temporal resolution. *Nature* **482**, 103-106.
- Cho M, Lee SH, Cho HT.** 2007. P-Glycoprotein4 Displays Auxin Efflux Transporter-Like Action in *Arabidopsis* Root Hair Cells and Tobacco Cells. *The Plant Cell Online* **19**, 3930-3943.
- Crawford NM, Glass ADM.** 1998. Molecular and physiological aspects of nitrate uptake in plants. *Trends in Plant Science* **3**, 389-395.
- Drew MC.** 1975. Comparison of the effects of a localised supply of phosphate, nitrate, ammonium and potassium on the growth of the seminal root system, and the shoot in barley. *New Phytologist* **75**, 479-490.
- Dubrovsky JG, Sauer M, Napsucialy-Mendivil S, Ivanchenko MG, Friml J, Shishkova S, Celenza J, Benkova E.** 2008. Auxin acts as a local morphogenetic trigger to specify lateral root founder cells. *Proc Natl Acad Sci U S A* **105**, 8790-8794.
- Forde BG.** 2002. Local and long-range signaling pathways regulating plant responses to nitrate. *Annual Reviews Plant Biology* **53**, 203-224.
- Forde BG.** 2014. Nitrogen signalling pathways shaping root system architecture: an update. *Current Opinion in Plant Biology* **21**, 30-36.
- Fukaki H, Okushima Y, Tasaka M.** 2007. Auxin-mediated lateral root formation in higher plants. *International Review Cytology* **256**, 111-137.
- Giehl RF, von Wiren N.** 2014. Root nutrient foraging. *Plant Physiology* **166**, 509-517.



- Giehl RFH, Lima JE, von Wirén N.** 2012. Localized iron supply triggers lateral root elongation in *Arabidopsis* by altering the AUX1-mediated auxin distribution. *The Plant Cell* **24**, 33-49.
- Gifford ML, Dean A, Gutierrez RA, Coruzzi GM, Birnbaum KD.** 2008. Cell-specific nitrogen responses mediate developmental plasticity. *Proc Natl Acad Sci U S A* **105**, 803-808.
- Gojon A, Krouk G, Perrine-Walker F, Laugier E.** 2011. Nitrate transceptor(s) in plants. *Journal of Experimental Botany* **62**, 2299-2308.
- Gojon A, Nacry P, Davidian JC.** 2009. Root uptake regulation: a central process for NPS homeostasis in plants. *Current Opinion in Plant Biology* **12**, 328-338.
- Guan P.** 2017. Dancing with Hormones: A Current Perspective of Nitrate Signaling and Regulation in *Arabidopsis*. *Frontiers in Plant Science* **8**, 1697.
- Gutierrez RA, Lejay LV, Dean A, Chiaromonte F, Shasha DE, Coruzzi GM.** 2007. Qualitative network models and genome-wide expression data define carbon/nitrogen-responsive molecular machines in *Arabidopsis*. *Genome Biology* **8**, R7.
- Hermans C, Hammond JP, White PJ, Verbruggen N.** 2006. How do plants respond to nutrient shortage by biomass allocation? *Trends in Plant Science* **11**, 610-617.
- Ho CH, Lin SH, Hu HC, Tsay YF.** 2009. CHL1 functions as a nitrate sensor in plants. *Cell* **138**, 1184-1194.
- Jacquot A, Li Z, Gojon A, Schulze W, Lejay L.** 2017. Post-translational regulation of nitrogen transporters in plants and microorganisms. *Journal of Experimental Botany* **68**, 2567-2580.
- Jenness MK, Carraro N, Pritchard CA, Murphy AS.** 2019. The *Arabidopsis* ATP-BINDING CASSETTE Transporter ABCB21 regulates auxin levels in cotyledons, the root pericycle, and leaves. *Frontiers in Plant Science* **10**, 806.
- Kiba T, Krapp A.** 2016. Plant Nitrogen Acquisition Under Low Availability: Regulation of Uptake and Root Architecture. *Plant and Cell Physiology* **57**, 707-714.
- Krouk G, Lacombe B, Bielach A, et al.** 2010. Nitrate-regulated auxin transport by NRT1.1 defines a mechanism for nutrient sensing in plants. *Developmental Cell* **18**, 927-937.
- Kubes M, Yang H, Richter GL, et al.** 2012. The *Arabidopsis* concentration-dependent influx/efflux transporter ABCB4 regulates cellular auxin levels in the root epidermis. *The Plant Journal* **69**, 640-654.
- Leran S, Munos S, Brachet C, Tillard P, Gojon A, Lacombe B.** 2013. *Arabidopsis* NRT1.1 is a bidirectional transporter involved in root-to-shoot nitrate translocation. *Molecular Plant* **6**, 1984-1987.
- Li B, Li Q, Su Y, Chen H, Xiong L, Mi G, Kronzucker HJ, Shi W.** 2011. Shoot-supplied ammonium targets the root auxin influx carrier AUX1 and inhibits lateral root emergence in *Arabidopsis*. *Plant, Cell and Environment* **34**, 933-946.
- Liu KH, Tsay YF.** 2003. Switching between the two action modes of the dual-affinity nitrate transporter CHL1 by phosphorylation. *The EMBO Journal* **22**, 1005-1013.
- Ljung K, Bhalerao RP, Sandberg G.** 2001. Sites and homeostatic control of auxin biosynthesis in *Arabidopsis* during vegetative growth. *The Plant Journal* **28**, 465-474.
- Lucas M, Kenobi K, von Wangenheim D, et al.** 2013. Lateral root morphogenesis is dependent on the mechanical properties of the overlaying tissues. *Proc Natl Acad Sci U S A* **110**, 5229-5234.
- Ma W, Li J, Qu B, He X, Zhao X, Li B, Fu X, Tong Y.** 2014. Auxin biosynthetic gene TAR2 is involved in low nitrogen-mediated reprogramming of root architecture in *Arabidopsis*. *The Plant Journal* **78**, 70-79.
- Malamy J, Benfey P.** 1997. Organization and cell differentiation in lateral roots of *Arabidopsis thaliana*. *Development (Cambridge, England)* **124**, 33-44.



- Marchant A, Bhalerao R, Casimiro I, Eklöf J, Casero PJ, Bennett M, Sandberg G.** 2002. AUX1 promotes lateral root formation by facilitating indole-3-acetic acid distribution between sink and source tissues in the Arabidopsis seedling. *The Plant Cell* **14**, 589-597.
- Miller AJ, Fan X, Orsel M, Smith SJ, Wells DM.** 2007. Nitrate transport and signalling. *Journal of Experimental Botany* **58**, 2297-2306.
- Mounier E, Pervent M, Ljung K, Gojon A, Nacry P.** 2014. Auxin-mediated nitrate signalling by NRT1.1 participates in the adaptive response of Arabidopsis root architecture to the spatial heterogeneity of nitrate availability. *Plant, Cell and Environment* **37**, 162-174.
- Mravec J, Kubes M, Bielach A, Gaykova V, Petrsek J, Skupa P, Chand S, Benkova E, Zazimalova E, Friml J.** 2008. Interaction of PIN and PGP transport mechanisms in auxin distribution-dependent development. *Development* **135**, 3345-3354.
- Munos S, Cazettes C, Fizames C, Gaymard F, Tillard P, Lepetit M, Lejay L, Gojon A.** 2004. Transcript profiling in the chl1-5 mutant of Arabidopsis reveals a role of the nitrate transporter NRT1.1 in the regulation of another nitrate transporter, NRT2.1. *The Plant Cell* **16**, 2433-2447.
- Nacry P, Bouguyon E, Gojon A.** 2013. Nitrogen acquisition by roots: physiological and developmental mechanisms ensuring plant adaptation to a fluctuating resource. *Plant and Soil* **370**, 1-29.
- Naz M, Luo B, Guo X, Li B, Chen J, Fan X.** 2019. Overexpression of Nitrate Transporter OsNRT2.1 Enhances Nitrate-Dependent Root Elongation. *Genes (Basel)* **10**.
- O'Brien JA, Vega A, Bouguyon E, Krouk G, Gojon A, Coruzzi G, Gutierrez RA.** 2016. Nitrate Transport, Sensing, and Responses in Plants. *Molecular Plant* **9**, 837-856.
- Olatunji D, Geelen D, Verstraeten I.** 2017. Control of Endogenous Auxin Levels in Plant Root Development. *International Journal of Molecular Sciences* **18**.
- Petrsek J, Friml J.** 2009. Auxin transport routes in plant development. *Development* **136**, 2675-2688.
- Porco S, Larrieu A, Du Y, et al.** 2016. Lateral root emergence in Arabidopsis is dependent on transcription factor LBD29 regulation of auxin influx carrier LAX3. *Development* **143**, 3340-3349.
- Remans T, Nacry P, Pervent M, Filleur S, Diatloff E, Mounier E, Tillard P, Forde BG, Gojon A.** 2006. The Arabidopsis NRT1.1 transporter participates in the signaling pathway triggering root colonization of nitrate-rich patches. *Proc Natl Acad Sci U S A* **103**, 19206-19211.
- Ruffel S, Krouk G, Ristova D, Shasha D, Birnbaum KD, Coruzzi GM.** 2011. Nitrogen economics of root foraging: transitive closure of the nitrate-cytokinin relay and distinct systemic signaling for N supply vs. demand. *Proc Natl Acad Sci U S A* **108**, 18524-18529.
- Santelia D, Vincenzetti V, Azzarello E, Bovet L, Fukao Y, Duchtig P, Mancuso S, Martinoia E, Geisler M.** 2005. MDR-like ABC transporter AtPGP4 is involved in auxin-mediated lateral root and root hair development. *FEBS Letters* **579**, 5399-5406.
- Sun CH, Yu JQ, Hu DG.** 2017. Nitrate: A Crucial Signal during Lateral Roots Development. *Frontiers in Plant Science* **8**, 485.
- Suzuki M, Yamazaki C, Mitsui M, Kakei Y, Mitani Y, Nakamura A, Ishii T, Soeno K, Shimada Y.** 2015. Transcriptional feedback regulation of YUCCA genes in response to auxin levels in Arabidopsis. *The Plant Cell Reports* **34**, 1343-1352.
- Swarup K, Benková E, Swarup R, et al.** 2008. The auxin influx carrier LAX3 promotes lateral root emergence. *Nature Cell Biology* **10**, 946-954.
- Swarup R, Kargul J, Marchant A, et al.** 2004. Structure-function analysis of the presumptive Arabidopsis auxin permease AUX1. *The Plant Cell* **16**, 3069-3083.
- Swarup R, Peret B.** 2012. AUX/LAX family of auxin influx carriers-an overview. *Frontiers in Plant Science* **3**, 225.

- Takato S, Kakei Y, Mitsui M, et al.** 2017. Auxin signaling through SCF(TIR1/AFBs) mediates feedback regulation of IAA biosynthesis. *Bioscience, biotechnology, and biochemistry* **81**, 1-7.
- Tsay YF, Schroeder JI, Feldmann KA, Crawford NM.** 1993. The herbicide sensitivity gene CHL1 of Arabidopsis encodes a nitrate-inducible nitrate transporter. *Cell* **72**, 705-713.
- Ulmasov T, Murfett J, Hagen G, Guilfole TJ.** 1997. Aux/IAA proteins repress expression of reporter genes containing natural and highly active synthetic auxin response element. *The Plant Cell* **9**, 1963-1971.
- Vidal EA, Alvarez JM, Gutierrez RA.** 2014. Nitrate regulation of AFB3 and NAC4 gene expression in Arabidopsis roots depends on NRT1.1 nitrate transport function. *Plant Signaling and Behavior* **9**, e28501.
- Vidal EA, Araus V, Lu C, Parry G, Green PJ, Coruzzi GM, Gutierrez RA.** 2010. Nitrate-responsive miR393/AFB3 regulatory module controls root system architecture in Arabidopsis thaliana. *Proc Natl Acad Sci U S A* **107**, 4477-4482.
- Vieten A, Sauer M, Brewer PB, Friml J.** 2007. Molecular and cellular aspects of auxin-transport-mediated development. *Trends in Plant Science* **12**, 160-168.
- Vieten A, Vanneste S, Wisniewska J, Benkova E, Benjamins R, Beeckman T, Luschnig C, Friml J.** 2005. Functional redundancy of PIN proteins is accompanied by auxin-dependent cross-regulation of PIN expression. *Development* **132**, 4521-4531.
- Walch-Liu P, Forde BG.** 2008. Nitrate signalling mediated by the NRT1.1 nitrate transporter antagonises L-glutamate-induced changes in root architecture. *The Plant Journal* **54**, 820-828.
- Walch-Liu P, Liu LH, Remans T, Tester M, Forde BG.** 2006. Evidence that L-glutamate can act as an exogenous signal to modulate root growth and branching in Arabidopsis thaliana. *Plant and Cell Physiology* **47**, 1045-1057.
- Wang R, Xing X, Wang Y, Tran A, Crawford NM.** 2009. A genetic screen for nitrate regulatory mutants captures the nitrate transporter gene NRT1.1. *Plant Physiology* **151**, 472-478.
- Wei J, Zheng Y, Feng H, Qu H, Fan X, Yamaji N, Ma JF, Xu G.** 2018. OsNRT2.4 encodes a dual-affinity nitrate transporter and functions in nitrate-regulated root growth and nitrate distribution in rice. *Journal of Experimental Botany* **69**, 1095-1107.
- Xu J, Scheres B.** 2005. Dissection of Arabidopsis ADP-RIBOSYLATION FACTOR 1 function in epidermal cell polarity. *The Plant Cell* **17**, 525-536.
- Zadnikova P, Petrasek J, Marhavy P, et al.** 2010. Role of PIN-mediated auxin efflux in apical hook development of Arabidopsis thaliana. *Development* **137**, 607-617.
- Zhang H, Forde BG.** 1998. An Arabidopsis MADS box gene that controls nutrient-induced changes in root architecture. *Science* **279**, 407-409.
- Zhang X, Cui Y, Yu M, Su B, Gong W, Baluska F, Komis G, Samaj J, Lin J.** 2019. Phosphorylation-Mediated Dynamics of Nitrate Transceptor NRT1.1 Regulate Auxin Flux and Nitrate Signaling in Lateral Root Growth. *Plant Physiology* **181**, 480-498.
- Zhao Y.** 2018. Essential Roles of Local Auxin Biosynthesis in Plant Development and in Adaptation to Environmental Changes. *Annual Review of Plant Biology* **69**, 417-435.

## FIGURE LEGENDS

### **Fig. 1: Relative mRNA accumulation of auxin transporters in roots of WT (Col) and *chl1-5* seedlings grown in presence or absence of $\text{NO}_3^-$ .**

Wild-type and *chl1-5* seedlings were grown for 8 days in vertical agar plates on basal medium supplemented with 1 mM  $\text{NO}_3^-$  or not. Transcript levels were quantified in the roots by quantitative RT-PCR using *CLATHRIN* as an internal standard. The ten auxin transporter genes investigated are grouped into three different classes, according to the response of their mRNA levels to  $\text{NO}_3^-$  and *NRT1.1* mutation:

- A) Auxin transporters not regulated by  $\text{NO}_3^-$
- B) Auxin transporters regulated by  $\text{NO}_3^-$  independently of *NRT1.1*.
- C) Auxin transporters regulated by  $\text{NO}_3^-$  in a *NRT1.1*-dependent way.

The data are the mean  $\pm$  se of 2 replicates from 3 independent experiments.

Statistically significant differences between genotypes or N treatments at  $P < 0.05$  are shown by different letters and ANOVA followed by a Turkey Test were used to show these significant differences.

### **Fig. 2: PIN3-GFP expression in lateral root primordia is repressed by N starvation both in WT (Col) and *chl1-5* genetic backgrounds.**

Seedlings expressing PIN3-GFP in Col (A) or in *chl1-5* (B) genetic background were grown for 8 days in vertical agar plates on basal medium deprived of N (left panel) or supplemented with either 0.5mM glutamine (Gln ; middle panel) or 1mM  $\text{NO}_3^-$  (right panel). Primary root portions with lateral root primordia at various developmental stages were observed under confocal microscope.

a = stage I-III, b = stage IV-V, c = stage VI-VII. White bar represents 100 $\mu\text{m}$ . The pictures shown are representative of >20 primordia from >10 plants of 3 independent experiments.

(C) Fluorescence quantification in different root tissues of 8-day-old plants expressing PIN3-GFP in *WT* or *chl1.5* lines in the absence or presence of  $\text{NO}_3^-$  or glutamine (Gln). Fluorescence was measured in unmerged primordia stage IV to VIII, and emerged primordia. Values are the mean of 5–10 plants from 2 independent experiments. P values were analyzed by Student's T Test and differences with N treatments are statistically significant at \*\*\*P , 0.001.

**Fig. 3: LAX3-YFP expression in tissues overlaying lateral root primordia is repressed in the absence of NO<sub>3</sub><sup>-</sup> in a *NRT1.1*-dependent manner.**

Seedlings expressing LAX3-YFP in Col (A) or in *chl1-5* (B) genetic background were grown for 8 days in vertical agar plates on basal medium deprived of N (left panel) or supplemented with either 0.5mM glutamine (middle panel) or 1mM NO<sub>3</sub><sup>-</sup> (right panel). Primary root portions with lateral root primordia at various developmental stages were observed under confocal microscope.

a = stage I-III, b = stage IV-V, c = stage VI-VII. White bar represents 100µm. The pictures shown are representative of >20 primordia from >10 plants of 3 independent experiments.

(C) Fluorescence quantification in different root tissues of 8-day-old plants expressing LAX3-YFP in *WT* or *chl1.5* lines in the absence or presence of NO<sub>3</sub><sup>-</sup> or glutamine (Gln). Fluorescence was measured in cortical/epidermal cells surrounding unmerged primordia stage IV to VIII. Values are the mean of 5–10 plants from 2 independent experiments. P values were analyzed by Student's T Test and differences with N treatments are statistically significant at \*\*\*\*P, 0.0001. (D) Relative mRNA accumulation of LBD29 in roots of Col and *chl1-5* plants grown for 12 days in agar plates on basal medium deprived of N or supplemented with 1 mM NO<sub>3</sub><sup>-</sup>. The expression was quantified by RT-qPCR and normalized by *CLATHRIN*.

The data are the mean ± se from 3 independent experiments. P values were analyzed by Student's T Test and differences between genotypes are statistically significant at \*P < 0.005.

**Fig.4: *TAR2* expression in the roots is repressed in the absence of NO<sub>3</sub><sup>-</sup> in a *NRT1.1*-dependent manner.**

Wild-type, *chl1-5* and *tar2-c* seedlings were grown for 12 days in vertical agar plates on basal medium deprived of N, or supplemented with either 1 mM NO<sub>3</sub><sup>-</sup>, or 0.5 mM L-glutamine.

(A) Relative accumulation of *TAR2* transcripts in roots. The data are the mean ± se of 2 replicates from 3 independent experiments. P values were analyzed by Student's T Test and differences between genotypes are statistically significant at \*\*P < 0.01.

(B) Histochemical staining of GUS activity in roots of plants expressing a *pTAR2::GUS* transgene in either wild type (Col) or *chl1-5* genetic background. Black bar represents 0.5 cm.

(C) Histochemical staining of GUS activity in the root stele and lateral root primordia of plants expressing a *pTAR2::GUS* transgene in either wild type (Col) or *chl1-5* genetic

background. The pictures shown display root portions with lateral root primordia at stages I-III. The pictures shown are representative of 10 primordia from >10 plants of two independent experiments. Black bar represents 0.5 cm.

**Fig. 5: Mutation of *TAR2* represses DR5::GFP expression in LRPs and in the stele under  $\text{NO}_3^-$  supply.**

(A) DR5::GFP expression in mature root zone of plants with WT (left panel) or *tar2c* (right panel) genetic background. The seedlings were grown for 12 days on basal medium supplemented with 1mM  $\text{NO}_3^-$ . White bars = 100  $\mu\text{m}$ .

(B) DR5::GFP expression in lateral root primordia of plants with WT (upper panel) or *tar2c* (lower panel) genetic background. The seedlings were grown for 12 days on basal medium supplemented with 1mM  $\text{NO}_3^-$ . The pictures display lateral root primordia at different developmental stages (from stage II to VI). The asterisk visualizes the location of the primordium. The pictures are representative of 10 plants of 2 independent experiments. White bars = 100  $\mu\text{m}$ .

(C) Fluorescence quantification in lateral root primordia at different developmental stages of 12-d-old plants expressing DR5::GFP in WT or *tar2c* background. Plants were grown on basal medium supplemented with 1mM of  $\text{NO}_3^-$ . Data represent mean  $\pm$  se of 2 independent experiments (n= 10). P values were analyzed by Student's T Test and differences between genotypes are statistically significant at \*,  $P < 0.05$ ; \*\*,  $P < 0.01$ ) \*\*\*,  $P < 0.001$ .

**Fig. 6: Regulation of lateral root development by  $\text{NO}_3^-$  and NRT1.1 relies on the TAR2 auxin biosynthesis enzyme.**

Root system architecture of Col, *chl1-5*, *tar2c* and *chl1-5tar2c* plants grown for 12 days in vertical agar plates on a basal nutrient medium deprived of N or supplemented with either 1 mM  $\text{NO}_3^-$  or 0.5 mM Gln.

(A) Representative pictures of Col, *chl1-5*, *tar2-c* and *tar2-c*  $\times$  *chl1-5* plants. White bar represents 0,5 cm.

(B) Mean length the primary root. Results (n=100) are representative of 5 independent experiments. P values were analyzed by Student's T Test and differences between mutant and wild-type genotypes are statistically significant at \*\*\*,  $P < 0.001$  (t test).

(C) Mean density of visible lateral roots ( $> 0,5$  mm). Results ( $n=100$ ) are representative of 5 independent experiments. P values were analyzed by Student's T Test and differences between mutant and wild-type genotypes are statistically significant at \*\*\*,  $P < 0.001$  (t test).

**Fig. 7: Schematic model of the integrated repressive role of NRT1.1 on lateral root development.**

NRT1.1 coordinately controls three separate auxin-related processes for repressing LRP development at low  $\text{NO}_3^-$  availability. First, NRT1.1 facilitates the basipetal transport of auxin out of the LRP (Krouk et al. 2010). Second NRT1.1 represses local auxin biosynthesis by downregulation the expression of the *TAR2* auxin biosynthetic gene, which reduces the acropetal flux of auxin to the LRP. These two actions are expected to lower auxin accumulation at the LRP apex that is required for normal LRP development and growth. Third, NRT1.1 maintains a mechanical resistance to the LRP outgrowth by repressing *LAX3* in the tissues overlying the LRP, thereby preventing cell wall remodeling in these tissues. All three repressive effects of NRT1.1 are suppressed by high  $\text{NO}_3^-$  availability ( $>0.5$  mM), leading to the local  $\text{NO}_3^-$ -induced stimulation of LRP development and emergence.






Figure 1.



Figure 2




Figure 3




Figure 4




Figure 5




Figure 6






Table 1. Primers used for qRT-PCR.

Description	AGI	Primer	Sequence (5'-3')
CLATHRIN	AT4G24550	Forward	AGCATACACTGCGTGCAAAG
		Reverse	TCGCCTGTGTACATATCTC
AUX1	AT2G38120	Forward	GAAGCCACCGTTCTTTATGC
		Reverse	GCTGCAGCTGGTTTACATTG
LAX3	AT1G77690	Forward	TTGTGAGGAGGGCATTGGTA
		Reverse	GGAGAGGCCACCGAGAGT
ABCB1	AT2G36910	Forward	CGACGGGAAAGACATAAGGA
		Reverse	CTGGTAGCGCGGATATGAAT
ABCB4	AT2G47000	Forward	CGTGTCGTGCTTTCTTTCT
		Reverse	ATCCAAAGGTGTTGCTGCTC
ABCB19	AT3G28860	Forward	CCAGCAAGAGGGTCGTAAAA
		Reverse	GAGACGATCCGTGGAGACAT
PIN1	AT1G73590	Forward	TACTCCGAGACCTTCCAACACTACG
		Reverse	TCCACCGCCACCACTTCC
PIN2	AT5G57090	Forward	CCTCGCCGCACTCTTTCTTTGG
		Reverse	CCGTACATCGCCCTAAGCAATGG
PIN3	AT1G70940	Forward	GAGGGAGAAGGAAGAAAGGGAAC
		Reverse	CTTGGCTTGTAATGTTGGCATCAG
PIN4	AT2G01420	Forward	TTGTCTCTGATCAACCTCGAAA
		Reverse	ATCAAGACCGCCGATATCAT
PIN7	AT1G23080	Forward	CGGCTGATATTGATAATGGTGTGG
		Reverse	GCAATGCAGCTTGAACAATGG
LDB29	AT3G58190	Forward	GCACATTGGCATTAGGCTTT
		Reverse	CTCAATCCCCACCTCAGCTA
TAR2	AT4G24670	Forward	CAGATTTGGCTTACTATTTGGCCACAG
		Reverse	GTCTTTCACCAAAGCCCATCCCAATC