



HAL
open science

Exogenous application of salicylic acid and gibberellic acid on biomass accumulation, antioxidant and anti-inflammatory secondary metabolites production in multiple shoot culture of *Ajuga integrifolia* Buch. Ham. ex D.Don

Bilal Haider Abbasi, Muhammad Asad Ullah, Muhammad Nadeem, Duangjai Tungmunnithum, Christophe Hano

► **To cite this version:**

Bilal Haider Abbasi, Muhammad Asad Ullah, Muhammad Nadeem, Duangjai Tungmunnithum, Christophe Hano. Exogenous application of salicylic acid and gibberellic acid on biomass accumulation, antioxidant and anti-inflammatory secondary metabolites production in multiple shoot culture of *Ajuga integrifolia* Buch. Ham. ex D.Don. *Industrial Crops and Products*, 2020, 145, 10.1016/j.indcrop.2020.112098 . hal-02619747

HAL Id: hal-02619747

<https://hal.inrae.fr/hal-02619747>

Submitted on 21 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution-NonCommercial 4.0 International License

1 Research Article for *Industrial Crops & Products*

2 **Exogenous application of salicylic acid and gibberellic acid on biomass**
3 **accumulation, antioxidant and anti-inflammatory secondary metabolites**
4 **production in multiple shoot culture of *Ajuga integrifolia* Buch. Ham. ex**
5 **D.Don.**

6 Bilal Haider Abbasi ^{a b c d}*, Muhammad Asad Ullah^a, Muhammad Nadeem^a, Duangjai Tungmunnithum^{b c}
7 ^c and Christophe Hano^{b c}

8 ^aDepartment of Biotechnology, Quaid-i -Azam University, Islamabad-45320, Pakistan

9 ^bLaboratoire de Biologie des Ligneux et des Grandes Cultures (LBLGC), INRA USC1328, Université d'Orléans,
10 45067 Orléans Cedex 2, France

11 ^cCOSM'ACTIFS, Bioactifs et Cosmétiques, CNRS GDR3711, 45067 Orléans Cedex 2, France

12 ^dEA2106 Biomolécules et Biotechnologies Vegetales, Université François-Rabelais de Tours, Tours, France

13 ^eDepartment of Pharmaceutical Botany, Faculty of Pharmacy, Mahidol University, Bangkok 10400, Thailand

14 *Correspondance: bhabbasi@qau.edu.pk

15

16

17

18 **Abstract**

19 *Ajuga integrifolia* Buch. Ham. ex D.Don is pharmacologically important plant, commonly
20 known as Kori Booti. Phytochemicals of this plant are used to treat variety of illnesses, ranging
21 from mild tooth ache to malaria and inflammatory diseases. In present study, exogenous role of
22 plant signaling molecules such as salicylic acid (SA) and gibberellic acid (GA) were investigated
23 for enhanced phytochemistry and biological activities in multiple shoot culture of *A. integrifolia*.
24 Shoots were exposed to various concentrations of SA and GA, and data was collected after 21
25 days. Significant increase in biomass (**dry weight (DW): 17.9 g/L**) was observed in response to
26 GA (**5.0 mg/L**) as compared with SA and control. High performance liquid chromatography was
27 employed to quantify eleven major secondary compounds. Results indicated that salicylic acid
28 promoted phytochemical production in shoot cultures whereas gibberellic acid down regulated
29 their biosynthesis. Salicylic acid (150 μ M) was found optimum for maximum accumulation of
30 majority of compounds, while rosmarinic acid and caffeic acid levels were higher in response to
31 SA (75 μ M) and SA (300 μ M), respectively. *In vitro* antioxidant and anti-inflammatory activities
32 were significantly increased under SA (150 μ M) treatment suggesting a direct correlation in
33 phytochemical biosynthesis and their respective biological activities.

34 **Keywords:** *Ajuga integrifolia*; Salicylic acid; shoot culture; Anti-inflammatory activity;
35 Phytochemicals; UPLC.

36

37

38 1. Introduction

39 Diseases are traced back since the origin of mankind on this planet. Humans are trying to combat
40 illnesses by exploitation of nature's primary resources such as microbes, animal and plants.
41 Plants have been a major source of effective substances to cure myriads of illnesses since
42 prehistoric times (Duraipandiyan et al., 2006). One of the many important medicinal plants
43 (MPs) is *Ajuga integrifolia*, commonly known as *Ajuga bracteosa* (Synonym) in Asian region,
44 belongs to family Lamiaceae. Both terminologies i.e. *bracteosa* and *integirfolia* are
45 interchangeable that are used to indicate same plant. *A. integrifolia* is a perennial herb that grows
46 up 5–50 cm tall and is usually found at high altitudes such as Himalayan region including
47 Pakistan, China and Afghanistan (Barkatullah et al., 2015; Jan et al., 2014). *A. integrifolia* plant
48 extracts have previously been studied for their enormous health benefit. Phytotoxic potential of
49 n-hexan mediated *A. integrifolia* extract showed herbicidal activity against *Lemna minor*
50 (Rehman et al., 2015) whereas anticancer activity of methanolic extracts was observed against
51 MCF-7 and Hep-2 cell lines (Pal et al., 2014). Multiple studies have demonstrated anti-
52 inflammatory and anti-plasmodial potential of *A. integrifolia* plant extracts by employing *in vivo*
53 mice models (Chandel and Bagai, 2010; Chandel and Bagai, 2011; Gautam et al., 2011;
54 Kaithwas et al., 2012).

55 Due to overexploitation of *Ajuga* plant on huge pharmaceutical demand, it is at the edge of
56 extinction as it has already been declared as endangered plant in Pakistan (Park et al., 2017; Rani
57 et al., 2017; Saeed et al., 2017). Use of biotechnological approaches such as *in vitro* cultures,
58 micro-propagation and organogenesis could really provide a suitable platform for conservation
59 of critically endangered plant (Park et al., 2017; Raghu et al., 2006; Shinwari, 2010).
60 Establishment of *in vitro* shoot organogenesis is considerably feasible method for conservation
61 of plants by producing whole plant in relatively short time irrespective to environmental
62 fluctuations (Gurel et al., 2011; Verma et al., 2011). Shoot culture of *A. integrifolia* was
63 previously established for optimum secondary metabolites production (Ali et al., 2018).

64 The present study was designed to determine the potential effects of salicylic acid and gibberellic
65 acid on enhanced production of pharmaceutically important secondary metabolites. The treated
66 shoot cultures were evaluated based on biomass accumulation (dry weight), phenolic and
67 flavonoids production, antioxidant, anti-inflammatory potential and special emphasize on

68 correlation of biological activities with secondary metabolites production upon treatment.
69 Quantification of shoot culture metabolites was performed using high performance liquid
70 chromatography (HPLC).

71 **2. Materials and Methods**

72 **2.1. Explant Collection and Multiple Shoot Culture Establishment**

73 *In vitro* shoot culture of *A. integrifolia* was carried out using leaf as explant source. Wild grown
74 plants were collected from Department of Biotechnology, Quaid-i-Azam University, Pakistan.
75 Leafs were separated from whole plants and surface sterilized for initiation of *in vitro* cultures.
76 Surface sterilization was done using 0.1% mercuric chloride solution for 40 sec, followed by 1
77 min treatment of 70% ethanol. Thereafter, plant leafs were washed three times with autoclaved
78 distilled water. Leaf sections (~0.5 cm²) were further inoculated on MS media (Murashige and
79 Skoog, 1962) containing 3% w/v sucrose and 0.8% w/v agar as solidifying agent. To promote
80 shooting, MS media was additionally supplemented with 1.0 mg/L BAP (6-benzylaminopurine)
81 and pH of media was set at 5.6 – 5.7 prior to autoclave. The culture was grown in 100 mL
82 vessels under dark conditions at 25 ± 2°C for maximum biomass production and constantly
83 transferred to fresh media after 4 weeks.

84 **2.2. Treatment of Salicylic Acid and Gibberellic Acid**

85 Basal culture of *A. integrifolia* established under dark conditions was used as inoculum to check
86 the effects of salicylic acid and gibberellic acid. Briefly, 1.0 g of shoots inoculum (3-4 shoots)
87 was added in previously used MS hormonal media additionally supplemented with various
88 concentrations of SA (75 µM, 150 µM, 300 µM) and GA (1.0 mg/L, 2.5 mg/L, 5.0 mg/L).
89 Cultures without any treatment of SA or GA were considered as control. Cultures were placed
90 under florescent light photoperiod conditions (16 h light/ 8 h dark) at 25 ± 2°C for optimum
91 shoot growth. Data was recorded after 21 days for maximum production of biomass and
92 secondary metabolites.

93 **2.3. Analytical Scheme**

94 After harvesting at 21 day, cultures were dried over night at 45°C for estimation of Dry weight
95 (DW). Dried shoot samples were subjected to extraction procedure according to (Zahir et al.,
96 2014). Briefly, 100 mg dried sample was dissolved in methanol (500 µl) by vortexing (5 min)
97 and sonication (30 min) at room temperature. The procedure was repeated twice, followed by
98 centrifugation at 15,000 rpm for 12 min to remove supernatant. Pallet was discarded and
99 supernatant was collected and stored at 4°C in eppendorf tube.

100 **2.4. Phytochemical Estimation**

101 **2.4.1. Phenolic and Flavonoids**

102 (Singleton and Rossi, 1965) protocol was employed for total phenolic content (TPC) estimation
103 using Folin–Ciocalteu’s reagent (FC) with minor adjustments. Briefly, extracted sample (20 µl)
104 from each treatment was thoroughly mixed with Na₂CO₃ (90 µl) and FC reagent solution (90 µl)
105 at room temperature in 96 well microplate. Mixture was incubated for 5 min prior to taking
106 absorbance using microplate reader at 630 nm. Gallic acid was employed as standard and results
107 were expressed as gallic acid equivalent. Phenolic production was calculated by following
108 formula,

$$109 \quad \text{Total phenolic production (mg/L)} = \text{DW (g/L)} \times \text{TPC (mg/g)}$$

110 Furthermore, total flavonoid contents (TFC) were calculated by (Ahmad et al., 2010) described
111 AlCl₃ colorimetric method. Sample extract (20 µl) was thoroughly mixed with potassium acetate
112 (10 µl) and aluminum chloride (10 µl) and dH₂O (160 µl) in 96 well microplate. Microplate
113 reader was used to measure absorbance of mixture at 415 nm after 30 min incubation time.
114 Quercetin was employed as standard and results were expressed as quercetin equivalent.
115 Flavonoid production was calculated by following formula,

$$116 \quad \text{Total flavonoid production (mg/L)} = \text{DW (g/L)} \times \text{TFC (mg/g)}$$

117 **2.4.2. Quantification of phytochemical via HPLC**

118 The separation, identification and quantification of the main metabolites accumulated in *A.*
119 *integrifolia* cultures were performed by HPLC. Separation was performed using the previously
120 established method allowing a good separation of caffeic acid derivatives and flavonoids, as

121 described in (Nazir et al., 2019). The separation was achieved using an Hypersil PEP 300 C18
122 column (250 × 4.6 mm, 5 μM) with a guard column Alltech (10 × 4.1 mm) at 35°C, by a Varian
123 high-performance liquid chromatography system (Varian Prostar 230 pump, Varian Prostar 410
124 autosampler and Varian Prostar 335 Photodiode Array Detector), driven by Galaxie version
125 1.9.3.2 software. The detection of compounds was initially set at 320 nm. The mobile phase
126 consisted of a mixture of two solvents: A composed of HCOOH/ H₂O, pH = 2.1 and B composed
127 of CH₃OH (HPLC-grade solvents). During the run, the mobile phase composition varied
128 according to the following nonlinear gradient: 8% B (0 min), 12% B (11 min), 30% B (17 min),
129 33% B (28 min), 100% B (30–35 min), 8% B (36 min) with a flow rate of 1mL/min. An
130 equilibration time of 10 min was applied after each individual run. The quantification was based
131 on the retention times and UV spectra of authentic commercial (Sigma-Aldrich). All samples
132 were analyzed in triplicate.

133 **2.5. Antioxidant Potential of *A. integrifolia***

134 **2.5.1. DPPH Scavenging Activity**

135 (Abbasi et al., 2010) protocol was used to determine free radical scavenging activity of shoot
136 extracts by employing DPPH reagent. Sample extract (20 μl) was thoroughly mixed with DPPH
137 solution (180 μl) in 96 well microplate, followed by 60 min incubation period in dark. The
138 final concentrations of ascorbic acid (5, 10, 20, 40 μg/ml) and DPPH (180 μl) with DMSO (20
139 μl) were taken as negative control. Absorbance was taken at 517 nm using microplate reader and
140 results were calculated using following formula,

$$141 \quad \text{Free radical scavenging activity (\%)} = 100 \times (1 - \text{AE}/\text{AD})$$

142 Herein, AE: Absorbance value with sample in mixture, AD: Absorbance value without sample in
143 mixture.

144 **2.5.2. Ferric Reducing Antioxidant Power (FRAP)**

145 (Benzie and Strain, 1996) method was employed to determine FRAP activity in shoot cultures of
146 *A. integrifolia*. Briefly, FRAP solution (190 μl) [composed of FeCl₃.6H₂O (20 mM); acetate
147 buffer (300 mM, pH 3.6) and TPTZ (10 mM); ratio 1:10:1 (v/v/v)] was mixed with sample

148 extract (10 μ l) from each treatment, followed by 15 min incubation time at room temperature.
149 Absorbance was taken at 630 nm using microplate reader. Results were expressed as TEAC
150 (Trolox C equivalent antioxidant capacity).

151 **2.5.3. Antioxidant ABTS Assay**

152 ABTS antioxidant activity of shoot cultures was measured according to (Tagliazucchi et al.,
153 2010) method. Briefly, potassium persulphate (2.45 mM) was mixed with ABTS salt (7 mM,
154 2,2-azinobis (3-ethylbenzthiazoline-6-sulphonic acid)) in equal proportion and incubated in dark
155 for 16 h to prepare ABTS solution. The absorbance was taken using microplate reader at 734 nm
156 and adjusted to 0.7. Extract was mixed in solution at room temperature and incubated in dark for
157 15 min, followed by absorbance at 734 nm. Results were expressed as TEAC (Trolox C
158 equivalent antioxidant capacity).

159 **2.6. Anti-inflammatory Activity of *A. integrifolia***

160 **2.6.1. COX-1 and COX-2 Inhibitory Activity**

161 COX-1 (Ovine) and COX-2 (human) assay kit was used according to manufacturer instructions
162 (701050, Cayman Chem. Co, Interchim, Montluçon, France) to determine inhibitory activity of
163 shoot extracts against COX-1 and COX-2. Arachidonic acid (1.1 mM) and ibuprofen (10 μ M)
164 were used as substrate and positive control respectively. The kit was used to measure COXs
165 peroxidase component. At 590 nm, Synergy II reader was used to check oxidized *N,N,N',N'*-
166 tetramethyl-*p*-phenylenediamine for 5 min in 96 well microplate.

167 **2.6.2. 15-LOX Inhibitory Activity**

168 Assay kit was used according to manufacturer instructions (760700, Cayman Chem. Co,
169 Interchim, Montluçon, France) to determine inhibitory activity of shoot extracts against 15-LOX.
170 Arachidonic acid (10 μ M) and nordihydroguariaretic acid (100 μ M) were used as substrate and
171 positive control inhibitor respectively. During lipooxygenation reaction, the hydroperoxides
172 concentration was measured by the kit using filtered soy 15-lipooxygenase standard in 10 mM
173 Tris-HCl buffer at 7.4 pH. At 940 nm, absorbance was taken using Synergy II reader in 96 well
174 microplate. After 5 min incubation of the inhibitor and enzyme, the absorbance was recorded

175 followed by incubation of 15 min after addition of the substrate and incubation for 5 min after
176 addition of chromogen.

177 **2.6.3. Secretory Phospholipase A2 (sPLA2) Inhibitory Activity**

178 Assay kit was used according to manufacturer instructions (10004883, Cayman Chem. Co,
179 Interchim, Montluçon, France) to determine inhibitory activity of shoot extracts against sPLA2.
180 Diheptanoyl thio-PC (1.44 mM) was employed as substrate whereas thiotheramide-PC (100 µM)
181 was considered, in this reaction, as positive control inhibitor. The cleavage of diheptanoyl thio-
182 PC ester releases free thiols which was measured by Synergy II reader (BioTek Instruments,
183 Colmar, France) at 420 nm in a 96-well microplate using DTNB (5-5'-dithio-bis-(2-nitrobenzoic
184 acid). Results were expressed as % inhibition by using following formula,

$$185 \quad \% \text{ Inhibition} = [(IA - \text{Inhibitor})/IA] \times 100$$

186 Here, Inhibitor: denotes enzymatic activity in the presence of inhibitor; IA: 100% enzymatic
187 activity without inhibitor presence.

188 **2.7. Statistical Analysis**

189 Whole experiment was conducted under controlled (temperature and light) conditions, performed
190 three times and repeated twice. Mean and standard deviations were calculated using Microsoft
191 Excel and graphs were prepared by Origin Pro 8.5.

192 **3. Results and Discussion**

193 **3.1. Biomass Production in *A. integrifolia* Shoot Culture**

194 Growth pattern in response to various concentrations of salicylic acid and gibberellic acid in *A.*
195 *integrifolia* shoots was measured by calculating respective biomass accumulation i.e. dry weight.
196 Data was recorded after 21 days treatment and compared with control (non-treated).
197 Morphological changes in shoot formation were also noted (Fig. 1). Salicylic acid produced
198 relatively mature and compact shoots compared with gibberellic acid and control. Overall,
199 gibberellic acid stimulated biomass accumulation rather than salicylic acid. Higher biomass
200 (DW: 17.9 g/L) was accumulated in GA (5.0 mg/L) treated shoot cultures as compared with
201 control (DW: 11.6 g/L) (Table 1). Effect of gibberellic acid concentration was found to be in

202 direct relation with biomass production whereas biomass was decreased with increase in salicylic
203 acid concentration. Maximum dry weight was observed at lowest concentration of salicylic acid
204 cultures as previously been reported (Sivanandhan et al., 2013) in *Withania somnifera* liquid
205 shoot culture. Similar trend was reported in salicylic acid treated shoot cultures of two
206 *Hypericum* species (Coste et al., 2011). Gibberellic acid mediated signaling in plants for shoot
207 formation and leaf shape has extensively been studied in the past (Davière and Achard, 2013;
208 Pramanik et al., 2018). The rapid cell division and cell elongation in response to gibberellic acid
209 could be the reason behind immature shooting after 21 days and enhanced biomass accumulation
210 (Gupta and Chakrabarty, 2013) (Fig. 1&Table 1). On the other hand, growth promoting effect of
211 exogenous salicylic acid has been studied in maize, wheat and soybean (Gunes et al., 2007;
212 Gutiérrez-Coronado et al., 1998; Shakirova et al., 2003). Lower concentrations of SA promoted
213 shooting and rooting in wheat and soybean plants whereas higher concentrations i.e. 250 µM SA
214 showed negative effect on chamomile plantlets (Kováčik et al., 2009).

215 **3.2. Phytochemical Production in *A. integrifolia* Shoot Cultures**

216 Plants tend to produce variety of biomolecules as protection mechanism against environmental
217 stimuli either biotic or abiotic. These phytochemicals mainly consists of phenolics and
218 flavonoids which help plants in smooth growth and development (Ali et al., 2006; Tan et al.,
219 2004). Current study involves the investigation of phytochemical's accumulation in response to
220 various concentrations of salicylic acid and gibberellic acid in *in vitro* derived multiple shoot
221 cultures of *A. integrifolia*. Highest phenolic contents (TPC: 11.08 mg/g) were observed against
222 SA 150 µM shoot cultures. Overall, salicylic acid promoted phenolics biosynthesis compared
223 with gibberellic acid except GA (2.5 mg/L) which showed relatively favorable results (TPC:
224 10.55 mg/g) compared with control. Since phenolic production was calculated based on dry
225 weight and TPC values, phenolic production was found optimum (TPP: 173.26 mg/L) in
226 gibberellic acid (GA: 5.0 mg/L) treated shoot cultures which is almost 1.5 fold higher than
227 control (TPP: 118.87 mg/L) (Fig. 2a). Optimum flavonoid contents (TFC: 1.85 mg/g) were noted
228 in SA 300 µM supplemented shoot cultures compared with control (TFC: 1.34 mg/g). Flavonoid
229 biosynthesis was observed to be dose dependent i.e. higher SA or GA concentration, higher
230 would be the flavonoid value. Similarly, flavonoid production (TFP: 28.5 mg/L) was found
231 optimum in GA (5.0 mg/L) shoot cultures compared with rest of treatments (Fig. 2b). Previously,

232 (Pérez-Tortosa et al., 2012) studied the effects of salicylic acid on enhanced production of
233 rosmarinic acid and antioxidant activity in shoots of *Thymus membranaceus*. Similarly, (Coste et
234 al., 2011) reported significant increase in hypericin and pseudohypericin production in
235 salicylic acid treated shoot cultures of *Hypericum hirsutum* and *Hypericum maculatum*.

236 **3.3. Antioxidant Evaluation in Multiple Shoot Culture of *A. integrifolia***

237 Plants produce cellular DNA damaging reactive oxygen species (ROS) such as H₂O₂, superoxide
238 and free radicals, when they endure environmental fluctuations or pathogenic attack (Moran et
239 al., 1994). Such stressed conditions could prevent plant's normal growth and development
240 processes through oxidative damage mechanism (Alexieva et al., 2001; Yu and Anderson, 1997).
241 To cope up with stress, plants produce large variety of secondary metabolites such as phenolics,
242 flavonoids, terpenoids etc. These plant biochemicals play vital role in mitigation of ROS by
243 acting as natural defense barrier (Ashry and Mohamed, 2011; Mittler, 2002; Samuolienė et al.,
244 2010). In current study, various *in vitro* based antioxidant assays were studied to investigate and
245 evaluate antioxidant capacity of *A. integrifolia* shoot extracts. These assays include DPPH (2,2-
246 diphenyl-1-picrylhydrazyle), FRAP (ferric reducing antioxidant power) and ABTS (2,2-azino-
247 bis-3-ethylbenzothiazoline-6-sulphonic acid). FRAP and ABTS works based on ET (electron
248 transfer) and HAT (hydrogen atom transfer) mechanism, respectively whereas DPPH is mixed
249 type assay, based on both mechanisms. Maximum DPPH activity (89.3%) was recorded in shoot
250 culture of *A. integrifolia* treated with salicylic acid (SA 150 µM) followed by SA 300 µM (87%)
251 compared with control (83.9%). Induction of gibberellic acid (GA) in shoot culture media
252 significantly reduced DPPH activity as compared with SA and control (Fig. 3a). Similarly, SA
253 150 µM treated shoot cultures showed optimum FRAP (532 µM) activity compared with rest of
254 treatments (Fig. 3b) whereas ABTS activity was found higher (435.5 µM) in SA 300 µM treated
255 shoots. Gibberellic acid negatively regulated antioxidant activities i.e. FRAP and ABTS in shoot
256 culture of *A. integrifolia*. (López-Orenes et al., 2013) studied the effects of exogenously applied
257 salicylic acid on antioxidant potential in shoots of *Cistus heterophyllus*. Results indicated that
258 lower concentration of salicylic acid was found effective in phenolics metabolism and
259 antioxidant activity, by triggering indigenous salicylic acid production in *C. heterophyllus*
260 shoots. Results of our study revealed a positive correlation in phytochemical biosynthesis and
261 antioxidant activity as previously studied (Pérez-Tortosa et al., 2012) in shoot cultures of *Thymus*

262 *membranaceus*. Higher level of phenolic contents in response to SA treatment could possibly be
263 the reason of optimum antioxidant activities in *A. integrifolia* shoot culture. Similar dependent
264 correlation of phytochemical and *in vitro* antioxidant activities have been observed in past
265 studies (Khan et al., 2018; Shah et al., 2019; Ullah et al., 2019a; Ullah et al., 2019b).

266 **3.4. Anti-Inflammatory Potential of *A. integrifolia* Multiple Shoot Culture**

267 Anti-inflammatory activity is mainly evaluated by means of % inhibition of different enzymes
268 responsible for inflammation process including cyclooxygenases (COX-1, COX-2), 15-
269 lipooxygenase (15-LOX) and secretory phospholipase A2 (sPLA2) (Mohale et al., 2014). These
270 enzymes are the key targets of non-steroidal anti-inflammatory drugs to reduce inflammation in
271 response to harmful stimuli. Plants ability to inhibit these enzymes through phytochemicals,
272 mainly flavonoids, shows their therapeutic role in development of alternative medicines
273 (Tungmunnithum et al., 2018). Anti-inflammatory activity of *A. integrifolia* shoot extracts was
274 measured using *in vitro* % inhibition assays of these enzymes. Optimum inhibitory activity was
275 recorded against sPLA2 ($39.51 \pm 1.04\%$), COX-1 ($34.44 \pm 0.93\%$), followed by 15-LOX (34.07
276 $\pm 0.37\%$) and COX-2 ($23.40 \pm 1.01\%$) in shoot extracts treated with salicylic acid (SA 150 μ M).
277 Overall, salicylic acid showed stimulatory effect on anti-inflammatory activity in *A. integrifolia*
278 shoots whereas gibberellic acid down regulated its activity compared with control. Anti-
279 inflammatory activity results in terms of % inhibition are shown in Table 2. Comparative
280 analysis of anti-inflammatory activities and phytochemical production in response to SA and GA
281 suggested that these phytochemicals play vital role in executing enzymatic inhibition that are
282 responsible for inflammation in the body (Guo et al., 2008; Lim et al., 2009; Szopa et al., 2018).
283 Anti-inflammatory potential of *A. integrifolia* has previously been studied using *in vitro* and *in*
284 *vivo* rat models (Hsieh et al., 2011; Kayani et al., 2016; Singh et al., 2012). Gautam et al
285 (Gautam et al., 2011) also reported COX-1 and COX-2 inhibition activity of *A. integrifolia* using
286 whole plant ethanol extract. This is the first study, evaluating enhanced anti-inflammatory
287 activities in *in vitro* derived *A. integrifolia* shoot culture in response to SA and GA.

288 **3.5. Phytochemical Quantification of *A. integrifolia***

289 For quantitative analysis of individual compounds present in shoots of *A. integrifolia*, an
290 effective HPLC method was employed. These phytochemicals actively form plant's defense

291 response against harsh environmental conditions to ensure smooth development (Balasundram et
292 al., 2006; Ullah et al., 2019a). In current study, 11 compounds were quantified in response to
293 salicylic acid and gibberellic acid treated shoot cultures. Overall, salicylic acid produced
294 optimum results whereas gibberellic acid down regulated phytochemicals production compared
295 with control. Optimum levels of chlorogenic acid (285.27 $\mu\text{g/g}$), apigenin (490 $\mu\text{g/g}$), luteolin
296 (493.61 $\mu\text{g/g}$), quercetin (542.3 $\mu\text{g/g}$), aucubin (187.09 $\mu\text{g/g}$), harpagoside (146.06 $\mu\text{g/g}$),
297 catalpol (91.67 $\mu\text{g/g}$) and 8-O-acetyl harpagoside (161.3 $\mu\text{g/g}$) were observed in shoot cultures in
298 response to SA 150 μM . On the contrary, higher levels of rosmarinic acid (569.8 $\mu\text{g/g}$) and
299 caffeic acid (332.9 $\mu\text{g/g}$) were accumulated against SA 75 μM and SA 300 μM treatments,
300 respectively (Table 3). Maximum production of harpagide (375.2 $\mu\text{g/g}$) was noted in untreated
301 (control) shoot cultures of *A. integrifolia*. Stimulatory effects of salicylic acid have previously
302 been reported in different *in vitro* derived cultures (Gadzovska et al., 2013; Mendoza et al., 2018;
303 Nadeem et al., 2019). (Largia et al., 2015) studied the enhanced bacoside A production in
304 salicylic acid treated *in vitro* derived shoot cultures of *Bacopa monnieri* L. Similar results were
305 reported by (López-Orenes et al., 2013) and (Pérez-Tortosa et al., 2012) in *Cistus heterophyllus*
306 and *Thymus membranaceus* shoot cultures against salicylic acid treatment. Negative effect of
307 gibberellic acid has been reported by Corbin et al (Corbin et al., 2013) in cell culture of *Linum*
308 *usitatissimum* L. Gibberellic acid significantly down regulated *LuPLR1* gene expression which is
309 responsible of biosynthesis of anti-cancerous lignan in flex. Phytochemicals, quantified in
310 current study, have huge pharmaceutical importance. They paly vital role as anticancer agents
311 through induction of apoptosis, oncogenes suppression and interruption of intracellular
312 communication (Mohammad Nabavi et al., 2015; White et al., 1989).

313 **4. Conclusion**

314 In current study, stimulatory effects of various concentrations of salicylic acid and gibberellic
315 acid on biomass and phytochemical synthesis were investigated in *in vitro* derived multiple shoot
316 culture of *Ajuga integrifolia*. Gibberellic acid showed maximum biomass accumulation by
317 stimulating shoot growth compared with salicylic acid and control. Salicylic acid proved to be
318 effect for total phenolic and flavonoid contents whereas phytochemical production was
319 significantly high in case of gibberellic acid treatment. *In vitro* based antioxidant and anti-
320 inflammatory activity were also found higher under the influence of salicylic acid which showed

321 positive correlation with plant secondary compounds. A significant increase in these
322 pharmaceutically important compounds was observed by quantification of high performance
323 liquid chromatography.

324 **Conflict of Interest**

325 The authors declare no conflict of interest.

326 **Acknowledgments**

327 B.H.A. acknowledges research fellowship of Le Studium-Institute for Advanced Studies, Loire
328 Valley, Orléans, France. D.T. gratefully acknowledges the support of the French government via
329 the French Embassy in Thailand in the form of Junior Research Fellowship Program 2018.

330 **References**

- 331 Abbasi, B.H., Khan, M.A., Mahmood, T., Ahmad, M., Chaudhary, M.F., Khan, M.A., 2010. Shoot
332 regeneration and free-radical scavenging activity in *Silybum marianum* L. Plant Cell Tiss. Org. (PCTOC)
333 101, 371-376.
- 334 Ahmad, N., Fazal, H., Abbasi, B.H., Rashid, M., Mahmood, T., Fatima, N., 2010. Efficient regeneration and
335 antioxidant potential in regenerated tissues of *Piper nigrum* L. Plant Cell Tiss. Org. (PCTOC) 102, 129-134.
- 336 Alexieva, V., Sergiev, I., Mapelli, S., Karanov, E., 2001. The effect of drought and ultraviolet radiation on
337 growth and stress markers in pea and wheat. Plant Cell Environ. t 24, 1337-1344.
- 338 Ali, H., Khan, M.A., Kayani, W.K., Khan, T., Khan, R.S., 2018. Thidiazuron regulated growth, secondary
339 metabolism and essential oil profiles in shoot cultures of *Ajuga bracteosa*. Ind. Crop Prod. 121, 418-427.
- 340 Ali, M.B., Khatun, S., Hahn, E.-J., Paek, K.-Y., 2006. Enhancement of phenylpropanoid enzymes and lignin
341 in *Phalaenopsis orchid* and their influence on plant acclimatisation at different levels of photosynthetic
342 photon flux. Plant Growth Regul. 49, 137-146.
- 343 Ashry, N.A., Mohamed, H.I., 2011. Impact of secondary metabolites and related enzymes in flax
344 resistance and or susceptibility to powdery mildew. World J. Agric. Sci 7, 78-85.
- 345 Balasundram, N., Sundram, K., Samman, S., 2006. Phenolic compounds in plants and agri-industrial by-
346 products: Antioxidant activity, occurrence, and potential uses. Food chem. 99, 191-203.
- 347 Barkatullah, I.M., Rauf, A., Hadda, T.B., Mubarak, M., Patel, S., 2015. Quantitative ethnobotanical survey
348 of medicinal flora thriving in Malkand Pass Hills, Khyber PakhtunKhwā Pakistan. J. Ethnopharmacology
349 169, 335-346.

350 Benzie, I.F., Strain, J.J., 1996. The ferric reducing ability of plasma (FRAP) as a measure of “antioxidant
351 power”: the FRAP assay. *Anal. Biochem.* 239, 70-76.

352 Chandel, S., Bagai, U., 2010. Antiplasmodial activity of *Ajuga bracteosa* against *Plasmodium berghei*
353 infected BALB/c mice. *Indian J. Med. Res.* 131, 440.

354 Chandel, S., Bagai, U., 2011. Screening of antiplasmodial efficacy of *Ajuga bracteosa* Wall ex. Benth.
355 *Parasitol. Res.* 108, 801-805.

356 Corbin, C., Renouard, S., Lopez, T., Lamblin, F., Lainé, E., Hano, C., 2013. Identification and
357 characterization of cis-acting elements involved in the regulation of ABA-and/or GA-mediated LuPLR1
358 gene expression and lignan biosynthesis in flax (*Linum usitatissimum* L.) cell cultures. *J. Plant Physiol.*
359 170, 516-522.

360 Coste, A., Vlase, L., Halmagyi, A., Deliu, C., Coldea, G., 2011. Effects of plant growth regulators and
361 elicitors on production of secondary metabolites in shoot cultures of *Hypericum hirsutum* and
362 *Hypericum maculatum*. *Plant Cell, Tiss. Org. (PCTOC)* 106, 279-288.

363 Davière, J.-M., Achard, P., 2013. Gibberellin signaling in plants. *Development* 140, 1147-1151.

364 Duraipandiyan, V., Ayyanar, M., Ignacimuthu, S., 2006. Antimicrobial activity of some ethnomedicinal
365 plants used by Paliyar tribe from Tamil Nadu, India. *BMC Complem. Altern. M.* 6, 35.

366 Gadzovska, S., Maury, S., Delaunay, A., Spasenoski, M., Hagège, D., Courtois, D., Joseph, C., 2013. The
367 influence of salicylic acid elicitation of shoots, callus, and cell suspension cultures on production of
368 naphthodianthrones and phenylpropanoids in *Hypericum perforatum* L. *Plant Cell, Tiss. Org. (PCTOC)* 113,
369 25-39.

370 Gautam, R., Jachak, S.M., Saklani, A., 2011. Anti-inflammatory effect of *Ajuga bracteosa* Wall Ex Benth.
371 mediated through cyclooxygenase (COX) inhibition. *J. Ethnopharmacology* 133, 928-930.

372 Gunes, A., Inal, A., Alpaslan, M., Eraslan, F., Bagci, E.G., Cicek, N., 2007. Salicylic acid induced changes on
373 some physiological parameters symptomatic for oxidative stress and mineral nutrition in maize (*Zea*
374 *mays* L.) grown under salinity. *J. Plant Physiol.* 164, 728-736.

375 Guo, L.Y., Hung, T.M., Bae, K.H., Shin, E.M., Zhou, H.Y., Hong, Y.N., Kang, S.S., Kim, H.P., Kim, Y.S., 2008.
376 Anti-inflammatory effects of schisandrin isolated from the fruit of *Schisandra chinensis* Baill. *Eur. J.*
377 *Pharmacol.* 591, 293-299.

378 Gupta, R., Chakrabarty, S., 2013. Gibberellic acid in plant: still a mystery unresolved. *Plant Sign. Behav.*
379 8, e25504.

380 Gurel, E., Yucesan, B., Aglic, E., Gurel, S., Verma, S.K., Sokmen, M., Sokmen, A., 2011. Regeneration and
381 cardiotonic glycoside production in *Digitalis davisiana* Heywood (*Alanya Foxglove*). *Plant Cell Tiss. Org.*
382 (PCTOC) 104, 217-225.

383 Gutiérrez-Coronado, M.A., Trejo-López, C., Larqué-Saavedra, A., 1998. Effects of salicylic acid on the
384 growth of roots and shoots in soybean. *Plant Physiol. Bioch.* 36, 563-565.

385 Hsieh, W.-T., Liu, Y.-T., Lin, W.-C., 2011. Anti-inflammatory properties of *Ajuga bracteosa* in vivo and in
386 vitro study and their effects on mouse model of liver fibrosis. *J. Ethnopharmacology* 135, 116-125.

387 Jan, M., Singh, S., Kaloo, Z., Maqbool, F., 2014. Medicinal importance of *Ajuga bracteosa* wall ex Benth.
388 A review. *Inter J Adv Res* 2, 389-394.

389 Kaithwas, G., Gautam, R., Jachak, S.M., Saklani, A., 2012. Antiarthritic effects of *Ajuga bracteosa* Wall ex
390 Benth. in acute and chronic models of arthritis in albino rats. *Asian Pac. J. Trop. Med.* 2, 185-188.

391 Kayani, W.K., Dilshad, E., Ahmed, T., Ismail, H., Mirza, B., 2016. Evaluation of *Ajuga bracteosa* for
392 antioxidant, anti-inflammatory, analgesic, antidepressant and anticoagulant activities. *BMC Complem.*
393 *Altern. M.* 16, 375.

394 Khan, T., Ullah, M.A., Garros, L., Hano, C., Abbasi, B.H., 2018. Synergistic effects of melatonin and
395 distinct spectral lights for enhanced production of anti-cancerous compounds in callus cultures of
396 *Fagonia indica*. *J. Photoch. Photobio. B*

397 Kováčik, J., Grúz, J., Bačkor, M., Strnad, M., Repčák, M., 2009. Salicylic acid-induced changes to growth
398 and phenolic metabolism in *Matricaria chamomilla* plants. *Plant Cell Rep.* 28, 135.

399 Largia, M.J.V., Pothiraj, G., Shilpha, J., Ramesh, M., 2015. Methyl jasmonate and salicylic acid synergism
400 enhances bacoside A content in shoot cultures of *Bacopa monnieri* (L.). *Plant Cell Tiss. Org. (PCTOC)* 122,
401 9-20.

402 Lim, Z.X., Ling, A.P.K., Hussein, S., 2009. Callus induction of *Ocimum sanctum* and estimation of its total
403 flavonoids content. *Asian J. Agri. Sci.* 1, 55-61.

404 López-Orenes, A., Martínez-Moreno, J.M., Calderón, A.A., Ferrer, M.A., 2013. Changes in phenolic
405 metabolism in salicylic acid-treated shoots of *Cistus heterophyllus*. *Plant Cell Tiss. Org. (PCTOC)* 113, 417-
406 427.

407 Mendoza, D., Cuaspud, O., Arias, J.P., Ruiz, O., Arias, M., 2018. Effect of salicylic acid and methyl
408 jasmonate in the production of phenolic compounds in plant cell suspension cultures of *Thevetia*
409 *peruviana*. *Biotech. Reports* 19, e00273.

410 Mittler, R., 2002. Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci.* 7, 405-410.

411 Mohale, D., Tripathi, A., Wahane, J., Chandewar, A., 2014. A Pharmacological Review on Cyclooxygenase
412 Enzyme. *Ind. J. Pharm. Pharmacol* 1, 46-58.

413 Mohammad Nabavi, S., Habtemariam, S., Daglia, M., Fazel Nabavi, S., 2015. Apigenin and breast cancers:
414 from chemistry to medicine. *Anti-Cancer Agents ME.* 15, 728-735.

415 Moran, J.F., Becana, M., Iturbe-Ormaetxe, I., Frechilla, S., Klucas, R.V., Aparicio-Tejo, P., 1994. Drought
416 induces oxidative stress in pea plants. *Planta* 194, 346-352.

417 Murashige, T., Skoog, F., 1962. A revised medium for rapid growth and bio assays with tobacco tissue
418 cultures. *Physiol. Plantarum* 15, 473-497.

419 Nadeem, M., Ahmad, W., Zahir, A., Hano, C., Abbasi, B.H., 2019. Salicylic acid-enhanced biosynthesis of
420 pharmacologically important lignans and neo lignans in cell suspension culture of *Linum ussitatissimum* L.
421 *Eng. Life Sci.* 19, 168-174.

422 Nazir, M., Tungmunnithum, D., Bose, S., Drouet, S., Garros, L., Giglioli-Guivarc'h, N., Abbasi, B.H., Hano,
423 C., 2019. Differential production of phenylpropanoid metabolites in callus cultures of *Ocimum basilicum*
424 L. with distinct in vitro antioxidant activities and in vivo protective effects against UV stress. *J. Agr. Food*
425 *Chem.*

426 Pal, A., Toppo, F.A., Chaurasiya, P.K., Singour, P.K., Pawar, R.S., 2014. In-vitro cytotoxicity study of
427 methanolic fraction from *Ajuga Bracteosa* wall ex. benth on MCF-7 breast adenocarcinoma and hep-2
428 larynx carcinoma cell lines. *Pharmacogn. Res.* 6, 87.

429 Park, H.Y., Kim, D.H., Sivanesan, I., 2017. Micropropagation of *Ajuga* species: a mini review. *Biotechnol.*
430 *Lett.* 39, 1291-1298.

431 Pérez-Tortosa, V., López-Orenes, A., Martínez-Pérez, A., Ferrer, M.A., Calderón, A.A., 2012. Antioxidant
432 activity and rosmarinic acid changes in salicylic acid-treated *Thymus membranaceus* shoots. *Food Chem.*
433 130, 362-369.

434 Pramanik, K., Pradhan, J., Sahoo, S.K., 2018. Role of auxin and gibberellins growth, yield and quality of
435 tomato: A review. *Int. J. Current Micro. App. Sci.* 6, 1624-1636.

436 Raghu, A., Geetha, S., Martin, G., Balachandran, I., Ravindran, P., 2006. Direct shoot organogenesis from
437 leaf explants of *Embelia ribes* Burm. f.: a vulnerable medicinal plant. *J. Forest Res-* 11, 57-60.

438 Rani, R., Khan, M.A., Kayani, W.K., Ullah, S., Naeem, I., Mirza, B., 2017. Metabolic signatures altered by
439 in vitro temperature stress in *Ajuga bracteosa* Wall. ex. Benth. *Acta physiol. Plant.* 39, 97.

440 Rehman, N.U., Begum, N., Ali, L., Al-Harrasi, A., Abbas, G., Ahmad, S., Khan, A.L., Shinwari, Z.K., Hussain,
441 J., 2015. Lipid peroxidation, Antiglycation, Cytotoxic, Phytotoxic, Antioxidant, Antiplatelet and
442 Antimicrobial activities of *Ajuga bracteosa* against various pathogens. *Pak J Bot* 47, 1195-1197.

443 Saeed, S., Ali, H., Khan, T., Kayani, W., Khan, M.A., 2017. Impacts of methyl jasmonate and phenyl acetic
444 acid on biomass accumulation and antioxidant potential in adventitious roots of *Ajuga bracteosa* Wall ex
445 Benth., a high valued endangered medicinal plant. *Physiol. Mol. Biol. Pla.* 23, 229-237.

446 Samuolienė, G., Brazaitytė, A., Urbonavičiūtė, A., Šabajevienė, G., Duchovskis, P., 2010. The effect of red
447 and blue light component on the growth and development of frigo strawberries. *Zemdirbyste-Agri.* 97,
448 99-104.

449 Shah, M., Ullah, M.A., Drouet, S., Younas, M., Tungmunnithum, D., Giglioli-Guivarc'h, N., Hano, C.,
450 Abbasi, B.H., 2019. Interactive Effects of Light and Melatonin on Biosynthesis of Silymarin and Anti-
451 Inflammatory Potential in Callus Cultures of *Silybum marianum* (L.) Gaertn. *Molecules* 24, 1207.

452 Shakirova, F.M., Sakhabutdinova, A.R., Bezrukova, M.V., Fatkhutdinova, R.A., Fatkhutdinova, D.R., 2003.
453 Changes in the hormonal status of wheat seedlings induced by salicylic acid and salinity. *Plant Sci.* 164,
454 317-322.

455 Shinwari, Z.K., 2010. Medicinal plants research in Pakistan. *J. Med. Plants Res.* 4, 161-176.

456 Singh, R., Patil, S., Pal, G., Ahmad, M., 2012. Evaluation of in vivo and in vitro anti-inflammatory activity
457 of *Ajuga bracteosa* Wall ex Benth. *Asian Pac. J. Trop. Dis.* 2, S404-S407.

458 Singleton, V.L., Rossi, J.A., 1965. Colorimetry of total phenolics with phosphomolybdic-phosphotungstic
459 acid reagents. *Am. J. Enol. Viticult.* 16, 144-158.

460 Sivanandhan, G., Rajesh, M., Arun, M., Jeyaraj, M., Dev, G.K., Arjunan, A., Manickavasagam, M.,
461 Muthuselvam, M., Selvaraj, N., Ganapathi, A., 2013. Effect of culture conditions, cytokinins, methyl
462 jasmonate and salicylic acid on the biomass accumulation and production of withanolides in multiple
463 shoot culture of *Withania somnifera* (L.) Dunal using liquid culture. *Acta physiol. Plant.* 35, 715-728.

464 Szopa, A., Dziurka, M., Warzecha, A., Kubica, P., Klimek-Szczykutowicz, M., Ekiert, H., 2018. Targeted
465 lignan profiling and anti-inflammatory properties of *Schisandra rubriflora* and *Schisandra chinensis*
466 extracts. *Molecules* 23, 3103.

467 Tagliacuzzi, D., Verzelloni, E., Bertolini, D., Conte, A., 2010. In vitro bio-accessibility and antioxidant
468 activity of grape polyphenols. *Food Chem.* 120, 599-606.

469 Tan, J., Bednarek, P., Liu, J., Schneider, B., Svatoš, A., Hahlbrock, K., 2004. Universally occurring
470 phenylpropanoid and species-specific indolic metabolites in infected and uninfected *Arabidopsis*
471 *thaliana* roots and leaves. *Phytochem.* 65, 691-699.

472 Tungmunnithum, D., Thongboonyou, A., Pholboon, A., Yangsabai, A., 2018. Flavonoids and other
473 phenolic compounds from medicinal plants for pharmaceutical and medical aspects: An overview.
474 *Medicines* 5, 93.

475 Ullah, M.A., Tungmunnithum, D., Garros, L., Drouet, S., Hano, C., Abbasi, B.H., 2019a. Effect of
476 Ultraviolet-C Radiation and Melatonin Stress on Biosynthesis of Antioxidant and Antidiabetic
477 Metabolites Produced in *In Vitro* Callus Cultures of *Lepidium sativum* L. *Int. J. Mol. Sci.* 20, 1787.

478 Ullah, M.A., Tungmunnithum, D., Garros, L., Hano, C., Abbasi, B.H., 2019b. Monochromatic lights-
479 induced trends in antioxidant and antidiabetic polyphenol accumulation in in vitro callus cultures of
480 *Lepidium sativum* L. *J. Photoch. Photobio. B.* 196, 111505.

481 Verma, S.K., YÜCESAN, B.B., CİNGÖZ, G., GÜREL, S., GÜREL, E., 2011. Direct shoot regeneration from leaf
482 explants of *Digitalis lamarckii*, an endemic medicinal species. *Turk. J. Bot.* 35, 689-695.

483 White, R.H., Worsham, A.D., Blum, U., 1989. Allelopathic potential of legume debris and aqueous
 484 extracts. *Weed Sci.* 37, 674-679.

485 Yu, T.-W., Anderson, D., 1997. Reactive oxygen species-induced DNA damage and its modification: a
 486 chemical investigation. *Mutat. Res-Fund. Mol. M.* 379, 201-210.

487 Zahir, A., Abbasi, B.H., Adil, M., Anjum, S., Zia, M., 2014. Synergistic effects of drought stress and
 488 photoperiods on phenology and secondary metabolism of *Silybum marianum*. *App. Biochem. Biotech.*
 489 174, 693-707.

490

491 **Figures & Tables:**

492

493 **Table 1.** Dry weight production in response to salicylic acid (SA) and gibberellic acid (GA) in multiple
 494 shoot culture of *A. integrifolia*.

Treatment	Concentration	dry weight (g/L)
Control	BAP 1.0 mg/L	11.66 ± 0.853
	SA 75 µM	14.21 ± 0.638
Salicylic Acid	SA 150 µM	14.90 ± 1.043
	SA 300 µM	13.66 ± 3.951
	GA 1.0 mg/L	15.25 ± 2.448
Gibberellic acid	GA 2.5 mg/L	16.27 ± 0.962
	GA 5.0 mg/L	17.97 ± 1.749

495

496

BAP: 6-benzylaminopurine

497

498

499

500 **Table 2.** Anti-inflammatory potential of *A. integrifolia* shoot cultures grown under different SA and GA
 501 treatments.

Treatment	% Inhibition			
	COX-1	COX-2	15-LOX	sPLA ₂
Control	32.37 ± 1.361	21.38 ± 0.997	29.20 ± 2.055	34.67 ± 0.443
SA 75 µM	31.10 ± 1.449	20.13 ± 1.204	32.35 ± 1.943	37.14 ± 1.394
SA 150 µM	34.44 ± 0.932	23.40 ± 1.015	34.07 ± 0.372	39.51 ± 1.044
SA 300 µM	31.77 ± 1.667	20.79 ± 1.743	33.58 ± 1.743	38.49 ± 1.493
GA 1.0 mg/L	17.83 ± 1.302	17.17 ± 0.846	23.59 ± 0.844	29.51 ± 1.943
GA 2.5 mg/L	16.22 ± 1.843	15.60 ± 1.563	22.71 ± 2.511	28.08 ± 1.338
GA 5.0 mg/L	32.37 ± 0.984	17.40 ± 1.337	22.68 ± 0.442	28.27 ± 1.437

502

503 SA: Salicylic Acid, GA: Gibberellic Acid, COX-1: Cyclooxygenase-1, COX-2: Cyclooxygenase-2, 15-LOX: 15-
 504 Lipooxygenase, sPLA₂: Secretory Phospholipase A2.

Table 3. HPLC based quantification of valuable metabolites in shoot culture of *A. integrifolia*.

Compounds	Treatments						
	BAP	Salicylic Acid (μM)			Gibberellic Acid (mg/L)		
	Control	75	150	300	1.0	2.5	5.0
Chlorogenic Acid	232.5 \pm 3.05	249.2 \pm 1.41	285.2 \pm 1.93	278.8 \pm 4.11	222.3 \pm 3.59	221.1 \pm 2.83	239.5 \pm 0.55
Caffeic Acid	196.2 \pm 6.04	269.4 \pm 3.95	316.2 \pm 4.22	332.9 \pm 2.01	193.8 \pm 1.87	230.2 \pm 8.55	185.4 \pm 2.79
Rosmarinic Acid	399.9 \pm 3.05	569.8 \pm 1.94	519.1 \pm 6.09	546.6 \pm 5.93	409.4 \pm 3.56	355.6 \pm 6.83	328.8 \pm 9.64
Apigenin	455.6 \pm 3.95	460.1 \pm 2.04	490.1 \pm 4.31	488.5 \pm 1.99	436.1 \pm 1.05	431.3 \pm 3.41	439.6 \pm 4.22
Luteolin	463.9 \pm 1.91	461.7 \pm 3.63	493.6 \pm 5.49	486.8 \pm 9.81	465.7 \pm 5.26	431.3 \pm 7.33	436.1 \pm 3.07
Quercetin	515.6 \pm 2.53	512.6 \pm 4.79	542.3 \pm 1.37	493.6 \pm 6.61	465.7 \pm 9.05	438.1 \pm 3.43	444.8 \pm 5.19
Harpagide	375.2 \pm 1.45	358.4 \pm 6.67	356.9 \pm 3.87	366.4 \pm 8.23	213.6 \pm 2.73	174.3 \pm 5.74	251.3 \pm 4.92
Aucubin	153.3 \pm 7.65	165.8 \pm 2.84	187.1 \pm 6.57	175.2 \pm 1.84	64.5 \pm 9.04	76.5 \pm 4.39	88.1 \pm 1.74
Harpagoside	129.5 \pm 5.54	112.1 \pm 8.51	146.1 \pm 1.84	117.4 \pm 7.48	81.9 \pm 3.82	71.2 \pm 5.44	57.8 \pm 1.63
Catalpol	74.9 \pm 3.10	73.5 \pm 7.45	91.6 \pm 2.29	65.8 \pm 4.84	39.6 \pm 8.85	46.2 \pm 3.48	29.4 \pm 9.57
8-O-Acetyl Harpagoside	153.3 \pm 2.04	141.6 \pm 4.51	161.3 \pm 1.23	144.9 \pm 5.43	88.4 \pm 8.74	75.6 \pm 3.66	67.9 \pm 6.73




Fig. 1. Effect of gibberellic acid (a: 1.0 mg/L, b: 2.5 mg/L, c: 5.0 mg/L) and salicylic acid (d: control, e: 75 μM, f: 150 μM, g: 300 μM) on biomass production in multiple shoot culture of *A. integrifolia*.




Fig. 2. Influence of SA and GA on a) Total phenolic content (TPC), Total phenolic production (TPP) b) Total flavonoid content (TFC), Total flavonoid production (TFP) in multiple shoot culture of *A. integrifolia*.




Fig. 3. *In Vitro* antioxidant activities in shoot culture of *A. integrifolia* in response to salicylic acid and gibberellic acid treatment **a)** DPPH activity, **b)** ABTS and FRAP activity. Values represent means \pm standard errors from triplicates.