



HAL
open science

Micro-food web interactions involving bacteria, nematodes, and mycorrhiza enhance tree P nutrition in a high P-sorbing soil amended with phytate

Mahafaka Ranoarisoa, Jean Trap, Anne-Laure Pablo, Damien Dezette, Claude Plassard

► To cite this version:

Mahafaka Ranoarisoa, Jean Trap, Anne-Laure Pablo, Damien Dezette, Claude Plassard. Micro-food web interactions involving bacteria, nematodes, and mycorrhiza enhance tree P nutrition in a high P-sorbing soil amended with phytate. *Soil Biology and Biochemistry*, 2020, 143, 10.1016/j.soilbio.2020.107728 . hal-02620477

HAL Id: hal-02620477

<https://hal.inrae.fr/hal-02620477>

Submitted on 7 Mar 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Micro-food web interactions involving bacteria, nematodes, and mycorrhiza enhance**
2 **tree P nutrition in a high P-sorbing soil amended with phytate**

3

4 **Authors**

5 Mahafaka Patricia Ranoarisoa¹, Jean Trap², Anne-Laure Pablo², Damien Dezette¹, Claude
6 Plassard^{1,*}

7

8 **Affiliations**

9 ¹Eco&Sols, INRAE, IRD, CIRAD, Montpellier SupAgro, Univ Montpellier, Montpellier,
10 France

11 ²Eco&Sols, IRD, INRAE, CIRAD, Montpellier SupAgro, Univ Montpellier, Montpellier,
12 France

13

14 *Corresponding author: Claude Plassard

15 Email address: claud.plassard@inra.fr

16 Full postal address: INRA, UMR Eco&Sols, 2 place Pierre Viala,
17 34060 Montpellier, Cedex 2, France

18

19 **Abstract**

20

21 Phytate is considered a poorly available plant P source but proved to be useful for particular
22 soil bacteria strains. In soil-free conditions, it has been shown that bacteria locked up the
23 mineralized phosphorus from phytate whereas bacterial grazers like nematodes were able to
24 deliver P to plants. Here, we aimed to determine if the interactions between phytate-
25 mineralizing bacteria, bacterial grazer nematodes, and mycorrhizal fungi could increase plant
26 P acquisition from phytate in high P-adsorbing soils. *Pinus pinaster* was grown in a Cambisol
27 supplemented with phytate. Plants, whether associated or not associated with the
28 ectomycorrhizal fungus *Hebeloma cylindrosporum*, were either inoculated or not inoculated
29 with the phytase-releasing bacteria *Bacillus subtilis* and the bacterial-feeding nematode
30 *Rhabditis* sp. After 100 days, the dual inoculation of bacteria and nematodes significantly
31 increased net plant P accumulation. We observed that, on average, mycorrhizal plants
32 accumulated more P in their shoots than non-mycorrhizal plants. However, the highest plant P
33 acquisition efficiency was found when the three soil organisms were present in the *P. pinaster*
34 rhizosphere. We conclude that, in a highly inorganic P-fixing soil, plant P acquisition from
35 phytate strongly depends on the grazing of phytate-mineralizing bacteria. Our results confirm
36 the importance of the soil microbial loop to improve plant P nutrition from phytate, which
37 should be considered a route to improve the utilization of this source of poorly available P by
38 plants.

39

40 **Keywords**

41 Phosphorus cycling; microbial loop; *Bacillus subtilis*; *Pinus pinaster*; *Hebeloma*
42 *cylindrosporum*; *Rhabditis* sp.

43

44

45 **1. Introduction**

46 Phosphorus (P) limits plant productivity in many terrestrial ecosystems (Holford, 1997;
47 Vitousek et al., 2010). Depending on the soil, P could be abundant as organic P (Po), with
48 compounds including monoesters, inositol phosphates, diesters and phosphonates. Although
49 depending on the environmental conditions, monoesters and diesters tend to be more labile
50 than the inositol phosphates which accumulate in soil (Darch et al., 2014; George et al.,
51 2018). A direct uptake of Po compounds by roots and mycorrhizal fungi has not been
52 documented excepted one report suggesting that non-mycorrhizal poplar (*Populus x*
53 *canescens*) and beech (*Fagus sylvatica*) roots might take up ADP and/or AMP by yet
54 unknown nucleotide transporters (Scheerer *et al.*, 2019). Hence, it is generally assumed that
55 plant roots take up P as inorganic P (Pi) in the form of free orthophosphate (HPO_4^{2-} or H_2PO_4^-
56 ; Bucher, 2007). The mineralization of Po is thus a key step in P cycling to supply available P
57 to plants (Marschner, 1995).

58 The soil Po pool can be mineralized by microbial enzymes, such as phosphatases and phytases
59 (Rodriguez & Fraga, 1999), into Pi potentially available for plants (Hayes, Simpson &
60 Richardson, 2000; Richardson, Hadobas & Hayes, 2000; Mullaney & Ullah, 2003). However,
61 large amounts of Pi released by microbial enzymes are immobilized in the microbial biomass
62 (Oberson et al., 2001; Pistocchi et al., 2018); they are strongly adsorbed by soil constituents
63 such as minerals (especially clays) and organic matter (Holford, 1997) or combined with
64 cations (Ca, Al, Fe). Therefore, only a small amount of Pi from organic P mineralization,
65 estimated to be in the 1–10 μM range (Hinsinger et al., 2011), will effectively supply the free
66 Pi pool. Consequently, two key ecological functions based on rhizosphere biological
67 interactions have evolved over time to increase the efficiency of plant P absorption: (i)
68 widespread mycorrhizal symbiosis between fungi and plant roots (Smith & Read, 2008) and

69 (ii) microbial loops involving interactions between bacteria and bacterial-feeders (protists and
70 nematodes) (Clarholm, 1985b).

71 In forest ecosystems, the mutualistic symbiosis between plants and ectomycorrhizal fungi is
72 one of the most important biological interactions involved in P acquisition by trees (Plassard
73 & Dell, 2010). The extraradical hyphae of ectomycorrhizal fungi are able to prospect a large
74 volume of soil, overcoming the Pi depletion zone created by plant roots (Becquer et al., 2014).
75 The fungal cells can also release carboxylates (Finlay, 2008) and enzymes (Colpaert & van
76 Laere, 1996; Courty et al., 2006; Louche et al., 2010) to increase P acquisition from soil
77 mineral and organic sources (Plassard et al., 2011). After uptake, P is transported along the
78 hyphae towards ectomycorrhizas as polyphosphates accumulated in mobile vacuoles (Nehls &
79 Plassard, 2018).

80 Among Po forms in soil, phytate (*myo*-inositol hexakisphosphate, $C_6H_{18}O_{24}P_6$) is considered
81 as the most abundant source (Alexander, 1977; Turner, 2007) but also as the poorest P source
82 to plants because most of the plant species are not able to release phytases (enzymes required
83 to release the Pi group from its ester bond) into their rhizosphere (Richardson et al., 2001). In
84 contrast, rhizosphere bacteria are able to produce phytases (Mullaney & Ullah, 2003; Turner,
85 2007) that could promote the release of Pi from phytate in the vicinity of plant roots.

86 Therefore, the inoculation of phytate-mineralizing bacteria in the rhizosphere of plants is
87 potentially highly beneficial for plant P nutrition (Richardson et al., 2001).

88 This effect could be strengthened when soil bacterial grazers are present in the rhizosphere of
89 plants as part of the soil microbial loop (Clarholm, 1985b). Bacterial feeders (mostly protists
90 and nematodes) release nutrients stored in the microbial biomass, increasing the activity of
91 decomposers (Kuikman et al., 1991, Jansen & Vanveen, 1991; Clarholm, 2005) and the
92 nutrients become available to plants (Bonkowski, 2004). Trap et al. (2016) showed that soil
93 bacterivores could significantly enhance plant P nutrition, up to 30%, through their grazing

94 activities. However, the increase in soil P availability induced by the interactions between
95 bacteria and bacterial-feeders has been reported to be slight (Anderson et al., 1978;
96 Darbyshire et al., 1994; Djigal et al., 2004a), high (Cole et al., 1978), or sometimes even null
97 (Griffiths, 1986). It is difficult to explain these discrepancies, mainly because the mechanisms
98 and the controlling factors by which bacterivores increase plant P acquisition have not been
99 identified clearly.

100 The experiment carried out by Irshad et al. (2012) in an agarose medium showed that the
101 presence of both bacterial-feeding nematodes and phytase-producing bacteria is required to
102 increase the P content in *Pinus pinaster* seedlings, probably by increasing the activity of
103 phytate-mineralizing bacteria. However, we do not know whether similar interactive effects
104 can be obtained in experiments using high P-sorbing soils rather than low-Pi-sorbing agarose.
105 In this study, we aimed to determine if the presence of the above-mentioned potential
106 rhizosphere mutualists in the rhizosphere, alone or together, promotes plant P acquisition
107 from phytate in a highly P-sorbing soil with a very low level of available P. For that purpose,
108 we grew *P. pinaster* seedlings, either associated or not associated with the ectomycorrhizal
109 fungus *Hebeloma cylindrosporum*, in a Cambisol amended with phytate, and either inoculated
110 or not inoculated with phytase-releasing bacteria (*Bacillus subtilis*) and bacterial-feeding
111 nematodes (*Rhabditis* sp.). We formulated four hypotheses. We first assumed that the plant
112 alone, without mycorrhiza nor free-living organisms, does not get P from phytate, as observed
113 in agarose (Irshad et al., 2012). Because mycorrhizal fungi are well-known to take up Pi not
114 available to roots, we expected higher plant P uptake when the ectomycorrhizal fungus is
115 inoculated, irrespective of the presence of free-living organisms (H2). We supposed that the
116 inoculation of free-living organisms will enhance phytate mineralization and plant P
117 availability (H3). Finally, we supposed that the plant P uptake from phytate would be
118 maximal when all potential rhizosphere mutualists are inoculated (H4).

119

120 **2. Materials and Methods**

121 **2.1. Plant and mycorrhizal synthesis**

122 Maritime pine (*P. pinaster*) seedlings were grown from seeds (Medoc, Landes-Sore-VG,
123 France) as described by Becquer et al. (2017). Briefly, the seeds were first soaked in
124 deionized water at 4 °C for 48 h before being sterilized in a 30% H₂O₂ solution for 30 minutes
125 and rinsed several times with 1 L of sterile deionized water. The seeds were deposited in a
126 solid medium (agar 1.5 g L⁻¹ and glucose 2 g L⁻¹) in Petri dishes (90 mm diameter) placed at
127 25 °C in the dark for 2–3 weeks. Petri dishes showing fungal or/and bacterial development
128 around the seeds were disposed of. Hence, we selected dishes containing actual sterile
129 germinated seeds to prepare plants in glass test tubes (see below and Figure 1A).

130 A dikaryotic strain of the ectomycorrhizal basidiomycete *H. cylindrosporum* (Debaud & Gay,
131 1987) was used to get ectomycorrhizal *P. pinaster*. The fungus was grown in the dark at 24
132 °C in a standard nitrate medium (Louche et al., 2010) for three weeks.

133 Glass test tubes (15 cm height) containing a filter paper (Whatman 1542–240 Hardened
134 Ashless Grade 542) inserted vertically inside the tube and 10 ml of a glucose solution (10 g L⁻¹)
135 were sterilized twice at an interval of 48 h (Becquer et al., 2017). The glucose solution of
136 each tube was then replaced by a nutritive solution containing 1 mM nitrate and 0.2 mM
137 phosphate (Irshad et al., 2011). Plants, whether inoculated or not with *H. cylindrosporum*,
138 were prepared with sterile germinated seeds having a root more than 2 cm long, but with their
139 external teguments still attached (Becquer et al., 2017). To prepare mycorrhizal (M) plants,
140 three small pieces of solid nitrate medium with *H. cylindrosporum* mycelium were placed on
141 the filter paper near the roots (Figure 1A and supplementary material 1). Non-mycorrhizal
142 plants (NM) were produced by placing germinated seedlings in the tubes without the fungus.
143 The tubes were kept for two months in a growth chamber (16/8 h light/dark cycle at 24/18 °C,

144 400 $\mu\text{mol m}^{-2} \text{s}^{-1}$, 400-700 nm, RH 80%). The level of the nutrient solution (15 ml per tube)
145 was topped up every week in sterile conditions.

146

147 **2.2. Bacterial strain and nematodes**

148 The bacterial strain used in the experiment was isolated from ectomycorrhizal roots collected
149 in a 15-year-old *P. pinaster* plot in the Landes Region, near Bordeaux (France). The strain
150 was identified as *Bacillus subtilis* (named 111b; Irshad et al., 2011) and was able to use
151 phytate as the sole source of P via the release of an alkaline phytase. For the inoculation
152 experiments, the bacteria were cultivated in a liquid medium, as described in Irshad *et al.*
153 (2011) for 4 d at 26 °C. The culture medium was centrifuged at 3500 g for 10 minutes and the
154 pellet was washed twice before being suspended in a volume of sterile deionized water
155 adjusted to get approximately 2 million bacteria per mL. Monoxenic populations of
156 nematodes (*Rhabditis* sp.) were isolated from the soil collected in the same plot (Landes,
157 France) and were cultured in *B. subtilis* (bacterial strain 111b) following the procedure
158 described previously (Irshad *et al.*, 2011). *Rhabditis* is a common, ubiquitous genus that is
159 classified in the first class along the colonizer-persistent gradient (cp) into the maturity index
160 (Bongers, 1990).

161

162 **2.3. Soil preparation and phytate addition**

163 The soil used for the co-inoculation experiment corresponded to the A layer of a chromic
164 cambisol collected in the south of France (Cazevieille, Hérault). It is a clay soil (48% clay, pH
165 in water = 6.9, total N = 2.5 g kg⁻¹, total P = 0.9 g kg⁻¹) with a very low concentration of
166 bicarbonate-extractable inorganic P (c.a. 3 mg kg⁻¹; Casarin et al., 2003). Before use, the soil
167 was dried at ambient temperature and sieved at 2 mm. Then, the soil was mixed with acid-
168 washed (HCl 1N) glass beads (50/50, w/w) to get homogenous humidification of the soil-bead

169 material during the experiment. The soil and glass bead mixtures were prepared for each
170 replicate by mixing them in a 50-ml polypropylene tube with 15 g of soil, 15 g of glass beads,
171 and 15 ml of a nutrient solution containing 9.3 mM KNO₃, 9.3 mM MgSO₄ 7H₂O, 18.6 mM
172 CaSO₄, 200 µg L⁻¹ thiamine hydrochloride, 2.3 ml L⁻¹ 1% Fe citrate, and 1 ml L⁻¹ solution of
173 micronutrients (Morizet & Mingeau, 1976). The soil was then autoclaved twice a week at 115
174 °C for 40 min.

175 A solution of phytate (9.24 g L⁻¹, inositol hexakisphosphate sodium salt from Sigma, ref
176 P0109) was prepared in sterile deionized water. Before sterilization by filtration (0.2 µm), the
177 pH of this solution was adjusted to 7 with HCl. After cooling the soil at room temperature,
178 each tube was opened in sterile conditions to add 46.2 mg of phytate (5 ml of the phytate
179 solution) and 350 µg of cholesterol (70 µl of cholesterol 5 mg L⁻¹, in ethanol). These amounts
180 of phytate and cholesterol were equal to those added to the agarose medium by Irshad et al.
181 (2012). The tubes were then shaken by inversion overnight to get a homogeneous soil
182 suspension.

183

184 ***2.4. Experimental design***

185 The experiment was carried out in sterile conditions in 12 x 12 cm square Petri dishes (Figure
186 1B-E and supplementary material 1). Each dish is first pierced with two holes, one in the lid
187 for watering and the other in a corner in order to allow the growth of the plant shoot outside
188 the dish and filled with the soil-bead-nutrient solution mixture from one polypropylene tube.
189 Two-month-old plants were removed from the test tubes and transferred to the Petri dishes.
190 Six biological treatments, corresponding to two levels of plant mycorrhizal status combined
191 with three inoculation treatments, were set up.
192 Each plant, associated (M) or not associated (NM) with the ectomycorrhizal basidiomycete *H.*
193 *cylindrosporum*, was (i) not inoculated with free-living organisms (None), (ii) inoculated with

194 the phytate-mineralizing bacteria *Bacillus subtilis* (+Bac), or (iii) inoculated with both the
195 bacteria and bacteria-feeding nematodes *Rhabditis* sp. (+Bac +Nem). The bacteria and the
196 nematodes were inoculated at a rate of approximately 10^6 bacteria and 300 nematodes per
197 dish, supplied in 0.5 ml solution each. Six replicates were set up for each biological treatment,
198 giving a total of 36 Petri dishes. The Petri dishes were placed horizontally in the same growth
199 chamber as described above and covered with aluminum foil to limit the light on the roots.
200 The plants were allowed to grow for 100 days. Each Petri dish was watered once a week with
201 sterile water to compensate for water loss during the experiment. At the beginning of the
202 experiment (time 0), 3 NM and 3 M plants were used to determine the initial values of root
203 and shoot dry weight and their P accumulation.

204

205 ***2.5. Plant and soil analysis***

206 At the end of the experiment, the shoots and the roots of each plant were separated. Intact root
207 systems were cleaned, then scanned and digitized using WinRHIZO
208 (<http://www.regentinstruments.com/products/rhizo/Rhizo.html>) to measure parameters such
209 as length, area, number of tips, number of forks, and number of crossings. The root and shoot
210 dry biomass were determined after oven-drying at 60 °C for 48 h. The total P content in the
211 shoots and roots was determined after crushing and mineralizing plant tissues with 36N
212 H₂SO₄ acid, as described by Aquino & Plassard (2004). The P concentration in the
213 mineralized solution was analyzed using colorimetry with malachite green (Ohno & Zibilske,
214 1991). Absorbance was measured at 630 nm. The P acquisition efficiency of the plants was
215 calculated by dividing the total plant P by the root length.
216 Furthermore, we collected as much soil-bead mixture as we could possibly could without
217 damaging the roots and used it (i) to check for the presence of living nematodes and (ii) to
218 quantify the abundance of 18S rDNA gene copies. Nematode survival rates were determined

219 by active filtering through fine cellulose wadding paper for 48 h and counted using a
220 binocular stereoscopic microscope (Southey, 1986). For each sample, microbial DNA was
221 extracted from 0.250 g of soil-bead mixture following the procedure developed by Tournier *et*
222 *al.* (2015). The quality of the extracted DNA was improved by adding 500 μ L of 5.5 M
223 guanidine thiocyanate during the seventh step of DNA extraction. Quantitative real-time PCR
224 was performed on 2.5 ng of extracted DNA to quantify fungal (18S) rDNA gene copies in
225 each sample. The PCR products were amplified with SYBR® Green Master Mix (BIORAD,
226 USA) using the primer pairs FR1 forward (AICCATTC AATCGGTAIT) and FF390 reverse
227 (CGATAACGAACGAGACCT) to estimate the number of 18S rDNA gene copies per gram
228 of soil. The thermocycling pattern used was as follows: 98 °C for 2 min; 39 cycles at 98 °C
229 for 5 s, 60 °C for 30 s, and 72 °C for 30 s; and a final elongation step at 55 °C for 5 s and 25
230 °C for 30 s.

231

232 ***2.6. Statistical analysis***

233 The mean and standard deviation ($n = 6$) for each measured variable were calculated for each
234 modality. Two-way analysis of variances was conducted to test the “mycorrhiza” and “free-
235 living organisms” factors and their interaction on plant variables. We conducted one-way
236 ANOVA followed by post hoc Tukey HSD test on all modalities as factors. The normality of
237 ANOVA residues was checked using the Shapiro-Wilk test. Using the Student t-test, we also
238 tested whether the final plant biomass and P amount values for each biological treatment were
239 different from the initial value. All statistical analyses were performed using the R software
240 (Team, 2016) with statistical significance at 5%.

241

242 **3. Results**

243 ***3.1. Plant biomass and root architecture***

244 At the end of the experiment, plant shoot and root dry biomasses were significantly higher
245 than those measured in two-month-old-plants (Figure 2). The two-way ANOVA (Table 1)
246 showed that the shoot and root dry biomasses were significantly affected by the presence of
247 the ectomycorrhizal fungus *H. cylindrosporum*, irrespective of the presence of free-living
248 organisms, with no interactive effect between the two main factors “mycorrhiza” and “free-
249 living organism” (Table 1). Taken as a whole, M treatment decreased shoot and root biomass
250 compared to NM treatment (Figure 2). The inoculation of free-living organisms did not
251 modify shoot biomasses of NM or M plants, nor the root biomasses of NM plants. However,
252 M plants not inoculated (None treatment) had lower root biomass (- 43 %) than NM plants,
253 whether not inoculated (None) or inoculated with bacteria and nematodes (+Bac +Nem).
254 Also, M plants inoculated with bacteria (+Bac) had lower root biomass (- 30 %) than NM
255 plants inoculated with bacteria and nematodes (+Bac +Nem). Only M plants inoculated with
256 nematodes and bacteria (+Bac +Nem) had the same root biomass as NM plants (Figure 2).
257 Regarding the root parameters (Table 2), the two-way ANOVA showed only a significant
258 effect of the M treatment and no significant interaction between the factors “mycorrhiza” and
259 “free-living organisms” for all the root variables (Table 3). The effect of the fungus was
260 confirmed by the one-way ANOVA (Table 2). In average, M treatment decreased all the root
261 parameters, with a stronger effect on root ramification (- 130% for the number of crossings, -
262 93 % for the number of forks, - 70 % for the number of tips) than on root length (- 47 %) and
263 surface areas (- 27 %) (Table 2).

264

265 **3.2. Plant P accumulation**

266 At harvest, only the shoots of NM plants inoculated with bacteria and nematodes (+Bac
267 +Nem), while those of all M plants, accumulated higher amounts of P than two-month-old
268 plants (Figure 3). In roots, the same trend was observed for NM plants, but only M plants

269 inoculated either with bacteria (+Bac) or with bacteria and nematodes (+Bac +Nem)
270 accumulated more P than two-month-old plants (Figure 3). The two-way ANOVA showed
271 that shoot P was significantly affected by the two factors “mycorrhiza” and “free-living
272 organism” while root P was affected only by the presence of free-living organisms, with no
273 interactive effect between these two main factors (Table 4). Taken as a group, M plants
274 accumulated more P in their shoots than the NM plants, irrespective of the presence of free-
275 living organisms (Figure 3). In contrast, P accumulation in roots did not differ between NM
276 and M plants. Finally, the calculation of total P accumulation per plant (Supplementary
277 Information 1) confirmed a different effect of free-living organisms in NM and M plants. In
278 NM plants, inoculation of bacteria alone (+Bac) did not modify the P accumulation whereas
279 the presence nematodes (+Bac, +Nem) was able to increase dramatically total P accumulation,
280 either by 88 % when compared to None treatment or by 98 % when compared to +Bac
281 treatment. In M plants, the inoculation of bacteria already increased by 40 % the P amounts
282 compared to None treatment. The addition of the nematodes increased by 66% the P amounts
283 compared to None treatment. However, M plants inoculated either with bacteria (+ Bac) or
284 with bacteria and nematodes (+Bac +Nem) accumulated the same amounts of total P per
285 plant.

286 We estimated the root P acquisition efficiency by dividing the total plant P amount by the root
287 length (Figure 4). The two-way ANOVA (Table 5) showed no significant interaction between
288 the factors “mycorrhiza” and “free-living organisms” on this parameter. In contrast, P
289 acquisition efficiency was highly affected by both the presence of the mycorrhizal fungus *H.*
290 *cylindrosporum* and the presence of free-living organisms. Within the same inoculation
291 treatment, M plants had always higher values of P acquisition efficiency than NM plants, with
292 increases of 78 % (None), 117 % (+Bac) and 47 % (+Bac +Nem). In NM plants, the presence
293 of *Rhabditis* sp. strongly increased P acquisition efficiency compared to the two other

294 treatments, by 90 % (None) and 70 % (+ Bac). In M plants, the presence of *Rhabditis* sp.
295 increased P acquisition efficiency only compared to plants not inoculated (None), by 60%.

296

297 ***3.3. Abundance of fungal DNA and nematodes***

298 To assess the soil colonization by *H. cylindrosporum* hyphae, we measured the number of 18S
299 rDNA gene copies of the fungus in soil DNA. As expected, the abundance of 18S rDNA
300 copies was significantly higher in soil with M plants than in soil with NM plants (Figure 5).
301 However, according to the two-way ANOVA (Table 6), the fungal rDNA gene copies number
302 was significantly affected by the interaction between the factors “mycorrhiza” and “free-
303 living organism. We did not find nematodes in the "None" and "+Bac" treatments. There were
304 ca 2000 nematodes per Petri dish in the soil in the "+Bac +Nem" treatments, irrespective of
305 the presence of the mycorrhizal fungus.

306

307 **4. Discussion**

308 ***4.1. Effects of rhizosphere organisms on plant growth***

309 We showed that (i) the presence of the mycorrhizal fungus had a slight negative impact on
310 shoot biomass and (ii) the presence of free-living organisms did not have any significant
311 effect on the accumulation of shoot biomass. At first glance, our results are opposed to those
312 found by Gyaneshwar et al. (2002), Djigal et al. (2004a), and Smith and Read (2008), who
313 showed that rhizosphere bacteria, bacterial-feeding nematodes, and mycorrhizal fungi
314 increased shoot biomass, respectively. However, this discrepancy can be explained by the
315 protocol used among these different studies. During the two-month pre-experiment required
316 for the mycorrhizal synthesis, plants were grown in a nutrient solution and accumulated P in
317 their tissues. Once placed in Petri dishes, they recycled their internal P despite the low P soil
318 availability in order to sustain shoot biomass production. Plants are able to vary their

319 stoichiometry to sustain shoot biomass production despite low soil mineral fertility (Gusewell,
320 2004) and this was shown for maritime pine (Ali et al., 2009; 2014). The negative effects of
321 the mycorrhizal fungus on shoot biomass can also be explained by the short duration of the
322 experiment. It has been shown that the effect of *H. cylindrosporum* on the rate of increase of
323 *P. pinaster* shoot biomass depends on the duration of the experiment (Aquino & Plassard,
324 2004). A period of 100 days does not appear to be long enough to detect significant effects of
325 the mycorrhizal fungus on shoot growth, especially with slow-growing plant species such as
326 *P. pinaster*. Despite these considerations, it was decided to stop the experiment after 100 days
327 because of the strong root development in the Petri dishes. It would be interesting to continue
328 the inoculation experiment for a longer period using larger containers in order to allow
329 sufficient space for root development.

330

331 ***4.2. Effect of organisms on plant P acquisition***

332 The analysis of the P content in the shoots and roots and the total P amount in the plants at the
333 end of the experiment showed that the presence of bacteria with bacterial-feeding nematodes
334 strongly increased the plants' P acquisition. The presence of *Bacillus subtilis* (alone or not) is
335 central for the mineralization of phytate by the extra-cellular enzyme (Jorquera et al., 2011;
336 Becquer et al., 2014), but the inoculation of bacteria alone in the non-mycorrhizal plants did
337 not significantly affect plant P nutrition. In other words, without the nematodes, the P
338 mineralization did not result in higher P availability. The inoculation of both bacteria and
339 nematodes is thus decisive for the net flow of P from phytate to the plant. This agrees with the
340 previous study carried out by Irshad et al. (2012) on agarose medium. Our results suggest that
341 the interaction between bacteria and bacterial-feeding nematodes improved soil P availability
342 for plants by stimulating the phytate-mineralizing bacterial activity and releasing microbial
343 biomass P (Trap et al., 2016). Similar patterns were found by Ranoarisoa et al. (2018) with

344 rice growing in Ferralsol from Madagascar with or without *Acrobelloides* sp., but only when
345 the soil pH was corrected with dolomite input. The promotion of the bacterial Po-mineralizing
346 activity by nematode inoculation has also been found by Djigal et al. (2004a; 2004b), who
347 showed that bacterial grazers increased soil phosphatase activity in the rhizosphere of maize
348 at all plant growth stages. From our data, we cannot decipher whether the nematodes
349 increased plant-available P through (i) higher microbial activity (excess of mineralized P not
350 immobilized in the microbial biomass) and/or (ii) excretion (P immobilized in the microbial
351 biomass and then released by the nematodes). The use of isotopic labeling (^{33}P) should
352 constitute the best way to achieve this goal (Ranoarisoa et al., 2018).

353 In contrast, the sole presence of the fungus did not increase plant P acquisition in “None”
354 treatment. The absence of plant response towards fungal inoculation is also probably due to
355 the duration of the experiment. Aquino and Plassard (2004) found that the same strain of *H.*
356 *cylindrosporum* had a beneficial effect on *P. pinaster* P nutrition after 6 months of growth, but
357 no effect after 4 months. However, it is important to note that the inoculation of bacteria in
358 ectomycorrhizal plants led to a significant effect on plant P amount (Supplementary
359 Information 1). This effect was clearly discernible when we calculated the plant P acquisition
360 efficiency, supposing that the Pi uptake rate of *P. pinaster* increased when both *B. subtilis* and
361 *H. cylindrosporum* were inoculated. As the fungus alone was shown to be poorly able to use
362 phytate in pure culture (Irshad et al., 2012), it could have been more efficient to use Pi
363 resulting from bacterial phytate mineralization. In this respect, the first hypothesis is that the
364 intense exploration of the soil by the hyphae would allow a higher ability of the mycorrhizal
365 roots to take up Pi released by the bacterial enzymes. Furthermore, we must not exclude the
366 possibility that, during the growth of the fungus, bacteria became attached to the hyphae and
367 move in the soil (Kohlmeier et al., 2005) promoting (i) the contact between the bacteria and
368 the substrate (phytate) and (ii) phytate mineralization. Finally, it is also possible that the high

369 expansion of fungal hyphae promotes carbon input into the soil away from the roots,
370 increasing bacterial activity in larger soil volumes. For instance, by manipulating the
371 endomycorrhizal fungus *Glomus intraradices* and protozoa in the rhizosphere of *Plantago*
372 *lanceolata*, Koller et al. (2013) showed that the presence of the bacterial grazers stimulated
373 the translocation of carbon from the plant via the fungus into the soil, fueling the microbial
374 loop (Clarholm, 1985a) and resulting in greater plant nutrition in comparison to plants without
375 protozoa. No such mechanism has been observed for ectomycorrhizal fungi, but our results
376 greatly encourage further studies focusing on this feature.

377

378 ***4.3. Effects of microorganisms on root growth and architecture***

379 The most notable effect was that *H. cylindrosporium* decreased the root biomass, irrespective
380 of the presence of free-living organisms. This agrees with a previous study carried out using
381 the same fungal isolate associated with *P. pinaster* (Aquino & Plassard, 2004). The reduced
382 root development, which was associated with ectomycorrhizal fungi, was attributed to the
383 high ability of the extraradical hyphae of the fungus to take up P from soil (Aquino &
384 Plassard, 2004; Plassard & Dell, 2010).

385 The analysis of the root architecture underlined the strong significant effect of the fungus on
386 the root compartment. In agreement with the findings of Jentschke et al. (1995) and
387 Bonkowski et al. (2001), the presence of the mycorrhizal fungus resulted in a less complex
388 root system. All root parameters were significantly reduced by the presence of the
389 ectomycorrhizal fungus. As discussed above, this pattern could be due to the high ability of
390 the ectomycorrhizal fungus to acquire P from the soil, thus shifting the plant carbon allocation
391 from the roots to the fungus (Brown *et al.*, 2013). In contrast to the ectomycorrhizal fungus,
392 bacteria and nematodes did not modify significantly the root architecture. Changes in the root
393 architecture in the presence of bacterial feeders (especially protozoa) have been interpreted as

394 hormonal non-nutritional effects on the lateral root production (Bonkowski, 2004; Bonkowski
395 & Clarholm, 2012). Here, the soil was inoculated with a population of *B. subtilis* (not with a
396 microbial community), which probably limited the ability of nematodes to induce non-
397 nutritional effects (Ranoarisoa et al., 2018).

398

399 ***4.4. Relevance of our simplified conditions to study micro-food web interactions***

400 In this study, we used an experimental system with simplified and artificial conditions to
401 study the effects of the micro-food web interactions on phytate mineralization as we worked
402 with sterilized soil-bead mixture and addition of selected free-living organisms. Compared to
403 *in situ* ecosystems presenting huge variations of temperatures and soil humidity, these two
404 environmental factors varied little during all the experiment, thus favoring enzyme (Rinkes et
405 al., 2013) and root activities for mineral uptake (Gessler et al., 1998). Although sterilized, our
406 soil-bead mixture enabled us to take into account the exchanges between mineral nutrients
407 and the soil as well as the rhizosphere changes able to modify greatly mineral nutrients
408 bioavailability for plants (Bravin et al., 2010). The control of biodiversity, although low in
409 comparison to field conditions, enabled us to quantify the potential contribution of each actor,
410 alone or in combination, to the mineralization of phytate and finally, to the bioavailability of P
411 released for the plant. Thus, as underlined by Mezeli et al. (2020), our experimental
412 conditions were suitable to reveal the effects of the soil microbial loop on phytate
413 mineralization and plant P bioavailability. Hence, the next step of this work will be to use
414 different richness levels of free-living organisms added in non-sterilized soil in order to take
415 into account better their interactions with soil biodiversity which might modify the final plant
416 P bioavailability.

417

418 **5. Conclusion**

419 We formulated four hypotheses. First, we supposed that the plant alone, without mycorrhiza
420 nor free-living organisms, is not able to uptake P from phytate (H1). This hypothesis was
421 validated since no P was acquired from phytate when no rhizospheric organisms were
422 inoculated. Second, we expected higher plant P uptake in the presence of *H. cylindrosporum*
423 in comparison to the plant alone (H2). This hypothesis was partially validated since the
424 presence of the fungus did not result in higher plant P nutrition in the “None” treatments, but
425 the inoculation of bacteria in ectomycorrhizal plants led to a significant effect on plant P
426 amount. The third hypothesis suggested that the presence of free-living organisms, bacteria,
427 and nematodes, would increase the plants’ P uptake. This hypothesis was fully validated and
428 constitutes the most remarkable result of the present study. The strong positive effect of the
429 presence of both phytate-mineralizing bacteria and bacterial-feeding nematodes on P
430 acquisition by plants growing in soil amended with phytate is central and suggests that
431 interactions between bacteria and bacterial-feeding nematodes are essential to increase soil Pi
432 availability for plants. Finally, the last hypothesis, supposing that plant P uptake from phytate
433 will be maximal when all potential rhizosphere mutualists are inoculated, was also validated.
434 The findings are particularly noteworthy because they show the importance of the rhizosphere
435 micro-food web role on plants’ P nutrition from a recalcitrant organic P source despite high Pi
436 adsorption onto soil components. Further investigations are required to assess better the
437 involved mechanisms. They could consist of the quantification of phosphatases and phytases
438 activities together with available Po and Pi in rhizosphere soil to determine the possible
439 relationships between soil micro-food web interactions and these parameters. Exploring the
440 benefits of rhizosphere interactions and identifying the potential drivers of these interactions
441 appear as pertinent goals to develop agricultural management practices that could take
442 advantage of the beneficial interactions of these organisms.

443

444 **Acknowledgments**

445

446 We would like to thank the three anonymous reviewers for their helpful suggestions that
447 helped us improve the manuscript. This project was part of the ANR project “UNLOCKP”
448 (ANR 11 BSV7 015 01; France) awarded to C. Plassard. The work by M.P. Ranoarisoa was
449 also funded by the UNLOCKP project. The work was also supported by the program
450 ‘Investments for the future’ (ANR-10- LABX-04-01) through the use of the Ecotrop platform
451 from CeMEB labEx. We are grateful to Pierre Trichet (ISPA, INRA Bordeaux) for providing
452 the soil samples for nematode isolation. We also thank Anne Jimenez and Cécile Villenave
453 (Elisol Environment) for their help and advice on the production of nematodes. We thank
454 Valentin Lauret, Guillaume Pugliese, Sebastian Mira, and Olivier Pacholick for their help
455 during the experiments.

456

457

458 **References**

459

460 Alexander, M., 1977. Introduction to soil microbiology. Wiley New York.

461 Ali, M.A., Louche, J., Duchemin, M., Plassard, C., 2014. Positive growth response of *Pinus pinaster*
462 seedlings in soils previously subjected to fertilization and irrigation. Forest Ecology and
463 Management 318, 62-70.

464 Ali, M.A., Louche, J., Legname, E., Duchemin, M., Plassard, C., 2009. *Pinus pinaster* seedlings and
465 their fungal symbionts show high plasticity in phosphorus acquisition in acidic soils. Tree
466 Physiology 29, 1587-1597.

467 Anderson, R.V., Elliott, E.T., McClellan, J.F., Coleman, D.C., Cole, C.V., Hunt, H.W., 1978. Trophic
468 interactions in soils as they affect energy and nutrient dynamics. 3. Biotic interactions of
469 bacteria, amebas, and nematodes. Microbial Ecology 4, 361-371.

470 Aquino, M., Plassard, C., 2004. Dynamics of ectomycorrhizal mycelial growth and P transfer to the
471 host plant in response to low and high soil P availability. FEMS Microbiology Ecology 48,
472 149-156.

473 Becquer, A., Torres-Aquino, M., Le Guernevé, C., Amenc, L.K., Trives-Segura, C., Staunton, S.,
474 Quiquampoix, H., Plassard, C., 2017. Establishing a Symbiotic Interface between Cultured
475 Ectomycorrhizal Fungi and Plants to Follow Fungal Phosphate Metabolism. Bio-Protocol 7,
476 e2577.

477 Becquer, A., Trap, J., Irshad, U., Ali, M.A., Plassard, C., 2014. From soil to plant, the journey of P
478 through trophic relationships and ectomycorrhizal association. Frontiers in plant science 5, 1-
479 7.

480 Bongers, T., 1990. The maturity index: an ecological measure of environmental disturbance based on
481 nematode species composition. Oecologia 83, 14-19.

482 Bonkowski, M., 2004. Protozoa and plant growth: the microbial loop in soil revisited. New
483 Phytologist 162, 617-631.

484 Bonkowski, M., Clarholm, M., 2012. Stimulation of plant growth through interactions of bacteria and
485 protozoa: Testing the auxiliary microbial loop hypothesis. *Acta Protozoologica* 51, 237-247.

486 Bonkowski, M., Jentschke, G., Scheu, S., 2001. Contrasting effects of microbial partners in the
487 rhizosphere: interactions between Norway Spruce seedlings (*Picea abies* Karst.), mycorrhiza
488 (*Paxillus involutus* (Batsch) Fr.) and naked amoebae (protozoa). *Applied Soil Ecology* 18,
489 193-204.

490 Bravin, M.N., Michaud, A.M., Larabi, B., Hinsinger, P., 2010. RHIZOtest: A plant-based biotest to
491 account for rhizosphere processes when assessing copper bioavailability. *Environmental*
492 *Pollution* 158, 3330–3337.

493 Brown, L.K., George, T.S., Barrett, G.E., Hubbard, S.F., White, P.J., 2013. Interactions between root
494 hair length and arbuscular mycorrhizal colonisation in phosphorus deficient barley (*Hordeum*
495 *vulgare*). *Plant and Soil* 372, 195-205.

496 Bucher, M., 2007. Functional biology of plant phosphate uptake at root and mycorrhiza interfaces.
497 *New Phytologist* 173, 11-26.

498 Casarin, V., Plassard, C., Souche, G., Arvieu, J.C., 2003. Quantification of oxalate ions and protons
499 released by ectomycorrhizal fungi in rhizosphere soil. *Agronomie* 23, 461–469.

500 Clarholm, M., 1985a. Interactions of bacteria, protozoa and plants leading to mineralization of soil-
501 nitrogen. *Soil Biology and Biochemistry* 17, 181-187.

502 Clarholm, M., 1985b. Interactions of bacteria, protozoa and plants leading to mineralization of soil
503 nitrogen. *Soil Biology and Biochemistry* 17, 181-187.

504 Clarholm, M. (2005) Soil protozoa: an under-researched microbial group gaining momentum. *Soil*
505 *Biology and Biochemistry* 37, 811-817.

506 Cole, C.V., Elliott, E.T., Hunt, H.W. & Coleman, D.C., 1978. Trophic interactions in soils as they
507 affect energy and nutrient dynamics. Phosphorus transformations. *Microbial Ecology* 4, 381-
508 387.

509 Colpaert, J.V., vanLaere, A., 1996. A comparison of the extracellular enzyme activities of two
510 ectomycorrhizal and a leaf-saprotrophic basidiomycete colonizing beech leaf litter. *New*
511 *Phytologist* 134, 133-141.

512 Courty, P.E., Pouysegur, R., Buée, M., Garbaye, J., 2006. Laccase and phosphatase activities of the
513 dominant ectomycorrhizal types in a lowland oak forest. *Soil Biology and Biochemistry* 38,
514 1219-1222.

515 Darbyshire, J.F., Davidson, M.S., Chapman, S.J., Ritchie, S., 1994. Excretion of nitrogen and
516 phosphorus by the soil Ciliate *Colpoda steinii* when fed the soil bacterium *Arthrobacter sp.*
517 *Soil Biology and Biochemistry* 26, 1193-1199.

518 Darch, T., Blackwell, M.S.A., Hawkins, J.M.B., Haygarth, P.M., Chadwick, D., 2014. A meta-analysis
519 of organic and inorganic phosphorus in organic fertilizers, soils, and water: implications for
520 water quality. *Critical Reviews in Environmental Science & Technology* 44, 2172–2202.

521 Debaud, J.C., Gay, G., 1987. *In vitro* fruiting under controlled conditions of the ectomycorrhizal
522 fungus *Hebeloma cylindrosporum* associated with *Pinus pinaster*. *New Phytologist* 105, 429-
523 435.

524 Djigal, D., Brauman, A., Diop, T.A., Chotte, J.L., Villenave, C., 2004a. Influence of bacterial-feeding
525 nematodes (*Cephalobidae*) on soil microbial communities during maize growth. *Soil Biology*
526 *and Biochemistry* 36, 323-331.

527 Djigal, D., Sy, M., Brauman, A., Diop, T.A., Mountport, D., Chotte, J.L., Villenave, C., 2004b.
528 Interactions between *Zeldia Punctata* (Cephalobidae) and bacteria in the presence or absence
529 of maize plants. *Plant and Soil* 262, 33-44.

530 Finlay, R.D., 2008. Ecological aspects of mycorrhizal symbiosis: with special emphasis on the
531 functional diversity of interactions involving the extraradical mycelium. *Journal of*
532 *Experimental Botany* 59, 1115–1126.

533 Gessler, A., Schneider, S., Von Sengbusch, D., Weber, P., Hanemann, U., Huber, C., Rothe, A.,
534 Kreutzer, K., Rennenberg, H., 1998. Field and laboratory experiments on net uptake of nitrate
535 and ammonium by the roots of spruce (*Picea abies*) and beech (*Fagus sylvatica*) trees. *New*
536 *Phytologist* 138, 275–285.

537 George, T.S., Giles, C.D., Menezes-Blackburn, D., Condon, L.M., Gama-Rodrigues, A.C., Jaisi, D.,
538 Lang, F., Neal, A.L., Stutter, M.I., Almeida, D.S., Bol, R., Cabugao, K.G., Celi, L., Cotner,
539 J.B., Feng, G., Goll, D.S., Hallama, M., Krueger, J., Plassard, C., Rosling, A., Darch, T.,

540 Fraser, T., Giesler, R., Richardson, A.E., Tamburini, F., Shand, C.A., Lumsdon, D.G., Zhang,
541 H., Blackwell, M.S.A., Wearing, C., Mezeli, M.M., Almás, Å.R., Audette, Y., Bertrand, I.,
542 Beyhaut, E., Boitt, G., Bradshaw, N., Brearley, C.A., Bruulsema, T.W., Ciais, P., Cozzolino,
543 V., Duran, P.C., Mora, M.L., de Menezes, A.B., Dodd, R.J., Dunfield, K., Engl, C., Frazão,
544 J.J., Garland, G., González Jiménez, J.L., Graca, J., Granger, S.J., Harrison, A.F., Heuck, C.,
545 Hou, E.Q., Johnes, P.J., Kaiser, K., Kjær, H.A., Klumpp, E., Lamb, A.L., Macintosh, K.A.,
546 Mackay, E.B., McGrath, J., McIntyre, C., McLaren, T., Mészáros, E., Missong, A.,
547 Mooshammer, M., Negrón, C.P., Nelson, L.A., Pfahler, V., Poblete-Grant, P., Randall, M.,
548 Seguel, A., Seth, K., Smith, A.C., Smits, M.M., Sobarzo, J.A., Spohn, M., Tawaraya, K.,
549 Tibbett, M., Voroney, P., Wallander, H., Wang, L., Wasaki, J., Haygarth, P.M., 2018. Organic
550 phosphorus in the terrestrial environment: a perspective on the state of the art and future
551 priorities. *Plant and Soil* 427, 191–208.

552 Griffiths, B.S., 1986. Mineralization of nitrogen and phosphorus by mixed cultures of the Ciliate
553 protozoan *Colpoda steinii*, the nematode *Rhabditis* sp and the bacterium *Pseudomonas*
554 *fluorescens*. *Soil Biology & Biochemistry* 18, 637-641.

555 Gusewell, S., 2004. N : P ratios in terrestrial plants: variation and functional significance. *New*
556 *Phytologist* 164, 243-266.

557 Gyaneshwar, P., Kumar, G.N., Parekh, L.J. & Poole, P.S., 2002. Role of soil microorganisms in
558 improving P nutrition of plants. *Plant and Soil* 245, 83-93.

559 Hayes, J., Simpson, R., Richardson, A., 2000. The growth and phosphorus utilisation of plants in
560 sterile media when supplied with inositol hexaphosphate, glucose 1-phosphate or inorganic
561 phosphate. *Plant and Soil* 220, 165-174.

562 Hinsinger, P., Brauman, A., Devau, N., Gerard, F., Jourdan, C., Laclau, J.-P., Le Cadre, E., Jaillard,
563 B., Plassard, C., 2011. Acquisition of phosphorus and other poorly mobile nutrients by roots.
564 Where do plant nutrition models fail? *Plant and Soil* 348, 29-61.

565 Holford, I.C.R., 1997. Soil phosphorus: Its measurement, and its uptake by plants. *Australian Journal*
566 *of Soil Research* 35, 227-239.

567 Irshad, U., Brauman, A., Villenave, C., Plassard, C., 2012. Phosphorus acquisition from phytate
568 depends on efficient bacterial grazing, irrespective of the mycorrhizal status of *Pinus pinaster*.
569 Plant and Soil 358, 148-161.

570 Irshad, U., Villenave, C., Brauman, A., Plassard, C., 2011. Grazing by nematodes on rhizosphere
571 bacteria enhances nitrate and phosphorus availability to *Pinus pinaster* seedlings. Soil Biology
572 and Biochemistry 43, 2121-2126.

573 Iyamuremye, F., Dick, R.P., Baham, J., 1996. Organic amendments and phosphorus dynamics: III.
574 Phosphorus speciation. Soil Science 161, 444-451.

575 Jentschke, G., Bonkowski, M., Godbold, D.L., Scheu, S., 1995. Soil protozoa and forest tree growth -
576 Non-nutritional effects and interaction with mycorrhizae. Biology and Fertility of Soils 20,
577 263-269.

578 Jorquera, M.A., Crowley, D.E., Marschner, P., Greiner, R., Fernández, M.T., Romero, D., Menezes-
579 Blackburn, D., De La Luz Mora, M., 2011. Identification of β -propeller phytase-encoding
580 genes in culturable *Paenibacillus* and *Bacillus* spp. from the rhizosphere of pasture plants on
581 volcanic soils. FEMS Microbiology Ecology 75, 163-172.

582 Kohlmeier, S., Smits, T.H., Ford, R.M., Keel, C., Harms, H., Wick, L.Y., 2005. Taking the fungal
583 highway: mobilization of pollutant-degrading bacteria by fungi. Environmental science &
584 technology 39, 4640-4646.

585 Koller, R., Rodriguez, A., Robin, C., Scheu, S., Bonkowski, M., 2013. Protozoa enhance foraging
586 efficiency of arbuscular mycorrhizal fungi for mineral nitrogen from organic matter in soil to
587 the benefit of host plants. New Phytologist 199, 203-211.

588 Kuikman, P.J., Jansen, A.G., Vanveen, J.A., 1991. N-15-nitrogen mineralization from bacteria by
589 protozoan grazing at different soil-moisture regimes. Soil Biology and Biochemistry 23, 193-
590 200.

591 Louche, J., Ali, M.A., Cloutier-Hurteau, B., Sauvage, F.-X., Quiquampoix, H., Plassard, C., 2010.
592 Efficiency of acid phosphatases secreted from the ectomycorrhizal fungus *Hebeloma*
593 *cylindrosporium* to hydrolyse organic phosphorus in podzols. FEMS Microbiology Ecology
594 73, 323-335.

595 Mezeli, M.M., Page, P., George, T.S., Neilson, N., Mead, A., Blackwell, M. S.A., Haygarth, P.M.,
596 2020. Using a meta-analysis approach to understand complexity in soil biodiversity and
597 phosphorus acquisition in plants. *Soil Biology and Biochemistry*, DOI:
598 10.1016/j.soilbio.2019.107695

599 Morizet, J., Mingeau, M., 1976. Influence des facteurs de milieu sur l'absorption hydrique. Etude
600 effectuee sur tomate decapitee en exsudation. i. facteurs nutritionnels. *Annales Agronomiques*
601 27, 333–342.

602 Mullaney, E.J., Ullah, A.H.J., 2003 The term phytase comprises several different classes of enzymes.
603 *Biochemical and Biophysical Research Communications* 312, 179-184.

604 Nehls, U., Plassard, C., 2018. Nitrogen and phosphate metabolism in ectomycorrhizas. *New*
605 *Phytologist* 220, 1047–1058.

606 Oberson, A., Friesen, D.K., Rao, I.M., Buhler, S., Frossard, E., 2001. Phosphorus Transformations in
607 an Oxisol under contrasting land-use systems: The role of the soil microbial biomass. *Plant*
608 *and Soil* 237, 197-210.

609 Ohno, T., Zibilske, L.M., 1991. Determination of low concentration of phosphorus in soil extracts using
610 malachite green. *Soil Science Society of America Journal* 55, 892-895.

611 Pistocchi, C., Mészáros, É., Tamburini, F., Frossard, E., Bünemann, E.K., 2018. Biological processes
612 dominate phosphorus dynamics under low phosphorus availability in organic horizons of
613 temperate forest soils. *Soil Biology and Biochemistry* 126, 64–75.

614 Plassard, C., Dell, B., 2010. Phosphorus nutrition of mycorrhizal trees. *Tree Physiology* 30, 1129-
615 1139.

616 Plassard, C., Louche, J., Ali, M.A., Duchemin, M., Legname, E., Cloutier-Hurteau, B., 2011. Diversity
617 in phosphorus mobilisation and uptake in ectomycorrhizal fungi. *Annals of Forest Science* 68,
618 33-43.

619 Ranoarisoa, M.P., Morel, C., Andriamananjara, A., Jourdan, C., Bernard, L., Becquer, T.,
620 Rabeharisoa, L., Rahajahalaza, K., Plassard, C., Blanchart, E., 2018. Effects of a
621 bacterivorous nematode on rice ³²P uptake and root architecture in a high P-sorbing ferrallitic
622 soil. *Soil Biology and Biochemistry* 122, 39-49.

623 Richardson, A.E., Hadobas, P.A., Hayes, J.E., 2000. Acid phosphomonoesterase and phytase activities
624 of wheat (*Triticum aestivum* L.) roots and utilization of organic phosphorus substrates by
625 seedlings grown in sterile culture. *Plant, Cell & Environment* 23, 397-405.

626 Richardson, A.E., Hadobas, P.A., Hayes, J.E., O'Hara, C.P., Simpson, R.J., 2001. Utilization of
627 phosphorus by pasture plants supplied with myo-inositol hexaphosphate is enhanced by the
628 presence of soil micro-organisms. *Plant and Soil* 229, 47-56.

629 Rinkes, Z., Sinsabaugh, R., Moorhead, D., Grandy, A., Weintraub, M., 2013. Field and lab conditions
630 alter microbial enzyme and biomass dynamics driving decomposition of the same leaf litter.
631 *Frontiers in Microbiology* 4, 260.

632 Rodriguez, H., Fraga, R., 1999. Phosphate solubilizing bacteria and their role in plant growth
633 promotion. *Biotechnology Advances* 17, 319-339.

634 Scheerer, U., Trube, N., Netzer, F., Rennenberg, H., Herschbach, C., 2019. ATP as Phosphorus and
635 Nitrogen Source for Nutrient Uptake by *Fagus sylvatica* and *Populus x canescens* Roots.
636 *Frontiers in Plant Science* 10, 378.

637 Smith, S.E., Read, D.J., 2008. *Mycorrhizal Symbiosis - Third Edition*. Academic Press, London.

638 Southey, J.F., 1986. *Laboratory methods for work with plant and soil nematodes*. Ministry of
639 Agriculture, Fisheries and Food, Reference Book 402, London.

640 Team, R.C. (2016) RStudio: Integrated Development for R [Internet]. [cited 2015 Nov 20].

641 Tournier, E., Amenc, L., Pablo, A.-L., Legname, E., Blanchart, E., Plassard, C., Robin, A., Bernard,
642 L., 2015. Modification of a commercial DNA extraction kit for safe and rapid recovery of
643 DNA and RNA simultaneously from soil, without the use of harmful solvents. *MethodsX* 2,
644 182-191.

645 Trap, J., Bonkowski, M., Plassard, C., Villenave, C., Blanchart, E., 2016. Ecological importance of
646 soil bacterivores for ecosystem functions. *Plant and Soil* 398, 1-24.

647 Turner, B.L., 2007. Inositol phosphates in soil: amounts, forms and significance of the phosphorylated
648 inositol stereoisomers. *Inositol phosphates: linking agriculture and the environment*. (eds B.L.
649 Turner, A.E. Richardson & E.J. Mullaney), pp. 186–206. CAB International, Wallingford,
650 UK.

651 Vitousek, P.M., Porder, S., Houlton, B.Z., Chadwick, O.A., 2010. Terrestrial phosphorus limitation:
652 mechanisms, implications, and nitrogen-phosphorus interactions. *Ecological Applications* 20,
653 5-15.

654

655 **Captions to figures**

656

657 **Figure 1. Schematic procedures to obtain the plants for inoculation experiments.**

658 A: ectomycorrhizal synthesis in glass tubes with *Pinus pinaster* and *Hebeloma*
659 *cylindrosporum*. Non mycorrhizal plants are obtained without fungal plugs added. The inset
660 diagram shows the home-made closing system of each Teflon tube. B-E: preparation of
661 square Petri dishes with soil-bead mixture. B: two holes for plant passage and watering are
662 first made with a hot tool. C: the sterile soil-bead-nutrient solution mixture is poured into the
663 dish. D: a two-month-old pine seedling is taken of the glass tube and placed on the soil-bead
664 mixture before addition of free-living organisms (*B. subtilis* and/or *Rhabditis* sp.). E: the hole
665 for watering and the dish are closed with pieces of sticky tape.

666

667 **Figure 2. Biomass accumulation in shoots and roots of *P. pinaster*, whether not**
668 **associated (NM) or associated with the ectomycorrhizal basidiomycete *H.***
669 ***cylindrosporum* (M) and different inoculation treatments.**

670 Two-month old plants were grown for 100 d in a cambisol supplemented with phytate with
671 different inoculation treatments that were: no addition of living-free organisms (None),
672 addition of a phytate-mineralizing bacteria *Bacillus subtilis* (+Bac) or addition of bacteria
673 together with bacteria-feeding nematodes *Rhabditis* sp. (+Bac +Nem). Bars are the means
674 (n=6) with standard deviation. Different capital letters (A, B) indicate significant differences
675 among mycorrhizal fungal status at $p < 0.05$ (two-way ANOVA). Different lowercase letters
676 (a, b, c) indicate significant differences among biological treatments at $p < 0.05$ (one-way
677 ANOVA followed by post hoc Tukey HSD test). Asterisk indicates a mean significantly
678 different from the initial value measured in two-month old plants at $p < 0.05$ (Student's t-test of
679 mean comparison)

680

681 **Figure 3. Total P amount in shoots and roots of *P. pinaster*, whether not associated (NM)**
682 **or associated with the ectomycorrhizal basidiomycete *H. cylindrosporium* (M).**

683 Two-month old plants were grown for 100 d in a cambisol supplemented with phytate with
684 different inoculation treatments that were: no addition of living-free organisms (None),
685 addition of a phytate-mineralizing bacteria *Bacillus subtilis* (+Bac) or addition of bacteria
686 together with bacteria-feeding nematodes *Rhabditis* sp. (+Bac +Nem). Bars are the means
687 (n=6) with standard deviation. Different capital letters (A, B) indicate significant differences
688 among mycorrhizal fungal status at $p < 0.05$ (two-way ANOVA). Different lowercase letters
689 (a, b, c) indicate significant differences among biological treatments at $p < 0.05$ (one-way
690 ANOVA followed by post hoc Tukey HSD test). Asterisk indicates a mean significantly
691 different from the initial value measured in two-month old plants at $p < 0.05$ (Student's t-test of
692 mean comparison).

693

694 **Figure 4. P acquisition efficiency of *P. pinaster*, whether not associated (NM) or**
695 **associated with the ectomycorrhizal basidiomycete *H. cylindrosporium* (M).**

696 Two-month old plants were grown for 100 d in a cambisol supplemented with phytate with
697 different inoculation treatments that were: no addition of living-free organisms (none),
698 addition of a phytate-mineralizing bacteria *Bacillus subtilis* (+Bac) or addition of bacteria
699 together with bacteria-feeding nematodes *Rhabditis* sp. (+Bac +Nem). Bars are the means
700 (n=6) with standard deviation. Different capital letters (A, B) indicate significant differences
701 among mycorrhizal fungal status at $p < 0.05$ (two-way ANOVA). Different lowercase letters
702 (a, b, c, d) indicate significant differences among biological treatments at $p < 0.05$ (one-way
703 ANOVA followed by post hoc Tukey HSD test).

704

705 **Figure 5. Abundance of fungal (18S) rDNA gene copies measured by qPCR in soil with**
706 ***P. pinaster*, whether not associated (NM) or associated with the ectomycorrhizal**
707 **basidiomycete *H. cylindrosporum* (M) after 100 d of growth in a cambisol supplemented**
708 **with phytate and with different inoculation treatments.**

709 Inoculation treatments were: no addition of living-free organisms (None), addition of a
710 phytate-mineralizing bacteria *Bacillus subtilis* (+Bac) or addition of bacteria together with
711 bacteria-feeding nematodes *Rhabditis* sp. (+Bac +Nem). Bars are the means (n=6) with
712 standard deviation. Different capital letters (A, B) indicate significant differences among
713 mycorrhizal fungal status at $p < 0.05$ (two-way ANOVA). Different lowercase letters (a, b, c)
714 indicate significant differences among biological treatments at $p < 0.05$ (one-way ANOVA
715 followed by post hoc Tukey HSD test).

716

717 **Supplementary information 1. Pictures of the devices used to produce plants for**
718 **inoculation experiments**

719 **Supplementary information 2. Plant total P amounts according to biological treatments.**




Figure 1. Schematic procedures to obtain the plants for inoculation experiments.

A: ectomycorrhizal synthesis in glass test-tubes with *Pinus pinaster* and *Hebeloma cylindrosporum*. Non mycorrhizal plants are obtained without fungal plugs added. The inset diagram shows the home-made closing system of each Teflon tube. B-E: preparation of square Petri dishes with soil-bead mixture. B: two holes for plant passage and watering are first made with a hot tool. C: the sterile soil-bead-nutrient solution mixture is poured into the dish. D: a two-month-old pine seedling is taken of the glass tube and placed on the soil-bead mixture before addition of free-living organisms (*B. subtilis* and/or *Rhabditis* sp.). E: the hole for watering and the dish are closed with pieces of sticky tape.




Figure 2. Biomass accumulation in shoots and roots of *P. pinaster*, whether not associated (NM) or associated with the ectomycorrhizal basidiomycete *H. cylindrosporium* (M) and different inoculation treatments.

Two-month old plants were grown for 100 d in a cambisol supplemented with phytate with different inoculation treatments that were: no addition of supplementary organisms (None), addition of a phytate-mineralizing bacteria *Bacillus subtilis* (+Bac) or addition of bacteria together with bacteria-feeding nematodes *Rhabditis* sp. (+Bac +Nem). Bars are the means (n=6) with standard deviation. Different capital letters (A, B) indicate significant differences among mycorrhizal fungal status at $p < 0.05$ (two-way ANOVA). Different lowercase letters (a, b, c) indicate significant differences among biological treatments at $p < 0.05$ (one-way ANOVA followed by post hoc Tukey HSD test). Asterisk indicates a mean significantly different from the initial value measured in two-month old plants at $p < 0.05$ (Student's t-test of mean comparison)




Figure 3. Total P amount in shoots and roots of *P. pinaster*, whether not associated (NM) or associated with the ectomycorrhizal basidiomycete *H. cylindrosporium* (M).

Two-month old plants were grown for 100 d in a cambisol supplemented with phytate with different inoculation treatments that were: no addition of supplementary organisms (None), addition of a phytate-mineralizing bacteria *Bacillus subtilis* (+Bac) or addition of bacteria together with bacteria-feeding nematodes *Rhabditis* sp. (+Bac +Nem). Bars are the means (n=6) with standard deviation. Different capital letters (A, B) indicate significant differences among mycorrhizal fungal status at $p < 0.05$ (two-way ANOVA). Different lowercase letters (a, b, c) indicate significant differences among biological treatments at $p < 0.05$ (one-way ANOVA followed by post hoc Tukey HSD test). Asterisk indicates a mean significantly different from the initial value measured in two-month old plants at $p < 0.05$ (Student's t-test of mean comparison).




Figure 4. P acquisition efficiency of *P. pinaster*, whether not associated (NM) or associated with the ectomycorrhizal basidiomycete *H. cylindrosporium* (M).

Two-month old plants were grown for 100 d in a cambisol supplemented with phytate with different inoculation treatments that were: no addition of supplementary organisms (none), addition of a phytate-mineralizing bacteria *Bacillus subtilis* (+Bac) or addition of bacteria together with bacteria-feeding nematodes *Rhabditis* sp. (+Bac +Nem). Bars are the means (n=6) with standard deviation. Different capital letters (A, B) indicate significant differences among mycorrhizal fungal status at $p < 0.05$ (two-way ANOVA). Different lowercase letters (a, b, c, d) indicate significant differences among biological treatments at $p < 0.05$ (one-way ANOVA followed by post hoc Tukey HSD test).




Figure 5. Abundance of fungal (18S) rDNA gene copies measured by qPCR in soil with *P. pinaster*, whether not associated (NM) or associated with the ectomycorrhizal basidiomycete *H. cylindrosporium* (M) after 100 d of growth in a cambisol supplemented with phytate and with different inoculation treatments.

Inoculation treatments were: no addition of supplementary organisms (None), addition of a phytate-mineralizing bacteria *Bacillus subtilis* (+Bac) or addition of bacteria together with bacteria-feeding nematodes *Rhabditis* sp. (+Bac +Nem). Bars are the means (n=6) with standard deviation. Different capital letters (A, B) indicate significant differences among mycorrhizal fungal status at $p < 0.05$ (two-way ANOVA). Different lowercase letters (a, b, c) indicate significant differences among biological treatments at $p < 0.05$ (one-way ANOVA followed by post hoc Tukey HSD test).

Table 1. Statistics (F and p-values) of two-way ANOVA performed on biomass accumulation in shoots and roots of *P. pinaster*.

Plants were grown for 100 days in a Cambisol supplemented with phytate, whether or not associated with the ectomycorrhizal fungus *H. cylindrosporum*, referred as "Mycorrhiza" (M) and whether or not inoculated with free-living microorganisms (bacteria and bacterial-feeding nematodes) referred as "Free-living organisms" (FO) as factors.

Factors	Dry biomass (g plant ⁻¹) in:	
	Shoot	Root
Mycorrhiza (M)	5.25 *	13.71 ***
Free-living organisms (FO)	1.81	2.56
M × FO	1.01	2.16

* p<0.05, ** p<0.01, *** p<0.001

Table 2. Root parameters measured in *P. pinaster* grown for 100 days in a Cambisol supplemented with phytate with different biological treatments.

Values are the means (n=6) with standard deviation between brackets. NM: non-mycorrhizal, M: mycorrhizal with *H. cylindrosporum*, None: no inoculation, +Bac: inoculation with *B. subtilis*, +Bac +Nem: inoculation with *B. subtilis* and *Rhabditis* sp.

Biological treatment		Root parameters (plant ⁻¹)				
		Length (cm)	Area (cm ²)	Tips (Number)	Forks (Number)	Crossings (Number)
NM	None	989 a (159)	150 a (19)	1851 a (387)	2726 ab (701)	428 a (143)
NM	+Bac	828 a (224)	133 a (30)	1745 ab (473)	2554 ab (930)	312 ab (125)
NM	+Bac +Nem	976 a (146)	154 a (23)	1768 ab (318)	3404 a (940)	472 a (161)
M	None	608 a (321)	107 a (50)	1200 bc (494)	1558 b (1267)	194 b (185)
M	+Bac	610 a (15)	112 a (5)	942 c (72)	1290 b (114)	142 b (20)
M	+Bac +Nem	647 a (90)	123 a (22)	1019 c (277)	1641 ab (407)	177 b (33)
NM vs M one-way ANOVA p-value		***	**	***	***	***

Different letters (a, b and c) indicate significant differences between biological treatments and stars (*p<0.05, **p<0.01, ***p<0.001) indicate significant differences between presence and absence of mycorrhiza according to one-way ANOVA followed by post hoc Tukey HSD test.

Table 3. Statistics (F and p-values) of two-way ANOVA performed on root parameters of *P. pinaster*.

Plants were grown for 100 days in a Cambisol supplemented with phytate, whether or not associated with the ectomycorrhizal fungus *H. cylindrosporum*, referred as "Mycorrhiza" (M) and whether or not inoculated with free-living microorganisms (bacteria and bacterial-feeding nematodes) referred as "Free-living organisms" (FO) as factors.

Factors	Length (cm)	Area (cm ²)	Tips (Number)	Forks (Number)	Crossings (Number)
Mycorrhiza (M)	24.86 ***	12.59 *	36.34 ***	32.03 ***	30.01 ***
Free-living organisms (FO)	0.87	1.12	0.79	1.90	2.04
M × FO	0.60	1.02	0.13	0.10	0.73

* p<0.05, ** p<0.01, *** p<0.001

Table 4. Statistics (F and p-values) of two-way ANOVA performed on P amount in shoots and roots of *P. pinaster*.

Plants were grown for 100 days in a Cambisol supplemented with phytate, whether or not associated with the ectomycorrhizal fungus *H. cylindrosporum*, referred as "Mycorrhiza" (M) and whether or not inoculated with free-living microorganisms (bacteria and bacterial-feeding nematodes) referred as "Free-living organisms" (FO) as factors.

Factors	Plant P (mg P plant ⁻¹) in:	
	Shoot	Root
Mycorrhiza (M)	4.42 *	1.76
Free-living organisms (FO)	7.42 **	23.94 ***
M × FO	1.58	4.08

* p<0.05, ** p<0.01, *** p<0.001

Table 5. Statistics (F and p-values) of two-way ANOVA performed on P acquisition efficiency of *P. pinaster*.

Plants were grown for 100 days in a Cambisol supplemented with phytate, whether or not associated with the ectomycorrhizal fungus *H. cylindrosporum*, referred as "Mycorrhiza" (M) and whether or not inoculated with free-living microorganisms (bacteria and bacterial-feeding nematodes) referred as "Free-living organisms" (FO) as factors.

Factors	P acquisition efficiency ($\mu\text{g P cm}^{-1}$ of roots)
Mycorrhiza (M)	51.61 ***
Free-living organisms (FO)	17.46 ***
M \times FO	1.4

* p<0.05, ** p<0.01, *** p<0.001

Table 6. Statistics (F and p-values) of two-way ANOVA performed on fungal (18S) rDNA gene copies number measured by qPCR in soil with *P. pinaster*.

Plants were grown for 100 days with the inoculation of ectomycorrhizal fungus referred as "Mycorrhiza" (M) and inoculation of free-living microorganisms (bacteria and bacterial-feeding nematodes) referred as "Free-living organisms" (FO) as factors.

Factors	18S rDNA gene copies number per gram of soil
Mycorrhiza (M)	194.71 ***
Free-living organisms (FO)	3.66 *
M × FO	3.36 *

* p<0.05, ** p<0.01, *** p<0.001