



**HAL**  
open science

## Adhesion to stainless steel surfaces and detection of viable but non cultivable cells of *Vibrio parahaemolyticus* and *Vibrio cholerae* isolated from shrimps in seafood processing environments: Stayin' alive?

Julia Mougin, Stéphanie Copin, Daline Bojolly, Virginie Ragueneau, Annick Robert-Pillot, Marie-Laure Quilici, Graziella Midelet-Bourdin, Thierry Grard, Maryse Bonnin-Jusserand

### ► To cite this version:

Julia Mougin, Stéphanie Copin, Daline Bojolly, Virginie Ragueneau, Annick Robert-Pillot, et al.. Adhesion to stainless steel surfaces and detection of viable but non cultivable cells of *Vibrio parahaemolyticus* and *Vibrio cholerae* isolated from shrimps in seafood processing environments: Stayin' alive?. *Food Control*, 2019, 102, pp.122-130. 10.1016/j.foodcont.2019.03.024 . hal-02620700

**HAL Id: hal-02620700**

**<https://hal.inrae.fr/hal-02620700>**

Submitted on 22 Oct 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Adhesion to stainless steel surfaces and detection of Viable But Non Cultivable cells of**  
2 ***Vibrio parahaemolyticus* and *Vibrio cholerae* isolated from shrimps in seafood processing**  
3 **environments: stayin' alive?**

4 Julia Mouglin<sup>a,b,c,d,e,1</sup>, Stéphanie Copin<sup>f,1</sup>, Daline Bojolly<sup>a</sup>, Virginie Raguene<sup>f</sup>, Annick Robert-  
5 Pillot<sup>g</sup>, Marie-Laure Quilici<sup>g</sup>, Graziella Midelet-Bourdin<sup>f</sup>, Thierry Grard<sup>a\*</sup>, Maryse Bonnin-  
6 Jusserand<sup>a</sup>

7 <sup>a</sup> *Univ. Littoral Côte d'Opale, convention ANSES, EA 7394 – ICV – Institut Charles Viollette,*  
8 *F-62321 Boulogne-sur-Mer, France*

9 <sup>b</sup> *INRA, France*

10 <sup>c</sup> *Univ. Lille F – 59000 Lille, France*

11 <sup>d</sup> *ISA, F – 59000 Lille, France*

12 <sup>e</sup> *Univ. Artois, F – 62000 Arras, France*

13 <sup>f</sup> *French Agency for Food, Environmental and Occupational Health & Safety (ANSES),*  
14 *Laboratory for Food Safety, Boulevard du Bassin Napoléon, 62200 Boulogne-sur-Mer,*  
15 *France*

16 <sup>g</sup> *Institut Pasteur, Paris, Unité des Bactéries Pathogènes Entériques, Centre National de*  
17 *Référence des Vibrions et du Choléra*

18 E-mail address of each author:

19 Julia Mouglin: [julia.mouglin@univ-littoral.fr](mailto:julia.mouglin@univ-littoral.fr)

20 Stéphanie Copin: [stephanie.copin@anses.fr](mailto:stephanie.copin@anses.fr)

21 Daline Bojolly: [daline.bojolly@gmail.com](mailto:daline.bojolly@gmail.com)

22 Virginie Raguene: [virginie.raguene@anses.fr](mailto:virginie.raguene@anses.fr)

23 Annick Robert-Pillot: [annick.robert-pillot@pasteur.fr](mailto:annick.robert-pillot@pasteur.fr)

24 Marie-Laure Quilici: marie-laure.quilici@pasteur.fr

25 Graziella Midelet-Bourdin: graziella.bourdin@anses.fr

26 Thierry Grard\*: thierry.grard@univ-littoral.fr

27 Maryse Bonnin-Jusserand: maryse.bonnin@univ-littoral.fr

28

29 \*Corresponding author: Thierry Grard

30 Tel: +33 3 21 99 25 08. E-mail address: thierry.grard@univ-littoral.fr

31 <sup>1</sup> J. Mougin and S. Copin share co-authorship of this article.

## 32 **Key words**

33 *Vibrio parahaemolyticus*; *Vibrio cholerae*; biofilm; shrimps; VBNC; PMA-qPCR; seafood  
34 industries

## 35 **Abstract**

36 A single strain of *Vibrio parahaemolyticus* and a single stain of *Vibrio cholerae* were isolated  
37 from shrimps sourced from seafood processing plants. The isolated strains were evaluated for  
38 their ability to adhere to stainless steel surfaces, under conditions encountered in seafood  
39 industries. In this way, the impact of different environmental factors such as temperature  
40 (8 °C or 37 °C) and culture media (tryptic soy broth (TSB) 2% NaCl, artificial sea water  
41 (ASW) or industrial brine) were studied. The viability of these strains was analyzed by three  
42 different methods: the enumeration of cultivable bacteria on agar media, and two PCR  
43 methods, i.e. a qPCR and a PMA-qPCR method in order to distinguish between viable and  
44 dead cells. The proportion of viable but non-cultivable (VBNC) cells was evaluated by  
45 comparing the results of propidium monoazide real-time PCR (PMA-qPCR) with the  
46 enumeration results. Cultivable bacteria were not detected in brine, regardless of the  
47 conditions tested. However, *V. cholerae* in the VBNC state was detected in brine at 8 °C until  
48 48h of incubation. *V. cholerae* exhibited higher viability at 8 °C compared to 37 °C. *V.*  
49 *cholerae* also exhibited higher viability compared to *V. parahaemolyticus*. In contrast, for *V.*  
50 *parahaemolyticus*, temperature and media (except industrial brine) had little influence on cell  
51 viability and adhesion behavior.

## 52 **1. Introduction**

53 Seafood can be contaminated by Gram-negative bacteria of the genus *Vibrio* present in coastal  
54 areas. In particular, *Vibrio parahaemolyticus* and *Vibrio cholerae* are two species potentially  
55 pathogenic for humans (Robert-Pillot, Copin, Himber, Gay, & Quilici, 2014). *V. cholerae* can  
56 be divided into two major groups: the cholera-causing strains of serogroups O1 and O139, and

57 non-O1/non-O139 *V. cholerae*. The pathogenicity of the strains depends on virulence factors,  
58 the most important of which is the cholera toxin (CT) encoded by the *ctxA* and *ctxB* genes.  
59 The non-O1/non-O139 strains rarely possess these cholera toxin genes but may also cause  
60 human illness. *V. parahaemolyticus* has been implicated in more than 45% of all outbreaks of  
61 bacterial gastroenteritis worldwide, and this proportion is increasing with the emergence of  
62 pandemic strains spreading over large geographic areas (Nair, et al., 2007). Typical *V.*  
63 *parahaemolyticus* illness results in mild to moderate acute gastroenteritis. As foodborne  
64 pathogens, *V. parahaemolyticus* and *V. cholerae* are responsible for vibriosis outbreaks and  
65 represent a risk to consumer health, especially in case of ingestion of raw, undercooked or  
66 mishandled seafood (Bonnin-Jusserand, et al., 2017); (M. F. R. Mizan, Jahid, & Ha, 2015).  
67 The risk associated with consumption of crustaceans, such as shrimps is well-documented for  
68 *V. parahaemolyticus* in several studies (Zarei, Borujeni, Jamnejad, & Khezzadeh, 2012);  
69 (Abdullah Sani, Ariyawansa, Babji, & Hashim, 2013); (Xu, et al., 2014) and for both *V.*  
70 *parahaemolyticus* and *V. cholerae* in others (Rosec, Causse, Cruz, Rauzier, & Carnat, 2012);  
71 (Yano, et al., 2014). *Vibrio* spp. can adopt specific survival strategies in harsh environments.  
72 They are found either free-living or associated with sediments, suspended particles and  
73 plankton (Thomas, Joseph, Raveendran, & Nair, 2006); (Vezzulli, et al., 2010); (Vezzulli,  
74 Colwell, & Pruzzo, 2013)). Their ability to adhere to the surface of the exoskeleton of animals  
75 composed of chitin, some zooplankton organisms, especially copepods (Zidour, et al., 2017),  
76 and to plastic marine debris (Zettler, Mincer, & Amaral-Zettler, 2013) allows them to survive  
77 in seawater longer than in a free-living state. *Vibrio* spp. have the ability to form organized  
78 structures called biofilms on various biotic and abiotic surfaces (Watnick & Kolter, 1999). It  
79 has been shown that *Vibrio* spp. are able to form biofilms on aquaculture equipment such as  
80 larval rearing tanks (Bourne, Høj, Webster, Swan, & Hall, 2006) as well as on several other  
81 substrates, including glass, polystyrene, polyethylene and polyvinyl chloride (Snoussi, et al.,

82 2009). *Vibrio* spp. biofilms have also been detected on fishing industry equipment (Bagge-  
83 Ravn, et al., 2003). Furthermore, biofilms represent a well-known hazard in the food industry  
84 (Shi & Zhu, 2009); (Bridier, et al., 2015), and many environmental factors influence biofilm  
85 formation. For example, recent studies on the role of additives currently used in seafood  
86 processing, like NaCl and glucose, on the biofilm formation of *V. parahaemolyticus* on  
87 shrimp surfaces have been carried out (M. F. Mizan, et al., 2016). Within the biofilm, cells  
88 persist in different physiological stages, including in the viable but non cultivable (VBNC)  
89 state. The VBNC state is defined as a form of resistance of bacteria to adverse environmental  
90 conditions. The cells conserve their metabolic activity but cultivability is lost. Therefore,  
91 bacteria cannot be detected on agar Petri dishes (Oliver & Bockian, 1995). *V.*  
92 *parahaemolyticus* (H. Zhong, et al., 2017); (Su, Jane, & Wong, 2013); (Mizunoe, Wai,  
93 Ishikawa, Takade, & Yoshida, 2000) and *V. cholerae* (Rowan, 2004) can enter the VBNC  
94 state. Importantly, the ISO standard 21872-1 applied to detect potentially pathogenic *Vibrio* in  
95 seafood cannot detect bacteria in the VBNC state. This could lead to underestimation of the  
96 bacterial population.

97 This study focused on the biofilm-forming ability on stainless steel of two different  
98 potentially pathogenic for humans *Vibrio* species (*V. cholerae* and *V. parahaemolyticus*)  
99 isolated from shrimps. The processing of shrimps in the seafood industry follows different  
100 steps (GBPH, 2016). Considering the possibility of a recontamination risk after cooking, brine  
101 could play a role by transporting *Vibrio*. Interestingly, there is no further critical control point  
102 (CCP) making it possible to limit the microbial population level after this step. Moreover,  
103 brine is recycled. Therefore, pathogen or spoilage flora could be present for instance in the  
104 VBNC state if renewal is not sufficiently frequent. VBNC cells could hardly be detected by  
105 classical microbiological methods and thus represent a potential risk for consumers.-In order  
106 to estimate the VBNC proportion of *V. parahaemolyticus* and *V. cholerae* within a biofilm

107 closest to industrial conditions, in our study, the experiments required the comparison of three  
108 different techniques: qPCR, PMA-qPCR (propidium monoazide real-time PCR) ((Q. Zhong,  
109 Tian, Wang, Fang, & Liao, 2018); (Zhu, Li, Jia, & Song, 2012); (Wu, Liang, & Kan, 2015))  
110 and enumeration on agar plates. The influence of environmental parameters such as  
111 temperature and media, including industrial brine, has been assessed. The aim of this study  
112 consists of assessing the adhesion to stainless steel surfaces and detecting Viable But Non  
113 Cultivable cells of *Vibrio parahaemolyticus* and *Vibrio cholerae* isolated from shrimps in  
114 seafood processing environments.

## 115 **2. Materials and methods**

### 116 **2.1 Bacterial strains and culture conditions**

117 Two environmental strains, isolated by the National Reference Laboratory (NRL: *Vibrio* sp.  
118 in fishery products), ANSES Boulogne-sur-mer, from seafood industries, were used in this  
119 study. The first one, *Vibrio parahaemolyticus* (ANSES collection 14-B3PA-0046) was  
120 isolated from shrimps imported from Nigeria. The second one, *Vibrio cholerae* (ANSES  
121 collection 09-B3PA-2350) was isolated from shrimps imported from Madagascar. The strains  
122 were maintained in marine broth with 10% glycerol and stored at - 80 °C. Prior to the  
123 experiments, bacteria were initially streaked on nutrient agar supplemented with 2% NaCl and  
124 incubated overnight at 37 °C. The strains were then transferred into tryptic soy broth (TSB  
125 2% NaCl) and incubated overnight at 37 °C.

126 Subsequently, cells were centrifuged at 5,000 g for 10 min and resuspended in TSB 2% NaCl  
127 or artificial sea water (ASW) or industrial brine (4 m<sup>3</sup> brine bath composed of water with a  
128 salinity of  $1.17 \pm 0.02$  (>1.10), pH 4). This step was repeated twice. Bacterial cell  
129 concentrations were adjusted to 10<sup>8</sup> CFU/mL by spectrophotometry (LibraS11, BioChrom  
130 Ltd., Cambridge, UK). In order to check the bacterial concentrations, suitable dilutions of  
131 suspensions were made and plated on salted nutrient agar using an automatic spiral plater  
132 (easySpiral<sup>®</sup>, Interscience, St-Nom-la-Bretèche, France). The plates were incubated at 37 °C

133 for 24 h. Cell numbers were then quantified using an automatic colony counter (Scan500<sup>®</sup>,  
134 Interscience).

## 135 **2.2 Preparation of 316 2B stainless steel coupons**

136 AISI 316 2B (Fe/Cr18/Ni10/Mo3) stainless steel coupons (37x16x1 mm) were purchased  
137 from Goodfellow SARL (Lille, France). First of all, 316 2B stainless steel coupons were  
138 rubbed with paper, soaked in a 50% acetone/50% ethanol (v/v) solution. Coupons were then  
139 immersed in a 20% alkali detergent solution (v/v) (RBS<sub>50</sub>, cleaning agent; Chemical Products  
140 R. Borghgraef, Brussels, Belgium) at 50 °C under agitation for 10 min. Coupons were rinsed  
141 in a 50 °C water bath under agitation for 20 min. Five successive rinses of 1 min each were  
142 carried out in ultra-pure water, while stirring at room temperature. Coupons were autoclaved  
143 for 15 min, at 121 °C.

## 144 **2.3 Biofilm formation on stainless steel coupons**

145 The bacterial suspension (11 mL) was transferred to a small sterile Petri dish (67 mm x 15  
146 mm) containing a sterile coupon. In order to create a humid atmosphere, the Petri dishes were  
147 then placed in bigger Petri dishes (140 mm x 20 mm) containing paper towels soaked with 25  
148 mL of sterilized distilled water. Sterile medium was used as a negative control. The Petri  
149 dishes were inoculated at 8 °C or 37 °C for 3 h, 24 h and 48 h.

## 150 **2.4 Quantification of viable, VBNC and dead cells attached to stainless steel coupons** 151 **after 3 h, 24 h and 48 h of incubation at either 37 °C or 8 °C**

152 After incubation, cell suspensions were removed and coupons were washed with 25 mL of  
153 sterile physiological water (0.9 % NaCl) in order to remove non-adhered bacteria. Washed  
154 coupons were then transferred to sterile Petri dishes.

155 Two coupons were used per experiment. Each coupon was swabbed twice with the same  
156 swab. This swab was vortexed into a tube containing 2 mL of physiological water. A second



157 swab, after undergoing the same process, was added to the same tube. This tube containing  
158 the two swabs was then vortexed for 20 sec. The contents of the tube were divided into three  
159 fractions. Each fraction underwent one of the following three techniques:

160 - First fraction: the proportion of viable cultivable bacteria was obtained by plating serial  
161 dilutions on salted nutrient agar using an automatic spiral plater (easySpiral<sup>®</sup>, Interscience).  
162 The plates were incubated at 37 °C for 24 h. Cell numbers were then quantified using an  
163 automatic colony counter (Scan500<sup>®</sup>, Interscience).

164 - Second fraction: qPCR was used to amplify DNA of viable, VBNC and dead bacteria.  
165 Genomic DNA from cell suspensions (495 µL) was extracted using the DNeasy<sup>®</sup> Blood &  
166 Tissue Kit 250 (Qiagen, Courtabœuf, France), following the manufacturer's instructions.  
167 Then, *V. parahaemolyticus* DNA was quantified by qPCR following the protocol described by  
168 Robert-Pillot, et al. (2014), with modifications. First at all, the LightCycler<sup>®</sup> 480 (Roche  
169 Diagnostics, France) was used for qPCR. The Qiagen Master Mix QuantiTect<sup>®</sup> Probe PCR Kit  
170 was used. This method was optimized for *V. cholerae*. Concentrations of reagents were  
171 adjusted to reach 4 mM Mg<sup>2+</sup> (Qiagen) and 6% glycerol using the Master Mix Light Cycler<sup>®</sup>  
172 480 Probes Master (Roche). The PCR cycling conditions were adjusted by including initial  
173 denaturation at 95 °C for 15 sec, followed by annealing at 61 °C for 1 min, extension at 72 °C  
174 for 15 sec.

175 - Third fraction: PMA-qPCR was used to amplify DNA of viable and VBNC bacteria.  
176 Propidium monoazide, PMA (Biotium Inc., Hayward, CA, USA) dissolved in ultra-pure water  
177 was added to 495 µL of cell suspension to reach a final concentration of 50 mM. Following an  
178 incubation period of 5 min in the dark, samples were light-exposed using a photo-activation  
179 system (PhAst Blue, GenIUL, Barcelona, Spain), with an intensity of 80 for 10 min. DNA  
180 from cell suspensions was extracted using the DNeasy<sup>®</sup> Blood & Tissue Kit 250 (Qiagen),  
181 following the manufacturer's instructions. Then, *V. parahaemolyticus* and *V. cholerae* DNA

182 was quantified by qPCR following the protocol described above for the second fraction. The  
183 genome equivalents (GEs) of the *V. parahaemolyticus* and *V. cholerae* strains were calculated  
184 based on a *V. parahaemolyticus* genome of  $5.0 \cdot 10^6$  bp (Makino, et al., 2003) and on a *V.*  
185 *cholerae* genome of  $4.0 \cdot 10^6$  bp (Heidelberg, et al., 2000).

## 186 **2.5 Microscopy observation**

187 A coupon was used to image biofilm samples on stainless steel. The biofilm was stained for  
188 15 min in the dark with 200  $\mu$ L taken from a LIVE/DEAD<sup>®</sup> BacLight<sup>™</sup> bacterial viability kit  
189 (Invitrogen, Carlsbad, CA, USA). Residual LIVE/DEAD<sup>®</sup> BacLight<sup>™</sup> solution was removed.  
190 Bacteria attached to the coupon were then examined in a wet state under an epifluorescence  
191 microscope (Imager.Z1, Zeiss, Marly-le-Roi, France) connected to a CCD camera (Axiocam -  
192 MRm, Zeiss) with apotome (x 40 magnification). A minimum of five observations were taken  
193 for each observed coupon.

## 194 **2.6 Statistical analysis**

195 Biofilm experiments were carried out in triplicate (three independent experiments). Data were  
196 analyzed by one-way analysis of variance (ANOVA) using R3.3.3 software (R Core Team,  
197 2017). The ANOVA model was verified using a Shapiro-Wilk normality test and Bartlett test  
198 of homogeneity of variances. If the normality and homogeneity of variances of distribution  
199 were not verified, a Kruskal-Wallis test was carried out, followed by a post hoc test. Post hoc  
200 tests were performed with Bonferroni correction and Fisher's least significant difference  
201 (LSD) criterion, using the agricolae 1.2-8 package (De Mendibru, 2017). The results were  
202 expressed as means  $\pm$  standard error. Statistical significance was considered at  $p < 0.05$ .

## 203 **3. Results**

204 *Vibrio parahaemolyticus* ANSES collection 14-B3PA-0046

205 *Total population (viable cultivable, VBNC and dead bacteria; qPCR)*

206 In TSB 2% NaCl and ASW broths, there was no impact of the incubation time at either 37 °C  
207 (Fig. 1) or 8 °C (Fig. 1) on the total population.

208 *Viable cultivable and VBNC bacteria (PMA-qPCR)*

209 In TSB 2% NaCl and ASW broths, the same proportion of bacteria was observed for 3 h (Fig.  
210 1.a, 1.b), 24 h (Fig. 1.c, 1.d) and 48 h (Fig. 1.e, 1.f) at either 37 °C or 8 °C.

211 No additional viable bacteria were detected in brine, regardless of the conditions (Fig. 1).

212 *Viable cultivable bacteria (enumeration on Petri dishes)*

213 A significant decrease in bacteria numbers was observed in TSB 2% NaCl broth depending on  
214 biofilm formation kinetics (3 h, 24 h and 48 h), regardless of the temperature ( $p<0.05$ ) (Fig.  
215 1). However, in ASW broth, there were more adherent bacteria at time point 24 h (Fig. 1.c,  
216 1.d) compared to 3 h (Fig. 1.a, 1.b) and 48 h (Fig. 1.e, 1.f), regardless of the temperature  
217 ( $p<0.05$ ). No viable cultivable bacteria were detected in brine at either 37 °C (Fig. 1) or 8 °C  
218 (Fig. 1).

219 *Vibrio cholerae* ANSES collection 09-B3PA-2350

220 *Total population (viable cultivable, VBNC and dead bacteria; qPCR)*

221 In TSB 2% NaCl broth, the total population had decreased at 37 °C at time point 48 h  
222 ( $p<0.05$ ) (Fig. 2.e). Otherwise, the total population was almost equivalent at 8 °C at 3 h (Fig.  
223 2.b), 24 h (Fig. 2.d) and 48 h (Fig. 2.f). In ASW broth, there were no differences regardless of  
224 the incubation time or the temperature. In brine, equivalent bacteria proportions were detected  
225 at 8 °C, regardless of the incubation time (Fig. 2.b, 2.d, 2.f). In contrast, bacterial populations  
226 had decreased at time point 48 h compared to 3 h and 24 h at 37 °C ( $p<0.05$ ) (Fig. 2.a, 2.c,  
227 2.e).

228 *Viable cultivable and VBNC bacteria (PMA-qPCR)*

229 In TSB 2% NaCl and ASW broths, the proportion of viable bacteria was equivalent regardless  
230 of the incubation time and temperature (Fig. 2). In brine, bacteria in the VBNC state were

231 detected at 8 °C for each incubation time, including 48 h. (Fig. 2.b, 2.d, 2.f). This means that  
232 the VBNC *V. cholerae* cells remained in brine at 8°C even for a long period of time.  
233 However, at 37 °C, at time point 48 h, viable bacteria were no longer detected (Fig. 2.e).  
234 Indeed, results showed no amplification curve of PMA-qPCR for an incubation time of 48 h  
235 at 37 °C in brine only. Thus, all *V. cholerae* cells were dead (Fig. 2.e).

#### 236 *Viable cultivable bacteria (enumeration on Petri dishes)*

237 In TSB 2% NaCl, the proportion of viable cultivable bacteria was not significant regardless of  
238 the incubation time and the temperature (Fig. 2). In ASW, cultivable *V. cholerae* were not  
239 detected at 37 °C regardless of the incubation time (Fig. 2.a, 2.c, 2.e). Furthermore, cultivable  
240 *V. cholerae* were not detected at 8 °C at time point 3 h (Fig. 2.b), but the proportion of  
241 bacteria was equivalent at 8 °C at 24 h (Fig. 2.d) and 48 h (Fig. 2.f) of incubation. No  
242 additional viable cultivable *V. cholerae* were detected in brine, regardless of the incubation  
243 time and temperature (Fig. 2).

#### 244 *Comparisons between V. cholerae and V. parahaemolyticus in brine for an incubation time of* 245 *48 h*

246 In Brine, for an incubation time of 48 h, temperature had significant effect on the adhesion of  
247 total bacteria regardless *Vibrio* species. Indeed, more bacteria were detected at 8 °C compared  
248 to 37 °C (Fig. 3.1).

249 VBNC cells of *V. parahaemolyticus* were no longer detected either at 37 °C or 8 °C, while  
250 VBNC cells of *V. cholerae* were detected at 8 °C only. According to these results, the  
251 adhesion of VBNC cells could be strain-dependent. Moreover, these results demonstrated that  
252 temperature has a significant effect on the adhesion of VBNC *V. cholerae* cells in brine (Fig.  
253 3.2).

254 Viable cultivable bacteria were not detected either at 37 °C or 8 °C on Petri dishes (Fig. 3.3).

#### 255 *Observations of adherent cells*

256 Micro-colonies were observed at 37 °C and 8 °C for *V. parahaemolyticus* at time points 3 h,  
257 24 h and 48 h (Fig. 4). In brine, aggregates were observed from the early stages of adhesion (3  
258 h), regardless of the temperature (Fig. 4.g, 4.p). These observations clearly showed a mix of  
259 live and dead bacteria during the establishment of these aggregates. Regarding *V. cholerae*  
260 (Fig. 5), dense cell clusters were observed from 3 h at 37 °C in TSB 2% NaCl (Fig. 5.a).  
261 These clusters decreased at 48 h, suggesting dispersal of the biofilm (Fig. 5.c). However, at  
262 8 °C at time point 48 h (Fig. 5.l), clusters especially composed of dead cells were still present.  
263 In ASW broth, minor changes in the structure of the biofilm were observed, regardless of the  
264 temperature and the incubation time (Fig. 5.d, 5.e, 5.f, 5.m, 5.n, 5.o). In brine, small  
265 aggregates were observed regardless of the temperature and time (Fig. 5.g, 5.h, 5.i, 5.p, 5.q,  
266 5.r). No further viable *V. cholerae* were observed in brine at 37 °C at time point 48 h (Fig.  
267 5.i). Viable bacteria could be VBNC cells not detected by the molecular methods used or Petri  
268 dish enumeration. This could be explained by a detection threshold level of the techniques  
269 used that may be too low.

### 270 **3. Discussion**

271 This study provides a deeper understanding of the viability of bacteria inside the biofilm.  
272 qPCR and PMA-qPCR methods were developed for both *Vibrio* species, in addition to cell  
273 enumeration on agar. Importantly, the combined findings of these three techniques and their  
274 comparison provided the proportions of viable cultivable, VBNC and dead bacteria.  
275 Additionally, microscopy using a LIVE/DEAD<sup>®</sup> staining kit coupled with Apotome  
276 technology enabled us to supplement the results obtained by the methods mentioned above. In  
277 this way, we were able to observe bacterial adhesion on stainless steel coupons. These  
278 observations were corroborated by results obtained by enumeration on agar plates and  
279 molecular biology (qPCR and PMA-qPCR methods).

280 Many factors influence biofilm formation: the bacteria themselves, that is to say their own  
281 genetics and physiology, the type of attachment surface, and the environment (pH,  
282 temperature, nutrient availability) (Whitehead & Verran, 2015). Among these factors,  
283 temperature seems to play a key role in *Vibrio* biofilm formation. In this study, small  
284 differences were observed regarding *V. parahaemolyticus* and *V. cholerae* biofilms at either  
285 37 °C or 8 °C, regardless of the incubation time (3 h, 24 h, 48 h). Indeed, the only significant  
286 influence of temperature determined was in brine but no influence was detected in TSB or  
287 ASW broths. Nonetheless, Ahmed, et al. (2018) showed that the biofilm formation of *V.*  
288 *parahaemolyticus* and *V. cholerae* was significantly higher at 37 °C and 25 °C than at 4 °C. In  
289 the study carried out by Beshiru and Igbinsosa (2018), *V. cholerae* showed better abilities to  
290 form biofilm at 21 °C and 37 °C, while the highest biofilm production of *V. parahaemolyticus*  
291 was observed at 30 °C in TSB under dynamic culture. Similarly, Song, et al. (2017) observed  
292 better biofilm production of *V. parahaemolyticus* in TSB 3% NaCl at 25 °C compared to  
293 15 °C and 37 °C. Han, Mizan, Jahid, and Ha (2016) showed that *V. parahaemolyticus* biofilm  
294 formation was better at 30 °C than 4 °C on stainless steel coupons immersed in fresh water.  
295 Furthermore, H.-C. Wong, Chung, and Yu (2002) showed that bacterial adhesion on stainless  
296 steel surfaces decreased when *V. parahaemolyticus* was incubated at 4 °C. All these results  
297 showed better biofilm formation for high temperatures (between 21 °C to 37 °C) compared to  
298 4 °C, contrary to our findings.

299 Adhesion is also media-dependent. In this study, we did not observe the same behavior on  
300 either TSB 2 % NaCl or ASW between *Vibrio* spp. *V. parahaemolyticus* exhibited the same  
301 behavior in TSB or ASW, regardless of the incubation temperature. However, in ASW for 24  
302 h and 48 h, better survival was observed for *V. cholerae* at 8 °C compared to 37 °C due to the  
303 presence of viable cultivable bacteria. The amount of viable cultivable bacteria at 37 °C in  
304 ASW was not higher than in TSB. H.-C. Wong, et al. (2002) showed that *V. parahaemolyticus*

305 exhibited better adhesion on stainless steel when cultured in chemically defined MM9  
306 medium than TSB. Regarding *V. cholerae*, this behavior was dependent on the temperature of  
307 incubation. At 8 °C for 24 h and 48 h, there were no differences in adhesion on either TSB  
308 2% NaCl or ASW. The fact that a nutrient-poor medium promotes adhesion has already been  
309 shown especially for *L. monocytogenes* (Kadam, et al., 2013). The composition of ASW  
310 medium is close to that of the marine environment, which is the natural habitat of *Vibrio*.  
311 Therefore, better biofilm formation ability could be expected in this medium, which was not  
312 the case in our study. Moreover, Beshiru and Igbinosa (2018) showed that *V. cholerae* and *V.*  
313 *parahaemolyticus* biofilm formation was not medium-dependent (even when the medium was  
314 rich or poor). Both tested strains also exhibited different behaviors in brine. Indeed, in this  
315 study, only dead bacteria were detected regardless of the temperature and incubation time for  
316 *V. parahaemolyticus*. However, VBNC bacteria of *V. cholerae* were detected at 37 °C for 3 h  
317 and 24 h. VBNC cells were detected for each incubation time at 8 °C. These results indicate a  
318 greater survival capacity of *V. cholerae* at 8 °C than at 37 °C in poor media. The induction of  
319 the VBNC state is triggered by a low temperature, more often 4 °C, and under starvation  
320 conditions (Q. Zhong, et al., 2018); (Su, et al., 2013); (H. C. Wong, Wang, Chen, & Chiu,  
321 2004). This could explain the presence of VBNC cells of *V. cholerae* in brine at 8 °C.

322 As for temperature and media, bacterial adhesion depends on the food contact surface.  
323 Stainless steel coupons were used in this study to mimic the conditions encountered in  
324 seafood industries. Also, Han, et al. (2016) studied *V. parahaemolyticus* biofilm formation on  
325 stainless steel, crab and shrimp. In addition to surface hydrophoby, cell hydrophoby must also  
326 be taken into account: *Vibrio* are strongly hydrophilic (Beshiru & Igbinosa, 2018). However,  
327 it is difficult to compare studies of *Vibrio* biofilms with one another because, most of the  
328 time, biofilm formation was studied *via* the microtiter plate assay with crystal violet staining.  
329 This assay remains a useful screening tool, but does not reflect real biofilm formation

330 conditions; in particular, the surface type is not considered. Moreover, *Vibrio* biofilm  
331 formation is strain-dependent (Song, et al., 2017); (Odeyemi & Ahmad, 2017); (M. F. Mizan,  
332 et al., 2016); (H.-C. Wong, et al., 2002). To supplement our results, different *V.*  
333 *parahaemolyticus* and *V. cholerae* strains of different origins (isolated from different seafood  
334 products or different countries), including pathogenic strains (carrying pathogenicity factors  
335 such as *ctxA/ctxB* genes for *V. cholerae* and *tdh* or *trh* genes for *V. parahaemolyticus*) should  
336 be assessed. Clearly, biofilm formation may be different depending on the strain's origin. The  
337 formation of mixed biofilm, for example *Vibrio* in association with *Listeria monocytogenes*,  
338 could also be evaluated. In addition to developing survival strategies (VBNC, biofilm), strains  
339 are constantly adapting to their environment. More and more studies point to the extent of  
340 antibiotic resistance for *Vibrio* (Yano, et al., 2014). This characteristic should therefore be  
341 taken into account in the biofilm survival state. Elexson, et al. (2014) tested several detergents  
342 against biofilms of antibiotic-resistant *V. parahaemolyticus* isolated from seafood and found  
343 hygiene issues and cross-contamination hazards in the seafood industries.

#### 344 **4. Conclusion**

345 For the first time, the cell viability was taken into account in order to better understand the  
346 survival strategies of potentially pathogenic human *Vibrio*. The aim of this study was to get as  
347 close as possible to the conditions encountered in seafood industries, even if this test remained  
348 *in vitro* for the standardization and exploitation purposes of the results. This scoping study  
349 highlighted the behavior of both *Vibrio* strains under laboratory conditions for different  
350 environmental parameters. Despite the fact that there are few data on the risk for consumer  
351 health of *V. cholerae* from shrimps, the strain tested in this study showed a greater survival  
352 capacity than the *V. parahaemolyticus* strain. Indeed, VBNC *V. cholerae* cells were detected  
353 at 8 °C in brine for each incubation time, including 48 h. Therefore, it could be useful to  
354 monitor *V. cholerae* contamination in shrimps. Seafood processing companies need to be



355 aware of and take into account the *Vibrio* risk in their hazard analysis critical control point  
356 (HACCP) approach. The possible presence of VBNC cells cannot be ruled out. The use of  
357 molecular biology methods like PMA-qPCR is recommended to ensure the safety of seafood  
358 products.

### 359 **Acknowledgements**

360 Julia Mougin would like to thank the “Hauts-de-France” regional council for its financial  
361 support of her PhD studies. This work was funded by the French government and the region  
362 Hauts-de-France in the framework of the CPER 2014-2020 MARCO project. This work was  
363 supported by a grant from FranceAgriMer (2014-0813) as part of the SURVIB project. The  
364 authors also thank the CITPPM (*Confédération des industries de traitement des produits des*  
365 *pêches maritimes et de l'aquaculture*) as a contributor to the SURVIB project. The authors are  
366 also grateful to Thomas Brauge for his scientific assistance, to Cindy Ducrocq and Marie-  
367 Ange Huchin for technical assistance and to Alexandre Dehaut for advices in statistical  
368 analyses.

### 369 **References**

- 370 Abdullah Sani, N., Ariyawansa, S., Babji, A. S., & Hashim, J. K. (2013). The risk assessment of  
371 *Vibrio parahaemolyticus* in cooked black tiger shrimps (*Penaeus monodon*) in Malaysia.  
372 *Food control*, 31(2), 546-552.
- 373 Ahmed, H. A., El Bayomi, R. M., Hussein, M. A., Khedr, M. H. E., Abo Remela, E. M., & El-  
374 Ashram, A. M. M. (2018). Molecular characterization, antibiotic resistance pattern and  
375 biofilm formation of *Vibrio parahaemolyticus* and *V. cholerae* isolated from crustaceans  
376 and humans. *Int. J. Food Microbiol.*, 274, 31-37.
- 377 Bagge-Ravn, D., Ng, Y., Hjelm, M., Christiansen, J. N., Johansen, C., & Gram, L. (2003). The  
378 microbial ecology of processing equipment in different fish industries - Analysis of the  
379 microflora during processing and following cleaning and disinfection. *Int. J. Food*  
380 *Microbiol.*, 87(3), 239-250.
- 381 Beshiru, A., & Igbiosa, E. O. (2018). Characterization of extracellular virulence properties and  
382 biofilm-formation capacity of *Vibrio* species recovered from ready-to-eat (RTE) shrimps.  
383 *Microb. Pathog.*, 119, 93-102.
- 384 Bonnin-Jusserand, M., Copin, S., Le Bris, C., Brauge, T., Gay, M., Brisabois, A., Grard, T., &  
385 Midelet-Bourdin, G. (2017). *Vibrio* species involved in seafood-borne outbreaks (*Vibrio*  
386 *cholerae*, *V. parahaemolyticus* and *V. vulnificus*): review of microbiological versus recent  
387 molecular detection methods in seafood products. *Crit. Rev. Food Sci. Nutr.*, 1-14.
- 388 Bourne, D. G., Høj, L., Webster, N. S., Swan, J., & Hall, M. R. (2006). Biofilm development within  
389 a larval rearing tank of the tropical rock lobster, *Panulirus ornatus*. *Aquaculture*, 260(1-  
390 4), 27-38.

391 Bridier, A., Sanchez-Vizuete, P., Guilbaud, M., Piard, J.-C., Naïtali, M., & Briandet, R. (2015).  
392 Biofilm-associated persistence of food-borne pathogens. *Food Microbiol.*, *45*, 167-178.

393 De Mendibru, F. A. (2017). Statistical procedures for agricultural research. R package version  
394 1.2-4. 2016. In.

395 Elexson, N., Afsah-Hejri, L., Rukayadi, Y., Soopna, P., Lee, H. Y., Tuan Zainazor, T. C., Nor Ainy,  
396 M., Nakaguchi, Y., Mitsuaki, N., & Son, R. (2014). Effect of detergents as antibacterial  
397 agents on biofilm of antibiotics-resistant *Vibrio parahaemolyticus* isolates. *Food control*,  
398 *35*(1), 378-385.

399 GBPH. (2016). *Guides de bonnes pratiques d'hygiène : crustacés cuits* (Vol. 5961).

400 Han, N., Mizan, M. F. R., Jahid, I. K., & Ha, S.-D. (2016). Biofilm formation by *Vibrio*  
401 *parahaemolyticus* on food and food contact surfaces increases with rise in temperature.  
402 *Food control*, *70*, 161-166.

403 Heidelberg, J. F., Eisen, J. A., Nelson, W. C., Clayton, R. A., Gwinn, M. L., Dodson, R. J., Haft, D.  
404 H., Hickey, E. K., Peterson, J. D., & Umayam, L. (2000). DNA sequence of both  
405 chromosomes of the cholera pathogen *Vibrio cholerae*. *Nature*, *406*(6795), 477.

406 Kadam, S. R., den Besten, H. M., van der Veen, S., Zwietering, M. H., Moezelaar, R., & Abee, T.  
407 (2013). Diversity assessment of *Listeria monocytogenes* biofilm formation: impact of  
408 growth condition, serotype and strain origin. *Int. J. Food Microbiol.*, *165*(3), 259-264.

409 Makino, K., Oshima, K., Kurokawa, K., Yokoyama, K., Uda, T., Tagomori, K., Iijima, Y., Najima,  
410 M., Nakano, M., Yamashita, A., Kubota, Y., Kimura, S., Yasunaga, T., Honda, T.,  
411 Shinagawa, H., Hattori, M., & Iida, T. (2003). Genome sequence of *Vibrio*  
412 *parahaemolyticus*: a pathogenic mechanism distinct from that of *V cholerae*. *Lancet*,  
413 *361*(9359), 743-749.

414 Mizan, M. F., Jahid, I. K., Kim, M., Lee, K. H., Kim, T. J., & Ha, S. D. (2016). Variability in  
415 biofilm formation correlates with hydrophobicity and quorum sensing among *Vibrio*  
416 *parahaemolyticus* isolates from food contact surfaces and the distribution of the genes  
417 involved in biofilm formation. *Biofouling*, *32*(4), 497-509.

418 Mizan, M. F. R., Jahid, I. K., & Ha, S.-D. (2015). Microbial biofilms in seafood: A food-hygiene  
419 challenge. *Food Microbiol.*, *49*, 41-55.

420 Mizunoe, Y., Wai, S. N., Ishikawa, T., Takade, A., & Yoshida, S.-i. (2000). Resuscitation of viable  
421 but nonculturable cells of *Vibrio parahaemolyticus* induced at low temperature under  
422 starvation. *FEMS Microbiol. Lett.*, *186*(1), 115-120.

423 Nair, G. B., Ramamurthy, T., Bhattacharya, S. K., Dutta, B., Takeda, Y., & Sack, D. A. (2007).  
424 Global dissemination of *Vibrio parahaemolyticus* serotype O3: K6 and its serovariants.  
425 *Clin. Microbiol. Rev.*, *20*(1), 39-48.

426 Odeyemi, O. A., & Ahmad, A. (2017). Population dynamics, antibiotics resistance and biofilm  
427 formation of *Aeromonas* and *Vibrio* species isolated from aquatic sources in Northern  
428 Malaysia. *Microb. Pathog.*, *103*, 178-185.

429 Oliver, J. D., & Bockian, R. (1995). In vivo resuscitation, and virulence towards mice, of viable but  
430 nonculturable cells of *Vibrio vulnificus*. *Appl. Environ. Microbiol.*, *61*(7), 2620-2623.

431 R Core Team. (2017). R: A language and environment for statistical computing (Version 3.4.  
432 2)[Computer software]. Vienna, Austria: R Foundation for Statistical Computing.

433 Robert-Pillot, A., Copin, S., Himber, C., Gay, M., & Quilici, M.-L. (2014). Occurrence of the three  
434 major *Vibrio* species pathogenic for human in seafood products consumed in France using  
435 real-time PCR. *Int. J. Food Microbiol.*, *189*, 75-81.

436 Rosec, J.-P., Causse, V., Cruz, B., Rauzier, J., & Carnat, L. (2012). The international standard  
437 ISO/TS 21872-1 to study the occurrence of total and pathogenic *Vibrio parahaemolyticus*  
438 and *Vibrio cholerae* in seafood: ITS improvement by use of a chromogenic medium and  
439 PCR. *Int. J. Food Microbiol.*, *157*, 189-194.

440 Rowan, N. J. (2004). Viable but non-culturable forms of food and waterborne bacteria: Quo Vadis?  
441 *Trends Food Sci. Technol.*, *15*(9), 462-467.

442 Shi, X., & Zhu, X. (2009). Biofilm formation and food safety in food industries. *Trends Food Sci.*  
443 *Technol.*, *20*(9), 407-413.

444 Snoussi, M., Noumi, E., Hajlaoui, H., Usai, D., Sechi, L. A., Zanetti, S., & Bakhrouf, A. (2009).  
445 High potential of adhesion to abiotic and biotic materials in fish aquaculture facility by  
446 *Vibrio alginolyticus* strains. *J. Appl. Microbiol.*, *106*(5), 1591-1599.

- 447 Song, X., Ma, Y., Fu, J., Zhao, A., Guo, Z., Malakar, P. K., Pan, Y., & Zhao, Y. (2017). Effect of  
448 temperature on pathogenic and non-pathogenic *Vibrio parahaemolyticus* biofilm  
449 formation. *Food control*, *73*, 485-491.
- 450 Su, C. P., Jane, W. N., & Wong, H. C. (2013). Changes of ultrastructure and stress tolerance of  
451 *Vibrio parahaemolyticus* upon entering viable but nonculturable state. *Int. J. Food*  
452 *Microbiol.*, *160*(3), 360-366.
- 453 Thomas, K. U., Joseph, N., Raveendran, O., & Nair, S. (2006). Salinity-induced survival strategy  
454 of *Vibrio cholerae* associated with copepods in Cochin backwaters. *Mar. Pollut. Bull.*,  
455 *52*(11), 1425-1430.
- 456 Vezzulli, L., Colwell, R. R., & Pruzzo, C. (2013). Ocean warming and spread of pathogenic *Vibrios*  
457 in the aquatic environment. *Microb. Ecol.*, *65*(4), 817-825.
- 458 Vezzulli, L., Previati, M., Pruzzo, C., Marchese, A., Bourne, D. G., & Cerrano, C. (2010). *Vibrio*  
459 infections triggering mass mortality events in a warming Mediterranean Sea. *Environ.*  
460 *Microbiol.*, *12*(7), 2007-2019.
- 461 Watnick, P. I., & Kolter, R. (1999). Steps in the development of a *Vibrio cholerae* El Tor biofilm.  
462 *Mol. Microbiol.*, *34*(3), 586-595.
- 463 Whitehead, K. A., & Verran, J. (2015). Formation, architecture and functionality of microbial  
464 biofilms in the food industry. *Current Opinion in Food Science*, *2*, 84-91.
- 465 Wong, H.-C., Chung, Y.-C., & Yu, J.-A. (2002). Attachment and inactivation of *Vibrio*  
466 *parahaemolyticus* on stainless steel and glass surface. *Food Microbiol.*, *19*(4), 341-350.
- 467 Wong, H. C., Wang, P., Chen, S. Y., & Chiu, S. W. (2004). Resuscitation of viable but non-  
468 culturable *Vibrio parahaemolyticus* in a minimum salt medium. *FEMS Microbiol. Lett.*,  
469 *233*(2), 269-275.
- 470 Wu, B., Liang, W., & Kan, B. (2015). Enumeration of viable non-culturable *Vibrio cholerae* using  
471 propidium monoazide combined with quantitative PCR. *J. Microbiol. Methods*, *115*, 147-  
472 152.
- 473 Xu, X., Wu, Q., Zhang, J., Cheng, J., Zhang, S., & Wu, K. (2014). Prevalence, pathogenicity, and  
474 serotypes of *Vibrio parahaemolyticus* in shrimp from Chinese retail markets. *Food*  
475 *control*, *46*, 81-85.
- 476 Yano, Y., Hamano, K., Satomi, M., Tsutsui, I., Ban, M., & Aue-umneoy, D. (2014). Prevalence and  
477 antimicrobial susceptibility of *Vibrio* species related to food safety isolated from shrimp  
478 cultured at inland ponds in Thailand. *Food control*, *38*, 30-36.
- 479 Zarei, M., Borujeni, M. P., Jamnejad, A., & Khezzadeh, M. (2012). Seasonal prevalence of *Vibrio*  
480 species in retail shrimps with an emphasis on *Vibrio parahaemolyticus*. *Food control*, *25*,  
481 107-109.
- 482 Zettler, E. R., Mincer, T. J., & Amaral-Zettler, L. A. (2013). Life in the "plastisphere": microbial  
483 communities on plastic marine debris. *Environ. Sci. Technol.*, *47*(13), 7137-7146.
- 484 Zhong, H., Zhong, Y., Deng, Q., Zhou, Z., Guan, X., Yan, M., Hu, T., & Luo, M. (2017). Virulence of  
485 thermolabile haemolysin *tlh*, gastroenteritis related pathogenicity *tdh* and *trh* of the  
486 pathogens *Vibrio Parahemolyticus* in Viable but Non-Culturable (VBNC) state. *Microb.*  
487 *Pathog.*, *111*, 352-356.
- 488 Zhong, Q., Tian, J., Wang, J., Fang, X., & Liao, Z. (2018). iTRAQ-based proteomic analysis of the  
489 viable but nonculturable state of *Vibrio parahaemolyticus* ATCC 17802 induced by food  
490 preservative and low temperature. *Food control*, *85*, 369-375.
- 491 Zhu, R. G., Li, T. P., Jia, Y. F., & Song, L. F. (2012). Quantitative study of viable *Vibrio*  
492 *parahaemolyticus* cells in raw seafood using propidium monoazide in combination with  
493 quantitative PCR. *J. Microbiol. Methods*, *90*(3), 262-266.
- 494 Zidour, M., Chevalier, M., Belguesmia, Y., Cudennec, B., Grard, T., Drider, D., Souissi, S., &  
495 Flahaut, C. (2017). Isolation and characterization of bacteria colonizing *Acartia tonsa*  
496 copepod eggs and displaying antagonist effects against *Vibrio anguillarum*, *Vibrio*  
497 *alginolyticus* and other pathogenic strains. *Front. Microbiol.*, *8*(1919).
- 498

499 **Figures caption:**

500 **Figure 1:** *Vibrio parahaemolyticus* ANSES collection 14-B3PA-0046 at 37 °C (a,c,e) and  
501 8 °C (b,d,f): enumeration of total bacteria: viable cultivable, VBNC, and dead bacteria  
502 (qPCR: black squares); viable cultivable and VBNC bacteria (PMA-qPCR: white squares);  
503 viable cultivable bacteria (on Petri dishes: grey squares) for 3 h, 24 h and 48 h of incubation  
504 on TSB (2% NaCl), ASW and brine media. Values are expressed by means,  $\pm$  95 %  
505 confidence interval, of three independent experiments.

506 **Figure 2:** *Vibrio cholerae* ANSES collection 09-B3PA-2350 at 37 °C (a,c,e) and 8 °C (b,d,f):  
507 enumeration of total bacteria (qPCR: black squares); viable and VBNC bacteria (PMA-qPCR:  
508 white squares); viable cultivable bacteria (on Petri dishes: grey squares) for 3 h, 24 h and 48 h  
509 of incubation on TSB (2% NaCl), ASW and brine media. Values are expressed by means,  $\pm$   
510 95 % confidence interval, of three independent experiments.

511 **Figure 3:** *Vibrio cholerae* ANSES collection 09-B3PA-2350 (hatched square) and *Vibrio*  
512 *parahaemolyticus* ANSES collection 14-B3PA-0046 (spotted square) in brine for 48 h, at  
513 37 °C and 8 °C; Enumeration of (1) total bacteria (qPCR); (2) Viable and VBNC bacteria  
514 (PMA-qPCR); (3) Viable cultivable bacteria (on Petri dishes). Values are expressed by  
515 means,  $\pm$  95 % confidence interval, of three independent experiments. The letters a and b  
516 represent differences between treatments with two temperatures, according to post hoc tests.  
517 The “\*” represents differences between strains at the same temperature, according to post hoc  
518 tests.


519 **Figure 4:** *Vibrio parahaemolyticus* ANSES collection 14-B3PA-0046 at 37 °C  
520 (a,b,c,d,e,f,g,h,i) and 8 °C (j,k,l,m,n,o,p,q,r): relevant microscopic photographs of stainless  
521 steel coupons after LIVE/DEAD<sup>®</sup>BacLight<sup>™</sup> staining for 3 h, 24 h and 48 h of incubation on  
522 TSB (2% NaCl), ASW and brine media.

523 Experiments were repeated three times and a minimum of five images were taken for each  
524 observed stainless steel coupon.


525 **Figure 5:** *Vibrio cholerae* ANSES collection 09-B3PA-2350 at 37 °C (a,b,c,d,e,f,g,h,i) and  
526 8 °C (j,k,l,m,n,o,p,q,r): relevant microscopic photographs of stainless steel coupons after  
527 LIVE/DEAD®BacLight™ staining for 3 h, 24 h and 48 h of incubation on TSB (2% NaCl),  
528 ASW and brine media.

529 Experiments were repeated three times and a minimum of five images were taken for each  
530 observed stainless steel coupon.

*Vibrio parahaemolyticus* (ANSES collection 14-B3PA-0046)



*Vibrio cholerae* (ANSES collection 09-B3PA-2350)




■ Viable cultivable, VBNC and dead bacteria (qPCR)


□ Viable cultivable and VBNC bacteria (PWA-qPCR)

■ Viable cultivable bacteria (Enumeration on agar medium)

## Brine, 48 h





*Vibrio parahaemolyticus* (ANSES collection 14-B3PA-0046)

*Vibrio cholerae* (ANSES collection 09-B3PA-2350)