



HAL
open science

Repellent activity of plants from the genus *Chenopodium* to *Ostrinia nubilalis* larvae

Dariusz Piesik, Didier Rochat, Jan Bocianowski, Frédéric Marion-Poll

► **To cite this version:**

Dariusz Piesik, Didier Rochat, Jan Bocianowski, Frédéric Marion-Poll. Repellent activity of plants from the genus *Chenopodium* to *Ostrinia nubilalis* larvae. *Plant Protection Science*, 2018, 54 (4), pp.265 - 271. 10.17221/143/2017-PPS . hal-02621436

HAL Id: hal-02621436

<https://hal.inrae.fr/hal-02621436v1>

Submitted on 26 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Repellent Activity of Plants from the Genus *Chenopodium* to *Ostrinia nubilalis* Larvae

DARIUSZ PIESIK^{1*}, DIDIER ROCHAT², JAN BOCIANOWSKI³ and FRÉDÉRIC MARION-POLL^{2,4}

¹Department of Entomology and Molecular Phytopathology, UTP University of Science and Technology, Bydgoszcz, Poland; ²UMR N°1272 INRA/Université Paris 6/AgroParisTech, Physiologie de l'Insecte: Signalisation et Communication, INRA Centre de Versailles-Grignon, Versailles Cedex, France; ³Department of Mathematical and Statistical Methods, Poznań University of Life Sciences, Poznań, Poland; ⁴AgroParisTech, Université Paris-Saclay, Paris, France

*Corresponding author: piesik@utp.edu.pl

Abstract

Piesik D., Rochat D., Bocianowski J., Marion-Poll F. (2018): Repellent activity of plants from the genus *Chenopodium* to *Ostrinia nubilalis* larvae. Plant Protect. Sci., 54: 265–271.

The olfactory choices expressed by naïve neonate European corn borer (ECB) larvae were tested by recording their locomotor activities in response to odours coming from plants of the genus *Chenopodium* (producing phytoecdysteroids or not). ECB larvae were found to be repelled from phytoecdysteroid-positive species, except *C. album* and *C. polyspermum*. On the contrary, they were mildly attracted or mildly repelled by phytoecdysteroid-negative species, except *C. botrys* (which emits a rancid odour). These observations indicate that neonate ECB larvae clearly differentiate plant odours and suggest that well-defended plants may emit repellent odours.

Keywords: European corn borer; odours; larvae; orientation; behaviour

Ecdysteroids are known insect moulting hormones that promote growth, moulting and metamorphosis (COLL *et al.* 2007; GELMAN *et al.* 2007). A number of plants synthesise and accumulate such hormones, which contribute to defend them against phytophagous insects (SIMON *et al.* 2004) and nematodes (SORIANO *et al.* 2004). Phytoecdysteroids (PEs) constitute a family of plant steroidal analogues of invertebrate steroid hormones, which include 503 molecules (LAFONT *et al.* 2002). They are detectable in 5–6% of higher plant species, frequently in leaves and flowers, but less so in stems, roots and seeds (DINAN *et al.* 2001). Their levels vary enormously between species and the distribution of ecdysteroid-containing species is only partly understood (DINAN *et al.* 2001; TARKOWSKA & STRNAD 2016).

The most common phytoecdysteroids are 20-hydroxyecdysone (BLACKFORD *et al.* 1996; DINAN *et al.* 2002), polypodine B, makisterone A, and ponasterone A (VOIGT *et al.* 2001). It has been shown that phytoecdysteroid concentrations increase in response to mechanical damage, insect herbivory, and application of methyl jasmonate (SCHMELZ *et al.* 1999, 2002). Phytoecdysteroids are believed to deter invertebrate predators and they play a defence role against phytophagous insects (DINAN 1998). In the tolerant insect species there is a rapid detoxification (as C-22 long-chain fatty acyl esters) and/or excretion of ingested ecdysteroids (BLACKFORD & DINAN 1997a,b).

The question asked here is if phytoecdysteroid-producing plants would emit specific deterring or repellent odours that would deter non-tolerant phy-

Supported by the Marie-Curie Foundation (Intra-European Fellowships), Grant No. MEIF-CT-2005-514931 (D. Piesik and F. Marion-Poll).

doi: 10.17221/143/2017-PPS

tophagous herbivores at distance. As an example of a non-tolerant phytophagous insect, European corn borer (*Ostrinia nubilalis* Hubner) was selected whose individuals are deterred from feeding by phytoecdysteroids at the larval stage (MARION-POLL & DESCOINS 2002) and from laying eggs at the adult stage (CALAS *et al.* 2007). It is reasoned that such an insect would benefit from an early detection of plant-producing ecdysteroids, in particular through sensing odours emitted by such plants. Plants from the family *Chenopodiaceae* which include plants that produce PEs and plants which do not produce these substances were selected (DINAN 1995; VOLODIN *et al.* 2002).

In this study, volatiles released by phytoecdysteroid-positive *Chenopodium* plants were analysed in order to find repellent compounds. The hypothesis that odours from such plants would repel European corn borer larvae when phytoecdysteroid-negative plants would be less repellent was also tested.

MATERIAL AND METHODS

Plant culture. Experiments were performed at the Plant Growth Centre, INRA, Versailles, France. Seeds for this study were obtained from the National Botanic Garden of Belgium. *Chenopodium album*, *C. ambrosioides*, *C. glaucum*, *C. polyspermum*, *C. schraderanum*,

C. urbicum, *C. amaranticolor*, *C. botrys*, *C. ficifolium*, *C. murale*, *C. vulvaria*, and *C. oleracea* were tested. The plants (3 per pot) were sown and grown in individual pots in a greenhouse. Daytime temperature was $22 \pm 2^\circ\text{C}$, overnight temperature was $18 \pm 2^\circ\text{C}$. According to DINAN (1992), there is a clear relationship between the amount of phytoecdysteroids in the vegetative parts and the levels found in the flowers and seeds, so that it is assumed that the tests performed on the seeds reflect the presence of phytoecdysteroids in the leaves (Table 1). Three green plants of each species were tested after 6 weeks of vegetation.

Insects. Behavioural experiments were performed on freshly hatched European corn borer (ECB) larvae not exposed to the food. The adult males and females (provided by INRA le Magneraud) were maintained in a cage at $25 \pm 0.5^\circ\text{C}$ and 70–80% relative humidity in a climatic chamber (16-h day/8-h night) and fed on sugared water (20%). Females laid eggs on wetted filter paper and put in glass vials (\varnothing 12 mm ID; 7.5 mm long). Each vial was sealed with a cotton plug, and put in a plastic box (25 × 15 × 10 cm). Vial plugs were lightly wetted daily with distilled water to prevent desiccation. The temperature was maintained at $25 \pm 0.5^\circ\text{C}$ and 70–80% relative humidity. Neonate larvae hatched from these papers were collected immediately after hatching and used for experiments. Each larva was used only once to avoid learning.

Table 1. Distribution of phytoecdysteroids in relation to the taxonomy of the plants of the genus *Chenopodium* (DINAN *et al.* 1998)

<i>Chenopodiaceae</i>	RIA ($\mu\text{g E. eg. g seed}$)	Against bioassay
<i>S. oleracea</i> L.	1185/1125/1340/2546 1033/1087/1204/1104	+++ / +++ / +++ / +++ +++ / +++ / +++ / +++
<i>Ch. album</i> L.	25/58/1370/1753/1294 428/957/790/499	+ / +++ / +++ / +++ / +++ ++ / +++ / +++ / +++
<i>Ch. amaranticolor</i> H.J. Coste and A. Reynie	1362	+++
<i>Ch. ficifolium</i> Sm.	481/975/1033	+++ / +++ / +++
<i>Ch. murale</i> L.	885/1297/169/117	+++ / +++ / +++ / +++
<i>Ch. vulvaria</i> L.	708/1258/984	+++ / +++ / +++
<i>Ch. ambrosioides</i> L.	–	–
<i>Ch. botrys</i> L.	–	–
<i>Ch. glaucum</i> L.	–	–
<i>Ch. polyspermum</i> L.	1975/2/1672/0/1477/2016	+++ / – / +++ / – / +++ / +++
<i>Ch. schraderanum</i> L.	–	–
<i>Ch. urbicum</i> L.	–	–

RIA ($\mu\text{g E. eg. g seed}$) – ecdysteroid-specific radioimmunoassay ($\mu\text{g ecdysone equivalents g seed}$); (–) below detection limit (4.3 μg 20-hydroecdysone equivalents/g; (+) 4.3–43 μg 20-hydroecdysone equivalents/g; (++) 43–430 μg 20-hydroecdysone equivalents/g; (+++) 430–4300 μg 20-hydroecdysone equivalents/g; (++++) more than 4300 μg 20-hydroecdysone equivalents/g

Locomotion compensator

Locomotion compensator. The locomotion and direction of each larva were recorded by placing it onto a Syntech TrackSphere LC-100 locomotion compensator (Syntech, Löptin, Germany). The locomotion of each larva was observed during 5 min, using 3 plants of 12 species on which 10 larvae and 10 neonates for the control were tested.

Odours were obtained by pushing 40 ml/min of humidified and charcoal-filtered air through a bag of Nalophan[®] (Scentroid, Whitchurch-Stouffville, Canada) plastic enclosing one *Chenopodiaceae* plant, using a Stimulus Air Controller CS-55 (Syntech) which mixed it with another stream of pure air so that the total airflow directed onto the insect was 80 ml/minute. The stimulus was pulsed at 0.1 Hz (5/5 s – 5 s of continuous pure and humidified air and 5 s of pulsed air with tested plant).

Volatile collection system. Volatiles were collected from *Chenopodiaceae* plants enclosed into a Nalophan[®] bag. A volatile collector trap (6.35-mm OD, 76-mm long glass tube; Analytical Research Systems, Micanopy,

USA) containing 30 mg of Super-Q adsorbent (Alltech Associates, Deerfield, USA) was inserted into each of 4 Tygon tubes (connection between airflow meter and collector trap). Purified, humidified air was delivered at a rate of 1.0 l/min over the plants, and a vacuum pump sucked 20% less (0.8 l/min) to avoid collecting odours from any gap of the system. Volatiles were collected from the whole plant. The volatile collection sequence (two-hour collections) was initiated after 6 weeks of growing. Six plants were taken for collection experiments plus two blanks (odours collected from empty Nalophan bags only). The results are presented in ng per 1 minute.

Chemical analyses. Volatiles were eluted from the Super-Q adsorbent with 225 µl of hexane and after this 7 ng of decane (both 95% of purity, Sigma-Aldrich, Poznań, Poland) were added as an internal standard. Volatiles were analysed by coupled gas chromatography-mass spectrometry (GC-MS). The GC-MS was a Perkin Elmer AutoSystem XL (Poland) instrument fitted with a 30-m DB-5MS capillary column (0.25-mm ID, 0.25 µm film thickness). The




Figure 1. Orientation of neonate ECB larvae to a source of odour delivery

doi: 10.17221/143/2017-PPS




Figure 2. Mean values for (Z)-3-HAL, (Z)-3-HAC, LIN, and β-CAR

identification of volatiles was verified with authentic standards purchased from commercial sources that had the same GC retention times and mass spectra.

Track analyses. The following parameters were used to quantify the tracks: walking speed (mm/s), straightness of walking (max. 1.00, straight line), angles (x , y), and upward length (mm). For each experimental condition the mean angles by Rayleigh Z tests using the procedure described by BATSCHLET (1981) (U tests) were tested. Calculations and statistical tests on circular data were done using the Oriana software (Kovach Computing Services, Anglesey, UK).

Statistical analysis. All the analyses were conducted using the GenStat v17 statistical software package. Multivariate analysis of variance (MANOVA) was performed on the basis of a MANOVA procedure in GenStat v17 (RENCHER 1992). Mahalanobis distance was suggested as a measure of “poly-VOCs” species similarity (SEIDLER-ŁOŻYKOWSKA & BOCIANOWSKI 2012), whose significance was verified by means of critical value D_α called “the least significant distance” (MAHALANOBIS 1936).

RESULTS

The servosphere was used to observe how ECB first instar larvae orient towards or downwind of the stimulus source, depending on the tested plants and the pulsing regime. None of the tested plants of the genus *Chenopodium* attracted the larvae. Attraction/repulsion was observed in response to odours from *Chenopodium album*, *C. ambrosioides*, *C. glaucum*, *C. polyspermum*, *C. schraderanum*, and *C. urbicum* (Figure 1). A clear repellency was recorded in response to odours from *C. amaranticolor*, *C. botrys*, *C. ficifolium*, *C. murale*, *C. vulvaria*, and *C. oleracea* (Figure 1) – Track Sphere Locomotion Compensator software.

Results of MANOVA (Wilk's $\lambda = 0.0001798$; $F_{10,55} = 43.39$; $P < 0.0001$) indicate that 4 volatile organic compounds (VOCs) were released in large amounts. Results of the analysis of variance for all VOCs confirm the variability of tested species at the significance level $\alpha = 0.01$ (except for LIN, $P = 0.489$).

Mean values for observed VOCs for both factors are presented in Figure 2 [repellent on the space (Yes) or

Tab. 2. Mahalanobis distance between species

Species	No	1	2	3	4	5	6	7	8	9	10	11
<i>Spinacia oleracea</i> L.	1	0	7.81	12.85	3.18	1.54	6.14	8.79	9.79	8.82	8.50	8.44
<i>Ch. album</i> L.	2		0	14.72	8.69	8.49	9.13	1.54	13.62	1.30	1.01	0.93
<i>Ch. amaranticolor</i>	3			0	11.12	11.83	11.73	15.55	13.89	15.46	15.28	15.23
<i>Ch. ficifolium</i> Sm.	4				0	1.80	6.42	9.76	10.80	9.75	9.41	9.34
<i>Ch. murale</i> L.	5					0	6.09	9.52	9.96	9.54	9.21	9.15
<i>Ch. vulvaria</i> L.	6						0	10.59	5.52	10.39	10.12	10.04
<i>Ch. ambrosioides</i> L.	7							0	14.92	0.41	0.56	0.65
<i>Ch. botrys</i> L.	8								0	14.76	14.53	14.47
<i>Ch. glaucum</i> L.	9									0	0.35	0.43
<i>Ch. polyspermum</i> L.	10										0	0.09
<i>Ch. schraderanum</i> L.	11											0

 $D_{0.05} = 7.73$




Figure 3. Distribution of ten *Chenopodiaceae* species and *Spinacia oleracea* L. in the space of the first two canonical variables

not active on the space (No) (left figure) and plants with ecdysteroids by Dinan (Green) or ecdysteroid-free plants by Dinan (Black) (right figure)]. The high mean values were observed in case of the repellent presence on the space for all four VOCs. Plants with ecdysteroids characterised higher mean values of all four VOCs than ecdysteroid-free plants (Figure 2).

The greatest variability in terms of all the analysed VOCs expressed jointly with the greatest Mahalanobis distance was recorded for *Ch. amaranticolor* and *Ch. ambrosioides* L. (the Mahalanobis distance was equal to 15.55). In turn, the greatest similarity was observed for *Ch. polyspermum* L. and *Ch. schraderanum* L. (the Mahalanobis distance for this pair was 0.09) (Table 2).

Multivariate analysis made it possible to compare tested species in terms of all four VOCs (Figure 3). The total variation explained by the first canonical variable was 62.03%, while for the second variable it was 23.55%.

DISCUSSION

The orientation of neonate ECB larvae to undamaged *Chenopodiaceae* plants which showed that these larvae clearly detect odours from such plants was recorded. WITZGALL *et al.* (2012), BENGTTSSON *et al.* (2014), and GONZALEZ *et al.* (2015) claim that many plants constitutively release VOCs, but quantities are often affected by stress, where volatiles have

various functions in plant development including defence against pathogens.

While our previous work showed that larvae of the same age were unconditionally attracted to maize odours (their host plants) or to green leaf volatiles (PIESIK *et al.* 2009, 2013), here, it was observed that ECB larvae were strongly repelled by odours from ecdysteroid-positive plants (Figure 2) and to a lesser extent by ecdysteroid-negative plants (Figure 1). *C. album* and *C. polyspermum* stand out in the group of plant odours being attractive/repellent. According to DINAN *et al.* (1998), these two species show important variations between samples: in *C. album*, the amounts varied between no detection to 1 to 3 levels, while for *C. polyspermum*, two of six samples were below the detection limit. In the group of repellent plants, all of them are ecdysteroid-positive except *C. botrys*.

These chemicals may be generally occurring compounds. In this work (*Z*)-3-hexenal, (*Z*)-3-hexen-1-yl acetate as well as linalool and β -caryophyllene were released in greater amounts in ecdysteroid-positive plants than in ecdysteroid-negative ones. These chemicals have been reported as repellent in different plants. KOSCHIER *et al.* (2002) found linalool and eugenol (main compounds of *O. majorana* and *O. gratissimum*) inhibiting feeding of onion thrips (*Thrips tabaci*). ZHENG *et al.* (2005) found (*Z*)-3-hexenal as a key component contributing to the aroma of omija (*Schizandra chinensis*) leaves. Moreover, (*Z*)-3-hexenal has been identified as the most abundant volatile compound in tomato (BUTTERY *et al.* 1987) and in orange juice

doi: 10.17221/143/2017-PPS

(BUETTNER & SCHIEBERLE 2001). FUHRMANN and GROSCH (2002) found (*Z*)-3-hexenal as one of the main odours of the apple cultivar Elstar. FUKUSHIMA *et al.* (2002) reported that (*E*)-2-hexenal, (*Z*)-3-hexen-1-ol, (*Z*)-3-hexen-1-yl acetate, β -myrcene, and linalool could be released not only from infested or artificially damaged plants, but also from undamaged ones. PEACOCK *et al.* (2001) found (*Z*)-3-hexen-1-yl acetate and (*Z*)-3-hexenol emitted from undamaged *S. dasyclados* (Wimm) leaves.

References

- Batschelet E. (1981): Circular Statistics in Biology. London, Academic Press.
- Bengtsson J.M., Gonzalez F., Cattaneo A.M., Montagné N., Walker W.B., Bengtsson M., Anfora G., Ignell R., Jacquini-Joly E., Witzgall P. (2014): A predicted sex pheromone receptor of codling moth *Cydia pomonella* detects the plant volatile pear ester. *Frontiers in Ecology and Evolution*, 2: 1–11.
- Blackford M., Clarke B., Dinan L. (1996): Tolerance of the Egyptian cotton leafworm *Spodoptera littoralis* (Lepidoptera: Noctuidae) to ingested phytoecdysteroids. *Journal of Insect Physiology*, 42: 931–936.
- Blackford M.J.P., Dinan L. (1997a): The effects of ingested ecdysteroid agonists (20-hydroxyecdysone, RH5849 and RH5992) and an ecdysteroid antagonist (cucurbitacin B) on larval development of two polyphagous lepidopterans (*Acherontia atropos* and *Lacanobia oleracea*). *Entomologia Experimentalis et Applicata*, 83: 263–276.
- Blackford M.J.P., Dinan L. (1997b): The effects of ingested 20-hydroxyecdysone on the larvae of *Aglais urticae*, *Inachis io*, *Cynthia cardui* (Lepidoptera: Nymphalidae) and *Tyria jacobaeae* (Lepidoptera: Arctiidae). *Journal of Insect Physiology*, 43: 315–327.
- Buettner A., Schieberle P. (2001): Evaluation of aroma differences between hand-squeezed juices from Valencia late and Navel oranges by quantitation of key odorants and flavor reconstitution experiments. *Journal of Agricultural and Food Chemistry*, 49: 2387–394.
- Buttery R.G., Teranishi R., Ling L.C. (1987): Fresh tomato aroma volatiles: a quantitative study. *Journal of Agricultural and Food Chemistry*, 35: 540–544.
- Calas D., Berthier A., Marion-Poll F. (2007): Do european corn borer females detect and avoid laying eggs in the presence of 20-hydroxyecdysone? *Journal of Chemical Ecology*, 33: 1393–1404.
- Coll J., Tandrón Y.A., Zeng X. (2007): New phytoecdysteroids from cultured plants of *Ajuga nipponensis* Makino. *Steroids*, 72: 270–277.
- Dinan L. (1992): The association of phytoecdysteroids with flowering in fat hen, *Chenopodium album*, and other members of the Chenopodiaceae. *Experientia*, 48: 305–308.
- Dinan L. (1995): A strategy for the identification of ecdysteroid receptor agonist and antagonists from plants. *European Journal of Entomology*, 92: 271–283.
- Dinan L. (1998): A strategy towards the elucidation of the contribution made by phytoecdysteroids to the deterrence of invertebrate predators on plants. *The Russian Journal of Plant Physiology*, 45: 296–305.
- Dinan L., Savchenko T., Whiting P. (2001): On the distribution of phytoecdysteroids in plants. *Cellular and Molecular Life Sciences*, 58: 1121–1132.
- Dinan L., Savchenko T., Whiting P. (2002): Chemotaxonomic significance of ecdysteroid agonists and antagonists in the Ranunculaceae: phytoecdysteroids in the genera *Helleborus* and *Hepatica*. *Biochemical Systematics and Ecology*, 30: 171–182.
- Fuhrmann E., Grosch W. (2002): Character impact odorants of the apple cultivars Elstar and Cox Orange. *Nahrung/Food*. 46: 187–193.
- Fukushima J., Kainoh Y., Honda H., Takabayashi J. (2002): Learning of herbivore-induced and nonspecific plant volatiles by a parasitoid, *Cotesia kariyai*. *Journal of Chemical Ecology*, 28: 579–586.
- Gelman D.B., Pszczolkowski M.A., Blackburn M.B., Ramaswamy S.B. (2007): Ecdysteroids and juvenile hormones of whiteflies, important insect vectors for plant viruses. *Journal of Insect Physiology*, 53: 274–284.
- Gonzalez F., Bengtsson J.M., Walker W.B., Rodrigues Sousa M.F., Cattaneo A.M., Montagné N., de Fouchier A., Anfora G., Jacquini-Joly E., Witzgall P., Ignell R., Bengtsson M. (2015): A conserved odorant receptor detects the same 1-indanone analogs in a tortricid and a noctuid moth. *Frontiers in Ecology and Evolution*, 3: 1–12.
- Koschier E.H., Sedy K.A., Novak J. (2002): Influence of plant volatiles on feeding damage caused by the onion thrips *Thrips tabaci*. *Crop Protection*, 21: 419–425.
- Lafont R., Harmatha J., Marion-Poll F., Dinan L., Wilson I.D. (2002): Ecdybase (The Ecdysone Handbook. 3rd Ed., on-line). Available at <http://ecdybase.org> (accessed 2002).
- Mahalanobis P.C. (1936): On the generalized distance in statistics. *Proceedings on the National Institute of Science of India*, 12: 49–55.
- Marion-Poll F., Descoins C. (2002): Taste detection of phytoecdysteroids in larvae of *Bombyx mori*, *Spodoptera littoralis* and *Ostrinia nubilalis*. *Journal of Insect Physiology*, 48: 467–476.
- Peacock L., Lewis M., Herrick S. (2001): Factors influencing the aggregative response of the blue willow beetle,

- Phratora vulgatissima*. Entomologia Experimentalis et Applicata, 98: 195–201.
- Piesik D., Rochat D., van der Pers J., Marion-Poll F. (2009): Pulsed odors from maize or spinach elicit orientation in european corn borer neonate larvae. Journal of Chemical Ecology, 35: 1032–1042.
- Piesik D., Rochat D., Delaney K.J., Marion-Poll F. (2013): Orientation of european corn borer first instar larvae to synthetic green leaf volatiles. Journal of Applied Entomology, 137: 234–240.
- Rencher A.C. (1992): Interpretation of canonical discriminant functions, canonical variates, and principal components. American Statistical Association, 46: 217–225.
- Schmelz E.A., Grebenok R.J., Galbraith D.W., Bowers W.S. (1999): Insect-induced synthesis of phytoecdysteroids in spinach, *Spinacia oleracea*. Journal of Chemical Ecology, 25: 1739–1757.
- Schmelz E.A., Grebenok R.J., Ohnmeiss T.E., Bowers W.S. (2002): Interactions between *Spinacia oleracea* and *Bradysia impatiens*: a role for phytoecdysteroids. Archives of Insect Biochemistry and Physiology, 51: 204–221.
- Seidler-Łożykowska K., Bocianowski J. (2012): Evaluation of variability of morphological traits of selected caraway (*Carum carvi* L.) genotypes. Industrial Crops and Products, 35: 140–145.
- Simon A., Pongrácz Z., Tóth G., Mák M., Máthé I., Báthori M. (2004): A new ecdysteroid with unique 9 β -OH and four other ecdysteroids from *Silene italica* ssp. *nemoralis*. Steroids, 69: 389–394.
- Soriano I.R., Riley I.T., Potter M.J., Bowers W.S. (2004): Phytoecdysteroids: a novel defense against plant-parasitic nematodes. Journal of Chemical Ecology, 30: 1885–1899.
- Tarkowska D., Strnad M. (2016): Plant ecdysteroids: plant sterols with intriguing distributions, biological effects and relations to plant hormones. Planta, 244: 545–555.
- Voigt B., Whiting P., Dinan L. (2001): The ecdysteroid agonist/antagonist and brassinosteroid-like activities of synthetic brassinosteroid/ecdyteroid hybrid molecules. Cellular and Molecular Life Sciences, 58: 1133–1140.
- Volodin V., Chadin I., Whiting P., Dinan L. (2002): Screening plants of European North-East Russia for ecdysteroids. Biochemical Systematics and Ecology, 30: 525–578.
- Witzgall P., Proffit M., Rozpedowska E., Becher P.G., Andreadis S., Coracini M., Lindblom T.U.T., Ream L.J., Haggman A., Bengtsson M., Kurtzman C.P., Piskur J., Knight A. (2012): “This is not an apple”-yeast mutualism in codling moth. Journal of Chemical Ecology, 38: 949–957.
- Zheng C.-H., Kim K.-H., Kim T.-H., Lee H.-J. (2005): Analysis and characterization of aroma-active compounds of *Schizandra chinensis* (omija) leaves. The Journal of the Science of Food and Agriculture, 85: 161–166.

Received: 2017–11–02

Accepted after corrections: 2017–01–29

Published online: 2018–02–04