



HAL
open science

Evaluation of the impact of a Herd Health and Production Management programme in organic dairy cattle farms: a process evaluation approach

Julie E. Duval, Nathalie Bareille, Aurélien Madouasse, Manon de Joybert, Karin Sjöström, Ulf Emanuelson, Florence Bonnet-Beaugrand, Christine Fourichon

► To cite this version:

Julie E. Duval, Nathalie Bareille, Aurélien Madouasse, Manon de Joybert, Karin Sjöström, et al.. Evaluation of the impact of a Herd Health and Production Management programme in organic dairy cattle farms: a process evaluation approach. *Animal*, 2018, 12 (7), pp.1475-1483. 10.1017/S1751731117002841 . hal-02621523

HAL Id: hal-02621523

<https://hal.inrae.fr/hal-02621523v1>

Submitted on 2 Apr 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Evaluation of the impact of a Herd Health and Production Management programme in organic dairy cattle farms: a process evaluation approach

J. E. Duval^{1†}, N. Bareille¹, A. Madouasse¹, M. de Joybert¹, K. Sjöström², U. Emanuelson², F. Bonnet-Beaugrand¹ and C. Fourichon¹

¹BIOEPAR, INRA, Oniris, 44307 Nantes, France; ²Department of Clinical Sciences, Division of Ruminant Medicine and Veterinary Epidemiology, Swedish University of Agricultural Sciences, PO Box 7054, SE-750 07 Uppsala, Sweden

(Received 28 February 2017; Accepted 11 September 2017; First published online 6 November 2017)

Animal health planning activities are not always providing a satisfactory positive impact on herd health and welfare. Moreover, evaluating the impact of advisory programmes is complex due to multiple interacting elements that influence its outcome. Therefore, measuring solely health outcomes is not sufficient: the whole process of the implementation and use of such programmes should be evaluated. In order to evaluate the impact of an intervention with a Herd Health and Production Management (HHPM) programme a process evaluation framework was designed and used. The intervention involved 20 organic dairy cattle farmers and their advisors, in both France and Sweden. In both countries 20 organic dairy farms were selected as control herds. The evaluation of the HHPM programme was based on: (a) the compliance to the programme; (b) the programme's functions influencing herd health management practices and stimulating dialogue between farmers and advisors; (c) its effectiveness in terms of improving herd health compared with control farms. Complete compliance to the programme was fulfilled by 21 out of 40 farmers–advisors. Results from a questionnaire showed that the programme functioned as intended (e.g. by allowing early identification of herd health problems), stimulated change in farmers' herd health management practices and farmer–advisor dialogue. Even though the majority of the users perceived that the programme contributed to herd health improvements, no significant differences in health outcomes were found when compared with control farms 12 months after the start of the intervention. The programme allowed creating an environment promoting the exchange of information between farmers and advisors, necessary to define pertinent advice in a farm-specific situation. Future research should aim at improving methods for the evaluation of the effect of advisory programmes, by identifying early indicators for effective advice and developing methods to evaluate the quality of advisory situations without interfering with them.

Keywords: dairy cattle, animal health planning, advisors, extension services, complex interventions

Implications

The results of this study show that when evaluating dairy herd health advisory programmes, looking beyond compliance levels and impact on herd health is valuable. It allows identifying the impact on factors that in theory lead to improved herd health advisory services. Furthermore, it provides a more detailed understanding of how Herd Health and Production Management (HHPM) programmes can be used in the field, as the amount of published information available on this subject is scarce. This study provides detailed information for advisors in herd health management and policy makers who would like to develop these programmes.

Introduction

The organic production principles and regulations aim for high levels of animal health and welfare on organic farms (International Federation of Organic Agriculture Movements, 2005; Anonymous, 2007). However, the large variation in production disease levels found on organic dairy cattle farms within and across European countries suggests that there is room for improvement (Krieger *et al.*, 2017). Since the 1980s, HHPM programmes were developed to support veterinarians in their role of advisors in disease prevention and health promotion. They require a holistic approach of the farm to be able to optimize animal health and productivity, whilst ensuring animal welfare, food safety, sustainable production and profitability. A HHPM programme is an iterative process

[†] E-mail: julie.duval@oniris-nantes.fr

involving: (1) setting objectives, (2) herd health monitoring activities, (3) implementation of practices and (4) evaluation of the outcome (Brand *et al.*, 2001). Improving the use of farm-specific animal health planning processes has been identified as a promising way to improve animal health and welfare on organic dairy farms (Vaarst *et al.*, 2011). In theory, HHPM programmes allow to design farm-specific and farmer-centred health programmes and promote the farmer–advisor dialogue. Duval *et al.* (2016b) described a participatory approach that allowed designing farm-specific monitoring tools which farmers intended to use in a HHPM programme. However, so far no information was available on how these indicators were used in a HHPM programme setting. In general, relatively little published information describes HHPM programmes’ actual use in field conditions.

Implementing a HHPM programme on farms can be regarded as a complex intervention. As defined by Craig *et al.* (2008), complex interventions contain several interacting components, involve different organizational levels, demand a number and certain complex behaviours by the persons receiving or delivering it, allow a certain adaptability of the intervention and finally can have a variation of results. HHPM programmes require a certain amount of flexibility to be adapted to farm-specific situations, and are influenced by decision-making processes from both farmer and adviser. Many factors influence the perceived pertinence of advice and ultimately farmers’ decision-making processes whether or not to adopt advice. Preventive health behaviour will, for example, be influenced by the perceived threat of disease and benefits of taking action (Janz and Becker, 1984) or the perceived ability to implement practices (Garforth, 2011). Outcomes of HHPM programmes are depending on the heterogeneity of farms and advisors. Complex interventions may be challenging to evaluate and outcomes of evaluation studies can be difficult to interpret and reproduce. It is thus important to evaluate not only the outcomes, but the

complete intervention process (Moore *et al.*, 2015). Process evaluation approaches aim at: understanding how the intervention under study was implemented, identifying causal mechanisms impacting the outcome and contextual factors explaining variation in implementation and results (Oakley *et al.*, 2006). Although process evaluation approaches are already used for the evaluation of public health interventions (Craig *et al.*, 2008), it has not been used previously in the context of intervention aiming at promoting dairy herd health and production. Therefore, the objectives of this study were to evaluate the use and impact of a HHPM programme targeting production diseases in dairy cattle in two different advisory contexts using a process evaluation framework.

Material and methods

General approach and study design

An intervention study was performed on 20 certified organic dairy cattle farms in France and on 20 farms in Sweden, with the objective to improve herd health by implementing a HHPM programme. In addition, in both countries 20 organic dairy farms were selected as control herds where no actions were taken. The countries were chosen to represent different contexts in terms of existing organic dairy farming systems and herd health advisory contexts. In Sweden, a mandatory and more standardized character of herd health monitoring and advisory services exists for organic dairy farms, which is non-existent in France (described in detail in Duval *et al.*, 2016b).

The impact of the HHPM programme was assessed using an adapted form of the process evaluation framework as described by Moore *et al.* (2015) (Figure 1). The framework allows evaluation of elements that we hypothesized could influence the impact of the programme on herd health, namely participants’ compliance to the HHPM programme, the programme’ ability to fulfil its intended uses, its ability to influence farmers’ animal health monitoring and disease




Figure 1 Process evaluation framework. Different elements can explain the impact of a Herd Health and Production Management (HHPM) programme on herd health advisory services and herd health. The elements that were evaluated during this study are surrounded by thick grey boxes and the origin of the data is described in italics.

prevention practices and to stimulate the interaction between farmers and advisors. The effectiveness of the programme in terms of improving herd health in intervention farms was compared with control farms and herd health situations before the intervention.

The Herd Health and Production Management programme

The main health disorders that are considered to have a negative influence on cows' health, welfare and production were identified as targets of the HHPM programme. Five health topics were selected: reproductive health and performance, udder health, calf health, locomotor disorders and metabolic disorders.

During an initial meeting (meeting 0) a researcher explained the HHPM programme to the participants. The nature of the tested HHPM programme conserved the general iterative concept of a HHPM programme; meeting 0 was used to set objectives and during the other meetings farmers and advisors were expected to (1) monitor herd health, (2) identify and implement disease preventive practices, if deemed necessary and (3) evaluate them (Supplementary Material S1). However, adaptations were made to the general concept of a HHPM programme by the research team. First, the indicators to be used for herd health monitoring were not imposed and standardized across farms. As described by Duval *et al.* (2016b), during meeting 0 a participatory approach was followed allowing farmers and their advisors to adapt the monitoring tool by choosing farm-specific indicators and alert thresholds to identify herd health problems targeting the five health topics. This resulted in the design of 40 farm-specific monitoring tools (paper-based or Excel[®]-sheets). Second, the tools provided for disease prevention by the research team contained lists of objectives to attain, rather than a detailed list of standard recommendations of good management practices that need to be implemented. The implementation of the HHPM programme was expected to lead to an improvement of herd health through: (1) herd health monitoring, (2) early identification of problems, (3) advisors providing farmers with adapted advice and (4) farmers' compliance with the advice.

Farmers and their advisors were asked to implement the HHPM programme and have at least three meetings per farm in Sweden and four in France over a period of 12 months. Budget limitations prevented to do the same number of meetings in both countries. These meetings were conducted without the researcher, with the objective to mimic as much as possible a real advisory situation.

Participant selection

Organic dairy farmers involved in an ongoing research project were invited to participate in the current study. Inclusion criteria were that farms had to have been certified as organic for at least 1 year before the study, participate in the official milk recording scheme and farmers had to give their consent to participate in the study and share their farm data. Farmers were promised that the data would be treated anonymously. Farmers chose any person that they deemed appropriate as

their advisor in herd health (veterinarian or other) to implement together the HHPM programme. In both countries, most often veterinarians were chosen.

Advisors could accompany several farmers, as was the case for one veterinarian in France and for five veterinarians in Sweden. Advisors were paid 1000 euros/farmer from the project budget to accompany farmers in implementing the HHPM programme and to report to the research teams on the farm meetings.

In total, 20 intervention farms were selected in western (the departments Loire-Atlantique and Morbihan) and eastern France (Lorraine region). In Sweden, 20 farms were selected from an area covering almost half of Sweden and with a relatively high density of dairy farms. The 20 French control farms were certified organic dairy farms located in the same geographic areas, with feeding practices, herd size and milk production level comparable with the intervention farms. In total, 20 Swedish control farms were randomly selected out of the certified organic farms present in Sweden. More details of the selection procedure can be found in Duval *et al.* (2016b).

Herd characteristics, advisor characteristics and a description of existing monitoring activities before the start of the study on the selected intervention farms were described in Duval *et al.* (2016b). Herd size was on average greater in Sweden than in France. Before the start of the study, herd health monitoring activities were non-existing in 15 out of 20 French farms. Out of 20 Swedish farmers 13 reported to monitor several herd health topics using a different approach than the one proposed in this study. Whether this was done with an advisor and/or in a herd health programme setting was not recorded.

The process evaluation framework and data collection

Evaluation of the compliance to the Herd Health and Production Management programme. The advisors reported after each meeting to the research teams using standardized templates for reporting (Supplementary Material S2). Participants' compliance to the HHPM programme was evaluated based on the number of meetings and the implementation of monitoring activities as planned, the use of the disease prevention tools after a herd health alert, the proposition of recommendations to improve a deteriorated health situation, the presence of a discussion on recommendations made during the previous meetings and implementation of recommended measures of previous meetings if there were any.

Evaluation of the users' opinion on the Herd Health and Production Management programme and its functions. The opinion of the participants of the HHPM programme was collected at the end of the intervention period. Each farmer and advisor that had performed at least one meeting after meeting 0 was asked by email to fill in a questionnaire using a web (Netigate[®]) or a paper form. Only one farmer used the paper format. Questions were related to the different types of possible use that could be made of the monitoring and the prevention tool, possible limits in its use, the value of

having regular farm meetings, the influence of the HHPM programme on the relationship between farmer and advisor, the perceived effectiveness on herd health, future use and cost of the HHPM programme (the questionnaire is available upon request from the corresponding author). Mostly, questions with closed answers were used. Answers to questions, where a Likert scale (1 to 6) was used, were transformed into agree or disagree answers for the analysis. The scores 1 to 3 were converted into disagree and scores of 4 to 6 were converted into agree. A maximum of three reminders to fill in the questionnaire were sent. Reasons for not responding to the questionnaire were not recorded.

Evaluation of the Herd Health and Production Management programme' impact on herd health. To assess the effectiveness of the HHPM programme in terms of improvement in herd health, changes in health situations before and after the intervention were evaluated and compared with the herd health situation in control farms from the corresponding countries. Subsequently, differences between French and Swedish farms were assessed. Farm data from the national recording systems were retrieved to calculate herd health and production indicators in a harmonized way, using the methodology described by Krieger *et al.* (2017). Data were obtained from the official milk recording schemes, artificial insemination databases and the animal identification and registration databases. The national recording systems are not harmonized and record keeping is different across countries, including the amount of information that is recorded. Therefore, the choice of herd health indicators was determined by data availability in both countries. For example, in contrast to Sweden, in France there is no information available on lameness. As a consequence, this health disorder was excluded from the evaluation even though lameness was one of the five health domains targeted by the HHPM programme.

Nine indicators were retained to calculate herd health and production performances in both countries. For milk production, the average daily milk yield produced per cow per herd was used. Udder health was assessed by using the prevalence of high somatic cell count (SCC), calculated as the proportion of records with a SCC-value $>200\,000$ cells/ml, and the apparent new udder infection risk defined as the proportion of cows with a SCC-value $<200\,000$ cells/ml that changed to $>200\,000$ cells/ml between consecutive test days. Reproductive health and performance indicators were calving interval, defined as the median interval between the last and the previous calving date, and calving to first artificial insemination interval, defined as the median duration of the interval between the last calving and the first artificial insemination after calving. For metabolic disorders, the prevalence of fat/protein ratios >1.5 between 30 and 100 days in milk was used as an indicator of an increased risk of ketosis; the prevalence of fat/protein ratios <1.0 was used as an indicator of an increased risk for sub-acute ruminal acidosis. On-farm mortality of cows, defined as the number of cows, that is after first calving, that died or

were euthanized on farm divided by the sum of days at risk of dying, was also used. Finally, calf mortality was measured by the number of calves that died between 1st day and 30 days of life, divided by the sum of days at risk of dying. Regarding both mortality indicators, sold animals were censored at the day of leaving the herd. The herd health and production performance indicators were calculated for two distinct periods. Period 1 is the reference period, from 13 October 2012 to 13 October 2014. Period 2 covers the implementation of the HHPM programme from the median date of meeting 1 (10 December 2014) to 10 December 2015. Between periods 1 and 2, meeting 0 were performed to introduce the HHPM programme on the farms.

Data were randomly missing for certain farms for the calculation of certain indicators in both countries, even though data availability was a selection criterion. In France, for 15 control herds and five herds in the intervention group, milk recording data were missing. In Sweden this data were missing for 13 control herds. For the calculation of the calving interval data from one intervention farm was missing in France, in Sweden this was the case for 17 control farms and one intervention farm. To calculate the indicator interval calving-first artificial insemination, data from one control and one intervention farm was missing in France, in Sweden data were missing for three control farms. To calculate calf mortality, the data of four and two control farms was missing in France and Sweden, respectively.

Statistical analysis

The impact of the HHPM programme on herd health was assessed using linear models (R, r-project.org). Models were specified as follows: $Y_i = \text{intercept} + \beta_1 \times \text{country}_i + \beta_2 \times \text{period}_i + \beta_3 \times \text{intervention}_i + \beta_4 \times \text{intervention}_i \times \text{period}_i + \varepsilon_i$ where Y_i was the health outcome of interest and country was the country in which the herd was located (0 = France or 1 = Sweden). The period was the period in which the health outcome was measured (0 = before intervention or meeting 0; 1 = after start HHPM or meeting 0), intervention was a variable indicating whether the herd was an intervention farm (0 = no and 1 = yes) and the interaction between period and intervention. β_1 to β_4 were coefficients measuring the strength of the associations between these explanatory variables and the outcomes. In these models, the value of β_4 was of interest. The normal distribution of the residuals was visually checked. Significance indicated a significant association between intervention and the outcome of interest. Fisher's tests were performed to compare the results on compliance, health impact and users' opinion.

Results

Compliance to the Herd Health and Production Management programme

Meeting 0 was implemented in all intervention farms. However, in both countries on two farms no further meetings were implemented. The reason stated by the involved advisors was their lack of time to implement the HHPM

programme. Out of the 40 farms 21 performed the number of meetings as planned (Table 1). There was no significant difference in the number of farmers who completed the full number of planned meetings in France (13/20, ≥ 4 meetings), compared with Sweden (8/20, ≥ 3 meetings) (P value = 0.30).

Compliance to monitoring activities on the five health topics was high (Table 1). Across countries, udder health was the domain with the highest number of health alerts, followed by reproduction and calf health (Figure 2). Herd health alerts most often lead to the use of a preventive tool (Table 1). No significant differences were found between France and Sweden. On average during the meetings, 100% of the alerts led to recommendations in Sweden compared to 85% in France (P value = 0.48) (data not shown). Prevention tools could also be used without a herd alert, and this was

done on 21 farms (data not shown). Identified corrective actions to improve animal health situations were 100% implemented, according to 35% and 27% of the farmers in France and Sweden, respectively (data not shown). This difference was not significant.

Participants' opinion on the Herd Health and Production Management programme and its functions

Out of the 18 eligible French farmers 17 answered the questionnaire, in Sweden 11 out of 18 farmers did. All French advisors answered the questionnaire and in Sweden 8 out of 13 advisors did. General appreciation of the tool was evaluated by asking both farmers and advisors whether they were of the opinion that the implementation of the tool during the study had a positive impact on herd health (Table 2). Numerically, farmers were more positive than advisors but this difference was not significant. Not all the farmers were willing to pay advisors for these kinds of services. Swedish farmers were more often willing to pay than French farmers (P value = 0.04). Advisors were asked whether their remuneration to implement the HHPM programme corresponded to what they would ask of farmers for this kind of services; 17.6% and 12.5% of the French and Swedish advisors, respectively, replied that they would ask more. In France and Sweden, respectively, 47.1% and 25% would ask less. Although not all French participants expected an improvement of the herd health situation, the percentage of participants answering that they would keep using certain tools was higher than the percentage of participants estimating a positive effect on herd health.

Table 1 Compliance to the Herd Health and Production Management programme in terms of number of meetings performed after meeting 0, implementation of monitoring activities and the use of disease prevention tools

	France	Sweden	Total
Number of implemented meetings per farm			
1	1	3	4
2	2	4	6
3	2	8	10
4	12	0	12
5	1	0	1
No data	0	3	3
Percentage of the meetings during which 5 health topics were monitored			
Meeting 1	72	87	80
Meeting 2	94	91	93
Meeting 3	80	85	83
Meeting 4	100	–	–
Percentage of the meetings with herd health alerts that lead to the use of disease prevention tools			
Meeting 1	80	92	86
Meeting 2	86	78	82
Meeting 3	80	73	77
Meeting 4	64	–	–




Figure 2 Number of alerts per health domain per meeting during the implementation of the Herd Health Management and Production programme in France and Sweden. n = total number of farms for which the data were available.

Participants' opinion of the monitoring and prevention tools' functions

Participants were asked whether the monitoring tool allowed doing what it was intended for. The main country difference was found regarding the statement that the implementation of the tool was a way to have regular contact between farmer and advisor. Numerically this seemed more important in France than in Sweden (P value = 0.08) (Table 3). Allowing each farmer to choose the indicators considered appropriate for herd health monitoring in his/her farm, was done with the intention to improve the shared understanding by farmer and advisor on the herd health situation of the farm, farmers' focus areas regarding herd health and, the way the farmer monitors health. Differences between the two countries were observed, as well as within country as between farmers and advisors. In numbers, farmers were more positive about the effect of the tool on the shared understanding between farmer and advisor, but these differences were not significant. The only significant difference observed between French and Swedish advisors concerned the effect of the chosen indicators on improving their knowledge on the way farmers monitor health and farmers' focus areas. Participants were allowed to change monitoring indicators during the course of the study. In France six advisors answered 'no I didn't identify the need', eight changed indicators and two replied 'it might

Table 2 Participants' perception on the effect of the Herd Health and Production Management (HHPM) programme on herd health and possible future use of the HHPM programme' tools

	Farmers			Advisors		
	FR (%)	SE(%)	P value	FR (%)	SE (%)	P value
The implementation of the advisory service, as proposed, has contributed to improve herd health	65	90	ns	59	63	ns
I am ready to pay the advisor for this kind of service	47	90	^a	–	–	–
I will keep using the HHPM programme' tools						
Yes, both the monitoring and the prevention tool	59	80	ns	65	63	ns
Yes, but only the monitoring tool	6	0	ns	6	0	ns
Yes, but only the prevention tool	6	0	ns	6	25	ns
I would recommend the monitoring and/or prevention tools to colleagues	65	100	ns	71	88	ns

FR = France; SE = Sweden.

^aP < 0.05.

Table 3 Participants' agreement on the fulfilment of intended uses of the monitoring and prevention tools of the Herd Health and Production Management programme and regular farm meetings

Agreement with the following statements	Farmers			Advisors		
	FR (%)	SE (%)	P value	FR (%)	SE (%)	P value
The herd health monitoring was useful because						
It allows for the early identification of herd health problems	82	69	ns	88	75	ns
It allows to secure herd health	77	46	ns	77	75	ns
It is a way to have regular contact with my advisor/the farmer	77	46	ns	94	63	ns
It gave me a better idea of how I can use data for herd health monitoring/it gave me more access to herd health data of the farm	71	62	ns	59	63	ns
Choosing indicators adapted to the farm						
Changed my perception of the herd health situation of the herd	53	23	ns	–	–	–
Improved the advisor' understanding of the way the farmer monitors herd health	82	50	ns	82	25	^b
Improved the advisor' knowledge on the herd health situation of the farm	82	67	ns	94	63	ns
Improved the advisors/my knowledge on your/ the farmers focus areas regarding herd health	82	67	ns	94	25	^a
Led to a list of indicators that was appropriate for herd health monitoring on the farm	77	75	ns	77	75	ns
In general, when a herd health problem was identified, the prevention tool helped to identify relevant risk factors present on the farm	82	91	ns	59	88	ns
Using the prevention tool allowed to show the link between management practices and animal health outcome	82	80	ns	82	88	ns
In general, it was possible to identify correctives actions on the farm corresponding to risk factors identified with the advisor/farmer	88	91	ns	82	88	ns
The use of the prevention tool stimulated discussion farm management practices						
Yes, we discussed more than we usually did	70	40	ns	47	75	ns
Yes, but in the past we already discussed farm management practices	18	60	ns	35	25	ns
No	12	0	ns	18	0	ns
Having regular farm meetings (for reasons other than emergencies) during the year was an opportunity						
To take more time to discuss the animal health situation on the farm	100	80	ns	100	88	ns
To have more time to discuss the questions the farmers have on animal health to the advisor/veterinarian	100	80	ns	94	88	ns
To discuss animal health management practices	100	80	ns	100	88	ns

FR = France; SE = Sweden.

^aP < 0,001; ^bP < 0.01.

have been useful'. In Sweden 5 out of 10 advisors answering that question had changed indicators or replied that it might have been useful (data not shown).

The prevention part of the tool was designed to serve different purposes: the identification of risk factors of disease, showing the link between practices and health

outcome, identifying corrective measures to improve or secure health and to stimulate discussion between farmer and advisors on management practices. In general, according to the participants, the prevention tool met the functions it was intended to accomplish (Table 3). No significant differences in perception were found between countries or groups of participants.

In general, the meetings were considered, by farmers and advisors, as an opportunity to discuss the animal health situation of the farm and questions of the farmer on animal health and animal health management practices (Table 3). Furthermore, French and Swedish advisors acknowledged that during the HHPM programme topics were discussed which they had not discussed in the setting of their usual collaboration (Figure 3). French advisors reported to have learned most often about farmers' objectives and farmers' farm and animal health management practices. Swedish farmers learned most often about farmers' farm and animal health management practices and of certain herd health problems (Figure 4).




Figure 3 Farmers' and advisors' experiences of areas which were discussed in-depth for the first time during the implementation of the Herd Health and Production Management programme with the advisors and in which advisors learned new information, respectively. FR = France; SE = Sweden; n = total number of persons from which data were available.




Figure 4 Reasons stated by French and Swedish farmers for not implementing the recommendations (multiple answers were possible). FR = France; SE = Sweden; n = total number of farmers from which data were available.

The Herd Health and Production Management programme' impact on herd health

No significant effect of the HHPM programme on herd health in the intervention farms was found (Table 4, additional descriptive herd figures can be found in Supplementary Material S3). The statistical analysis showed no significant difference in the health status changes, before and after the start of the HHPM programme, between the HHPM and the control group or between countries. The average daily milk production was significantly higher in Sweden than in France. The prevalence and the apparent new udder infection risk was lower in Sweden than in France, as well as, calving interval and median calving to first artificial insemination interval. No significant differences were found between countries metabolic disorders' indicators. Finally, calf mortality was significantly higher in France than in Sweden in contrast to the situation for cow mortality, which in turn was lower in France, but not significantly.

Discussion

The value of a process evaluation framework

This study was innovative using a HHPM programme which was even more farm centric compared with previous studies, because farmers could chose to use it as suggested, modify it or replace it. In addition, a single-advisory method was evaluated across countries. Finally, the evaluation of the intervention of the HHPM programme, using a process evaluation framework, was an innovative approach in the field of animal health management. As health management interventions were often evaluated mainly based on compliance levels and/or changes in herd health indicators (e.g. Green *et al.*, 2007; Bell *et al.*, 2009; Ivemeyer *et al.*, 2009). The impact of a health intervention depends on numerous causal mechanism that need to be evaluated (Moore *et al.*, 2015). In this study, an evaluation solely based on compliance and health improvements would have concluded that the HHPM programme failed. However, in general, when meetings were included, the results showed that the HHPM programme was used as intended. Both farmers and advisors agreed that the monitoring and the prevention part of the tool fulfilled most of the functions it was expected to fulfil to ultimately promote health. Moreover, the HHPM programme appeared to be appreciated by most users of the tool, based upon the perceived effectiveness of the intervention on herd health, according to a majority of the respondents of the questionnaire, and their willingness to continue to use (elements) of the tool.

This study illustrates the value of a process evaluation approach in pilot studies, testing complex interventions in small and heterogeneous study populations. Several limits in the study design can explain why no significant herd health improvements were measured in the intervention farms. The initial herd health situations probably differed. If the only objective of the study had been to measure the effectiveness of the intervention on herd health it would have been more

Table 4 Results of linear models for herd health indicators in the study groups and their evolution after implementation of the Herd Health and Production Management programme in France and Sweden

Effects	Evolution of herd health indicators ¹																										
	Milk yield per cow per day (kg)			Prevalence of high somatic cell count (%)			Apparent new udder infection risk (%)			Ln calving interval (days)			Ln median calving to first artificial insemination interval (days)			Prevalence of fat/protein ratios >1.5 (%)			Prevalence of fat/protein ratios <1.0 (%)			Ln on-farm mortality adult cows (%)			Ln calf mortality 1 to 30 days after birth (%)		
	Estimate	P		Estimate	P		Estimate	P		Estimate	P		Estimate	P		Estimate	P		Estimate	P		Estimate	P		Estimate	P	
Intercept	+17.79	a		+32.04	a		+15.59	a	+6.01	a		+4.57	a	+0.12	a		+0.05	a	+1.24	a		+1.49	a				
Country SE	+9.33	a		-4.81	a		-2.26	b	-0.04	a		-0.13	b	-0.01	ns		0.00	ns	+0.18	ns		-0.61	a				
Intervention	+1.30	ns		+0.70	ns		-0.21	ns	-0.01	ns		+0.09	ns	0.00	ns		0.00	ns	-0.02	ns		-0.21	ns				
Study period 2	+0.91	ns		-0.75	ns		-0.79	ns	+0.01	ns		0.00	ns	0.00	ns		0.00	ns	-0.07	ns		-0.41	b				
Intervention × study period 2	-0.04	ns		-1.20	ns		+0.15	ns	-0.01	ns		-0.17	ns	-0.01	ns		-0.01	ns	+0.02	ns		+0.32	ns				

¹For all models the reference modality was the control group in France (first row, model intercept). For the indicators calving interval, median calving to first artificial insemination interval, on-farm mortality of adult cows and calf mortality from 1 to 30 days after birth the natural logarithm (Ln) of the indicator was modelled. For these indicators the coefficient estimates must be exponentiated to get back to the original scale, and the effects are multiplicative. For example, the calving interval in the control group in Sweden (SE) is exp. (6.03) × exp. (-0.03) = 403.4 days. ^aP < 0.001; ^bP < 0.01.

appropriate to test the tool in, for example, herds with severe udder health problems. Other studies reported more improvements in animal health in herds with a poorer health level at the start of the intervention (e.g. Green *et al.*, 2007; Ivemeyer *et al.*, 2009). In addition, because of the possible different health problems in different farms, the sample size was probably too small per health disorder, especially with the high amount of missing data, limiting the statistical power. Moreover, the testing period may have been too short to observe an effect of the programme based on the indicators chosen. In other intervention studies similar difficulties were identified (Bell *et al.*, 2009; Ivemeyer *et al.*, 2012). The methodological challenges encountered support thus the idea to perform a process evaluation; evaluating both the process and herd health impact. Especially, since large-scale and long intervention studies are often not feasible for financial and organizational reasons.

The Herd Health and Production Management programme' impact on the dialogue between farmer and advisor
The HHPM programme proved to have characteristics facilitating the development of advisory services that stimulated the farmer–advisor dialogue and thus possibly their mutual understanding. The transfer of information occurred in both directions from farmer to advisor and the other way around. Even in Sweden, both farmers and advisors learned from each other despite the existing advisory context that could lead to expect that Swedish advisors had already more often in-depth knowledge of the herd health situation and farm management than in France. The quality of the farmer–advisor dialogue is considered as a key to success of animal health planning activities (Vaarst *et al.*, 2011). The project imposed regular and frequent meetings, which could have been a step forward in itself to stimulate an advisory role for advisors in herd health management on organic dairy farms. French veterinarians have been found to be rarely invited to organic dairy farms and find it difficult to make their role in organic dairy farmers animal health management evolve from a therapeutic role towards an advisory role (Duval *et al.*, 2016a). This situation might be true for other countries, as has been reported in Denmark (e.g. Vaarst *et al.*, 2006).

Difficulties encountered when evaluating the advisory relationship between farmer and advisor
Information will only be of significance to the receiver if it is built upon his/her existing knowledge (Klerkx and Jansen, 2010), underlining thus the importance of the farmer–advisor dialogue to exchange information with the aim to build new knowledge (co-constructing advice) rather than only exchanging information (like a recipe). The results show that both farmers and advisors perceived that they have acquired knowledge that was new to them. However, the research strategy chosen does not allow to measure if and how that was used to adapt it to farm-specific advice. Research methods such as qualitative research interviews could have been more appropriate to understand that process (Malterud, 2001).

A need for adaptable herd health management tools?

Participants used the possibility to adapt the programme to farm-specific situation (Duval *et al.*, 2016b). It has been questioned whether adaptability would only lead to 'quick and easy solutions' (Beekhuis-Gibbon *et al.*, 2011). In this study, the freedom to choose indicators did not seem to lead to situations in which farmers set health standards that are relatively easy to achieve, as meetings without an alert were relatively rare.

Testing a tool under field-like conditions to improve its relevance to practice

The process evaluation framework chosen provided useful additional information on the context and implementation of the HHPM programme. A dialogue between designers and end-users that have tested a prototype creates a learning environment in which the response of the tool to 'field-like' situations can be discussed (Cerf *et al.*, 2012). Moreover, testing the tool and reporting on the context of its implementation should make reports more useful for future users and decision makers (Waters *et al.*, 2011).

Conclusion

Although compliance to the HHPM programme was not fully fulfilled and no significant effect of the intervention on herd health was demonstrated, the HHPM-tool triggered multiple factors that could promote the dialogue between a farmer and an advisor, and stimulate farmers' decisions to implement advice on animal health management. The methodological challenges encountered for the evaluation of the HHPM programs' impact on herd health consolidates the need for specific research approaches for the evaluation of such complex interventions.

Acknowledgements

The authors would like to thank the participating farmers and advisors for their extensive collaboration and feedback. This study received funding from the European Union's Seventh Framework Programme for research, technological development and demonstration under grant agreement number 311824 (IMPRO), and by the Region Pays de la Loire under grant agreement number 201309596.

Supplementary material

To view supplementary material for this article, please visit <https://doi.org/10.1017/S1751731117002841>

References

Anonymous 2007. Council regulation (EC) No 834/2007 on organic production and labelling of organic products and repealing Regulation (EEC) No 2092/91.

Beekhuis-Gibbon L, Devitt C, Whyte P, O'Grady L, More SJ, Redmond B and Doherty ML 2011. A HACCP-based approach to mastitis control in dairy herds. Part 2: implementation and evaluation. *Irish Veterinary Journal* 64, 7.

Bell NJ, Bell MJ, Knowles TG, Whay HR, Main DJ and Webster AJF 2009. The development, implementation and testing of a lameness control programme based on HACCP principles and designed for heifers on dairy farms. *The Veterinary Journal* 180, 178–188.

Brand A, Noordhuizen JPTM and Schukken YH 2001. Herd Health and Production Management in dairy practice. Wageningen Academic Publishers, Wageningen, The Netherlands.

Cerf M, Jeuffroy MH, Prost L and Meynard JM 2012. Participatory design of agricultural decision support tools: taking account of the use situations. *Agronomy for Sustainable Development* 32, 899–910.

Craig P, Dieppe P, Macintyre S, Health P, Unit S, Michie S and Petticrew M 2008. Developing and evaluating complex interventions: the new Medical Research Council Guidance. *British Medical Journal* 337, 1655.

Duval JE, Bareille N, Fourichon C, Madouasse A and Vaarst M 2016a. Perceptions of French private veterinary practitioners on their role in organic dairy farms and opportunities to improve their advisory services for organic dairy farmers. *Preventive Veterinary Medicine* 128, 10–21.

Duval JE, Fourichon C, Madouasse A, Sjöström K, Emanuelson U and Bareille N 2016b. A participatory approach to design monitoring indicators of production diseases in organic dairy farms. *Preventive Veterinary Medicine* 133, 12–22.

Garforth C 2011. Effective communication to improve udder health: can social science help? In *Udder health and communication* (ed. H Hogeveen and TJGM Lam), pp. 55–66. Wageningen Academic Publishers, Utrecht, The Netherlands.

Green MJ, Leach KA, Breen JE, Green LE and Bradley AJ 2007. National intervention study of mastitis control in dairy herds in England and Wales. *Veterinary Record* 160, 287–293.

International Federation of Organic Agriculture Movements 2005. Principles of organic agriculture. IFOAM. Retrieved on 13 February 2017 from <http://www.ifoam.bio/en/organic-landmarks/principles-organic-agriculture>.

Ivemeyer S, Smolders G, Brinkmann J, Gratzner E, Hansen B, Henriksen BIF and Walkenhorst M 2012. Impact of animal health and welfare planning on medicine use, herd health and production in European organic dairy farms. *Livestock Science* 145, 63–72.

Ivemeyer S, Walkenhorst M, Heil F, Notz C, Maeschli A, Butler G and Klocke P. 2009. Management factors affecting udder health and effects of a one year extension program in organic dairy herds. *Animal* 3, 1596–1604.

Janz NK and Becker MH 1984. The health belief model: a decade later. *Health Education Quarterly* 11, 1–47.

Klerkx L and Jansen J 2010. Building knowledge systems for sustainable agriculture: supporting private advisors to adequately address sustainable farm management in regular service contacts. *International Journal of Agricultural Sustainability* 8, 148–163.

Krieger M, Sjöström K, Blanco-Penedo I, Madouasse A, Duval JE, Bareille N, Fourichon C, Sundrum A and Emanuelson U. 2017. Prevalences of production diseases in European organic dairy herds. *Livestock Science* 198, 104–108.

Malterud K 2001. The art and science of clinical knowledge: evidence beyond measures and numbers. *The Lancet* 358, 397–400.

Moore GF, Audrey S, Barker M, Bond L, Bonell C, Hardeman W and Baird J 2015. Process evaluation of complex interventions: Medical Research Council guidance. *British Medical Journal* 350, h1258.

Oakley A, Strange V, Bonell C, Allen E, Stephenson J, and RIPPLE Study Team. 2006. Process evaluation in randomised controlled trials of complex interventions. *British Medical Journal* 332, 413–416.

Vaarst M, Bennedsgaard TW, Klaas I, Nissen TB., Thamsborg SM and Østergaard S 2006. Development and daily management of an explicit strategy of nonuse of antimicrobial drugs in twelve Danish organic dairy herds. *Journal of Dairy Science* 89, 1842–1853.

Vaarst M, Winckler C, Roderick S, Smolders G, Ivemeyer S, Brinkmann J and Huber J 2011. Animal health and welfare planning organic dairy cattle farms. *The Open Veterinary Science Journal* 5, 19–25.

Waters E, Hall BJ, Armstrong R, Doyle J, Pettman TL and De Silva-Sanigorski A 2011. Essential components of public health evidence reviews: capturing intervention complexity, implementation, economics and equity. *Journal of Public Health* 33, 462–465.