



HAL
open science

**Tick-borne pathogens in ticks (Acari: Ixodidae)
collected from various domestic and wild hosts in
Corsica (France), a Mediterranean island environment**

Sébastien Grech-Angelini, Frederic Stachurski, Muriel Vayssier Taussat,
Elodie Devillers, François Casabianca, Renaud Lancelot, Gerrit Uilenberg,
Sara Moutailler

► **To cite this version:**

Sébastien Grech-Angelini, Frederic Stachurski, Muriel Vayssier Taussat, Elodie Devillers, François Casabianca, et al.. Tick-borne pathogens in ticks (Acari: Ixodidae) collected from various domestic and wild hosts in Corsica (France), a Mediterranean island environment. *Transboundary and emerging diseases*, 2020, on-line, 13 p. 10.1111/tbed.13393 . hal-02621778

HAL Id: hal-02621778

<https://hal.inrae.fr/hal-02621778>

Submitted on 8 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **ORIGINAL ARTICLE**

2 **Tick-borne pathogens in ticks (Acari: Ixodidae) collected from various domestic**
3 **and wild hosts in Corsica (France), a Mediterranean island environment**

4

5 **Running title:** Eleven TBPs from six genera were found in ticks collected on Corsican animal hosts.

6 Sébastien Grech-Angelini^{1,*}, Frédéric Stachurski^{2,3}, Muriel Vayssier-Taussat⁴, Elodie Devillers⁴,

7 François Casabianca¹, Renaud Lancelot^{2,3}, Gerrit Uilenberg⁵, Sara Moutailler⁴

8 ¹INRA, UR045 Laboratoire de recherches sur le développement de l'élevage, Corte, France

9 ²CIRAD, UMR ASTRE, F-34398 Montpellier, France

10 ³ASTRE, CIRAD, INRA, Univ. Montpellier, Montpellier, France

11 ⁴UMR BIPAR, Animal Health Laboratory, ANSES, INRA, Ecole Nationale Vétérinaire d'Alfort,
12 Université Paris-Est, Maisons-Alfort, France

13 ⁵“A Surgente”, Route du Port, Cargèse, France

14 Correspondence:

15 Sébastien Grech-Angelini, GTV Corse, Ghisonaccia, France ; email: grech.angelini@gtvcorse.fr ;

16 Sara Moutailler, UMR BIPAR, Maisons-Alfort, France, sara.moutailler@anses.fr

17 **Present address:*

18 - *Groupement Technique Vétérinaire de Corse, Ghisonaccia, France*

19 - *UMR BIPAR, Animal Health Laboratory, ANSES, INRA, Ecole Nationale Vétérinaire d'Alfort,*

20 *Université Paris-Est, Maisons-Alfort, France*

21

22

23 ABSTRACT

24 Corsica is a touristic mountainous French island in the north-west of the Mediterranean Sea presenting
25 a large diversity of natural environments where many interactions between humans, domestic animals
26 and wild fauna occur. Despite this favourable context, tick-borne pathogens (TBPs) have not
27 systematically been investigated. In this study, a large number of TBPs were screened in ticks
28 collected during one year from domestic and wild hosts in Corsica. More than 1,500 ticks belonging
29 to nine species and five genera (*Rhipicephalus*, *Hyalomma*, *Dermacentor*, *Ixodes* and *Haemaphysalis*)
30 were analysed individually or pooled (by species, gender, host and locality). A real-time microfluidic
31 PCR was used for high-throughput screening of TBPs DNA. This advanced methodology permitted
32 the simultaneous detection of 29 bacterial and 12 parasitic species (including *Borrelia*, *Anaplasma*,
33 *Ehrlichia*, *Rickettsia*, *Bartonella*, *Candidatus* Neoehrlichia, *Coxiella*, *Francisella*, *Babesia* and
34 *Theileria*). CCHF virus was investigated individually in tick species known to be vectors or carriers of
35 this virus. In almost half of the tick pools (48%), DNA from at least one pathogen was detected and
36 eleven species of TBPs from six genera were reported. TBPs were found in ticks from all collected
37 hosts and were present in more than 80% of the investigated area. The detection of some pathogens
38 DNA confirmed their previous identification in Corsica, such as *Rickettsia aeschlimannii* (23% of
39 pools), *Rickettsia slovaca* (5%), *Anaplasma marginale* (4%) and *Theileria equi* (0.4%), but most
40 TBPs DNA was not reported before in Corsican ticks. This included *Anaplasma phagocytophilum*
41 (16%), *Rickettsia helvetica* (1%), *Borrelia afzelii* (0.7%), *Borrelia miyamotoi* (1%), *Bartonella*
42 *henselae* (2%), *Babesia bigemina* (2%) and *Babesia ovis* (0.5%). The important tick infection rate and
43 the diversity of TBPs reported in this study highlight the probable role of animal reservoir hosts for
44 zoonotic pathogens and human exposure to TBPs on Corsica.

45 **Keywords:** Ticks (Ixodidae), Tickborne pathogens, Domestic animals, Wild animals, Corsica, France

46

47

48 1. INTRODUCTION

49 The regional sanitary importance of ticks depends on the tick species and tick-borne pathogens (TBPs)
50 present in an area, and to a large extent on the local climate, management and breeds of livestock and
51 human activities (Jongejan and Uilenberg, 2004). The role of ticks as vectors of human pathogens is
52 second in importance to that of mosquitoes (Parola and Raoult, 2001) and they are worldwide the
53 most important vectors in the veterinary field (Nicholson et al., 2009). Ticks can transmit many
54 varieties of pathogens, including bacteria, parasites and viruses. Moreover, human tick-borne diseases
55 are usually zoonotic and asymptomatic for non-human vertebrate hosts which most often are the
56 reservoirs of pathogens causing human infection (Jongejan and Uilenberg, 2004).

57 Corsica is a French island in the western part of the Mediterranean area, situated 15 km north of
58 Sardinia and 90 km west of Tuscany in Italy. It is the fourth Mediterranean island in size and the most
59 mountainous and forested one. The island consists of two departments (Corse-du-Sud and Haute-
60 Corse) and 360 communes (the smallest administrative unit in France; Fig. 1). Tourism (three million
61 people annually, 320,000 permanent inhabitants), extensive farming (sheep, goats, pigs and cattle),
62 hunting and hiking are important activities in Corsica (Grech-Angelini et al., 2016b). Therefore, in
63 this context, permanent interactions occur between livestock, wildlife and humans in a small area,
64 which certainly favour the circulation of TBPs, including zoonotic ones. Corsica is also on the route
65 of migratory birds which create a natural link between Africa and Europe and could spread ticks
66 infected with TBPs (Hoffman et al., 2018).

67 Only scattered observations on the tick fauna of Corsica, mostly grouped together in a book on the
68 ticks of France (Pérez-Eid, 2007), were available before 2014. From May 2014 to May 2015, a large
69 tick survey (Grech-Angelini et al, 2016b) on domestic and a few wild animals led to the identification
70 of nine species: *Rhipicephalus bursa*, *Hyalomma marginatum*, *Dermacentor marginatus*, *Rh.*
71 *sanguineus* sensu lato, *Hy. scupense*, *Ixodes ricinus*, *Haemaphysalis punctata*, *Rh. (Bo.) annulatus*,
72 and *Ha. sulcata*. The diversity of the Corsican tick species, characterized by ticks usually collected in

73 humid environments (*I. ricinus*) and others in drier areas (*Hyalomma* spp., *Rh. bursa*), suggested a
74 potential high diversity of TBPs on the island.

75 Some of the TBPs occurring on Corsica have been reported more or less reliably in former and recent
76 studies. Various species of the genera *Anaplasma*, *Rickettsia* and *Ehrlichia* (ICTTD, 2000; Matsumoto
77 et al., 2004; Dahmani et al., 2017; Selmi et al., 2017; Cabezas-Cruz et al., 2019; Cicculli et al., 2019a,
78 2019b, 2019c), the genera *Babesia* and *Theileria* (ICTTD, 2000) have been identified, and *Borrelia*
79 *burgdorferi* sensu lato was recently reported (Cicculli et al., 2019c). It remained uncertain whether all
80 of the Corsican TBPs were known. The aim of this study was to have a large overview regarding the
81 TBPs carried in more than 1,500 ticks collected on different Corsican animal hosts, focusing on the
82 main pathogens of medical and veterinary importance known in the Mediterranean area, including
83 *Borrelia* spp., *Rickettsia* spp., *Anaplasma* spp., *Francisella* spp., *Ehrlichia* spp., *Coxiella* spp.,
84 *Theileria* spp., *Babesia* spp., *Bartonella* spp., *Candidatus* Neoehrlichia and CCHF virus.

85

86 **2. MATERIAL AND METHODS**

87 **2.1 Study area and tick collection.**

88 A large-scale tick collection has been conducted on domestic and a few wild animals in Corsica
89 (Grech-Angelini et al., 2016a and 2016b). Cattle were chosen as host model because of the extensive,
90 free-ranging livestock farming system, with a low frequency of acaricide treatments. Ticks were
91 collected over a year (May 2014 to May 2015) in the three cattle Corsican slaughterhouses. Ticks on
92 sheep, goats and horses were collected less systematically from May to August 2014 in three farms for
93 each host. Ticks from domestic carnivores were provided by practising veterinarians. Ticks from wild
94 boars, mouflons and deer were obtained respectively from hunters, from staff of the National Office
95 for Hunting and Wildlife (ONCFS) and of the Regional Natural Park of Corsica (PNRC). Ticks were
96 stored in 70 % ethanol at -20°C until their identification.

97 Ticks were identified on their morphology; when deemed necessary, some specimens were also
98 molecularly examined by sequencing mitochondrial *cox1* (cytochrome *c* oxidase subunit 1) and 16S
99 ribosomal RNA genes and ITS2 (internal transcribed spacer 2) (Grech-Angelini et al., 2016a and
100 2016b).

101 **2.2 Pools of ticks**

102 Tick were analysed individually or by pools consisting of two to five ticks. *Rhipicephalus bursa*, *Hy.*
103 *marginatum*, *Ha. punctata*, *Rh. sanguineus* s.l. and *D. marginatus*, found in large numbers on their
104 hosts, were grouped in pools by sex, host and locality. Other species found more rarely or with a
105 special sanitarian interest were systematically analysed individually: *I. ricinus*, *Hy. scupense*, *Ha.*
106 *sulcata* and *Rh. (Bo.) annulatus*.

107 **2.3 DNA and RNA extraction**

108 After washing once in 70% ethanol for 5 min and twice in distilled water for 5 min, pools of one to
109 five ticks were crushed in 300 µl of DMEM with 10% foetal calf serum and six steel balls using the
110 homogenizer Precellys® 24 Dual (Bertin, France) at 5500 rpm for 40s. DNA was then extracted using
111 100µl of the homogenate according to the Wizard genomic DNA purification kit (Promega, France)
112 following the manufacturer's instruction. Total DNA per sample was eluted in 50 µl of rehydration
113 solution and stored at -20°C until further use (Michelet et al, 2014). Using 100µl of tick homogenate
114 for *Hyalomma* spp. and *Rh. bursa* (species known to transmit or to carry CCHF virus), the Nucleospin
115 RNA II kit (Macherey-Nagel, Duren, Germany) was used for total RNA extraction following the
116 manufacturer's instruction (Gondard et al., 2018).

117 **2.4 Detection of tick-borne pathogen DNA or RNA**

118 Bacteria and parasites

119 Forty-one sets of primers and probes were used in this study to detect TBPs (29 bacterial and 12
120 parasitic species, Tab. 1). The BioMark™ real-time PCR system (Fluidigm, USA) was used for high-
121 throughput microfluidic real-time PCR amplification using 48.48 dynamic arrays (Fluidigm). These

122 chips dispense 48 PCR mixes and 48 samples into individual wells, after which on-chip microfluidics
123 assemble PCR reactions in individual chambers prior to thermal cycling resulting in 2,304 individual
124 reactions (Michelet et al., 2014).

125 Conventional PCR using primers targeting genes or regions other than those of the BioMark™ system
126 were used to confirm the presence of pathogenic DNA in the field samples. Amplicons were
127 sequenced by Eurofins MWG Operon (Germany), and then assembled using BioEdit software (Ibis
128 Biosciences, Carlsbad). An online BLAST (National Center for Biotechnology Information) was used
129 to compare results with published sequences listed in GenBank sequence databases (Michelet et al,
130 2014; Gondard et al., 2019).

131 Specific detections of CCHF virus and *Theileria* spp.

132 CCHF virus RNA was specifically investigated in tick species known to transmit, or at least able to
133 carry it: *Hy. marginatum*, *Hy. scupense*, and *Rh. bursa*. The presence of *Theileria* spp. DNA was also
134 investigated by specific real-time PCR in *Hy. scupense* and *Haemaphysalis* spp. because *Hy. scupense*
135 is an efficient vector of *T. annulata* and because *T. buffeli*, of which *Ha. punctata* is a vector, has
136 already been reported in Corsica (ICTTD, 2000) (for primers and probes see Gondard et al., 2018 and
137 2019).

138 **2.5 Data analysis**

139 The infection rate is given for all pathogens in each tick species and each host. The infection rate of
140 individually analysed tick species is the real one detected. The infection rate of pooled tick species (cf.
141 Methods, 2.4) means that in an infected pool there was at least one tick that gave a positive result. To
142 assess differences in tick species infection between development stage and sex a Kruskal-Wallis test
143 was used. Differences were considered as statistically significant if $p < 0.05$.

144

145

146

147 **3. RESULTS**

148 **3.1 Collected ticks and pool preparation**

149 More than three thousand ticks (3,134) from nine species and five genera were collected from May
150 2014 to May 2015 (Grech-Angelini et al., 2016b). Almost half of them (1,523) were analysed to
151 detect animal and human pathogens DNA or RNA (Tab. 2). A total of 569 samples were analysed
152 (consisting of one to five ticks): 269 adult females, 261 adult males and 39 nymphs. Nymphs of four
153 species were found: *Rh. bursa*, *Hy. scupense*, *Rh. sanguineus* group and *Rh. (Bo.) annulatus* (Tab. 2).
154 *Rhipicephalus bursa* ticks (4.8 ticks per pool on average), *Hy. marginatum* (4.1), *Dermacentor*
155 *marginatus* (4.1), *Rh. sanguineus* sensu lato (3.7) and *Ha. punctata* (2.5) were systematically pooled
156 whereas ticks from the other four species were analysed individually (*Rh. (Bo.) annulatus*, *Ha.*
157 *sulcata*, *Hy. scupense* and *Ixodes ricinus*).

158 **3.2 Detection of TBPs DNA or RNA in tick species**

159 As for each tick species, the proportion of infected pools was not significantly different between
160 development stage and sex ($p > 0,05$), the identified pathogens were presented by tick species, not
161 separately for the stage of development and the sex of adults. Almost half of the samples (48%) were
162 positive for at least one pathogen DNA (Tab. 2) and among them 12% were positive for two
163 pathogens DNA or more. The most infected tick species were *Hy. marginatum* (100% of pools), *I.*
164 *ricinus* (70% of the individual ticks) and *D. marginatus* (66% of pools). Pathogen DNA was found in
165 all tick species collected in Corsica except in *Ha. sulcata*, but only two specimens were analysed
166 (Tab. 2 and Tab. 3).

167 **3.3 Pathogens identified**

168 DNA of eleven pathogen species from six genera were identified in the ticks: *Borrelia* spp., *Rickettsia*
169 spp., *Anaplasma* spp., *Babesia* spp., *Bartonella* spp. and *Theileria* spp.

170 ***Borrelia* spp.**

171 *Borrelia* spp. DNA was detected in two tick species and two different *Borrelia* were identified. DNA
172 of *Borrelia miyamotoi* was found in *Ixodes ricinus* (2% of individual ticks) and in *Ha. punctata* (13%
173 of the pools). A sequence was obtained from *I. ricinus* (GenBank, accession number: MK732472),
174 and showed 100% homology with reference sequences from *B. miyamotoi* strain isolated in Austria
175 (AN: KP202177). DNA of *B. afzelii* was reported only in *I. ricinus* (4% of ticks). Unfortunately, the
176 sequence could not be obtained.

177 ***Rickettsia* spp.**

178 DNA of *Rickettsia* spp. was detected in six tick species (Tab. 3). Three species of *Rickettsia* were
179 identified. *Rickettsia aeschlimannii* was found in *Rh. bursa* (27% of pools), *Hy. marginatum* (100% of
180 pools), *Rh. sanguineus* s.l. (10% of pools) and *Hy. scupense* (5% of individual ticks). The four
181 sequences obtained from *Hy. marginatum* and *Hy. scupense* presented 100% identity. One sequence
182 was deposited in GenBank (AN: MK732478), and showed 99% homology with reference sequences
183 from *R. aeschlimannii* strain isolated in Egypt (HQ335153). *Rickettsia slovaca* was mostly identified
184 in *D. marginatus* (66% of pools). Three pools of *Rh. sanguineus* s.l. (7%) and one *Hy. scupense* tick
185 (0.7%) were also positive for this bacterium. The sequence obtained from *Hy. scupense* (GenBank,
186 AN: MK732480) showed 99% homology with reference sequences from *R. slovaca* strain isolated in
187 United States (AN: KR559551). *Rickettsia helvetica* was detected only in *I. ricinus* (6% of ticks) but
188 the sequence could not be obtained.

189 ***Anaplasma* spp.**

190 DNA of *Anaplasma* spp. was detected in eight tick species (Tab. 3) and two species were recorded: *A.*
191 *phagocytophilum* and *A. marginale*. *Anaplasma phagocytophilum* DNA was mainly found in *I. ricinus*
192 (63% of ticks), but it was also reported in *Rh. sanguineus* s.l. (7% of pools), *Rh. bursa* (7% of pools),
193 *Hy. marginatum* (1% of pools), *Hy. scupense* (2% of ticks), *D. marginatus* (3% of pools) and *Ha.*
194 *punctata* (7%). Unfortunately, the sequence of *A. phagocytophilum* could not be obtained. *Anaplasma*
195 *marginale* DNA was frequently detected in *Rh. (Bo.) annulatus* (18% of ticks) but it was also

196 identified in *Rh. bursa* (8% of pools), *Hy. marginatum* (6% of pools), *I. ricinus* (2% of ticks), *Ha.*
197 *punctata* (7% of pools) and *Rh. sanguineus* s.l. (7% of pools). The sequence obtained from *Rh. (Bo.)*
198 *annulatus* (GenBank, AN: MK732471) showed 99% homology with reference sequences from *A.*
199 *marginale* strain isolated in South Africa (AN: AF414873).

200 ***Babesia* spp.**

201 DNA of *Babesia* spp. was detected only in *Rh. bursa* and two species were identified. *Babesia*
202 *bigemina* was found in 9% of the pools of *Rh. bursa*. The two sequences obtained presented 100%
203 identity. One sequence was deposited in GenBank (AN: MK732475) and showed 100% homology
204 with reference sequences from *Ba. bigemina* strain from Virgin Islands (AN: EF458206). *Babesia ovis*
205 was identified in 3% of *Rh. bursa* pools. The sequence obtained (GenBank, AN: MK732477) showed
206 92% homology with reference sequences from *Ba. ovis* strain isolated in Sudan (AN: AY260171).

207 ***Bartonella* spp.**

208 *Bartonella henselae* was the only species of the genus *Bartonella* detected and its DNA was identified
209 in seven tick species (Tab. 3) but mainly in *Rh. bursa* (5% of pools), *I. ricinus* (3% of ticks), *Rh. (Bo.)*
210 *annulatus* (9% of ticks) and *Ha. punctata* (3% of pools). The five sequences obtained in *Rh. bursa* and
211 *I. ricinus* presented 100% identity. One sequence was deposited in GenBank (AN: MK732473) and
212 showed 100% homology with reference sequences from *Bar. henselae* strain Houston-1 (AN:
213 BX897699).

214 ***Theileria* spp.**

215 *Theileria equi* DNA was reported in one pool of *Rh. bursa* (1%) and in one *Rh. (Bo.) annulatus* (9%
216 of ticks) (Tab. 3). The sequence obtained from *Rh. bursa* (GenBank, AN: MK732476) showed 99%
217 homology with reference sequences from *T. equi* isolated in Brazil (AN: KJ573370). No *Theileria*
218 *annulata* DNA was detected by the BioMark™ real-time PCR system in the analysed ticks. Individual

219 PCR for *Theileria* spp. did not show the presence of any other species of *Theileria* in the tick species
220 analysed (*Hy. scupense* and *Haemaphysalis* spp.).

221

222 **Others pathogens and micro-organisms**

223 *Francisella tularensis* was not identified in this study but *Francisella*-like endosymbionts were
224 detected in five tick species: *Hy. marginatum* (90% of pools), *Hy. scupense* (7% of ticks), *I. ricinus*
225 (4% of ticks), *Rh. sanguineus* s.l. (10% of pools) and *Rh. (Bo.) annulatus* (27% of ticks). Neither
226 DNA of *Coxiella* spp. nor DNA of *Candidatus Neoehrlichia mikurensis* were identified in this study.
227 RNA of CCHF virus was not found in *Hy. marginatum* (89 pools; 362 ticks), *Rh. bursa* (108; 518) and
228 *Hy. scupense* (135 ticks).

229 For some *Rickettsiae*, the exact species could not be determined t. These *Rickettsiae* were reported in
230 seven tick species: *Rh. bursa* (8% of pools), *Hy. scupense* (2% of ticks), *I. ricinus* (9% of ticks), *Rh.*
231 *sanguineus* s.l. (12% of pools), *D. marginatus* (11% of pools), *Ha. punctata* (3% of pools) and *Rh.*
232 *(Bo.) annulatus* (9% of ticks). One sequence was obtained from *Rh. bursa* (GenBank, AN:
233 MK732479) and showed 99% homology with reference sequences from *Rickettsia*-like endosymbiont
234 strain 162 citrate synthase (gltA) gene (AN: JQ925589).

235 **3.4 Presence of pathogens in ticks collected on the different hosts**

236 Ticks were collected from different hosts: 404 samples (consisting to one to five ticks) originated
237 from cattle, 19 from goats, 12 from sheep, 24 from horses, 22 from dogs, four from cats (Tab. 4), 41
238 from wild boars, 26 from mouflons, 13 from deer and two from hedgehog and birds (Tab. 5). For
239 birds and deer, as the collected ticks were all *I. ricinus*, they were individually analysed.

240 *Borrelia* spp. DNA was found only in ticks from cattle; *Borrelia miyamotoi* and *Bo. afzelii* infected
241 respectively 2% and 1% of pools from this host.

242 *Rickettsia* spp. DNA was found in ticks from almost all collected hosts. *Rickettsia aeschlimannii* was
243 the most important pathogen infecting ticks from domestic ruminants and horses, reported in 23% of
244 pools from cattle, 37% from goats, 58% from sheep and 75% from horses. It was rarely found in pools
245 from dogs (5%), wild boars (7%) and mouflons (4%). *Rickettsia slovaca* DNA was mainly found in
246 pools from wild boars (59%) but it was also reported in pools from cattle (0.5%), horses (4%), dogs
247 (5%) and hedgehog (one positive pool). *Rickettsia helvetica* DNA was identified only in ticks
248 collected from cattle (2%).

249 *Anaplasma phagocytophilum* DNA was reported in ticks from cattle (18% of pools), goat (11% of
250 pools), cats (25% of pools), wild boars (2% of pools), deer (92% of individual ticks) and birds (two
251 positive ticks) whereas *A. marginale* DNA was found in pools from cattle (6%) and mouflons (4%).

252 *Babesia bigemina* DNA was detected mainly in ticks sampled on cattle (2% of pools) but one pool of
253 ticks from goats was also positive for this pathogen. *Babesia ovis* DNA was reported in two hosts, one
254 pool of ticks collected on cattle and two pools from mouflons were positive.

255 *Bartonella henselae* DNA was reported in pools from cattle (2%), goat (11%), horses (4%) and wild
256 boars (2%). *Theileria equi* DNA was found in one pool from cattle and one from wild boars.

257 **3.5 Geographical distribution of TBPs found in ticks**

258 Ticks from 82 municipalities were analysed and among them, pathogen DNA was found in 68 (Fig.
259 1). *Rickettsia* spp. were widespread (Fig. 1a) as they were reported in more than 80% of the
260 municipalities investigated. *Rickettsia slovaca* DNA was found in ticks collected from 13
261 municipalities, *R. aeschlimannii* DNA from 49 and *R. helvetica* DNA was reported from a single
262 municipality in the centre of Corsica (Haute-Corse). *Anaplasma* spp. (Fig. 1b) were also distributed in
263 a large part of the investigated area, found in 37% of the municipalities. *Anaplasma phagocytophilum*
264 DNA was reported in 19 municipalities and *A. marginale* DNA in 16. *Borrelia* spp. were found in five
265 municipalities (Fig. 1c). *Borrelia miyamotoi* DNA was identified in all these municipalities whereas
266 *B. afzelii* DNA was reported only in one from Haute-Corse. *Babesia* spp. were identified in 10

267 municipalities as well in Corse-du-Sud as in Haute-Corse (Fig. 1d). *Bartonella henselae* DNA was
268 identified in 12 municipalities whereas *T. equi* DNA was found in two municipalities near the eastern
269 coast in Haute-Corse (results not shown).

270

271 **4. DISCUSSION**

272 In this study, a method using multiple primers/probe sets was implemented to perform high-
273 throughput detection of TBPs DNA. This large-scale investigation has (i) enabled the detection of rare
274 pathogens, (ii) generated prevalence estimations for pathogens, thus creating a comprehensive
275 overview of the epidemiological situation for 29 bacteria and 12 parasites present or not found in ticks
276 from Corsica. A positive tick or a positive pool means it contains DNA (or RNA) sequences similar to
277 those of corresponding genes of TBPs, and does not necessarily mean that the TBP is present in the
278 tick. For the sake of commodity, the term “infection rate” is often used in this study but it means more
279 “positive rate for pathogen DNA”. As all collected ticks were removed from their hosts, potentially
280 infected by pathogens, the identification of pathogen DNA in a tick suggests its presence in Corsica
281 but the tick species is not necessarily a biological vector of the pathogen. The micro-organism or its
282 DNA may be simply present in the ingested blood meal, the host having been infected by a real vector
283 species of tick (Estrada-Peña et al., 2013). However, it has to be noted that in this survey, the main
284 pathogens were most often identified in their natural vector suggesting a higher link between one
285 pathogen and its natural vector than with other tick species. Almost 70% of the *R. aeschlimannii* were
286 detected in *Hy. marginatum*, 86% of the *R. slovaca* in *D. marginatus*, 80% of *A. phagocytophilum* in
287 *I. ricinus*, more than 60% of *A. marginale* in ticks from the genus *Rhipicephalus* and all the *Babesia*
288 spp. were identified in *Rh. bursa*.

289 **4.1 Tick-borne pathogens reported for the first time in Corsican ticks**

290 In almost half of the pools, DNA of at least one pathogen was reported and 11 species of pathogens
291 from six genera were identified. Among them, seven were reported for first time in Corsican ticks: *B.*
292 *afzelii*, *B. miyamotoi*, *R. helvetica*, *A. phagocytophilum*, *Ba. ovis*, *Ba. bigemina* and *Bar. henselae*.

293 *Borrelia afzelii* DNA, found in 4% of the Corsican *I. ricinus*, belongs to the *Borrelia burgdorferi* s.l.
294 group, the causative agents of Lyme disease, the most important zoonotic infection in Europe. The
295 presence of *B. burgdorferi* s.l. is linked to its natural vectors, ticks of the genus *Ixodes*, mainly *I.*
296 *ricinus* in the Mediterranean area. This pathogenic agent was believed to be rare or even absent from
297 Corsica, but a few cases of Lyme disease have been reported in recent years by the French National
298 Centre of *Borrelia* and Santé Publique France (Vandenesch et al., 2014). *Borrelia burgdorferi* s.l. was
299 reported in Continental France, continental Italy, Spain (Hubálek *et al.*, 1997) and also in northern
300 Africa (Tunisia, Algeria and Morocco; Benredjem et al., 2014) but has not been found in ticks from
301 Sardinia and Sicilia, two Mediterranean islands with a drier climate than Corsica. Recently, it was
302 reported in *I. ricinus* collected from rats in Corsica (Cicculli et al., 2019c), but without determining
303 the exact species. The identification of *B. afzelii* DNA in ticks confirmed the exposure of Corsican
304 population to Lyme disease.

305 DNA of *Borrelia miyamotoi*, first identified in 1995 in ticks from Japan (Fukunaga et al., 1995), was
306 also detected for the first time in Corsica. It belongs to the relapsing fever group of *Borrelia* and its
307 natural vectors are *Ixodes* ticks. *Borrelia miyamotoi* is currently considered as an emerging pathogen
308 affecting humans. Relapsing fever borreliosis is characterized by influenza-like illness and one or
309 more relapse episodes of bacteraemia and fever (Platonov *et al.*, 2011). *Borrelia miyamotoi* was
310 reported in 3% of ticks from continental France (Cosson et al., 2014) and 0.7% of northern Italy
311 (Ravagean *et al.*, 2018). Although no human case of *B. miyamotoi* was identified in Corsica and in the
312 Mediterranean area, the bacterium has been found in humans throughout Europe (Siński et al., 2016).

313 *Rickettsia helvetica* DNA was only found in *I. ricinus* which is a natural vector of this bacterium. It
314 was initially reported in 1979 in Switzerland from *I. ricinus* (Beati et al., 1993) and now the species
315 has been identified in ticks from many countries over the world. It is especially well spread in the

316 Mediterranean area including continental France (Parola *et al.*, 1998), Italy (Beninati *et al.*, 2002),
317 Spain (Fernández-Soto *et al.*, 2004), Croatia (Dobec *et al.*, 2009), Algeria (Kernif *et al.*, 2012) and
318 Tunisia (Sfar *et al.*, 2008). Human cases of rickettsiosis caused by *R. helvetica* have been reported in
319 Italy and continental France (Portillo *et al.*, 2015). *Rickettsia helvetica* is a member of the Spotted
320 Fever Group (SFG) and reportedly causes a self-limiting illness associated with headache and
321 myalgias (Parola *et al.*, 2013).

322 *Anaplasma phagocytophilum* DNA was mostly found in *I. ricinus*, an important vector of this
323 zoonotic agent responsible for human granulocytic anaplasmosis (HGA), tick-borne fever (or pasture
324 fever) of ruminants and equine anaplasmosis. In Corsica, this pathogen was relatively unknown, and
325 was previously only detected in a single sample of bovine blood (ICTTD, 2000). There is no HGA
326 case reported in Corsica, but in continental France the first one was identified in 2003 and other were
327 infrequently reported (Edouard *et al.*, 2012). Human cases occurred also in continental Italy (Ruscio
328 and Cinco, 2003). A recent study showed that, in the French Basque Country, 22.4% of collected ticks
329 contained *A. phagocytophilum* DNA (Dahmani *et al.*, 2017). It was also reported from Sardinia
330 (Alberti *et al.*, 2005), Sicily (Torina *et al.*, 2010) and Northern Africa (Dahmani *et al.*, 2015).

331 *Babesia* spp. are protozoan blood parasites with more than 100 described species. In this study, *Ba.*
332 *bigemina* and *Ba. ovis* DNA were found for the first time in ticks collected in Corsica. *Babesia*
333 *bigemina* is a causal agent for bovine babesiosis and it occurs in most areas of the world (Uilenberg,
334 2006). *Babesia ovis* is a causative agent of ovine babesiosis, with strains varying in pathogenicity and
335 is also widespread throughout the world (Uilenberg, 2006).

336 Ticks (especially *I. ricinus*) are highly suspected to be among the vectors of *Bartonella* species.
337 *Bartonella henselae* DNA was reported for the first time in ticks from Corsica and was found in seven
338 tick species. It causes an infection commonly encountered in cats (cat scratch disease) and potentially
339 dogs and humans worldwide (Álvarez-Fernández *et al.*, 2018). In Sardinia, it was found in at least
340 0.2% of the collected ticks (Chisu *et al.*, 2018).

341 **4.2 Other TBP identified in Corsica**

342 *Rickettsia* species were the pathogens with the highest infection rate found in the Corsican pools. They
343 are known as zoonotic agents and, if their pathogenicity and the reservoir role for animals are
344 discussed, most species of this genus cause important human diseases (Davoust et al., 2010). The real
345 impact of rickettsial diseases in Corsica is unknown, but some cases of Mediterranean spotted fever
346 (MSF, caused by *R. conorii*) are reported by local medical doctors and the French National Reference
347 Centre (CNR) for *Rickettsia* species. A former seroepidemiological study showed in 1985 that 4.8%
348 of people were exposed to these pathogens in Corse-du-Sud (Raoult et al., 1985). *Rickettsia*
349 *aeschlimannii* DNA was the most frequently detected, infecting 100% of *Hy. Marginatum* pools, one
350 of its main natural vectors (Matsumoto *et al*, 2004), confirming the important presence of this
351 bacterium in Corsica, already reported in 74% of *Hy. marginatum* collected by Matsumoto *et al.*
352 (2004). *Rickettsia aeschlimannii* was first isolated from *Hy. marginatum* in 1997 from Morocco (Beati
353 *et al.*, 1997) and is now reported in the whole Mediterranean area (Parola et al., 2013). *Rickettsia*
354 *aeschlimannii* infections in humans cause spotted fever; this has previously been confirmed in
355 northern Africa and South Africa, and in 2010 a first case occurred in Southern Europe, in a Greek
356 patient (Germanakis et al., 2013). Given the important infestation rate of this pathogen in Corsican
357 ticks, human exposure to *R. aeschlimannii* infection is high and human cases of tick-borne spotted
358 fever acquired in Corsica could often be due to *R. aeschlimannii*.

359 *Rickettsia slovaca* DNA was identified mainly in *D. marginatus* which is an important natural vector
360 of this bacterium. This report confirmed the results of Selmi et al. (2017) who identified *R. slovaca* in
361 *D. marginatus* collected on Corsican vegetation. *Rickettsia slovaca* was first isolated in 1968 in a *D.*
362 *marginatus* collected in Slovakia and is largely spread in the Mediterranean area. It was found in
363 Sardinia and continental Italy (Chisu et al., 2016 and Selmi et al, 2017), continental France (Michelet
364 et al., 2017), and Spain (Fernández-Soto et al., 2006). The first proven human case of *R. slovaca*
365 infection was reported in 1997 in continental France and this micro-organism is now known as a cause
366 of disease in various European Mediterranean countries (de Sousa et al., 2013). It is associated with

367 TIBOLA (tickborne lymphadenopathy) syndrome, characterized by lymph node enlargement and
368 scalp eschars (Parola et al., 2013). So far, the TIBOLA syndrome has not been reported in Corsica.
369 This study did not allow to identify DNA of other *Rickettsia* spp. previously reported in Corsica as
370 *Rickettsia felis*, a SFG *Rickettsia*, found in a flea (*Archaeopsylla erinacei*) collected from a fox (Marié
371 et al., 2012). Matsumoto et al. (2005) reported the species *R. massiliae* in *Rh. sanguineus* s.l. collected
372 on dogs in Corse-du-Sud and a recent study identified *R. africae* DNA in an *Amblyomma variegatum*,
373 tick species not established in Corsica but whose one adult was collected, certainly following
374 introduction of a nymph by migrating bird (Cicculli et al., 2019a). DNA of another species of
375 *Rickettsia*, Candidatus *Ri. barbariae*, was recently detected in ticks collected from Corsican cattle
376 (Cicculli et al., 2019c), but its pathogenicity remains unknown.

377 This study also showed that *Anaplasma* species are widespread in Corsican ticks. *Anaplasma*
378 *marginale* DNA was found in ticks from the genus *Rhipicephalus* which are its natural vectors but
379 also in *Hy. marginatum*, *I. ricinus* and *Ha. punctata*. *Anaplasma marginale* DNA was formerly
380 reported in blood from Corsican cattle (ICTTD, 2000) and two recent studies detected it in *Rh. bursa*
381 collected on cattle confirming its presence on the island (Dahmani et al., 2017 and Cicculli et al.,
382 2019c). *Anaplasma marginale* is distributed worldwide in tropical and subtropical regions. It is the
383 causative agent of erythrocytic bovine anaplasmosis that can affect various species of domestic and
384 wild ruminants (Aubry and Geale, 2011). From islands near to Corsica, it was identified in Sardinia
385 (Zobba et al, 2014) and Sicily (Torina et., 2010), but there is no report of its presence in continental
386 France. *Anaplasma ovis* DNA, not found in this study, was recently reported in blood (52%) and ticks
387 (*Rh. bursa*) collected from Corsican goats (Cabezas-Cruz et al., 2019). *Anaplasma bovis* and *A.*
388 *omatjenne* were formerly reported in cattle blood (ICTTD, 2000) and other potential new species of
389 *Anaplasma* were described from Corsican sheep blood (Dahmani et al., 2017).

390 Concerning *Babesia* species, *B. bovis*, not found in this study, was reported earlier from Corsican
391 cattle blood (ICTTD, 2000). *Babesia bovis* occurs in most subtropical and tropical regions of the
392 world (Uilenberg, 2006), and two of its proven vectors occur on Corsica (*Rh. bursa* and *Rh. (B.)*)

393 *annulatus*). Bovine babesiosis affecting cattle in Corsica is probably due either to *B. bigemina* or to *B.*
394 *bovis*.

395 No species from the genus *Ehrlichia* were found in this study although *E. minacensis*, a species
396 unknown so far from the Mediterranean area, has been detected recently in a *Hy. marginatum*
397 (Cicculli et al., 2019b).

398 *Theileria equi* DNA was identified in two pools. It is a causative agent of equine piroplasmosis, that
399 can be also due to *Babesia caballi* (not detected in this study). As the disease is largely spread in the
400 island, it is well known by local veterinary practitioners and horse owners. *Theileria annulata* DNA
401 has not been reported so far in the Corsican tick population. The high occurrence in Corsica of *Hy.*
402 *scupense*, one of its main natural vectors, highlights however the risk of transmission of *T. annulata* to
403 the local cattle population (Grech-Angelini et al., 2016a). *Theileria buffeli*, a benign cattle pathogen,
404 already detected in Corsica (ICTTD, 2000), was not found in this study.

405 A total of 332 pools (1,015 ticks) of three species (*Hy. marginatum*, *Hy. scupense* and *Rh. bursa*) were
406 tested for CCHF virus RNA and all were negative. The CCHF virus is mainly transmitted by ticks of
407 the genus *Hyalomma* and *Hy. marginatum* is its main natural vector in Europe. Crimean-Congo
408 haemorrhagic fever is now the most worldwide tick-borne viral infection of humans, and it can lead to
409 haemorrhagic manifestations and considerable mortality. It is also an important emerging zoonotic
410 disease in Turkey and south-eastern Europe (Dreshaj et al., 2016). Moreover, in 2016, the first
411 autochthonous human cases have been reported in Spain showing that the disease can occur in western
412 Europe (Negredo et al., 2017). Lately, CCHF RNA has been reported in a nymph of *Hy. rufipes* on a
413 migrating bird on the small Italian island of Ventotene, identifying migrating birds as possible
414 introduction pathway for CCHF virus into western Europe (Mancuso et al., 2019). *Hyalomma rufipes*,
415 an important natural vector of CCHF virus in subSaharan Africa, was ever found on migrating birds in
416 Corsica (Pérez-Eid, 2007). It is possible that CCHF virus was not detected in this study due to an
417 insufficient sample size of analysed ticks or a deterioration of the genetic material. Because of the
418 high frequency of *Hy. marginatum* in Corsica (the second most abundant tick species collected on

419 animal hosts, Grech-Angelini et al., 2016b), it would be necessary to closely monitor the possible
420 emergence of CCHF virus. More ticks, mainly *Hy. marginatum*, should be analysed and animal sera
421 should be collected to test the presence of antibodies against CCHF virus in domestic ruminants which
422 are known to be asymptomatic reservoir of CCHF virus (Bente et al., 2013).

423 **4.3 Potential Corsican animal reservoirs for zoonotic pathogens and human exposure**

424 These data concerning the presence of TBPs DNA on Corsica Island will be useful for studying
425 possible future emerging diseases and the role of animals as reservoirs of TBPs. This study
426 demonstrated the presence of expected pathogens DNA, as well as unexpected pathogens. Among the
427 eleven pathogens reported in ticks collected from animal hosts, seven are zoonotic: *B. miyamotoi*, *B.*
428 *afzelii*, *R. aeschlimannii*, *R. slovaca*, *R. helvetica*, *A. phagocytophilum* and *Bar. henselae*. Most of
429 them are asymptomatic, or their effects on animal health are unknown. *Rickettsia aeschlimannii* DNA
430 was reported in a high proportion in ticks from domestic ruminants and horses whereas *R. slovaca*
431 DNA was mostly identified in ticks from wild boars and *R. helvetica* DNA only in ticks from cattle.
432 All positive pools for *Borrelia* spp. consisted of ticks collected from cattle. These domestic or wild
433 animals could be specific reservoirs for the respective cited pathogens. *Anaplasma phagocytophilum*
434 was mainly found in *I. ricinus* on various hosts including migrating birds that showed the potential to
435 import pathogens on Corsica by this pathway. Even if no human cases of these pathogens have been
436 diagnosed in Corsica to date, the important interactions that occur between domestic animals, wild
437 fauna and humans highlight the risk for the human population. Some of these zoonotic pathogens
438 could circulate undetected in the Corsican human population. Such studies may help avert that more
439 efforts should be devoted to the surveillance of the animal reservoirs and the human exposure.
440 Pathogen DNA is not enough to be able to confirm the presence and the circulation of those pathogens
441 in Corsica. Nevertheless, this first step will allow to perform in-depth research on those interesting
442 pathogens to better characterize their epidemiological cycle. Indeed, serological survey in domestic
443 and wild animals and humans need to be carry out, as well as trying to isolate those pathogens from

444 infected ticks and/or infected animals to be able to conclude regarding the risk for human and animal
445 health.

446

447 **5. CONCLUSION**

448 In this study, 569 tick pools (1,523 ticks) collected from animals on Corsica were analysed to
449 investigate the presence of 27 bacteria and 12 parasites by a high throughput real-time microfluidic
450 PCR system. The CCHF virus and *Theileria* spp. were specifically investigated in their respective
451 potential vectors. DNA of eleven pathogens from six genera was identified in Corsican ticks and
452 among them seven were reported for first time in Corsican ticks: *B. miyamotoi*, *B. afzelii*, *R. helvetica*,
453 *A. phagocytophilum*, *Bar. henselae*, *Ba. bigemina* and *Ba. ovis*. These results also confirmed the
454 presence of four other TBPs in Corsica: *R. aeschlimannii*, *R. slovaca*, *A. marginale*, and *T. equi*. Many
455 of the pathogens found in this survey, as well as among those found by others, are mostly
456 asymptomatic or benign for animals, highlighting domestic and wild Corsican animals are probably an
457 important epidemiological reservoir increasing the human exposure to these zoonotic pathogens.
458 These findings highlight the importance to look deeper into TBPs epidemiological situation in Corsica
459 in the near future.

460

461 **ETHIC STATEMENT**

462 The authors obtained the agreement of all the owners to collect ticks from their domestic animals. The
463 cattle inspected were slaughtered for the human consumption and the wild boars collected were
464 legally hunted during the hunting season. The collected deer, mouflons and birds were captured by
465 PNRC and ONCF and they were all released. This study was approved by the veterinary Institutes
466 (DDCSPP of Corse-du-Sud and Haute-Corse) and the French Ministry of Agriculture, General
467 Directorate for Food (DGAl).

468 **ACKNOWLEDGEMENTS**

469 The authors are grateful to the staff of the slaughterhouse of Ponte-Leccia, Cuttoli and Porto-Vecchio
470 for their help in collecting ticks from cattle and to ONCFS, PNRC, hunters (especially Oscar
471 Maestrini, INRA, Corte, France) and practicing veterinarians for collecting ticks respectively from
472 mouflons, deer (and birds), wild boars and domestic carnivores.

473 **CONFLICT OF INTEREST**

474 The authors declare no conflict of interest.

475 **REFERENCES**

- 476 Alberti, A., Addis, M. F., Sparagano, O., Zobba, R., Chessa, B., Cubeddu, T., ... & Pittau, M. (2005).
477 *Anaplasma phagocytophilum*, Sardinia, Italy. *Emerging infectious diseases*, *11*(8), 1322.
- 478 Álvarez-Fernández, A., Breitschwerdt, E. B., & Solano-Gallego, L. (2018). Bartonella infections in
479 cats and dogs including zoonotic aspects. *Parasites & vectors*, *11*(1), 624.
- 480 Aubry, P., & Geale, D. W. (2011). A review of bovine anaplasmosis. *Transboundary and emerging*
481 *diseases*, *58*(1), 1-30.
- 482 Beati, L., Meskini, M., Thiers, B., & Raoult, D. (1997). Rickettsia aeschlimannii sp. nov., a new
483 spotted fever group rickettsia associated with Hyalomma marginatum ticks. *International Journal of*
484 *Systematic and Evolutionary Microbiology*, *47*(2), 548-554.
- 485 Beati, L., Péter, O., Burgdorfer, W., Aeschlimann, A., & Raoult, D. (1993). Confirmation that
486 Rickettsia helvetica sp. nov. is a distinct species of the spotted fever group of rickettsiae. *International*
487 *Journal of Systematic and Evolutionary Microbiology*, *43*(3), 521-526.
- 488 Beninati, T., Lo, N., Noda, H., Esposito, F., Rizzoli, A., Favia, G., & Genchi, C. (2002). First
489 detection of spotted fever group rickettsiae in Ixodes ricinus from Italy. *Emerging infectious diseases*,
490 *8*(9), 983.

- 491 Benredjem, W., Leulmi, H., Bitam, I., Raoult, D., & Parola, P. (2014). *Borrelia garinii* and *Rickettsia*
492 *monacensis* in *Ixodes ricinus* ticks, Algeria. *Emerging infectious diseases*, 20(10), 1776.
- 493 Bente, D.A., Forrester, N.L., Watts, D.M., McAuley, A.J., Whitehouse, C.A., Bray M. (2013)
494 Crimean-Congo hemorrhagic fever: history, epidemiology, pathogenesis, clinical syndrome and
495 genetic diversity. *Antiviral Research*, 100: 159–89.
- 496 Cabezas-Cruz, A., Gallois, M., Fontugne, M., Allain, E., Denoual, M., Moutailler, S., ... & Agoulon,
497 A. (2019). Epidemiology and genetic diversity of *Anaplasma ovis* in goats in Corsica, France.
498 *Parasites & Vectors*, 12(1), 3.
- 499 Chisu, V., Foxi, C., Mannu, R., Satta, G., & Masala, G. (2018). A five-year survey of tick species and
500 identification of tick-borne bacteria in Sardinia, Italy. *Ticks and tick-borne diseases*, 9(3), 678-681.
- 501 Chisu, V., Leulmi, H., Masala, G., Piredda, M., Foxi, C., & Parola, P. (2017). Detection of *Rickettsia*
502 *hoogstraalii*, *Rickettsia helvetica*, *Rickettsia massiliae*, *Rickettsia slovaca* and *Rickettsia aeschlimannii*
503 in ticks from Sardinia, Italy. *Ticks and tick-borne diseases*, 8(3), 347-352.
- 504 Cicculi, V., de Lamballerie, X., Charrel, R., & Charrel, R. (2019a). First molecular detection of
505 *Rickettsia africae* in a tropical bont tick, *Amblyomma variegatum*, collected in Corsica, France. *Exp*
506 *Appl Acarol*.
- 507 Cicculi, V., Masse, S., Capai, L., de Lamballerie, X., Charrel, R., Falchi, A. (2019b). First detection of
508 *Ehrlichia minasensis* in *Hyalomma marginatum* ticks collected from cattle in Corsica, France. *Vet*
509 *Med Sci*.
- 510 Cicculi, V., Capai, L., Quilichini, Y., Masse, S., Fernández-Alvarez, A., Minodier, L., ... & Falchi, A.
511 (2019c). Molecular investigation of tick-borne pathogens in ixodid ticks infesting domestic animals
512 (cattle and sheep) and small rodents (black rats) of Corsica, France. *Ticks and tick-borne diseases*, 10,
513 606-613.

- 514 Cosson, J. F., Michelet, L., Chotte, J., Le Naour, E., Cote, M., Devillers, E., ... & Moutailler, S.
515 (2014). Genetic characterization of the human relapsing fever spirochete *Borrelia miyamotoi* in
516 vectors and animal reservoirs of Lyme disease spirochetes in France. *Parasites & vectors*, 7(1), 233.
- 517 Dahmani, M., Davoust, B., Benterki, M. S., Fenollar, F., Raoult, D., & Mediannikov, O. (2015).
518 Development of a new PCR-based assay to detect Anaplasmataceae and the first report of *Anaplasma*
519 *phagocytophilum* and *Anaplasma platys* in cattle from Algeria. *Comparative immunology,*
520 *microbiology and infectious diseases*, 39, 39-45.
- 521 Dahmani, M., Davoust, B., Rousseau, F., Raoult, D., Fenollar, F., & Mediannikov, O. (2017). Natural
522 *Anaplasmataceae* infection in *Rhipicephalus bursa* ticks collected from sheep in the French Basque
523 Country. *Ticks and tick-borne diseases*, 8(1), 18-24.
- 524 Dahmani, M., Davoust, B., Tahir, D., Raoult, D., Fenollar, F., & Mediannikov, O. (2017). Molecular
525 investigation and phylogeny of *Anaplasmataceae* species infecting domestic animals and ticks in
526 Corsica, France. *Parasites & Vectors*, 10(1), 302.
- 527 Davoust, B., Mediannikov, O., Marié, J. L., Socolovschi, C., Parola, P., & Raoult, D. (2010). Les
528 animaux vertébrés sont-ils réservoirs de *Rickettsia*? Bull. Acad. Vét. France, Tome 163 - N°4/5
- 529 de Sousa, R., Pereira, B. I., Nazareth, C., Cabral, S., Ventura, C., Crespo, P., ... & da Cunha, S.
530 (2013). *Rickettsia slovaca* infection in humans, Portugal. *Emerging infectious diseases*, 19(10), 1627.
- 531 Dobec, M., Golubic, D., Punda-Polic, V., Kaeppli, F., & Sievers, M. (2009). *Rickettsia helvetica* in
532 *Dermacentor reticulatus* ticks. *Emerging infectious diseases*, 15(1), 98-100.
- 533 Dreshaj, S., Ahmeti, S., Ramadani, N., Dreshaj, G., Humolli, I., & Dedushaj, I. (2016). Current
534 situation of Crimean-Congo hemorrhagic fever in Southeastern Europe and neighboring countries: a
535 public health risk for the European Union? *Travel medicine and infectious disease*, 14(2), 81-91.

- 536 Dwużnik D., Mierzejewska EJ., Drabik P., Kloch A., Alzarraf M., Behnke JM., Bajer A. (2019). The
537 role of juvenile *Dermacentor reticulatus* ticks as vectors of microorganisms and the problem of “meal
538 contamination”. *Experimental and Applied Acarology*. <https://doi.org/10.1007/s10493-019-00380-6>.
- 539 ECDC (2013). Epidemiological situation of rickettsioses in EU/EFTA countries. Technical report.
540 www.ecdc.europa.eu.
- 541 Edouard, S., Koebel, C., Goehringer, F., Socolovschi, C., Jaulhac, B., Raoult, D., & Brouqui, P.
542 (2012). Emergence of human granulocytic anaplasmosis in France. *Ticks and tick-borne diseases*, 3(5-
543 6), 403-405.
- 544 Estrada-Peña, A., Gray, J. S., Kahl, O., Lane, R. S., & Nijhoff, A. M. (2013). Research on the ecology
545 of ticks and tick-borne pathogens—methodological principles and caveats. *Frontiers in cellular and*
546 *infection microbiology*, 3, 29.
- 547 Fernández-Soto, P., Pérez-Sánchez, R., Encinas-Grandes, A., & Sanz, R. Á. (2006). Rickettsia slovacica
548 in Dermacentor ticks found on humans in Spain. *European Journal of Clinical Microbiology and*
549 *Infectious Diseases*, 25(2), 129.
- 550 Fernández-Soto, P., Pérez-Sánchez, R., Encinas-Grandes, A., & Sanz, R. Á. (2004). Detection and
551 identification of Rickettsia helvetica and Rickettsia sp. IRS3/IRS4 in Ixodes ricinus ticks found on
552 humans in Spain. *European Journal of Clinical Microbiology & Infectious Diseases*, 23(8), 648-649.
- 553 Fukunaga, M., Takahashi, Y., Tsuruta, Y., Matsushita, O., Ralph, D., McClelland, M., & Nakao, M.
554 (1995). Genetic and phenotypic analysis of Borrelia miyamotoi sp. nov., isolated from the ixodid tick
555 Ixodes persulcatus, the vector for Lyme disease in Japan. *International Journal of Systematic and*
556 *Evolutionary Microbiology*, 45(4), 804-810.
- 557 Germanakis, A., Chochlakis, D., Angelakis, E., Tselentis, Y., & Psaroulaki, A. (2013). Rickettsia
558 aeschlimannii infection in a man, Greece. *Emerging infectious diseases*, 19(7), 1176-7.

- 559 Gondard, M., Michelet, L., Nisavanh, A., Devillers, E., Delannoy, S., Fach, P., ... & Moutailler, S.
560 (2018). Prevalence of tick-borne viruses in *Ixodes ricinus* assessed by high-throughput real-time PCR.
561 *Pathogens and disease*, 76(8).
- 562 Gondard, M., Delannoy, S., Pinarello, V., Aprelon, R., Devillers, E., Galon, C., ... & Moutailler, S.
563 (2019). Upscaling surveillance of tick-borne pathogens in the French Caribbean islands. *bioRxiv*,
564 532457.
- 565 Grech-Angelini, S., Stachurski, F., Lancelot, R., Boissier, J., Allienne, J. F., Gharbi, M., & Uilenberg,
566 G. (2016a). First report of the tick *Hyalomma scupense* (natural vector of bovine tropical theileriosis)
567 on the French Mediterranean island of Corsica. *Veterinary parasitology*, 216, 33-37.
- 568 Grech-Angelini, S., Stachurski, F., Lancelot, R., Boissier, J., Allienne, J. F., Marco, S., ... &
569 Uilenberg, G. (2016b). Ticks (Acari: Ixodidae) infesting cattle and some other domestic and wild
570 hosts on the French Mediterranean island of Corsica. *Parasites & Vectors*, 9(1), 582.
- 571 Habálek, Z., & Halouzka, J. (1997). Distribution of *Borrelia burgdorferi* sensu lato genomic groups in
572 Europe, a review. *European journal of epidemiology*, 13(8), 951-957.
- 573 Hoffman, T., Lindeborg, M., Barboutis, C., Erciyas-Yavuz, K., Evander, M., Fransson, T., ...
574 Salaneck, E. (2018). Alkhurma Hemorrhagic Fever Virus RNA in *Hyalomma rufipes* Ticks Infesting
575 Migratory Birds, Europe and Asia Minor. *Emerging infectious diseases*, 24(5), 879-882.
- 576 ICTTD (2000). Tick-borne infections of cattle on Corsica. Newsletter n° 14:13 (ISSN: 1385-3422).
- 577 Jongejan F, Uilenberg G. The global importance of ticks. *Parasitology*. 2004; 129 (Supplement): S3–
578 14.
- 579 Kernif, T., Messaoudene, D., Ouahioune, S., Parola, P., Raoult, D., & Bitam, I. (2012). Spotted fever
580 group rickettsiae identified in *Dermacentor marginatus* and *Ixodes ricinus* ticks in Algeria. *Ticks and*
581 *tick-borne diseases*, 3(5-6), 380-381.

- 582 Mancuso E., Toma L., Polci A., d'Alessio S.G., Di Luca M. *et al.* (2019). Crimean-Congo
583 haemorrhagic fever virus genome in tick from migratory bird, Italy. *Emerging infectious diseases*, 25:
584 1418-1420.
- 585 Marié, J. L., Davoust, B., Socolovschi, C., Mediannikov, O., Roqueplo, C., Beaucournu, J. C., ...,
586 Parola, P. (2012). Rickettsiae in arthropods collected from red foxes (*Vulpes vulpes*) in France.
587 *Comparative immunology, microbiology and infectious diseases*, 35(1), 59-62.
- 588 Matsumoto, K., Parola, P., Brouqui, P., Raoult, D. (2004). *Rickettsia aeschlimannii* in *Hyalomma*
589 ticks from Corsica. *European Journal of Clinical Microbiology and Infectious Diseases*, 23: 732-734.
- 590 Matsumoto, K., Ogawa, M., Brouqui, P., Raoult, D., & Parola, P. (2005). Transmission of *Rickettsia*
591 *massiliae* in the tick, *Rhipicephalus turanicus*. *Medical and veterinary entomology*, 19: 263-270.
- 592 Michelet, L., Delannoy, S., Devillers, E., Umhang, G., Aspan, A., Juremalm, M., ... Klitgaard, K.
593 (2014). High-throughput screening of tick-borne pathogens in Europe. *Frontiers in Cellular and*
594 *Infection Microbiology*, 4, 103.
- 595 Negrodo, A., de la Calle-Prieto, F., Palencia-Herrejón, E., Mora-Rillo, M., Astray-Mochales, J.,
596 Sánchez-Seco, M. P., ... & Keough-Delgado, E. (2017). Autochthonous Crimean–Congo Hemorrhagic
597 Fever in Spain. *New England Journal of Medicine*, 377(2), 154-161.
- 598 Nicholson WL, Sonenshine DE, Lane RS, Uilenberg G. 2009. Chapter 26. Ticks (Ixodida). In: Mullen
599 GR, Durden LA, editors. *Medical and veterinary entomology*. Second ed. Elsevier; Academic Press;
600 pp. 493–542.
- 601 Parola P, Raoult D. Ticks and tick-borne bacterial diseases in humans: an emerging infectious threat.
602 *Clin Infect Dis*. 2001; 32:897–928.

- 603 Parola, P., Beati, L., Cambon, M., & Raoult, D. (1998). First isolation of *Rickettsia helvetica* from
604 *Ixodes ricinus* ticks in France. *European Journal of Clinical Microbiology and Infectious Diseases*,
605 *17*(2), 95-100.
- 606 Parola, P., Paddock, C. D., Socolovschi, C., Labruna, M. B., Mediannikov, O., Kernif, T., ... &
607 Raoult, D. (2013). Update on tick-borne rickettsioses around the world: a geographic approach.
608 *Clinical microbiology reviews*, *26*(4), 657-702.
- 609 Parola, P., Rovero, C., Rolain, J. M., Brouqui, P., Davoust, B., & Raoult, D. (2009). *Rickettsia slovaca*
610 and *R. raoultii* in tick-borne rickettsioses. *Emerging infectious diseases*, *15*(7), 1105.
- 611 Pérez-Eid, C. (2007). *Les tiques: identification, biologie, importance médicale et vétérinaire*.
612 Lavoisier.
- 613 Platonov, A. E., Karan, L. S., Kolyasnikova, N. M., Makhneva, N. A., Toporkova, M. G., Maleev, V.
614 V., ... & Krause, P. J. (2011). Humans infected with relapsing fever spirochete *Borrelia miyamotoi*,
615 Russia. *Emerging infectious diseases*, *17*(10), 1816.
- 616 Portillo, A., Santibáñez, S., García-Álvarez, L., Palomar, A. M., & Oteo, J. A. (2015). Rickettsioses in
617 Europe. *Microbes and infection*, *17*(11-12), 834-838.
- 618 Raoult, D., Fournier, P. E., Abboud, P., & Caron, F. (2002). First documented human *Rickettsia*
619 *aeschlimanii* infection. *Emerging infectious diseases*, *8*(7), 748.
- 620 Raoult, D., Nicolas, D., De Micco, P. H., Gallais, H., & Casanova, P. (1985). Aspects
621 épidémiologiques de la fièvre boutonneuse méditerranéenne en Corse du Sud. *Bulletin de la Société*
622 *de Pathologie Exotique et Filiales*, *78*, 446-451.
- 623 Ravagnan, S., Tomassone, L., Montarsi, F., Krawczyk, A. I., Mastrorilli, E., Sprong, H., ... & Capelli,
624 G. (2018). First detection of *Borrelia miyamotoi* in *Ixodes ricinus* ticks from northern Italy. *Parasites*
625 *& vectors*, *11*(1), 130.

- 626 Ruscio, M., & Cinco, M. (2003). Human granulocytic ehrlichiosis in Italy. *Annals of the New York*
627 *Academy of Sciences*, 990(1), 350-352.
- 628 Selmi, M., Ballardini, M., Salvato, L., & Ricci, E. (2017). Rickettsia spp. in Dermacentor marginatus
629 ticks: analysis of the host-vector-pathogen interactions in a northern Mediterranean area. *Experimental*
630 *and Applied Acarology*, 72(1), 79-91.
- 631 Sfar, N., M'ghirbi, Y., Letaief, A., Parola, P., Bouattour, A., & Raoult, D. (2008). First report of
632 Rickettsia monacensis and Rickettsia helvetica from Tunisia. *Annals of Tropical Medicine &*
633 *Parasitology*, 102(6), 561-564.
- 634 Siński, E., Welc-Falęciak, R., & Zajkowska, J. (2016). Borrelia miyamotoi: A human tick-borne
635 relapsing fever spirochete in Europe and its potential impact on public health. *Advances in medical*
636 *sciences*, 61(2), 255-260.
- 637 Torina, A., Galindo, R. C., Vicente, J., Di Marco, V., Russo, M., Aronica, V., ... & Kocan, K. M.
638 (2010). Characterization of Anaplasma phagocytophilum and A. ovis infection in a naturally infected
639 sheep flock with poor health condition. *Tropical animal health and production*, 42(7), 1327-1331.
- 640 Uilenberg, G. (2006). Babesia—a historical overview. *Veterinary Parasitology*, 138(1-2), 3-10.
- 641 Uilenberg G., Gray J., Kahl O. (2018) Research on Piroplasmidora and other tick-borne agents: Are
642 we going the right way? *Ticks and Tick-Borne Diseases* 9: 860-863.
- 643 Vandenesch, A., Turbelin, C., Couturier, E., Arena, C., Jaulhac, B., Ferquel, E., Choumet, V.,
644 Saugeon, C., Coffinieres, E., Blanchon, T., Vaillant, V., Hanslik, T. (2014). Incidence and
645 hospitalisation rates of Lyme borreliosis, France, 2004 to 2012. *Euro Surveill.* 19 20883 [pii].
- 646 Zobba, R., Anfossi, A. G., Parpaglia, M. L. P., Dore, G. M., Chessa, B., Spezzigu, A., ... & Alberti, A.
647 (2014). Molecular investigation and phylogeny of Anaplasma spp. in Mediterranean ruminants reveal

648 the presence of neutrophil-tropic strains closely related to *A. platys*. *Appl. Environ. Microbiol.*, 80(1),
649 271-280.

650

651 **TABLES**

652

653 **Table 1. List of pathogens (and endosymbionts) detectable by the real-time PCR chip (Biomark™)**

Genus	Number of target pathogens/endosymbionts	Species
<i>Anaplasma</i>	6	<i>A. marginale</i> , <i>A. centrale</i> , <i>A. phagocytophilum</i> , <i>A. ovis</i> , <i>A. bovis</i> , <i>A. platys</i>
<i>Borrelia</i>	8	<i>B. burgdorferi</i> sensu stricto, <i>B. garinii</i> , <i>B. afzelii</i> , <i>B. valaisiana</i> , <i>B. lusitaniae</i> , <i>B. spielmanii</i> , <i>B. bissetti</i> , <i>B. miyamotoi</i>
<i>Bartonella</i>	2	<i>Bar. henselae</i> , <i>Bar. quintana</i>
<i>Ehrlichia</i>	3	<i>E. canis</i> , <i>E. chaffeensis</i> , <i>E. ruminantium</i>
<i>Francisella</i>	2	<i>F. tularensis</i> , <i>Francisella</i> -like endosymbionts
<i>Neoehrlichia</i>	1	<i>Candidatus Neoehrlichia mikurensis</i>
<i>Rickettsia</i>	5	<i>R. conorii</i> , <i>R. slovaca</i> , <i>R. massiliae</i> , <i>R. helvetica</i> , <i>R. aeschlimannii</i>
<i>Coxiella</i>	2	<i>Coxiella burnetii</i> , <i>Coxiella</i> -like organisms
<i>Babesia</i>	10	<i>Ba. divergens</i> , <i>Ba. microti</i> , <i>Ba. canis</i> , <i>Ba. vogeli</i> , <i>Ba. bovis</i> , <i>Ba. caballi</i> , <i>Ba. venatorum</i> , <i>Ba. bigemina</i> , <i>Ba. major</i> , <i>Ba. ovis</i>
<i>Theileria</i>	2	<i>T. equi</i> , <i>T. annulata</i>

654

655

656 **Table 2. Infection rates of individual ticks or pools of ticks collected in Corsica**

Tick species	Number of pools	Stade or sex of ticks			Total ticks analysed	Infection rate %
		Female	Male	Nymph		
<i>Rh. bursa</i>	108	42	51	15	518	43
<i>Hy. marginatum</i>	89	24	65	0	362	100
<i>Hy. scupense</i> *	135	60	56	19	135	8
<i>I. ricinus</i> *	115	79	36	0	115	70
<i>Rh. sanguineus</i> s.l.	41	22	16	3	150	24
<i>D. marginatus</i>	38	16	22	0	156	66
<i>Ha. punctata</i>	30	18	12	0	74	30
<i>Rh. (Bo.) annulatus</i> *	11	6	3	2	11	36
<i>Ha. sulcata</i> *	2	2	0	0	2	0

657 * Ticks individually analysed, the infection rate is the real tick infection rate

658

Tick species (pools; ticks analysed)						
	<i>Borrelia</i>	<i>Rickettsia</i>	<i>Anaplasma</i>	<i>Babesia</i>	<i>Bartonella</i>	<i>Theileria</i>
<i>Rh. bursa</i> (108;518)	0	27	15	12	5	1
<i>Hy. marginatum</i> (89;362)	0	100	7	0	1	0
<i>Hy. scupense</i> (135;135)*	0	6	2	0	0	0
<i>I. ricinus</i> (115; 115)*	5	6	64	0	2	0
<i>Ha. punctata</i> (30;74)	13	0	13	0	3	0
<i>Ha. sulcata</i> (2;2)*	0	0	0	0	0	0
<i>Rh. sanguineus</i> s.l. (41;150)	0	17	12	0	2	0
<i>Rh. (B.) annulatus</i> (11;11)*	0	0	18	0	9	9
<i>D. marginatus</i> (38;156)	0	66	3	0	3	0

659 **Table 3. Proportion (%) of infected pools (or ticks)**

660 *Ticks individually analysed.

661

662 **Table 4. Pathogen DNA detected in ticks collected from domestic animals**

Host	Analysed pools (ticks)	Tick species pools (or individual ticks*)	% infected pools (or ticks*)	Pathogens identified (% of infected pools or ticks*)
Cattle	404 (894)	<i>Rh. bursa</i> (54), <i>Hy. marginatum</i> (67), <i>Hy. scupense</i> (135), <i>I. ricinus</i> (98), <i>Rh. sanguineus</i> s.l. (9), <i>D. marginatus</i> (4), <i>Ha. punctata</i> (26), <i>Rh. (B.) annulatus</i> (11)	47	<i>Borrelia</i> spp. (3%), <i>Rickettsia</i> spp. (25%), <i>Anaplasma</i> spp. (22%), <i>Babesia</i> spp. (3%), <i>Bartonella</i> spp. (2%), <i>Theileria</i> spp. (0.3%)
Goat	19 (91)	<i>Rh. bursa</i> (18), <i>Ha. punctata</i> (1)	47	<i>Rickettsia</i> spp. (37%), <i>Anaplasma</i> spp. (2%), <i>Babesia</i> spp. (5%), <i>Bartonella</i> spp. (11%)
Sheep	12 (45)	<i>Rh. bursa</i> (9), <i>Hy. marginatum</i> (2), <i>Rh. sanguineus</i> s.l. (1)	58	<i>Rickettsia</i> spp. (58%)
Dogs	22 (86)	<i>Hy. marginatum</i> (1), <i>Rh. sanguineus</i> s.l. (21)	14	<i>Rickettsia</i> spp. (9%), <i>Anaplasma</i> spp. (4%),
Horses	24 (106)	<i>Rh. bursa</i> (6), <i>Hy. marginatum</i> (16), <i>Rh. sanguineus</i> s.l. (1), <i>D. marginatus</i> (1),	83	<i>Rickettsia</i> spp. (79%), <i>Bartonella</i> spp. (4%), <i>Theileria</i> spp. (4%)
Cats	4 (6)	<i>I. ricinus</i> (1), <i>Rh. sanguineus</i> s.l. (3)	25	<i>Anaplasma</i> spp. (25%)

663

664

665 **Table 5. Pathogen DNA detected in ticks collected from wild animals**

Host	Analysed pools	Total ticks	Tick species (pools or individual ticks*)	% infected pools (or ticks*)	Pathogens (% of infected pools or ticks*)
Wild boars	41	177	<i>Rh. bursa</i> (2), <i>Hy. marginatum</i> (3), <i>I. ricinus</i> (1), <i>Rh. sanguineus</i> s.l. (3), <i>D. marginatus</i> (32),	66	<i>Rickettsia</i> spp. (66%), <i>Anaplasma</i> spp. (2%), <i>Bartonella</i> spp. (2%)
Mouflons	26	98	<i>Rh. bursa</i> (19), <i>Rh. sanguineus</i> s.l. (1), <i>D. marginatus</i> (1), <i>Ha. punctata</i> (3), <i>Ha. sulcata</i> (2)	12	<i>Rickettsia</i> spp. (4%), <i>Anaplasma</i> spp. (4%), <i>Babesia</i> spp. (8%)
Deer	13	13	<i>I. ricinus</i> (13*)	92	<i>Anaplasma</i> spp. (92%*)
Hedgehogs	2	5	<i>Rh. sanguineus</i> s.l. (2)	50	<i>Rickettsia</i> spp. (50%)
Birds	2	2	<i>I. ricinus</i> (2*)	100	<i>Anaplasma</i> spp. (100%*)

666

667

668





669 **FIGURE LEGENDS**

670 **Figure 1. Distribution of the main DNA pathogens found in Corsican ticks. a) *Rickettsia* spp., b)**

671 ***Anaplasma* spp., c) *Borrelia* spp., d) *Babesia* spp.**

672

673



Area with analysed ticks

Pathogen not found

Presence of the pathogen

Municipalities not investigated

CORSE-DU-SUD

HAUTE-CORSE