

Author Correction: Intermediate degrees of synergistic pleiotropy drive adaptive evolution in ecological time

Léa Frachon, Cyril Libourel, Romain Villoutreix, Sébastien Carrère, Cédric Glorieux, Carine Huard-Chauveau, Miguel Navascués, Laurène Gay, Renaud Vitalis, Etienne Baron, Laurent Amsellem, Olivier Bouchez, Marie Vidal, Valérie Le Corre, Dominique Roby, Joy Bergelson and Fabrice Roux

Correction to: *Nature Ecology & Evolution* **1**, 1551–1561 (2017); published online 4 September 2017.

In the version of this Article previously published, there was a typographical error ('4' instead of '2') in the equations relating F_{ST} and effective population size (N_e) in the Methods section 'Genome-wide scan for selection based on temporal differentiation'. The correct equations are given below.

$$F_{ST} \approx \frac{T}{T+2} = \frac{\tau}{\tau + 2N_e}$$

$$\widehat{N}_e = \frac{\tau(1 - \widehat{F}_{ST})}{2\widehat{F}_{ST}}$$

All the calculations and results presented in the Article were obtained using these correct equations and need no amendment. The conclusions of our study are therefore in no way affected.

Published online: 5 December 2017

<https://doi.org/10.1038/s41559-017-0405-2>