



# Genome-wide identification and characterization of aquaporin gene family in beta vulgaris

Weilong Kong, Shaozong Yang, Yulu Wang, Mohammed Bendahmane, Xiaopeng Fu

## ► To cite this version:

Weilong Kong, Shaozong Yang, Yulu Wang, Mohammed Bendahmane, Xiaopeng Fu. Genome-wide identification and characterization of aquaporin gene family in beta vulgaris. PeerJ, 2017, 5, 10.7717/peerj.3747 . hal-02622429

**HAL Id: hal-02622429**

**<https://hal.inrae.fr/hal-02622429>**

Submitted on 26 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



# Genome-wide identification and characterization of aquaporin gene family in *Beta vulgaris*

Weilong Kong<sup>1</sup>, Shaozong Yang<sup>1</sup>, Yulu Wang<sup>1</sup>, Mohammed Bendahmane<sup>2</sup> and Xiaopeng Fu<sup>1</sup>

<sup>1</sup> College of Horticulture and Forestry Sciences, Huazhong Agricultural University, Key Laboratory of Horticultural Plant Biology, Ministry of Education, Wuhan, Hubei, China

<sup>2</sup> INRA-CNRS-Lyon1-ENS, Laboratoire Reproduction et Développement des Plantes, Ecole Normale Supérieure Lyon, France

## ABSTRACT

Aquaporins (AQPs) are essential channel proteins that execute multi-functions throughout plant growth and development, including water transport, uncharged solutes uptake, stress response, and so on. Here, we report the first genome-wide identification and characterization AQP (*BvAQP*) genes in sugar beet (*Beta vulgaris*), an important crop widely cultivated for feed, for sugar production and for bioethanol production. Twenty-eight sugar beet AQPs (*BvAQPs*) were identified and assigned into five subfamilies based on phylogenetic analyses: seven of plasma membrane (PIPs), eight of tonoplast (TIPs), nine of NOD26-like (NIPs), three of small basic (SIPs), and one of x-intrinsic proteins (XIPs). *BvAQP* genes unevenly mapped on all chromosomes, except on chromosome 4. Gene structure and motifs analyses revealed that *BvAQP* have conserved exon-intron organization and that they exhibit conserved motifs within each subfamily. Prediction of *BvAQPs* functions, based on key protein domains conservation, showed a remarkable difference in substrate specificity among the five subfamilies. Analyses of *BvAQPs* expression, by mean of RNA-seq, in different plant organs and in response to various abiotic stresses revealed that they were ubiquitously expressed and that their expression was induced by heat and salt stresses. These results provide a reference base to address further the function of sugar beet aquaporins and to explore future applications for plants growth and development improvements as well as in response to environmental stresses.

**Subjects** Agricultural Science, Bioinformatics, Genomics, Plant Science

**Keywords** Gene structure, Expression profile, *Beta vulgaris*, Aquaporins, Abiotic stress

## INTRODUCTION

AQPs are known to facilitate water transport and other small nutrients through cell membranes (Maurel *et al.*, 2009; Maurel *et al.*, 2008). Since the discovery of the first aquaporin (AQP1) in mammals, AQPs were identified in many microorganisms, plants and animals (Gomes *et al.*, 2009).

Plant aquaporin families are complex, and are composed of a large number of genes. For example, there are 35 AQPs in *Arabidopsis thaliana*, 31 in *Zea mays*, 34 in *Oryza sativa*, 55 in *Populus trichocarpa* and 66 AQPs in *Glycine max* (Johanson & Kjellbom, 2001; Nguyen,

Submitted 13 April 2017  
Accepted 8 August 2017  
Published 19 September 2017

Corresponding author  
Xiaopeng Fu,  
fuxiaopeng@mail.hzau.edu.cn

Academic editor  
David Hyten

Additional Information and  
Declarations can be found on  
page 17

DOI 10.7717/peerj.3747

© Copyright  
2017 Kong et al.

Distributed under  
Creative Commons CC-BY 4.0

OPEN ACCESS

[Moon & Jung, 2013](#); [Chaumont et al., 2001](#); [Gupta & R, 2009](#); [Zhang et al., 2013](#)). In plants AQPs play major roles in water and solute transport, in maintaining water homeostasis and in the response to environment stresses. AQP's roles in glycerol, urea, boric acid, silicic acid,  $H_2O_2$ ,  $NH_3$  and  $CO_2$  transport through cell membranes were reported to be important for cytoplasm homeostasis, seed germination, embolism recovery, petal and leaf movement, guard cells closure, fruit ripening and maintenance of cell turgor under various stresses ([Fitzpatrick & Reid, 2009](#); [Maurel et al., 2009](#); [Maurel et al., 2008](#); [Mitani-Ueno et al., 2011](#); [Heinen & Chaumont, 2009](#); [Maurel et al., 2008](#); [Prado & Maurel, 2013](#); [Uehlein & Kaldenhoff, 2008](#); [Wudick & Maurel, 2009](#)).

To date, AQPs are recognized into seven subfamilies: PIPs, TIPs, NIPs, SIPs, XIPs, GIPs, and HIPs ([Anderberg, Kjellbom & Johanson, 2012](#); [Danielson & Johanson, 2008](#)). Green plants usually contain four subfamilies: PIPs, TIPs, NIPs, and SIPs. However, members of XIPs subfamily were also found in some dicots, such as *Solanum lycopersicum* ([Venkatesh, Yu & Park, 2013](#)), *Populus trichocarpa* ([Gupta & Sankararamakrishnan, 2009](#)) and *Glycine max* ([Cheng et al., 2013](#)), but were absent in Brassicaceae and monocots ([Danielson & Johanson, 2008](#)). GIPs and HIPs subfamilies were reported in moss (*Physcomitrella patens*) and fern (*Selaginella moellendorffii*).

AQPs are highly conserved in all living organisms, consisting of six transmembrane domains (TM1–TM6) connected by five loops (LA–LE). NPA motifs, the ar/R selectivity and Froger's position are critical for AQP's functions. Two NPA motifs (Asn-Pro-Ala) are located on LB and LE, forming a central aqueous pore in the middle of the lipid bilayer involved in proton exclusion and substrate selectivity ([Bansal & Sankararamakrishnan, 2007](#)). The ar/R selectivity is formed by four residues from TM2 (H2), TM5 (H5), LE (LE1 and LE2) and acts as a size-exclusion barrier ([Hove & Bhawe, 2011](#); [Mitani-Ueno et al., 2011](#)). Froger's position consists of five conserved residues (P1–P5) and discriminations between AQP-type and GIP-type AQPs ([Froger et al., 1998](#)). Nine specificity-determining positions for non-aqua substrates (i.e., urea, boric acid, silicic acid, ammonia, carbon dioxide and hydrogen peroxide ( $H_2O_2$ )) were also proposed based on a comprehensive analysis of functionally characterized AQPs ([Hove & Bhawe, 2011](#)).

Sugar beet belongs to Caryophyllales, which lays on basal taxa of core dicots. It is an important crop in temperate climates region and provides nearly 30% of the world's annual sugar production ([Dohm et al., 2014](#)). It is also used as a source for animal feed and for bioethanol production.

To date, little is known about AQPs in sugar beet and in Caryophyllales. So far, only information on 26 AQPs, grouped in five subfamilies (eight PIPs, 11 TIPs, four NIPs, two SIPs and one XIP), was reported in carnation ([Morita et al., 2017](#)). More information on AQPs in Caryophyllales plants is therefore required to help understand their function and evolution.

Here we used the available high-quality genome sequence ([Dohm et al., 2014](#)) and RNA-seq datasets ([Minoche et al., 2015](#)) of sugar beet to identify and characterize the expression of BvAQPs. We report the distribution of BvAQPs on chromosomes, phylogeny analysis, gene structure, subcellular location, conserved residues and conserved elements.

We propose putative functions of sugar beet AQPs based on their detailed genes expression patterns analysis by means of RNA-seq.

## METHODS

### Identification and distribution of AQP genes in sugar beet

BvAQPs were identified by HMM (Hidden Markov Model) and BLAST homology search. The sugar beet predicted proteome was collected using the sugar beet genome RefBeet-1.2 (<http://bvseq.molgen.mpg.de/Genome/Download/index.shtml>). The Hidden Markov model (HMM) of the MIP domain (PF00230) was downloaded from the Sanger database (<http://pfam.xfam.org/family/PF00230>). PF00230 was then used to query the predicted Sugar beet proteome using HMMER 3.0 software (<http://hmmmer.org/>). 35 Arabidopsis AtAQPs were download from TAIR Database (<https://www.arabidopsis.org/browse/genefamily/Aquaporins.jsp>) and then used to search BvAQPs with BLASTp tool using NCBI sugar beet genome (<https://www.ncbi.nlm.nih.gov/genome/?term=Beta+vulgaris>) and sugar beet genome (<http://www.genomforschung.uni-bielefeld.de/en/projects/annobeet>) (Baranwal, Negi & Khurana, 2016) with cut-off  $E$ -value of  $e^{-5}$ . All no-redundant gene sequences were analyzed by SMART (<http://smart.embl-heidelberg.de/>) and Pfam (<http://pfam.xfam.org/search/sequence>). Sequences encoding complete MIP domain and two NPA motifs were considered as putative AQP genes.

Additionally, the molecular weight (MW) and isoelectric point (PI) of BvAQPs were calculated by ExPASy (<http://www.expasy.org/>); transmembrane helical domains (TMHs) were assessed by TMHMM Sever v.2.0 (<http://www.cbs.dtu.dk/services/TMHMM/>); the subcellular localization of BvAQPs were predicted using Plant-mPLOC (<http://www.csbio.sjtu.edu.cn/bioinf/plant-multi/>) and Wolf PSORT (<http://www.genscript.com/wolf-psort.html>). The position of the AQP genes on the sugar beet chromosomes were identified based on position information from the sugar beet genome database and the distribution graph of AQP genes was drawn by MapInspect software (<http://mapinspect.software.informer.com/>).

### Phylogenetic analyses and sequence alignments

BvAQPs were aligned with AtAQPs from Arabidopsis (<https://www.arabidopsis.org/>) DcAQPs from *Dianthus caryophyllus* (Morita et al., 2017; <http://carnation.kazusa.or.jp/blast.html>), by using Clustal W. Phylogenetic tree was built by MEGA6.0 (<http://www.megasoftware.net/history.php>) using the neighbor-joining (NJ) method, with 1,000 times bootstrap replicates. BvAQPs were named based on their sequence homology and phylogenetic analyses. Greek letters ( $\alpha$ ,  $\beta$ ) were used to denote the transcripts derived from the same gene.

BvAQPs and two *S. tuberosum* AQPs (Venkatesh, Yu & Park, 2013) were aligned by DNAMAN (<http://dnaman.software.informer.com/>) with default parameters. Two NPA motifs, ar/R selectivity filter and Froger's position were inferred from the multiple sequence alignment result from DNAMAN. Specificity-determining positions (SDP1-SDP9) from alignments with the structure resolved *Spinacia oleracea* PIP2;1 and functionally characterized AQPs as being collected by Hove and Bhavé (Hove & Bhavé, 2011).



**Table 1** Identification of *BvAQP* genes using sugar beet genome data.


Family	Gene	Gene ID	Gene code	Protein length (aa)	MW (kDa)	pI	TMHs	Plant-mPLoc	WoLF PSORT
PIP	BvPIP1;1	fpur	Bv1g004510_fpur.t1	289	31.1	8.84	5	plas	cyto
	BvPIP1;2	shpz	Bv1g004520_shpz.t1	285	30.67	9.05	6	plas	plas
	BvPIP1;3	gqok	Bv2g024120_gqok.t1	286	30.74	9.14	6	plas	plas
	BvPIP2;1	cqnr	Bv7g163390_cqnr.t1	284	30.32	8.31	6	plas	plas
	BvPIP2;2	ixem	Bv9g210030_ixem.t1	288	31.09	7.08	6	plas	plas
	BvPIP2;3	yige	Bv9g210020_yige.t1 (XP_010689549.1) <sup>a</sup>	274	29.64	8.97	6	plas	plas
	BvPIP2;4	reke	Bv9g216070_reke.t1	281	30.13	8.84	6	plas	plas
TIP	BvTIP1;1	kzkq	Bv7ug180930_kzkq.t1	254	26.06	5.38	6	vacu	plas
	BvTIP1;2	iuuk	Bv2g037380_iuuk.t1	252	26.3	5.92	6	vacu	vacu
	BvTIP1;3	ynzf	Bv7g176430_ynzf.t1	248	25.46	5.13	6	vacu	vacu
	BvTIP2;1	dkzm	Bv9g223310_dkzm.t1	247	25.26	5.6	6	vacu	plas
	BvTIP2;2	xunf	Bv5g104980_xunf.t1	249	25.09	4.7	7	vacu	vacu
	BvTIP3;1	dreg	Bv8g190600_dreg.t1	257	27.13	7.07	5	vacu	nucl/mito/vacu
	BvTIP4;1	yjno	Bv2g032200_yjno.t1	247	26.1	6.57	6	vacu	cyto/vacu
NIP	BvNIP5;1	gghp	Bv3ug068240_gghp.t1	255	26.54	8.47	6	plas	chlo
	BvNIP1;1	ughi	Bv8ug202570_ughi.t1	292	30.87	8.91	6	plas	plas
	BvNIP4;1	aejh	Bv2g027680_aejh.t1	273	29.3	8.87	6	plas	vacu
	BvNIP4;2	xash	Bv2g027660_xash.t1	299	32.62	8.27	6	plas	plas
	BvNIP5;1	gkiq	Bv6TE021760_gkiq.t1 (XP_010680949.1) <sup>a</sup>	261	27.61	8.96	6	plas	vacu
	BvNIP5;2	oani	Bv6g139140_oani.t1	298	30.9	8.73	6	plas	plas
	BvNIP6;1	hmzo	Bv9g225280_hmzo.t1	306	31.75	7.66	6	plas	plas
SIP	BvNIP6;2	zkgo	Bv5g108450_zkgo.t1 (XP_010677474.1) <sup>a</sup>	266	27.46	7.82	6	plas	plas
	BvNIP6;3	jecw	Bv5g108440_jecw.t1 (XP_010677699.1) <sup>a</sup>	327	35.08	9.03	6	plas	plas
	BvNIP7;1	kqew	Bv3ug070540_kqew.t1	289	30.72	7.13	7	plas	plas
SIP	BvSIP1;1 α	zywx	Bv2g035790_zywx.t1 (XP_010669579.1) <sup>a</sup>	250	26.46	9.46	5	plas	plas
	BvSIP1;1 β	fzwq	Bv2g035780_fzwq.t1	247	26.39	9.74	6	plas/vacu	vacu
	BvSIP2;1	qzqg	Bv3g064810_qzqg.t1 (XP_010673209.1) <sup>a</sup>	236	26.02	9.56	5	plas	vacu
XIP	BvXIP1;1	iwpe	Bv9g217040_iwpe.t1	312	34.13	8.38	7	plas	plas

**Notes.** Note1: Best possible cell localization prediction by the Plant-mPLoc and WoLF PSORT tool (Chlo: chloroplast; Cyto: cytosol; Cysk: cytoskeleton; E.R: endoplasmic reticulum; Extr: extracellular; Golg: Golgi apparatus; Lyso: lysosome; Mito: mitochondria; Nucl: nuclear; Pero: peroxisome; Plas: plasma membrane; Vacu: vacuolar membrane).  
<sup>a</sup>sequences were splicing errors in AnnoBeet, and were corrected by NCBI Beta vulgaris (ID 221)—Genome. These were verified by PCR amplification.

## Genes mapping on the sugar beet chromosomes, gene duplications, gene structure and alternative splicing

Twenty-eight *AQP* genes were unevenly mapped on eight chromosomes (Fig. 2). Seven *BvAQP* genes (25%) mapped on chromosome (Chr) 2, 6 *BvAQP* genes (21%) mapped on Chr9. Chr3, Chr5 and Chr7 each had three *BvAQP* genes (10%) and Chr1, Chr6 and Chr8







**Figure 1** Multiple alignments and phylogenetic analysis of *Bv*AQPs *A. thaliana* AtAQPs and *D. caryophyllus* DcAQPs. Multiple alignments were performed using the default parameter of Clustal W. Phylogenetic dendrogram was generated by MEGA 6 using neighbor-joining (NJ) method with 1,000 bootstrap replicates.

each had two *BvAQP* genes. No putative AQP was found on Chr4. Tandem duplication events were found on Chr1, Chr2, Chr5, Chr6 and Chr8 (i.e., *BvPIP1;1* and *BvPIP1;2*, *BvNIP4;1* and *BvNIP4;2*, *BvNIP6;2* and *BvNIP6;3*, *BvNIP5;1* and *BvNIP5;2*, *BvPIP2;2* and *BvPIP2;3*).

The exon-intron structures play crucial roles during plant evolution. The sugar beet AQPs exon-intron structures are showed in Fig. 3. Most PIP subfamily genes have four exons, while *BvPIP1;3* had 5 and *BvPIP2;4* had three. TIP subfamily genes had three exons; specifically, the third exon of *BvTIP1;1* had a noncoding exon and codes a long 5' translated region (5' UTR). NIP subfamily genes had five exons with the exception *BvNIP5;2*, *BvNIP6;2* and *BvNIP6;3* had four exons. Most SIP subfamily genes have three exons but *BvXIP1;1* only had two exons, which were similar to situations reported in common bean, tomato, potato, and so on. (Ariani & Gepts, 2015; Reuscher et al., 2013a;



**Figure 2** Distribution of *BvAQP* genes on the nine *B. vulgaris* chromosomes. Note: Black lines represent tandem gene duplications. Red lines represent the existence of different transcripts from single gene.




**Figure 3** *BvAQP* genes structure analysis.

Venkatesh, Yu & Park, 2013). These results suggested that *BvAQP* gene structure is globally conserved in sugar beet. The conserved exon-intron structure provided an additional proof to support the classification results (Fig. 1).

To seek further insights into the gene structure, the splicing pattern of sugar beet *BvAQP* pre-mRNA sequence were analyzed. The splicing analysis revealed that *BvSIP1;1 α* is the result of second exon skipping (Fig. 4). The amino acid sequence of *BvSIP1;1α* and *BvSIP1;1β* shows a 74.8% similarity.





**Figure 4** Alternative splicing of *SIP1;1* generate two different transcripts: *SIP1;1α* and *SIP1;1β*.

## Conserved residues and conserved motifs distribution

To further understand the possible physiological role and substrate specificity of BvAQPs, they were aligned and TMHs and conserved residues (NPA motifs, ar/R selectivity filter and Froger's position) were analyzed (Table 2). The data revealed that all BvAQPs had six characteristic TMHs for AQPs, required for transport function (Fig. S1). However, BvPIP1;1 showed partial loss of TM6. The resulting missing protein domains may cause dominant negative effect on protein functions, although this need to be confirmed. All PIPs and TIPs showed the dual typical NPA motifs, but some members of NIPs such as BvNIP5;1, BvNIP5;2 and BvNIP6;1 showed that the Alanine (A) in the third residue of the second NPA motif was replaced by a Valine (V). All SIPs and XIPs showed various third residues in the first NPA motif, in which Alanine (A) was replaced by Threonine (T) or by a Leucine (L). PIPs' ar/R selectivity filter and Froger's position showed an apparent family-specificity compared to NPA motifs, with ar/R filter configuration typical for water-transporting AQPs (F-H-T-R) and Q/M-S-A-F-W residues in Froger's position. TIPs contained the H-I-A/G-V/R residues in the ar/R selectivity filter and T-S/A-A-Y-W residues in Froger's position, but BvTIP5;1 showed that the N-V-G-Y in the ar/R selectivity filter was different from other TIPs and formed a single-gene clade with the TIPs. In NIPs, the ar/R selectivity filter and Froger's position had multiple types. BvNIP1;1, BvNIP4;1 and BvNIP4;2 showed W-V-A-R in the ar/R selectivity filter and F/L-S-A-Y-L/I in the Froger's position. BvNIP5;1 and BvNIP5;2 showed A-I-G/A-R in the ar/R selectivity filter and F-T-A-Y-M in the Froger's position. BvNIP6;1, BvNIP6;2 and BvNIP6;3 showed S-I-G/A-R in the ar/R selectivity filter and Y-T-A-Y-F/M/L in the Froger's position. BvNIP7;1 showed A-V-G-R in the ar/R selectivity filter and F-S-A-Y-F in the Froger's position. SIP1s and SIP2s showed distinct difference in the ar/R selectivity filter and the Froger's position. BvSIP1;1  $\alpha$  and BvSIP1;1  $\beta$  showed I/V-V-P-N ar/R selectivity filter and M-A-A-Y-W in the Froger's position, but BvSIP2;1 showed S-N-G-S ar/R selectivity filter and F-V-A-Y-W in the Froger's position. BvXIP1;1 showed V-S-A-R ar/R selectivity filter and F-C-A-F-W in the Froger's position.


**Table 2** Conserved dual NPA motifs, ar/R (H2, H5, LE1 and LE2), Froger's positions (P1–P5) analysis of BvAQPs.

Subfamily	Gene	NPA		Ar/R selectivity filter				Froger's position				
		LB	LE	H2	H5	LE1	LE2	P1	P2	P3	P4	P5
PIP	BvPIP1;1	NPA	NPA	F	H	T	R	Q	S	A	F	W
	BvPIP1;2	NPA	NPA	F	H	T	R	Q	S	A	*	*
	BvPIP1;3	NPA	NPA	F	H	T	R	Q	S	A	F	W
	BvPIP2;1	NPA	NPA	F	H	T	R	Q	S	A	F	W
	BvPIP2;2	NPA	NPA	F	H	T	R	Q	S	A	F	W
	BvPIP2;3	NPA	NPA	F	H	T	R	Q	S	A	F	W
	BvPIP2;4	NPA	NPA	F	H	T	R	M	S	A	F	W
TIP	BvTIP1;1	NPA	NPA	H	I	A	V	T	S	A	Y	W
	BvTIP1;2	NPA	NPA	H	I	A	V	T	S	A	Y	W
	BvTIP1;3	NPA	NPA	H	I	A	V	T	S	A	Y	W
	BvTIP2;1	NPA	NPA	H	I	G	R	T	S	A	Y	W
	BvTIP2;2	NPA	NPA	H	I	G	R	T	S	A	Y	W
	BvTIP3;1	NPA	NPA	H	I	A	R	T	A	A	Y	W
	BvTIP4;1	NPA	NPA	H	I	A	R	T	S	A	Y	W
NIP	BvTIP5;1	NPA	NPA	N	V	G	Y	T	S	A	Y	W
	BvNIP1;1	NPA	NPA	W	V	A	R	F	S	A	Y	L
	BvNIP4;1	NPA	NPA	W	V	A	R	F	S	A	Y	I
	BvNIP4;2	NPS	NPA	W	A	A	R	L	S	A	Y	I
	BvNIP5;1	NPS	NPV	A	I	G	R	F	T	A	Y	M
	BvNIP5;2	NPS	NPV	A	I	A	R	F	T	A	Y	M
	BvNIP6;1	NPS	NPV	S	I	G	R	F	T	A	Y	F
	BvNIP6;2	NPA	NPA	S	I	G	R	Y	T	A	Y	M
SIP	BvNIP6;3	NPA	NPA	S	I	A	R	Y	T	A	Y	L
	BvNIP7;1	NPA	NPA	A	V	G	R	F	S	A	Y	F
	BvSIP1;1 $\alpha$	NPT	NPA	I	V	P	N	M	A	A	Y	W
SIP	BvSIP1;1 $\beta$	NPT	NPA	V	V	P	N	M	A	A	Y	W
	BvSIP2;1	NPL	NPA	S	N	G	S	F	V	A	Y	W
XIP	BvXIP1;1	NPT	NPA	V	S	A	R	F	C	A	F	W

To explore further the diversity in each group, the conserved motifs were predicted (Fig. 5). Motifs 5 and 6 appeared specifically in PIPs; Motif 8, in NIPs; Motif 9 in TIPs and XIPs.

### Subcellular localization

Plant-mPLOC prediction was used to predict the BvAQPs subcellular localization (Table 1). PIPs, NIPs, TIPs and SIPs (except BvSIP1;1  $\beta$ ) were predicted to localize to plasma membrane and all TIPs were also predicted to localize on vacuolar membrane. However, subcellular localizations predicted by WoLF PSORT were diverse and not always in agreement with that predicted by Plant-mPLOC (Table 1). For example, most PIPs were predicted to localize on plasma membrane. TIPs were predicted to localize on vacuolar membrane, plasma membrane as well as in the nucleus, mitochondria, cytosol and chloroplast. NIPs, SIPs and XIPs were predicted to localize on vacuolar membrane




**Figure 5** BvAQPs protein motifs identified by MEME. Using the complete amino acid sequences of BvAQPs. Combined *p*-values are indicated in (A) and different motifs were shown by different colors and numbered from 1 to 10 in (B).

and plasma membrane. Although these predications represent a starting base, more experiments, i.e., with tagged proteins, are will be useful better characterize the subcellular localization of BvAQPs.


## Expression analysis of sugar beet AQP genes in different organs and in response to salt & heat stress

We used the available RNA-seq data from different organs and young leaf under heat, salt stress treatment of *B. vulgaris* (Dohm et al., 2014; Minoche et al., 2015) to evaluate the expression of BvAQPs. 27 BvAQP genes (96%) were expressed in at least one tissue (Fig. 6, Table S1) and one BvAQP gene (NIP6;3) did not show any expression in all tested plant organs. Eighteen BvAQP genes (64%) expressed in all organs, including six PIP, four TIP, four NIP, three SIP and one XIP. Twenty-six BvAQP genes (92%) expressed in taproot, with BvPIP1;3, BvTIP1;1 and BvPIP2;4 exhibiting the highest expression levels. Twenty-six AQP genes (92%) were expressed in inflorescence, among which BvPIP2;4, BvPIP1;3 and BvTIP1;1 exhibiting the highest expression levels. Twenty-two AQP genes (79%) were expressed in seed (with BvTIP3;1, BvPIP2;1, BvPIP1;3 and BvTIP1;1 showing the highest expression levels). Only 18 AQP genes (64%) were expressed in leaf (BvTIP1;1, BvPIP1;3 and BvTIP2;1 showed the highest expression levels). Most genes were expressed constitutively in the tested tissues, while one gene (BvTIP5;1) showed tissue-specific expression in the inflorescence. In addition, the transcripts of BvTIP1;2 and BvNIP7;1 were also shown to be relatively abundant in inflorescence and BvTIP2;2 was highly expressed in taproot and seeding.

The expression patterns of sugar beet AQP genes were found to fluctuate under different stress conditions. Eleven AQP genes (44%) expression were down-regulated both under salt and heated stress. These include BvPIP1;1, BvPIP1;2, BvPIP2;3, BvNIP4;1, BvNIP4;2,



**Figure 6** Expression profiles of the 28 *BvAQP* genes in different plant organs and tissues.



**Figure 7** Expression pattern of the 23 *BvAQP* genes in young leaf under abiotic stress.

*BvNIP5;2*, *BvNIP6;1*, *BvNIP7;1*, *BvSIP1;1 α*, *BvSIP1;1 β* and *BvSIP2;1* (Fig. 7). In contrast, the expression of *BvPIP2;4*, *BvTIP1;1* and *BvTIP2;1* (12%) were up-regulated both under salt and heat stress. Interestingly, nine AQP genes showed different expression in response to salt or heat treatments. Eight among these AQP genes (32%; *BvPIP1;3*, *BvPIP2;1*, *BvPIP2;2*, *BvTIP1;3*, *BvTIP4;1*, *BvNIP1;1*, *BvNIP5;1* and *BvXIP1;1*) were up-regulated under heat treatment but down-regulated under salt treatment. Notably, members of PIP genes and XIP genes (*BvPIP1;3*, *BvPIP2;1*, *BvPIP2;2*, *BvXIP1;1*) were highly up-regulated under heat stress but slightly down-regulated under salt stress. Only *BvTIP2;2* (4%) was up-regulated under salt but down-regulated under heat stress.

Interestingly, *BvNIP6;3* wasn't expressed in all organs, neither in young leaf under salt and heat stress (Fig. 6; Table S1), and *BvNIP6;3* formed a single-gene clade within the NIP6s, with the highest MV (35.08kDa) and longest protein length (Table 1). In addition, gene duplication analysis revealed *BvNIP6;3* and *BvNIP6;2* were tandem duplication (Fig. 2) and suggested that *BvNIP6;3* was originated from *BvNIP6;2* and was pseudo-genes.

Other genes, such as *BvTIP1;2*, *BvTIP3;1*, *BvNIP6;2* and *BvNIP6;3* were not expressed in non-treated or in stressed leaf. *BvTIP5;1* was not expressed in control or heat stressed young leaf, but showed only very expression in young leaf under salt stress (Table S1).

## DISCUSSION

Many studies have confirmed that AQP genes were involved in plant water transport, in regulating growth and development and that they are widely distributed in animals and plants. Most functional studies of AQP genes were mainly conducted using model plants, such as *A. thaliana*, *N. tabacum* L., *Z. mays* L. and *O. sativa* L. Almost nothing or very little is known about *B. vulgaris* AQPs. Here we report the first investigation of AQPs in sugar beet. Twenty-eight AQP genes were identified, including seven PIPs, eight TIPs, nine NIPs, three SIPs and one XIP.

The number of AQP genes in *B. vulgaris* is similar to that of *D. caryophyllus* (Morita et al., 2017), a closely related species belonging to the same family. *B. vulgaris* contains less AQP genes compared to plants that did undergo 1-3 times of whole genome duplication (WGD), fragment duplication (FD) and/or tandem duplication (TD), such as *G. max* L (Zhang et al., 2013), *Brassica rapa* (Tao et al., 2014), *Z. mays* L. (Chaumont et al., 2001), *O. sativa* L (Nguyen, Moon & Jung, 2013) and *P. trichocarpa* (Gupta & R, 2009). *B. vulgaris* contains more AQP genes than basal plants such as *P. patens*, *S. moellendorffii* (Bowers et al., 2003; Doyle et al., 2008; Gupta & R, 2009; Yu & Yang, 2002; Zhang et al., 2013). Almost all subgroups of PIPs, TIPs, SIPs and XIPs are present in the Caryophyllaceae plants *B. vulgaris* and *D. caryophyllus* (Morita et al., 2017).

The relatively small number of AQP genes in sugar beet, compared to other higher plant species, is likely due to the fact that *B. vulgaris* did not undergo WGD. Moreover, the rapid expansion of NIP subfamily which led to the production of new subgroups, might have occurred after the divergence of the basal core dicot and core dicot. The divergence and proliferation of NIP subfamily may be an adaptive response to an ever-changing environment, playing crucial role in plants disease resistance and multi-stress

(Liu & Zhu, 2010). Some studies also suggested that NIPs allow larger solutes, such as silicic acid, to permeate. *O. sativa* L genes *OsNIP2;1* (*Lsi1*), *OsNIP2;2* (*Lsi6*), and *Hordeum vulgare* L. *HvNIP2;1* (*HvLsi1*) transport silicon across the biomembrane and enhance the resistance of plants to biotic and abiotic stress (Chiba et al., 2009; Ma et al., 2006; Yamaji & Ma, 2009; Yamaji, Mitatni & Ma, 2008), and *O. sativa* L *OsNIP2;1* could also be permeable to water, urea, boric acid, arsenite (Ma et al., 2008; Mitani, Yamaji & Ma, 2008).

To analyze the evolutionary relationship and BvAQPs putative functions, an unrooted phylogenetic tree was constructed using aquaporins from *A. thaliana* (in which a complete set of AtAQP genes is well known) and *D. caryophyllus*, closely related species to *B. vulgaris* (Johanson & Kjellbom, 2001; Morita et al., 2017). During the process of cluster analysis, two carnation AQPs (DcSIP2;1 and DcNIP5;1) aroused our attention and made us confused in classification at once. Previously, the classification of DcSIP2;1 and DcNIP5;1 was not well resolved (Morita et al., 2017). In the case of DcSIP2;1, it shared the highest similarity of 67% with BvSIP1;1 and 51% with AtSIP1;1, and also shared the same NPA motifs, Ar/R selectivity filter and Froger's position with BvSIP1;1. It clustered closer to the SIP1 subgroup. In the case of DcNIP5;1, it farther clustered to the NIP5 subgroup and clustered closer to the NIP6 subgroup. The closest homolog of DcNIP5;1 is BvNIP6;3 (similarity is 51%) and both proteins share the same Ar/R selectivity filter (S-I-A-R) with BvNIP6;3 and BvNIP6;1, with a similar Ar/R selectivity filter (A-I-A-R) to DcNIP6;1. Therefore, DcNIP5;1 belongs to NIP6 subgroup.

Alternative splicing is a mechanism by which genes can produce multiple transcript variants protein products, which allows in turn to increase the diversity of gene functions. Alternative splicing plays an important role in various processes such as development, response to pathogen and to various abiotic stresses (Bove et al., 2008; Gassmann, 2008; Jang et al., 2009; Reddy & Golovkin, 2010). A pair of splice variants (*BvSIP1;1*  $\alpha$  and *BvSIP1;1*  $\beta$ ) were found in BvAQPs. *BvSIP1;1*  $\alpha$  resulted from second exon skipping. First exon and second exon encoded the same 'NPT' type NPA motif, so *BvSIP1;1*  $\alpha$  and *BvSIP1;1*  $\beta$  shared similar amino acid sequence and protein topology. These two genes had a similar expression pattern in different organs. It is possible that *BvSIP1;1*  $\alpha$  and *BvSIP1;1*  $\beta$  may have redundant function. Exon skipping splicing were also found in *S. tuberosum*, while *St-SIP1;1*  $\alpha$  and *St-SIP1;1*  $\beta$  also resulted from second exon skipping and shared similar expression pattern (Venkatesh, Yu & Park, 2013). However, the identified alternative splicing events found for BvAQPs genes were not reported in other plant species, suggesting that selective splicing may be species-specific.

NPA motifs, ar/R selectivity filter and Froger's position were reported involved in substrate selection and transport activity. BvAQPs functions could be conferred based on the comparison of these residues with other plants AQPs. BvPIPs showed typical NPA motif, F-H-T-R highly conserved ar/R selectivity filter and Q-S-A-F-W Froger's position, this composition is highly conserved in PIPs of *A. thaliana* (Johanson & Kjellbom, 2001), *Z. mays* (Chaumont et al., 2001), *S. lycopersicum* (Reuscher et al., 2013b), *P. trichocarpa* (Gupta & R, 2009), *G. max* (Zhang et al., 2013) and *B. rapa* (Tao et al., 2014). This composition of PIPs was reported to likely regulate root and leaf hydraulics, facilitate the CO<sub>2</sub> diffusion, affect photosynthesis (Gupta & R, 2009). Therefore, *B. vulgaris* homologous PIPs may have



similar roles in regulating water absorption, plant hydraulics and CO<sub>2</sub> diffusion. Based on the SDPs analysis proposed by Hove and Bhavé ([Hove & Bhavé, 2011](#)) ([Table 3](#); [Fig. S2](#)), all *B. vulgaris* PIPs represented urea-type SDPs, BvPIP1;1 and BvPIP1;2 represented boric acid-type SDPs, all PIP2s represented H<sub>2</sub>O<sub>2</sub>-type SDPs, thus supporting conserved functions. In addition, BvPIP1;2, BvPIP1;3 and BvPIP2;1 seemed to represent novel CO<sub>2</sub>-type SDPs (I/L/V-M-C-A-I/V-D/H/K-W-D-W) with the substitution of Ile for Met in SDP2 and the substitution of D to H/k in SDP6. Although the ar/R selectivity filter varied highly, plant TIPs were shown to transport water as efficiently as PIPs ([Zhi et al., 2015](#)). Additionally, BvTIP1;1, BvTIP1;2, BvTIP1;3, BvTIP3;1 and BvTIP4;1 contained dual conserved NPA motifs, H-I-A-V/R ar/R selectivity filter and T-S-A-Y-W Froger's position, which had the same composition of *Citrus sinensis* AQPs, such as CsTIP1s, CsTIP3s showed to transport urea and H<sub>2</sub>O<sub>2</sub> ([Hove & Bhavé, 2011](#)). Compared to other TIPs, BvTIP2;1 and BvTIP2;2 had H-I-G-R ar/R selectivity filter, and this type ar/R selectivity filter was experimentally proven to transport formamide ([Hove & Bhavé, 2011](#)), suggesting BvTIPs play a crucial role in transporting a wide range molecules. All TIPs (exception of BvTIP5;1) represented urea-type SDPs, indicating transporting urea function. BvTIP1;1 and BvTIP3;1 represented H<sub>2</sub>O<sub>2</sub>-type SDPs, indicating transporting H<sub>2</sub>O<sub>2</sub> potential capacity.

Substrate selection specificity of NIPs is largely determined by two pores formed by NPA motifs and ar/R selectivity filter respectively. 'W-V-A-R' type ar/R selectivity filter were proven to be permeable to uncharged molecules i.e., glycerol, formamide and water, implying that BvNIP1;1 and BvNIP4;1 may have similar functions. 'A/S/T-V/I-A/G-R' type ar/R selectivity filter were reported can transport glycerol, formamide and larger solutes, like urea, boric, but were impermeable to water. Besides, NPS/NPV aqueous pore and A-I-G-R ar/R selectivity filter was known as a boric acid transporter in *A. thaliana* AtNIP5;1, and orthologs BvNIP5;1 can be involved in boron transport in *B. vulgaris*, suggesting that NIPs could be involved in the transport of larger solutes. Based on SDPs analysis, all NIPs (except BvNIP7;1) are potential urea transporters, while BvNIP4;2 is a potential H<sub>2</sub>O<sub>2</sub> transporters, BvNIP5;2 and BvNIP6;1 are potential boric acid transporters and finally BvNIP1;1 could represent a novel NH<sub>3</sub>-type SDPs with H replaced K/L/N/V at SDP2. It should be noticed that we were unable to find NIP2 subgroup and silicic acid-type SDP AQPs in sugar beet, suggesting that other types of SDPs AQPs could be involved in absorption of silicic during plant growth and development, in sugar beet. Sugar beet SIPs and XIPs exhibited great variety in ar/R selectivity filter and Froger's position, with a variable first NPA motif compared to other AQPs subfamilies. It has been reported that *A. thaliana* SIPs (AtSIP1;1 and AtSIP1;2) localize to endoplasmic reticulum (ER) and facilitate water transport. SDPs analysis suggest that BvXIP1;1 can transport boric acid, it is thus possible that BvXIP has different role in *B. vulgaris*.

Sequence alignment revealed that *BvPIP1;1* codes for a truncated protein that lacks TM6 domain. The absence of TM6 domain possibly affects subcellular localization, proper folding, and oligomerization and transport activity of this truncated AQP in *B. vulgaris*. Similarly truncated AQP TdPIP2;1 has been reported to affect water transport activity in wheat ([He et al., 2012](#); [Hu, 2012](#)). RNA-seq analysis revealed that AQP genes were expressed in all examined tissues of *B. vulgaris*, thus similar to previously reported

**Table 3** Identified typical SDPs in BvAQPs.

Aquaporin	Specificity-determining positions								
	SDP1	SDP2	SDP3	SDP4	SDP5	SDP6	SDP7	SDP8	SDP9
<b>Typical NH<sub>3</sub> transporter</b>	F/T	K/L/N/V	F/T	V/L/T	A	D/S	A/H/L	E/P/S	A/R/T
BvNIP1;1	F	H	F	T	A	D	L	E	T
<b>Typical Boric Acid transporter</b>	T/V	I/V	H/I	P	E	I/L	I/L/T	A/T	A/G/K/P
BvPIP1;2	T	I	H	P	E	L	L	T	P
BvPIP1;3	T	I	H	P	E	L	L	T	P
BvNIP5;2	T	I	H	P	E	L	L	A	P
BvNIP6;1	T	I	H	P	E	L	L	A	P
BvXIP1;1	T	I	H	P	E	L	L	T	P
<b>Typical CO<sub>2</sub> transporter</b>	I/L/V	I	C	A	I/V	D	W	D	W
BvPIP1;2	V	M	C	A	I	D	W	H	W
BvPIP1;3	V	M	C	A	I	H	W	D	W
BvPIP2;1	I	M	C	A	V	K	W	D	W
<b>Typical H<sub>2</sub>O<sub>2</sub> transporters</b>	A/S	A/G	L/V	A/F/L/T/V	I/L/V	H/I/L/Q	F/Y	A/V	P
BvPIP2;1	A	G	V	F	I	H	F	V	P
BvPIP2;2	A	G	V	F	I	H	F	V	P
BvPIP2;3	A	G	V	F	I	H	F	V	P
BvPIP2;4	A	G	V	F	I	H	F	V	P
BvTIP1;1	S	A	L	A	I	H	Y	V	P
BvTIP3;1	A	A	L	T	I	H	Y	V	P
BvNIP4;2	S	A	L	L	I	L	Y	V	P
<b>Typical silicic acid transporters</b>	C/S	F/Y	A/E/L	H/R/Y	G	K/N/T	R	E/S/T	A/K/P/T
Not found									
<b>Typical urea transporters</b>	H	P	F/I/L/T	A/C/F/L	L/M	A/G/P	G/S	G/S	N
BvPIP1;1	H	P	F	L	L	P	G	G	N
BvPIP1;2	H	P	F	L	L	P	G	G	N
BvPIP1;3	H	P	L	F	L	P	G	G	N
BvPIP2;1	H	P	F	L	L	P	G	G	N
BvPIP2;2	H	P	F	F	L	P	G	G	N
BvPIP2;3	H	P	F	F	L	P	G	G	N
BvPIP2;4	H	P	F	F	L	P	G	G	N
BvTIP1;1	H	P	F	F	L	A	G	S	N
BvTIP1;2	H	P	F	F	L	P	G	S	N
BvTIP1;3	H	P	L	F	L	A	G	S	N
BvTIP2;1	H	P	F	A	L	P	G	S	N
BvTIP2;2	H	P	F	A	L	P	G	S	N
BvTIP3;1	H	P	F	L	L	P	G	S	N
BvTIP4;1	H	P	L	A	L	L	G	S	N
BvNIP1;1	H	P	I	A	L	P	G	S	N
BvNIP4;1	H	P	L	A	L	P	G	S	N
BvNIP4;2	H	P	I	A	L	T	G	S	N

(continued on next page)

Table 3 (continued)

Aquaporin	Specificity-determining positions								
	SDP1	SDP2	SDP3	SDP4	SDP5	SDP6	SDP7	SDP8	SDP9
BvNIP5;1	H	P	I	A	L	P	G	S	N
BvNIP5;2	H	P	I	A	L	P	G	S	N
BvNIP6;1	H	P	I	A	L	P	G	S	N
BvNIP6;2	H	P	L	A	L	P	G	S	N
BvNIP6;3	H	P	I	A	L	P	G	S	N

data in maize ([Chaumont et al., 2001](#)), *A. Arabidopsis* ([Quigley, 2001](#)), tomato ([Reuscher et al., 2013a](#)) and potato ([Venkatesh, Yu & Park, 2013](#)). Total transcript abundance in the different examined tissues showed high expression levels in organs involved in water absorption, transport and evaporation, such as taproot and leaf, suggesting the main role of AQPs in water and nutrients transport. The RNA-seq data suggest that BvAQPs may play a key role in radial and axial water transportation in sugar beet, thus in agreement with [Amodeo et al. \(1999\)](#). Many AQPs show similar expression patterns, suggesting that they may act synergistically in some tissues. Co-expression of tandem PIP2-PIP1 dimers in *Xenopus* oocytes showed that they can form PIP2-PIP1 hetero-tetramers and synergistically increase water transport capacity ([Bellati et al., 2010](#); [Jozefkowicz et al., 2013](#); [Jozefkowicz et al., 2015](#)). *BvPIP1;3*, *BvTIP1;1* and *BvPIP2;4* were considerably abundant in taproot and are likely involved in water and nutrients transport in root. *BvPIP2;4*, *BvPIP1;3* and *BvTIP1;1* were abundant in inflorescence and seeding, and could be involved in monitoring the water balance in young tissues. According to their expression profiles, *BvTIP1;1*, *BvPIP1;3* and *BvTIP2;1* could play key roles in leaf hydraulics. *BvTIP3;1*, *BvPIP2;1*, *BvPIP1;3* and *BvTIP1;1* were considerably abundant in seed and therefore could be involved in water balance in seed. *BvPIP1;3* and *BvTIP1;1* show high expression levels in all examined tissues, while *BvTIP3;1* were only expressed in abundance in seed, and showed very low expression in other organs, suggesting a key role in seed development. *BvTIP3;1* closest homolog *Arabidopsis AtTIP3;1* and castor bean *RcTIP3;1* also shared similar expression pattern. *AtTIP3;1* was reported to be seed- and embryo-specific AQP ([Johnson, Herman & Chrispeels, 1989](#)). *RcTIP3;1* expressed preferentially in endosperm of developing seeds and considerably low in germinating seed ([Zou et al., 2015](#)). Moreover, *BvPIP2;2* and *BvTIP2;2* were highly expressed in seeding where they may control the water balance. In total, several AQP genes (like *BvTIP3;1*, *BvPIP2;1*, and *BvPIP1;3*) play a constitutive role and some AQP genes (like *BvTIP3;1*, *BvPIP2;2* and *BvTIP2;2*) play a specialized function in specific plants organs. In addition, it is noteworthy that several putative non-aqua transporter-encoding genes (i.e., *BvNIP7;1*, *BvNIP4;1*, *BvXIP1;1*, *BvNIP5;2*, *BvNIP6;1* and *BvSIP2;1*) were shown to be relatively high abundant in certain tissues. Compared to others tissues, transcript level of *BvNIP4;1*, *BvNIP5;2* and *BvNIP6;1* was considerably high in taproot. Similar to *BvNIP7;1*, *Arabidopsis AtNIP7;1* (specifically expressed in anthers) encode for a less efficient boric acid transporter, compared to *AtNIP5;1* and *AtNIP6;1*. It is possible that *BvNIP7;1* plays a role in B absorption and balance in inflorescence tissue. *BvXIP1;1* had relatively

high expression level in leaf. Similarly, *BvXIP1;1* the closest homolog to *RcXIP1;1*, also showed high expression levels in leaf, but their exact functions remain unclear and further functional investigation are required.

In this study, a large number of *BvAQP* genes showed transcriptional changes when exposed to salt and heat stresses. For example, *BvTIP1;1* and *BvTIP2;1* showed strong induction ( $\text{Log}_2$ -based value  $>1$ ) after salt stress; *BvPIP2;2*, *BvTIP1;1*, *BvTIP2;1* and *BvXIP1;1* showed strong induction ( $\text{Log}_2$ -based value  $>1$ ) after heat stress, suggesting an extensive response of *BvAQP* genes to abiotic stress and a potential function for improving sugar beet resistance to abiotic stress. The results were consistent with previously reported data that showed that *BvPIP2;2* expression was up-regulated in response to high salt stress (Skorupaklaput et al., 2015). Biochemical and genetic evidence has demonstrated that some AQP genes (like *TaAQP7*, *TaAQP8*, *TaNIP*, *MaPIP1;1*, *MusaPIP2;6* and *NtAQP1*) improve plants resistance to abiotic stress (Gao et al., 2010; Hu, 2012; Sade, 2010; Sreedharan, Shekhawat & Ganapathi, 2015; Xu et al., 2014; Zhou et al., 2012). Additionally, we noticed that *BvTIP1;1* and *BvTIP2;1* were strongly induced by both salt and heat stresses, suggesting that these two AQPs may play key roles in *B. vulgaris* adaption to environmental changes, e.g., heat and salt stresses.

## CONCLUSIONS

Twenty-eight *BvAQP* genes were identified based on genome data, chromosome distribution, phylogenetic, protein characteristics, gene structural, gene duplication, conserved motifs and RNA-seq expression analysis of *BvAQP* genes were further researched. The results suggested that AQPs had multiple functions in different tissues; several *BvAQP* genes responded to abiotic stresses and improved the plants' resistance to abiotic stresses. The potential functions of *BvAQPs* were predicted and discussed based on NPA motifs, ar/R selectivity filter, Froger's positions and SDP positions analysis. This study provide a useful resource for identifying and characterizing *BvAQPs* and a base for the breeding and genetic engineering of sugar beet.

## ACKNOWLEDGEMENTS

The authors appreciate those contributors who make the sugar beet genome and transcriptome data accessible in public databases.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This study was supported by the National Natural Science Foundation of China (31000918) and the Fundamental Research Funds for the Central Universities (2662015PY052 and 2662016PY041). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

## Grant Disclosures

The following grant information was disclosed by the authors:

National Natural Science Foundation of China: 31000918.

Fundamental Research Funds for the Central Universities: 2662015PY052, 2662016PY041.

## Competing Interests

The authors declare there are no competing interests.

## Author Contributions

- Weilong Kong conceived and designed the experiments, performed the experiments, analyzed the data, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.
- Shaozong Yang and Yulu Wang performed the experiments.
- Mohammed Bendahmane and Xiaopeng Fu conceived and designed the experiments, contributed reagents/materials/analysis tools.

## Data Availability

The following information was supplied regarding data availability:

The raw data has been supplied as [Supplementary Files](#).

## Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.3747#supplemental-information>.

## REFERENCES

- Amodeo G, Dorr R, Vallejo A, Sutka M, Parisi M. 1999.** Radial and axial water transport in the sugar beet storage root. *Journal of Experimental Botany* **50**(333):509–516 DOI [10.1093/jxb/50.333.509](https://doi.org/10.1093/jxb/50.333.509).
- Anderberg HI, Kjellbom P, Johanson U. 2012.** Annotation of *Selaginella moellendorffii* major intrinsic proteins and the evolution of the protein family in terrestrial plants. *Frontiers in Plant Science* **3**:Artn 33 DOI [10.3389/fpls.2012.00033](https://doi.org/10.3389/fpls.2012.00033).
- Ariani A, Gepts P. 2015.** Genome-wide identification and characterization of aquaporin gene family in common bean (*Phaseolus vulgaris* L.). *Molecular Genetics and Genomics* **290**:1771–1785 DOI [10.1007/s00438-015-1038-2](https://doi.org/10.1007/s00438-015-1038-2).
- Bansal A, Sankararamakrishnan R. 2007.** Homology modeling of major intrinsic proteins in rice, maize and Arabidopsis: comparative analysis of transmembrane helix association and aromatic/arginine selectivity filters. *BMC Structural Biology* **7**:1–17 DOI [10.1186/1472-6807-7-27](https://doi.org/10.1186/1472-6807-7-27).
- Baranwal VK, Negi N, Khurana P. 2016.** Genome-wide identification and structural, functional and evolutionary analysis of WRKY components of Mulberry. *Scientific Reports* **6**:30794 DOI [10.1038/srep30794](https://doi.org/10.1038/srep30794).
- Bellati J, Alleva K, Soto G, Vitali V, Jozefkowicz C, Amodeo G. 2010.** Intracellular pH sensing is altered by plasma membrane PIP aquaporin co-expression. *Plant Molecular Biology* **74**(1):105–118 DOI [10.1007/s11103-010-9658-8](https://doi.org/10.1007/s11103-010-9658-8).

- Bove J, Kim CY, Gibson CA, Assmann SM. 2008.** Characterization of wound-responsive RNA-binding proteins and their splice variants in Arabidopsis. *Plant Molecular Biology* 67:71–88 DOI 10.1007/s11103-008-9302-z.
- Bowers JE, Chapman BA, Rong J, Paterson AH. 2003.** Unravelling angiosperm genome evolution by phylogenetic analysis of chromosomal duplication events. *Nature* 422:433–438 DOI 10.1038/nature01521.
- Chaumont F, Barrieu F, Wojcik E, Chrispeels MJ, Jung R. 2001.** Aquaporins constitute a large and highly divergent protein family in maize. *Plant Physiology* 125:1206–1215 DOI 10.1104/pp.125.3.1206.
- Cheng F, Mandáková T, Wu J, Xie Q, Lysak MA, Wang X. 2013.** Deciphering the diploid ancestral genome of the mesohexaploid *Brassica rapa*. *The Plant Cell* 25:1541–1554 DOI 10.1105/tpc.113.110486.
- Chiba Y, Mitani N, Yamaji N, Ma JF. 2009.** HvLsi1 is a silicon influx transporter in barley. *Plant Journal* 57:810–818 DOI 10.1111/j.1365-313X.2008.03728.x.
- Danielson JAH, Johanson U. 2008.** Unexpected complexity of the aquaporin gene family in the moss *Physcomitrella patens*. *BMC Plant Biology* 8:Article 45 DOI 10.1186/1471-2229-8-45.
- Dohm JC, Minoche AE, Holtgräwe D, Capellagutiérrez S, Zakrzewski F, Tafer H, Rupp O, Sørensen TR, Stracke R, Reinhardt R. 2014.** The genome of the recently domesticated crop plant sugar beet (*Beta vulgaris*). *Nature* 505:546–549 DOI 10.1038/nature12817.
- Doyle JJ, Paterson AH, Soltis DE, Wendel JF. 2008.** Evolutionary genetics of genome merger and doubling in plants. *Genetics* 42:443–461 DOI 10.1146/annurev.genet.42.110807.091524.
- Fitzpatrick KL, Reid RJ. 2009.** The involvement of aquaglyceroporins in transport of boron in barley roots. *Plant Cell & Environment* 32:1357–1365 DOI 10.1111/j.1365-3040.2009.02003.x.
- Froger A, Tallur B, Thomas D, Delamarche C. 1998.** Prediction of functional residues in water channels and related proteins. *Protein Science* 7:1458–1468 DOI 10.1002/pro.5560070623.
- Gao Z, He X, Zhao B, Zhou C, Liang Y, Ge R, Shen Y, Huang Z. 2010.** Overexpressing a putative aquaporin gene from wheat, TaNIP, enhances salt tolerance in transgenic Arabidopsis. *Plant & Cell Physiology* 51:767–775 DOI 10.1093/pcp/pcq036.
- Gassmann W. 2008.** Alternative splicing in plant defense. *Current Topics in Microbiology & Immunology* 326:219–234.
- Gentleman RC, Carey VJ, Bates DM, Bolstad B, Dettling M, Dudoit S, Ellis B, Gautier L, Ge Y, Gentry J. 2004.** Bioconductor: open software development for computational biology and bioinformatics. *Genome Biology* 5:1–16 DOI 10.1186/gb-2004-5-10-r80.
- Gomes D, Agasse A, Thiébaud P, Delrot S, Gerós H, Chaumont F. 2009.** Aquaporins are multifunctional water and solute transporters highly divergent in living organisms. *Biochimica Et Biophysica Acta* 1788:1213–1228 DOI 10.1016/j.bbamem.2009.03.009.
- Gu Z, Cavalcanti A, Chen FC, Bouman P, Li WH. 2002.** Extent of gene duplication in the genomes of drosophila, nematode, and yeast. *Molecular Biology and Evolution* 19:256–262 DOI 10.1093/oxfordjournals.molbev.a004079.



- Gupta A, R S. 2009.** Genome-wide analysis of major intrinsic proteins in the tree plant *Populus trichocarpa*: characterization of XIP subfamily of aquaporins from evolutionary perspective. *BMC Plant Biology* 9:134 DOI 10.1186/1471-2229-9-134.
- Gupta AB, Sankararamakrishnan R. 2009.** Genome-wide analysis of major intrinsic proteins in the tree plant *Populus trichocarpa*: characterization of XIP subfamily of aquaporins from evolutionary perspective. *BMC Plant Biology* 9:9 DOI 10.1186/1471-2229-9-134.
- He H, Dong Q, Shao Y, Jiang H, Zhu S, Cheng B, Xiang Y. 2012.** Genome-wide survey and characterization of the WRKY gene family in *Populus trichocarpa*. *Plant Cell Reports* 31:1199–1217 DOI 10.1007/s00299-012-1241-0.
- Heinen RB, Chaumont F. 2009.** Role of aquaporins in leaf physiology. *Journal of Experimental Botany* 60:2971–2985 DOI 10.1093/jxb/erp171.
- Hove RM, Bhavé M. 2011.** Plant aquaporins with non-aqua functions: deciphering the signature sequences. *Plant Molecular Biology* 75:413–430 DOI 10.1007/s11103-011-9737-5.
- Hu W. 2012.** Overexpression of a wheat aquaporin gene, TaAQP8, enhances salt stress tolerance in transgenic tobacco. *Plant & Cell Physiology* 53:2127–2141 DOI 10.1093/pcp/pcs154.
- Jang YH, Lee JH, Park HY, Kim SK, Lee BY, Suh MC, Kim JK. 2009.** OsFCA transcripts show more complex alternative processing patterns than its Arabidopsis counterparts. *Journal of Plant Biology* 52:161–166 DOI 10.1007/s12374-009-9018-x.
- Johanson U, Kjellbom P. 2001.** The complete set of genes encoding major intrinsic proteins in Arabidopsis provides a framework for a new nomenclature for major intrinsic proteins in plants. *Plant Physiology* 126:1358–1369 DOI 10.1104/pp.126.4.1358.
- Johnson KD, Herman EM, Chrispeels MJ. 1989.** An abundant, highly conserved tonoplast protein in seeds. *Plant Physiology* 91:1006–1013 DOI 10.1104/pp.91.3.1006.
- Jozefkowicz C, Rosi P, Sigaut L, Soto G, Pietrasanta LI, Amodeo G, Alleva K. 2013.** Loop A is critical for the functional interaction of two *Beta vulgaris* PIP aquaporins. *PLOS ONE* 8(3):e57993 DOI 10.1371/journal.pone.0057993.
- Jozefkowicz C, Sigaut L, Scochera F, Soto G, Ayub N, Pietrasanta LI, Amodeo G, Flecha González FL, Alleva K. 2015.** PIP water transport and its pH dependence are regulated by tetramer stoichiometry. *Biophysical Journal* 110(6):1312–1321 DOI 10.1016/j.bpj.2016.01.026.
- Liu QP, Zhu ZJ. 2010.** Functional divergence of the NIP III subgroup proteins involved altered selective constraints and positive selection. *BMC Plant Biology* 10:Article 256 DOI 10.1186/1471-2229-10-256.
- Ma JF, Tamai K, Yamaji N, Mitani N, Konishi S, Katsuhara M, Ishiguro M, Murata Y, Yano M. 2006.** A silicon transporter in rice. *Nature* 440:688–691 DOI 10.1038/nature04590.
- Ma JF, Yamaji N, Mitani N, Xu XY, Su YH, McGrath SP, Zhao FJ. 2008.** Transporters of arsenite in rice and their role in arsenic accumulation in rice grain. *Proceedings of the National Academy of Sciences of the United States of America* 105:9931–9935 DOI 10.1073/pnas.0802361105.

- Maurel C, Santoni V, Luu DT, Wudick MM, Verdoucq L. 2009.** The cellular dynamics of plant aquaporin expression and functions. *Current Opinion in Plant Biology* 12:690–698 DOI 10.1016/j.pbi.2009.09.002.
- Maurel C, Verdoucq L, Luu DT, Santoni V. 2008.** Plant aquaporins: membrane channels with multiple Integrated functions. *Plant Biology* 59:595–624 DOI 10.1146/annurev.arplant.59.032607.092734.
- Minoche AE, Dohm JC, Schneider J, Holtgräwe D, Viehöver P, Montfort M, Sörensen TR, Weisshaar B, Himmelbauer H. 2015.** Exploiting single-molecule transcript sequencing for eukaryotic gene prediction. *Genome Biology* 16:1–13 DOI 10.1186/s13059-015-0729-7.
- Mitani N, Yamaji N, Ma JF. 2008.** Characterization of substrate specificity of a rice silicon transporter, Lsi1. *Pflügers Archiv-European Journal of Physiology* 456:679–686 DOI 10.1007/s00424-007-0408-y.
- Mitani-Ueno N, Yamaji N, Zhao FJ, Ma JF. 2011.** The aromatic/arginine selectivity filter of NIP aquaporins plays a critical role in substrate selectivity for silicon, boron, and arsenic. *Journal of Experimental Botany* 62:4391–4398 DOI 10.1093/jxb/err158.
- Morita S, Sugiyama S, Tateishi A, Satoh S. 2017.** Identification and characterization of plasma membrane intrinsic protein (PIP) aquaporin genes in petals of opening carnation flowers. *Horticulture Journal* 86:78–86 DOI 10.2503/hortj.MI-127.
- Nguyen MX, Moon S, Jung KH. 2013.** Genome-wide expression analysis of rice aquaporin genes and development of a functional gene network mediated by aquaporin expression in roots. *Planta* 238:669–681 DOI 10.1007/s00425-013-1918-9.
- Prado K, Maurel C. 2013.** Regulation of leaf hydraulics: from molecular to whole plant levels. *Frontiers in Plant Science* 4:58–60 DOI 10.3389/fpls.2013.00255.
- Quigley F. 2001.** From genome to function: the Arabidopsis aquaporins. *Genome Biology* 3:RESEARCH0001 DOI 10.1186/gb-2001-3-1-research0001.
- Reddy ASN, Golovkin MV. 2010.** *Nuclear pre-mRNA processing in plants*. Berlin: Springer, 326.
- Reuscher S, Akiyama M, Mori C, Aoki K, Shibata D, Shiratake K. 2013a.** Genome-wide identification and expression analysis of aquaporins in tomato. *PLOS ONE* 8:e79052 DOI 10.1371/journal.pone.0079052.
- Reuscher S, Akiyama M, Mori C, Aoki K, Shibata D, Shiratake K. 2013b.** Genome-wide identification and expression analysis of aquaporins in tomato. *PLOS ONE* 8:615–629 DOI 10.1371/journal.pone.0079052.
- Sade N. 2010.** The role of tobacco Aquaporin1 in improving water use efficiency, hydraulic conductivity, and yield production under salt stress. *Plant Physiology* 152:245–254 DOI 10.1104/pp.109.145854.
- Skorupakłaput M, Szczepanek J, Kurnik K, Tretyn A, Tyburski J. 2015.** The expression patterns of plasma membrane aquaporins in leaves of sugar beet and its halophyte relative, Beta vulgaris ssp. maritima, in response to salt stress. *Biologia* 70(4):467–477 DOI 10.1515/biolog-2015-0056.
- Sreedharan S, Shekhawat UKS, Ganapathi TR. 2015.** Constitutive and stress-inducible overexpression of a native aquaporin gene (MusaPIP2;6) in transgenic banana

- plants signals its pivotal role in salt tolerance. *Plant Molecular Biology* **88**:41–52 DOI [10.1007/s11103-015-0305-2](https://doi.org/10.1007/s11103-015-0305-2).
- Stracke R, Holtgräwe D, Schneider J, Pucker B, Sörensen TR, Weisshaar B. 2014.** Genome-wide identification and characterisation of R2R3-MYB genes in sugar beet (*Beta vulgaris*). *BMC Plant Biology* **14**:1–17 DOI [10.1186/s12870-014-0249-8](https://doi.org/10.1186/s12870-014-0249-8).
- Tao P, Zhong X, Li B, Wang W, Yue Z, Lei J, Guo W, Huang X. 2014.** Genome-wide identification and characterization of aquaporin genes (AQPs) in Chinese cabbage (*Brassica rapa* ssp. *pekinensis*). *Molecular Genetics and Genomics* **289**:1131–1145 DOI [10.1007/s00438-014-0874-9](https://doi.org/10.1007/s00438-014-0874-9).
- Uehlein N, Kaldenhoff R. 2008.** Aquaporins and plant leaf movements. *Annals of Botany* **101**:1–4 DOI [10.1093/aob/mcm278](https://doi.org/10.1093/aob/mcm278).
- Venkatesh J, Yu JW, Park SW. 2013.** Genome-wide analysis and expression profiling of the *Solanum tuberosum* aquaporins. *Plant Physiology & Biochemistry* **73C**:392–404 DOI [10.1016/j.plaphy.2013.10.025](https://doi.org/10.1016/j.plaphy.2013.10.025).
- Wudick MM, Maurel C. 2009.** A look inside: localization patterns and functions of intracellular plant aquaporins. *New Phytologist* **184**:289–302 DOI [10.1111/j.1469-8137.2009.02985.x](https://doi.org/10.1111/j.1469-8137.2009.02985.x).
- Xu Y, Hu W, Liu J, Zhang J, Jia C, Miao H, Xu B, Jin Z. 2014.** A banana aquaporin gene, MaPIP1;1, is involved in tolerance to drought and salt stresses. *BMC Plant Biology* **14**:59 DOI [10.1186/1471-2229-14-59](https://doi.org/10.1186/1471-2229-14-59).
- Yamaji N, Ma JF. 2009.** A transporter at the node responsible for intervascular transfer of silicon in rice. *The Plant Cell* **21**:2878–2883 DOI [10.1105/tpc.109.069831](https://doi.org/10.1105/tpc.109.069831).
- Yamaji N, Mitatni N, Ma JF. 2008.** A transporter regulating silicon distribution in rice shoots. *The Plant Cell* **20**:1381–1389 DOI [10.1105/tpc.108.059311](https://doi.org/10.1105/tpc.108.059311).
- Yu J, Yang H. 2002.** A draft sequence of the rice genome (*Oryza sativa* L. ssp. *indica*). *Science* **296**:1937–1942 DOI [10.1126/science.1068037](https://doi.org/10.1126/science.1068037).
- Zhang DY, Ali Z, Wang CB, Xu L, Yi JX, Xu ZL, Liu XQ, He XL, Huang YH, Khan IA. 2013.** Genome-wide sequence characterization and expression analysis of major intrinsic proteins in soybean (*Glycine max* L.). *PLOS ONE* **8**:e56312 DOI [10.1371/journal.pone.0056312](https://doi.org/10.1371/journal.pone.0056312).
- Zhi Z, Gong J, Feng A, Xie G, Wang J, Mo Y, Yang L. 2015.** Genome-wide identification of rubber tree (*Hevea brasiliensis* Muell. Arg.) aquaporin genes and their response to ethephon stimulation in the laticifer, a rubber-producing tissue. *BMC Genomics* **16**:1–18 DOI [10.1186/s12864-015-2152-6](https://doi.org/10.1186/s12864-015-2152-6).
- Zhou S, Hu W, Deng X, Ma Z, Chen L, Huang C, Wang C, Wang J, He Y, Yang G. 2012.** Overexpression of the wheat aquaporin gene, TaAQP7, enhances drought tolerance in transgenic tobacco. *PLOS ONE* **7**:e52439 DOI [10.1371/journal.pone.0052439](https://doi.org/10.1371/journal.pone.0052439).
- Zou Z, Gong J, Huang Q, Mo Y, Yang L, Xie G. 2015.** Gene structures, evolution, classification and expression profiles of the aquaporin gene family in castor bean (*Ricinus communis* L.). *PLOS ONE* **10**:e0141022 DOI [10.1371/journal.pone.0141022](https://doi.org/10.1371/journal.pone.0141022).