

Complete genome sequence of an *Ehrlichia minasensis* strain isolated from cattle

Daniel M. Aguiar, João P. Araujo, Luciano Nakazato, Emilie Bard, Alejandro Cabezas Cruz

► To cite this version:

Daniel M. Aguiar, João P. Araujo, Luciano Nakazato, Emilie Bard, Alejandro Cabezas Cruz. Complete genome sequence of an *Ehrlichia minasensis* strain isolated from cattle. *Microbiology Resource Announcements*, American Society for Microbiology, 2019, 8 (15), 10.1128/MRA.00161-19. hal-02622602

HAL Id: hal-02622602

<https://hal.inrae.fr/hal-02622602>

Submitted on 28 Sep 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Complete Genome Sequence of an *Ehrlichia minasensis* Strain Isolated from Cattle

Daniel M. Aguiar,^a João P. Araujo, Jr.,^b Luciano Nakazato,^c Emilie Bard,^d Alejandro Cabezas-Cruz^e

^aLaboratory of Virology and Rickettsiosis, Veterinary Hospital (HOVET), Faculty of Veterinary Medicine (FAVET), Federal University of Mato Grosso (UFMT), Cuiabá, MT, Brazil

^bInstitute for Biotechnology, São Paulo State University (UNESP), Botucatu, SP, Brazil

^cLaboratory of Molecular Biology, Veterinary Hospital (HOVET), Faculty of Veterinary Medicine (FAVET), Federal University of Mato Grosso (UFMT), Cuiabá, MT, Brazil

^dEPIA, INRA, VetAgro Sup, Saint Genès Champanelle, France

^eUMR BIPAR, INRA, ANSES, École Nationale Vétérinaire d'Alfort, Université Paris-Est, Maisons-Alfort, France

ABSTRACT *Ehrlichia minasensis* is a tick-borne pathogen affecting cattle, cervids, and dogs, and it is closely related to the monocytotropic pathogen *Ehrlichia canis*. Here, we announce the draft genome sequence of *Ehrlichia minasensis* strain Cuiabá, isolated from a naturally infected calf from Santo Antônio do Leverger, Mato Grosso, Brazil.

The genus *Ehrlichia* is composed of tick-borne obligate intracellular Gram-negative alphaproteobacteria of the family *Anaplasmataceae* that primarily infect dogs (*Ehrlichia canis*, *Ehrlichia chaffeensis*, and *Ehrlichia ewingii*), mice (*Ehrlichia muris*), and ruminants (*Ehrlichia ruminantium*) (1). Some *Ehrlichia* species, however, have also been found to infect humans (i.e., *E. canis*, *E. ruminantium*, *E. chaffeensis*, and *E. ewingii*) (2, 3). Recently, we reported the *in vitro* culture (4), molecular (5), and morphological (6) characterization of *Ehrlichia minasensis* strain UFMG-EV^T (7), a novel *Ehrlichia* species that appears to have evolved from a highly variable strain of *E. canis* (8). Another *E. minasensis* strain (UFMT-BV) was found to be pathogenic for cattle in Brazil (9). The geographical distribution of *E. minasensis* is not restricted to the American continent as previously thought (5, 9, 10), and recent reports found these bacteria in Corsica, France (11), Pakistan (12), Ethiopia (13), and Israel (14).

The 16S rRNA phylogenetic tree shows that *Ehrlichia* is a diverse genus (7), and despite recent improvements in bacterial genome sequencing technologies, only a few genomes are available for *Ehrlichia* species, including those of *E. chaffeensis*, *E. ruminantium*, *E. canis*, *E. muris*, and *E. minasensis* UFMG-EV^T (15). Here, we report the genome sequence of *E. minasensis* strain Cuiabá, isolated from whole blood of an 8-month-old naturally infected male Holstein calf. The bacteria were grown in DH82 cells. The sample was treated with 2.5 units of Benzonase (Sigma-Aldrich, MO, USA) and 2 units of Turbo DNase (Thermo Fisher Scientific, Waltham, MA, USA) following total DNA extraction using a QIAamp DNA kit (Qiagen, Hilden, Germany) and quantified using Qubit fluorometric quantification (Thermo Fisher Scientific). The library was made starting from 5 ng of DNA using the Nextera XT library preparation kit (Illumina, San Diego, CA, USA), and the mean insert size of the final library was 450 bp. The library was then sequenced with a NextSeq 550 system using sequencing by synthesis (SBS) chemistry (Illumina). Read quality was controlled using Illumina's cloud service BaseSpace. A total of 37 million reads passed the quality filter; reads averaged 150 bp in length and showed an average quality score above Q30 in more than 90% of the bases. *De novo* assembly was performed with all quality reads using SPAdes (16). Prior

Citation Aguiar DM, Araujo JP, Jr, Nakazato L, Bard E, Cabezas-Cruz A. 2019. Complete genome sequence of an *Ehrlichia minasensis* strain isolated from cattle. *Microbiol Resour Announc* 8:e00161-19. <https://doi.org/10.1128/MRA.00161-19>.

Editor Julie C. Dunning Hotopp, University of Maryland School of Medicine

Copyright © 2019 Aguiar et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Emilie Bard, emilie.bard@inra.fr, or Alejandro Cabezas-Cruz, cabezasalejandrocruz@gmail.com.

Received 13 February 2019

Accepted 18 March 2019

Published 11 April 2019

to assembly, the overlapping reads were merged with Fast Length Adjustment of SHort reads (FLASH) (17). The contigs were aligned with Mauve (18) to the *E. minasensis* UFMG-EV^T genome (GenBank accession number [CDGH00000000](https://doi.org/10.1093/nar/gkz000)). The assembly step returned 59 contigs and 1,361,734 bases with an N_{50} value of 694,769 and 29.72% G+C content. The genome annotation was run with Prokka (19) and revealed the presence of 1,020 genes, including 978 coding sequences (CDSs) and 42 RNA genes (rRNA, transfer-messenger RNA [tmRNA], and tRNA). Among these, 346 genes are related to proteins with enzymatic activity (with defined Enzyme Commission [EC] codes), 93 are involved in cellular process and signaling with 15 related to membrane proteins, 162 in information storage and processing with 38 related to replication, recombination, and repair, and 226 in metabolism with 56 related to energy production and conversion.

E. minasensis genomes provide an invaluable tool for studying the relation between “host shift” (i.e., the capacity of some pathogens to infect novel hosts) and vector promiscuity within the *Anaplasmataceae*. This idea is supported by two facts. First, while the common tick vector of *E. canis* is almost exclusively *Rhipicephalus sanguineus*, *E. minasensis* is mainly found in *Rhipicephalus microplus* (5, 20) but has been detected also in *Hyalomma marginatum* (11) and *Hyalomma anatolicum* (12). Second, while *E. canis* is mainly pathogenic for dogs, *E. minasensis* infects not only bovines (9, 10, 13) and cervids (21) but also dogs (14).

Data availability. The raw sequence reads have been submitted to the NCBI SRA under the accession number [PRJNA478569](https://www.ncbi.nlm.nih.gov/sra/PRJNA478569), and the whole-genome shotgun project of the *Ehrlichia minasensis* strain Cuiabá has been deposited in DDBJ/EMBL/GenBank under the accession number [QOHL00000000](https://www.ncbi.nlm.nih.gov/genbank/COHL00000000). Genome assembly data are available under the accession number [GCA_004181775](https://www.ncbi.nlm.nih.gov/genbank/GCA_004181775).

REFERENCES

- Saito TB, Walker DH. 2016. Ehrlichioses: an important One Health opportunity. *Vet Sci* 3:20. <https://doi.org/10.3390/vetsci3030020>.
- Perez M, Bodor M, Zhang C, Xiong Q, Rikihisa Y. 2006. Human infection with *Ehrlichia canis* accompanied by clinical signs in Venezuela. *Ann N Y Acad Sci* 1078:110–117. <https://doi.org/10.1196/annals.1374.016>.
- Thomas RJ, Dumler JS, Carlyon JA. 2009. Current management of human granulocytic anaplasmosis, human monocytic ehrlichiosis and *Ehrlichia ewingii* ehrlichiosis. *Expert Rev Anti Infect Ther* 7:709–722. <https://doi.org/10.1586/eri.09.44>.
- Zweygarth E, Schöl H, Lis K, Cabezas-Cruz A, Thiel C, Silaghi C, Ribeiro MFB, Passos LMF. 2013. *In vitro* culture of a novel genotype of *Ehrlichia* sp. from Brazil. *Transbound Emerg Dis* 60 Suppl 2:86–92. <https://doi.org/10.1111/tbed.12122>.
- Cabezas-Cruz A, Zweygarth E, Ribeiro MFB, da Silveira JAG, de la Fuente J, Grubhoffer L, Valdés JJ, Passos LMF. 2012. New species of *Ehrlichia* isolated from *Rhipicephalus (Boophilus) microplus* shows an ortholog of the *E. canis* major immunogenic glycoprotein gp36 with a new sequence of tandem repeats. *Parasit Vectors* 5:291. <https://doi.org/10.1186/1756-3305-5-291>.
- Cabezas-Cruz A, Vancová M, Zweygarth E, Ribeiro MFB, Grubhoffer L, Passos LMF. 2013. Ultrastructure of *Ehrlichia mineirensis*, a new member of the *Ehrlichia* genus. *Vet Microbiol* 167:455–458. <https://doi.org/10.1016/j.vetmic.2013.08.001>.
- Cabezas-Cruz A, Zweygarth E, Vancova M, Broniszewska M, Grubhoffer L, Passos LMF, Ribeiro MFB, Alberdi P, de la Fuente J. 2016. *Ehrlichia minasensis* sp. nov., isolated from the tick *Rhipicephalus microplus*. *Int J Syst Evol Microbiol* 66:1426–1430. <https://doi.org/10.1099/ijsem.0.000895>.
- Cabezas-Cruz A, Valdés JJ, de la Fuente J. 2014. The glycoprotein TRP36 of *Ehrlichia* sp. UFMG-EV and related cattle pathogen *Ehrlichia* sp. UFMG-EV evolved from a highly variable clade of *E. canis* under adaptive diversifying selection. *Parasit Vectors* 7:584. <https://doi.org/10.1186/PREACCEPT-1737878488143835>.
- Aguiar DM, Ziliani TF, Zhang X, Melo ALT, Braga IA, Witter R, Freitas LC, Rondelli ALH, Luis MA, Sorte ECB, Jaune FW, Santarém VA, Horta MC, Pescador CA, Colodel EM, Soares HS, Pacheco RC, Onuma SSM, Labruna MB, McBride JW. 2014. A novel *Ehrlichia* genotype strain distinguished by the TRP36 gene naturally infects cattle in Brazil and causes clinical manifestations associated with ehrlichiosis. *Ticks Tick Borne Dis* 5:537–544. <https://doi.org/10.1016/j.ttbdis.2014.03.010>.
- Gajadhar AA, Lobanov V, Scandrett WB, Campbell J, Al-Adhami B. 2010. A novel *Ehrlichia* genotype detected in naturally infected cattle in North America. *Vet Parasitol* 173:324–329. <https://doi.org/10.1016/j.vetpar.2010.06.034>.
- Ciculli V, Masse S, Capai L, de Lamballerie X, Charrel R, Falchi A. 2019. First detection of *Ehrlichia minasensis* in *Hyalomma marginatum* ticks collected from cattle in Corsica, France. *Vet Med Sci*. <https://doi.org/10.1002/vms3.140>.
- Rehman A, Conraths FJ, Sauter-Louis C, Krücken J, Nijhof AM. 2019. Epidemiology of tick-borne pathogens in the semi-arid and the arid agro-ecological zones of Punjab province, Pakistan. *Transbound Emerg Dis* 66:526–536. <https://doi.org/10.1111/tbed.13059>.
- Hailemariam Z, Krücken J, Baumann M, Ahmed JS, Clausen P-H, Nijhof AM. 2017. Molecular detection of tick-borne pathogens in cattle from southwestern Ethiopia. *PLoS One* 12:e0188248. <https://doi.org/10.1371/journal.pone.0188248>.
- Thomson K, Yaaran T, Belshaw A, Curson L, Tisi L, Maurice S, Kiddle G. 2018. A new TaqMan method for the reliable diagnosis of *Ehrlichia* spp. in canine whole blood. *Parasit Vectors* 11:350. <https://doi.org/10.1186/s13071-018-2914-5>.
- Cabezas-Cruz A, Zweygarth E, Broniszewska M, Passos LMF, Ribeiro MFB, Manrique M, Tobes R, de la Fuente J. 2015. Complete genome sequence of *Ehrlichia mineirensis*, a novel organism closely related to *Ehrlichia canis* with a new host association. *Genome Announc* 3:e01450-14. <https://doi.org/10.1128/genomeA.01450-14>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Mago T, Salzberg SL. 2011. FLASH: Fast Length Adjustment of Short reads to improve genome assemblies. *Bioinformatics* 27:2957–2963. <https://doi.org/10.1093/bioinformatics/btr507>.
- Rissman AI, Mau B, Biehl BS, Darling AE, Glasner JD, Perna NT. 2009.

- Reordering contigs of draft genomes using the Mauve aligner. *Bioinformatics* 25:2071–2073. <https://doi.org/10.1093/bioinformatics/btp356>.
19. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.
 20. Carvalho ITS, Melo ALT, Freitas LC, Verçoza RV, Alves AS, Costa JS, Chitarra CS, Nakazato L, Dutra V, Pacheco RC, Aguiar DM. 2016. Minimum infection rate of *Ehrlichia minasensis* in *Rhipicephalus microplus* and *Amblyomma sculptum* ticks in Brazil. *Ticks Tick Borne Dis* 7:849–852. <https://doi.org/10.1016/j.ttbdis.2016.04.004>.
 21. Lobanov VA, Gajadhar AA, Al-Adhami B, Schwantje HM. 2012. Molecular study of free-ranging mule deer and white-tailed deer from British Columbia, Canada, for evidence of *Anaplasma* spp. and *Ehrlichia* spp. *Transbound Emerg Dis* 59:233–243. <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1865-1682.2011.01259.x>.