



## **Author correction: A genome scan for milk production traits in dairy goats reveals two new mutations in Dgat1 reducing milk fat content**

Pauline Martin, Isabelle Palhiere, Cyrielle Maroteau, Philippe Bardou, Kamila Canale-Tabet, Julien Sarry, Florent Woloszyn, Justine Bertrand-Michel, Ines Racke, Hueseyin Besir, et al.

### **► To cite this version:**

Pauline Martin, Isabelle Palhiere, Cyrielle Maroteau, Philippe Bardou, Kamila Canale-Tabet, et al.. Author correction: A genome scan for milk production traits in dairy goats reveals two new mutations in Dgat1 reducing milk fat content. Scientific Reports, 2017, 8, 10.1038/s41598-018-22118-x . hal-02622800

**HAL Id: hal-02622800**

**<https://hal.inrae.fr/hal-02622800>**

Submitted on 8 Dec 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

# SCIENTIFIC REPORTS

OPEN

## Author Correction: A genome scan for milk production traits in dairy goats reveals two new mutations in *Dgat1* reducing milk fat content

Pauline Martin<sup>1</sup>, Isabelle Palhière<sup>1</sup>, Cyrielle Maroteau<sup>1,2</sup>, Philippe Bardou<sup>1,3</sup>, Kamila Canale-Tabet<sup>1</sup>, Julien Sarry<sup>1</sup>, Florent Woloszyn<sup>1</sup>, Justine Bertrand-Michel<sup>4</sup>, Ines Racke<sup>5</sup>, Hüseyin Besir<sup>5</sup>, Rachel Rupp<sup>1</sup> & Gwenola Tosser-Klopp<sup>1</sup> 

Correction to: *Scientific Reports* <https://doi.org/10.1038/s41598-017-02052-0>, published online 12 May 2017

The Acknowledgements section in this Article is incomplete.

“The authors thank Pierre Martin and all the Capgenes breeding organizations for providing the data and their helpful contributions. This work was supported by grants from French organizations through “PhénoFinlait” (a research program including INRA, APIS-GENE, ALLICE (formerly UNCEIA), CAPGENES and FCEL) and EC (FP7/2007–2013), grant no. 245140, “3SR”, Sustainable Solutions for Small Ruminants (<http://www.3sr-breeding.eu/>). The authors thank all the partners involved. The authors also thank Cécile Albenne for her precious help in enzymology, Stéphane Fabre, Marc Teisser and Julie Demars for their involvement and help with softwares and molecular part, and Aurélie Tircazes for her help with genotype calling. Lipidomic analyses were performed on the Toulouse MetaToul-Lipidomique Core Facility (I2MC, Inserm 1048, Toulouse, France), MetaboHUB-ANR-11-INBS-0010.”

should read:

“The authors thank Pierre Martin and all the Capgenes breeding organizations for providing the data and their helpful contributions. This work was supported by grants from French organizations through “PhénoFinlait” (a research program including INRA, APIS-GENE, ALLICE (formerly UNCEIA), CAPGENES and FCEL) and EC (FP7/2007–2013), grant no. 245140, “3SR”, Sustainable Solutions for Small Ruminants (<http://www.3srbreeding.eu/>). The authors thank all the partners involved. The authors also thank Cécile Albenne for her precious help in enzymology, Stéphane Fabre, Marc Teisser and Julie Demars for their involvement and help with softwares and molecular part, and Aurélie Tircazes for her help with genotype calling. Lipidomic analyses were performed on the Toulouse MetaToul-Lipidomique Core Facility (I2MC, Inserm 1048, Toulouse, France), MetaboHUB-ANR-11-INBS-0010. Pauline Martin received a doctoral research grant from “Région Midi-Pyrénées”, Animal Genetics Department from INRA and GENOMCAP research program including INRA, APIS-GENE, ALLICE (formerly UNCEIA), CAPGENES and FCEL). Cyrielle Maroteau received a CIFRE grant from ALLICE”.

<sup>1</sup>GenPhySE, Université de Toulouse, INRA, INPT, ENVT, Castanet Tolosan, France. <sup>2</sup>Division of Molecular and Clinical Medicine, School of Medicine, University of Dundee, Ninewells Hospital and Medical School, Dundee, UK. <sup>3</sup>INRA, Sigénac, Castanet-Tolosan, France. <sup>4</sup>MetaToul-Lipidomic Core Facility, MetaboHUB, INSERM U 1048, Toulouse, France. <sup>5</sup>Protein Expression and Purification Core Facility, EMBL Heidelberg, Heidelberg, Germany. Rachel Rupp and Gwenola Tosser-Klopp contributed equally to this work. Correspondence and requests for materials should be addressed to G.T. (email: [gwenola.tosser@inra.fr](mailto:gwenola.tosser@inra.fr))



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018