



**HAL**  
open science

## First Report of Sclerotium stem rot caused by Sclerotium rolfsii on Stevia rebaudiana in southwestern France

Zoé Le Bihan, Jonathan Gaudin, Fanny Robledo-Garcia, Patrick Cosson,  
Cecile Hastoy, Dominique Rolin, Valerie Schurdi-Levraud

### ► To cite this version:

Zoé Le Bihan, Jonathan Gaudin, Fanny Robledo-Garcia, Patrick Cosson, Cecile Hastoy, et al.. First Report of Sclerotium stem rot caused by Sclerotium rolfsii on Stevia rebaudiana in southwestern France. *Plant Disease*, 2020, 104 (2), pp.584-584. 10.1094/PDIS-04-19-0696-PDN . hal-02623266

**HAL Id: hal-02623266**

**<https://hal.inrae.fr/hal-02623266>**

Submitted on 24 Feb 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## First Report of *Sclerotium* stem rot caused by *Athelium rolfsii* on *Stevia rebaudiana* in southwestern France

Z. Le Bihan, INRA, Université de Bordeaux, UMR 1332 Biologie du Fruit et Pathologie, Villenave d'Ornon, France and Oviatis SA, Lacaussade, France; J. Gaudin, INRA, UMR 1065 Santé et Agroécologie du Vignoble, ISVV, Villenave d'Ornon, France; F. Robledo-Garcia, INRA, UMR 1065 Santé et Agroécologie du Vignoble (SAVE), ISVV, Villenave d'Ornon, France ; P. Cosson, INRA, Université de Bordeaux, UMR 1332 Biologie du Fruit et Pathologie, Villenave d'Ornon, France; C. Hastoy, Oviatis SA, Lacaussade, France; D. Rolin INRA, Université de Bordeaux, UMR 1332 Biologie du Fruit et Pathologie 1332, Villenave d'Ornon, France and V. Schurdi-Levraud, INRA, Université de Bordeaux, UMR 1332 Biologie du Fruit et Pathologie 1332, Villenave d'Ornon, France.

*Stevia rebaudiana* is a perennial species that accumulates steviol glycosides in its leaves. These natural compounds produce an intense sweet taste but are non-caloric and are hence in demand by consumers to reduce daily sugar consumption. The cultivation of *S. rebaudiana* has recently become popular in France. In August 2017, disease symptoms similar to a sudden wilting were observed in a *S. rebaudiana* 0.5 ha commercial field in Sainte-Livrade (N44° 23' 48"; E0°35'13"), France. Symptoms first appeared on the aerial parts, resulting in wilting and drying of the entire plant. Additional symptoms included lesions on the stem at the collar base with abundant white mycelium and round-white to dark-brown small globoid sclerotia were observed on diseased plants. These symptoms were also observed on nearby pepper crops. The outbreaks of infection were increasing with time. Disease incidence was estimated at approximately 17%. To isolate the pathogen, the collar of one diseased plant was collected and incubated in moist chamber for 2 days at 24°C. Mycelia was removed from the diseased plant parts and placed on MAC (malt extract agar amended with 50 mg/liter chloramphenicol) and incubated at 22°C in a growth chamber with a 12 h photoperiod. After 7 days, immature sclerotia were abundantly formed, and after ten days small globoid, white sclerotia were approximately 1 to 3 mm diam. and became dark brown with age. The freshly isolated pathogenic fungus was examined microscopically. The white mycelia had clamp connection typical of that described for *A. rolfsii* (Mordue 1974). No sexual reproductive structures were observed. Genomic DNA of one isolate (17SCL\_STEV1) was extracted as described (Hastoy et al. 2019). The Large ribosomal subunit locus (LSU rRNA) was amplified and sequenced with primers LROR / LR5 (Stielow et al. 2015). The length of the amplicon was 884 bp (GenBank accession MK680087). Phylogenetic analysis was done with the neighbor-joining method in MEGA7. Isolate 17SCL\_STEV1 was identified as *A. rolfsii* and presented 99% sequence similarity with *A. rolfsii* isolate DGADY14 from *Capsicum annum*. To conduct pathogenicity testing, mycelium was cultivated on MA (Malt-Agar) medium incubated at 22°C for 6 days in the growth chamber. Colonized MA plugs (0.5 diam.) were removed and four were applied to the stems of 5 two-month-old *S. rebaudiana* cv. Shoutian III plants following injury by a scalpel. For the control, stems were wounded and inoculated with sterile MA plugs with the same method. Plants were kept in a growth chamber at 22°C with a 12-h photoperiod and 85% humidity. After 2 days, lesions appeared on the collar. White mycelium and incipient sclerotia were observed on the lesion and soil surface and turned into typical brownish sclerotia. Inoculated plants wilted and died after 5 days. Three re-isolation were performed once by sampling mycelium from symptomatic plants. Isolates were confirmed as *A. rolfsii* based on morphological and molecular characteristics. Although there are previous reports of *A. rolfsii* on *S. rebaudiana* in India (Kamalakaran et al. 2007), in the USA (Koehler and Shew 2014), Italy (Carrieri et al. 2015) and Greece (Chatzivassiliou et al. 2015), to the best of our knowledge, this is the first report of *A. rolfsii* causing *Sclerotium* stem rot of *S. rebaudiana* in France. This is of primary importance given the damage to yield and the difficulty of controlling *A. rolfsii* (Punja 1985).

## References

- Carrieri, R., Cozzolino, E., Tarantino, P., Cerrato, D., Lahoz, E., 2015. First Report of Southern Blight on Candyleaf (*Stevia rebaudiana*) Caused by *Sclerotium rolfsii* in Italy. *Plant Disease* 100, 220. <https://doi.org/10.1094/PDIS-03-15-0324-PDN>
- Chatzivassiliou, E.K., Giakountis, A., Testa, A., Kienle, U., Jungbluth, T., 2015. Natural Infection of *Stevia rebaudiana* by Cucumber mosaic virus in Spain and by *Sclerotium rolfsii* in Greece. *Plant Disease* 100, 1029–1029. <https://doi.org/10.1094/PDIS-10-15-1149-PDN>
- Hastoy, C., Le Bihan, Z., Gaudin, J., Cosson, P., Rolin, D., Schurdi-Levraud, V., 2019. First report of *Septoria* sp. infecting *Stevia rebaudiana* in France and screening of *Stevia rebaudiana* genotypes for host resistance. *Plant Disease*. <https://doi.org/10.1094/PDIS-10-18-1747-RE>
- Kamalakannan, A., Valluvarparidasan, V., Chitra, K., Rajeswari, E., Salaheddin, K., Ladhakshmi, D., Chandrasekaran, A., 2007. First report of root rot of stevia caused by *Sclerotium rolfsii* in India. *Plant Pathology* 56.
- Koehler, A., Shew, H., 2014. First Report of Stem and Root Rot of *Stevia* Caused by *Sclerotium rolfsii* in North Carolina. *Plant Disease* 98, 1005–1005. <https://doi.org/10.1094/PDIS-12-13-1238-PDN>
- Mordue, E. 1974. Page 1 in: *CMI Descriptions of Pathogenic Fungi and Bacteria*, No. 410. Commonwealth Mycological Institute, Kew, U.K.
- Punja, Z.K., 1985. The Biology, Ecology, and Control of *Sclerotium Rolfsii*. *Annu. Rev. Phytopathol.* 23, 97–127. <https://doi.org/10.1146/annurev.py.23.090185.000525>
- Stielow, J.B., Lévesque, C.A., Seifert, K.A., Meyer, W., Irinyi, L., Smits, D., Renfurm, R., Verkley, G.J.M., Groenewald, M., Chaduli, D., Lomascolo, A., Welti, S., Lesage-Meessen, L., Favel, A., Al-Hatmi, A.M.S., Damm, U., Yilmaz, N., Houbraeken, J., Lombard, L., Quaedvlieg, W., Binder, M., Vaas, L.A.I., Vu, D., Yurkov, A., Begerow, D., Roehl, O., Guerreiro, M., Fonseca, A., Samerpitak, K., van Diepeningen, A.D., Dolatabadi, S., Moreno, L.F., Casaregola, S., Mallet, S., Jacques, N., Roscini, L., Egidi, E., Bizet, C., Garcia-Hermoso, D., Martín, M.P., Deng, S., Groenewald, J.Z., Boekhout, T., de Beer, Z.W., Barnes, I., Duong, T.A., Wingfield, M.J., de Hoog, G.S., Crous, P.W., Lewis, C.T., Hambleton, S., Moussa, T.A.A., Al-Zahrani, H.S., Almaghrabi, O.A., Louis-Seize, G., Assabgui, R., McCormick, W., Omer, G., Dukik, K., Cardinali, G., Eberhardt, U., de Vries, M., Robert, V., 2015. One fungus, which genes? Development and assessment of universal primers for potential secondary fungal DNA barcodes. *Pers - Int Mycol J* 35, 242–263. <https://doi.org/10.3767/003158515X689135>

e-Xtra 1

Corresponding author: [valerie.schurdi-levraud@inra.fr](mailto:valerie.schurdi-levraud@inra.fr)

e-Xtra 2

Z. Le Bihan and C. Hastoy were supported by ANRT n°2017/0489 and ANRT n°2014/0915 funding and Oviatis SA, France. The Nouvelle-Aquitaine Region supported the work through specific CIFRE funding. Work was partly supported by the Research Federation on Integrative Biology and Ecology (Bordeaux University, France).

e-Xtra 3



Figure S1. Sclerotium rot on *Stevia rebaudiana* and mycological characteristics of the pathogenic fungus, *Athelia rolfsii*. A and B typical symptoms of plant wilting and drying in the field; C white mycelium formed on the collar of infected plant from the field after 1 day of incubation in a moist chamber; D colony of *A. rolfsii* with mycelial mat and sclerotia grown on MAC medium after 12 days of incubation; E clamp connections structure formed on the fungus hyphae (arrow); F symptoms (wilt) and signs (white mycelium on the stem and near the soil line) appeared 2 days after artificial inoculation (left; control on the right).



Figure S2. Phylogenetic tree of isolate 17SCL\_STEV1 and related sequences from pepper and from NCBI were constructed by MEGA7. Branch stability was estimated by bootstrapping with 1000 replicates, support values are shown at the branch points. The analysis involved 16 nucleotide sequences. All ambiguous positions were removed for each sequence pair. There were a total of 446 positions in the final dataset.