Variability of free and glycosylated volatiles from strawberries destined for the fresh market and for processing, assessed using direct enzymatic hydrolysis

Stéphane Gaborieau, Aurélie Cendres, David Page, Christian Ginies,
Catherine M.G.C. Renard

► To cite this version:

Stéphane Gaborieau, Aurélie Cendres, David Page, Christian Ginies, Catherine M.G.C. Renard. Variability of free and glycosylated volatiles from strawberries destined for the fresh market and for processing, assessed using direct enzymatic hydrolysis. *LWT - Food Science and Technology*, 2018, 98, pp.187-196. 10.1016/j.lwt.2018.08.026 . hal-02623748

HAL Id: hal-02623748

<https://hal.inrae.fr/hal-02623748>

Submitted on 26 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Variability of free and glycosylated volatiles from strawberries destined for the fresh market and for processing, assessed using direct enzymatic hydrolysis

Stephane Gaborieau, Aurelie Cendres, David Page, Christian Ginies, Catherine M.G.C. Renard

PII: S0023-6438(18)30679-0

DOI: [10.1016/j.lwt.2018.08.026](https://doi.org/10.1016/j.lwt.2018.08.026)

Reference: YFSTL 7338

To appear in: *LWT - Food Science and Technology*

Received Date: 1 February 2018

Revised Date: 26 June 2018

Accepted Date: 13 August 2018

Please cite this article as: Gaborieau, S., Cendres, A., Page, D., Ginies, C., Renard, C.M.G.C., Variability of free and glycosylated volatiles from strawberries destined for the fresh market and for processing, assessed using direct enzymatic hydrolysis, *LWT - Food Science and Technology* (2018), doi: 10.1016/j.lwt.2018.08.026.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Comment citer ce document :

Gaborieau, S. (Auteur de correspondance), Cendres, A., Page, D., Ginies, C., Renard, C. (2018). Variability of free and glycosylated volatiles from strawberries destined for the fresh market and for processing, assessed using direct enzymatic hydrolysis. *LWT - Food Science and Technology*, 98, 187-196. . DOI : 10.1016/j.lwt.2018.08.026

Title: Variability of free and glycosylated volatiles from strawberries destined for the fresh market and for processing, assessed using direct enzymatic hydrolysis

Authors:

Stephane Gaborieau^{a,b}, Aurelie Cendres^b, David Page^a, Christian Ginies^a, Catherine M.G.C. Renard^a

^a UMR408 Sécurité et Qualité des Produits d'Origine Végétale, INRA, Avignon University F-84000 Avignon, France

^b Atelier du fruit, 7 rue des Pommerets, 21 600 Longvic-FRANCE

Corresponding author :

Mr. Stephane Gaborieau

INRA Centre de recherche PACA, Domaine Saint-Paul, Site Agroparc

228 route de l'Aérodrome

CS 40509 F-84914 Avignon Cedex 9 France

Tel.: +33603861398

E-mail: stephane.gaborieau@inra.fr

Abstract: Free- and glycosylated-volatile profiles of 14 strawberry varieties, 9 for industrial processing ('Darselect', 'Clery', 'Honey', 'Honeoye', 'Siabel', 'FCMO060', 'Fraise19', 2 'Senga Sangana') and 5 for fresh market ('Gariguet', 'Charlotte', 'CIR121', 2 'Clery' (full-field and hydroponic)), were compared. All volatiles were analysed by GC-MS. Volatiles from glycosides were first released by direct enzymatic hydrolysis. The extraction method was optimised for furaneol, a key component of strawberry aroma. More than 60 volatile compounds were identified, the most abundant being butyl acetate (average: 17 mg/kg), furaneol (average: 2 mg/kg) and free hexanoic acid (average: 3 mg/kg). Free-volatile profiles showed a split between fresh market strawberries, distinguished by esters and carbonyl molecules like isobutyl acetate or hexanal, and strawberries for processing, distinguished by molecule like 3-penten-2-one and 1-butanol. The three 'Clery' profiles were different notably in their hexanal, 4-vinylguaiacol and 3-penten-2-one concentrations. The glycosylated volatile profiles were similar among most strawberry varieties with, as major glycosylated volatiles, hexanoic acid (average: 1.7 mg/kg), benzyl alcohol (average: 0.5 mg/kg), gamma-decalactone (average: 0.5 mg/kg) and coumaran (average: 2.5 mg/kg). The potential for volatile enhancement by deconjugation was different. Potentially fresh market strawberries had a volatile increase of 6% against 50% for strawberries for processing.

Keywords: glycoside, glycosidase, GC-MS, aroma, *Fragaria x ananassa*

1. Introduction

Over 4.3 millions of tons of strawberries are produced each year, out of which 80% to 90% are destined for the fresh market (López-Aranda et al., 2011) and the rest for processing. Strawberries are found in juices, in jams and in semi-processed food products, which can be incorporated in yogurt, ice cream or pastry. Strawberry is also the most appreciated fruit by French consumers and 92% of them (Bhat, Geppert, Funken, & Stamminger, 2015) like strawberries because of their specific aroma. Therefore, aromatic potential of strawberries for processing interest for future process optimisation to preserve or enhance the aromatic power of strawberries.

The molecules most impacting strawberry aroma (Nuzzi, Lo Scalzo, Testoni, & Rizzolo, 2008) are furaneol, mesifuran, ethyl butanoate, ethyl hexanoate, β -linalool and hexanal. They are responsible for “caramel”, “apple-like”, “fruity” and “green” notes, respectively, in strawberries. Several molecules are not ubiquitous but are important for the aroma of strawberries in which they are found. This is the case of gamma-decalactone with “peach-like” note found in ‘Elsanta’ or ‘Senga Sangana’ varieties or trans-nerolidol with “floral” note found in ‘Elvira’ or ‘Pandera’ varieties (Larsen, Poll, & Olsen, 1992). Despite its complexity, scientists agree that furaneol is the key molecule (Wein et al., 2002; Ulrich et al., 1997). This furanone is obtained by several successive enzymatic transformations of D-fructose-1,6-diphosphate, the last one of which is a reduction by a quinone oxidoreductase (Raab, 2006).

Various parameters can influence the aroma of strawberries (Perez, Rios, Sanz, & Olias, 1992; Miszczak, Forney, & Prange, 1995). The first one is genetic. Indeed Aharoni (2004) compared the aroma profile of wild strawberries (*Fragaria vesca*) and cultivated strawberries (*Fragaria x ananassa*). Nerolidol dominates among terpenes in cultivated strawberries, while, in wild strawberries, monoterpenes like β -myrcene and α -pinene,

responsible of the characteristic musky aroma of this species, are more prevalent (Bianchi, Lovazzano, Lanubile, & Marocco, 2014). Maturity also impacts the strawberry aroma: Forney, Kalt, & Jordan (2000) showed an increase of methyl butanoate and methyl hexanoate with maturity and Darbellay, Luisier, Villettaz, Amadò, & Azodanlou (2004) a decrease of (Z)-3-hexenyl acetate and an increase of ethyl hexanoate during ripening in 'Carezza', 'Darselect' and 'Marmolada' varieties. The atmosphere during packaging also impacts volatile production: Nielsen & Leufven (2008) showed an increase of ethyl acetate (fermented molecules) with CO₂-enriched atmosphere in the 'Korona' variety.

However not all volatiles exist as such in the intact raw fruit. Some are formed during eating or processing by endogenous enzymes acting on precursors. A well-known reaction cascade is that of lipoxygenase leading to "green" notes from fatty acids (Pérez, Sanz, Olías, & Olías, 1999). Another major route is that of glycosides, where the flavour-bearing molecules (aglycones which become volatiles if liberated) are bound to sugars for storage. The sugar moiety can be a mono- or disaccharide and more rarely a trisaccharide, the first aglycone-bound sugar always being β -D-glucose (Sarry & Gunata, 2004). These precursors are present in all fruits, including strawberries. They were studied by Stahl-Biskup, Intert, Holthuijzen, Stengele, & Schulz (1993) in apples, apricots, blackberries, mangoes, plums and tomatoes or Gunata, Bayonove, Baumes, & Cordonnier (1985) in grapes. In strawberries, furaneol, mesifuran, benzyl alcohol, 2-phenylethyl alcohol, hydroxylinalool, 2-methylbutanoic acid or hexanoic acid can be found as glycosides (Groyne, Lognay, & Marlier, 1999). For example furaneol is predominantly conjugated to malonyl-D-glucose (Raab, 2006).

These glycosides are traditionally identified and quantified after isolation by solid phase extraction (SPE) (Young & Paterson, 1995). SPE consists in injecting the fruit extract on an XAD-2 or a C18-column, which binds the aglycone moieties of the glycosides. The

sugars and free volatiles are first eluted by water and dichloromethane, respectively; the glycosides are thereafter eluted with another solvent (generally methanol) (Ubeda et al., 2012; Roscher, Herderich, Steffen, Schreier, & Schwab, 1996). This method allows to eliminate matrix interactions. However, it is not adapted to compare free volatiles to bound volatiles because of the discrimination that is generated by affinity extraction, and because the natural evolution of volatiles in fruit matrices is not considered. Indeed, Hampel, Robinson, Johnson, & Ebeler (2014) showed that this pre-treatment leads to a loss and a discrimination of many compounds, and can not be used to compare free volatile to glycoconjugated volatiles. They advised direct hydrolysis to compare quantitatively free and bound volatiles.

The aim of this study is to establish direct hydrolysis by AR2000 glycosidase as a method to compare aroma profiles and aromatic potentials of processing and fresh market strawberries. Fourteen strawberry batches, 9 for processing (obtained from industrial sources) and 5 for fresh market (obtained from CIREF (Centre Interrégional de Recherche et d'Expérimentation de la Fraise, Douville (France))) were characterised. Direct enzymatic hydrolysis was combined with GC-MS analysis and descriptive statistics to evaluate aromatic potential of these strawberries.

2. Material & method

2.1. Plant material

Strawberries for processing were commercial samples harvested in 2015, individually quick frozen (IQF), and stored at -20°C. Among them, 'FCMO' and 'Fraise19' are a mix of several strawberry varieties used by the French food industry. Fresh market strawberries were from CIREF, harvested in 2016, and were frozen at -20°C (Table 1).

2.2. Reagents and solvents

Pentane (Normapur purity $\geq 99.9\%$) and Ethyl acetate (purity $\geq 99.9\%$) were purchased from VWR International (Darmstadt, Germany). Enzymatic kits to measure D-glucose/fructose (ref: E 0139106), D-glucose (ref: E 0716251), L-malic acid (ref: E 0139 068) and citric acid (ref: E 0139076) were from R-Biopharm (Darmstadt, Germany). Hydromatrix™ (diatomaceous earth) was from Agilent Technologies (Les Ulys, France). The internal standard was 4-nonanol (purity > 97 %) from Merck Schuchardt (Hohenbrunn, Germany). The AR2000 enzyme to measure the aromatic potential came from Oenobrand (Montferrier-sur-Lez, France) as freeze-dried powder (6.17 nkat/mg of powder). For retention index (RI) calculations in GC-MS, a solution of C7-C30 saturated alkane 1000 $\mu\text{g/mL}$ (ref: 49451-U) from Supelco (Bellefonte, Pennsylvania, USA) was used.

2.3. Fruit characterisation

2.3.1. Brix and pH

Strawberries (about 100 g) were ground and used for measurement of Brix degree using a digital refractometer PR-101 from Agato (Tokyo, Japan) and of pH using pHmeter FE20 FireEasy from Mettler Toledo (Viroflay, France).

2.3.2 Sugars and Acids

Sugars (glucose, fructose, sucrose) and main organic acids (citric acid and malic acid) were quantified by colorimetric enzymatic measurements with kits for food analysis from R-Biopharm (Darmstadt, Germany) and expressed in g.kg^{-1} FW. The measurements were performed on a 96 well-microplate with a SAFAS (FLX-Xenius, SAFAS, Monaco) equipped with an automatic injection device (Garcia & Renard, 2014).

2.4 Characterization of Volatiles Compounds

2.4.1 Sample preparation:

A dozen of frozen strawberries (around 150 g) were mashed for 1 min in a blender (Waring-Nova, Grosseron, St. Herblain, France) (Figure 1). An aliquot (40 g) was sampled for enzymatic hydrolysis with 40 mg of AR2000. Sodium chloride was added to the rest of the mashed strawberries (75 g/100 g of strawberry) to inhibit endogenous enzymes. After another mixing, this was divided: one aliquot of 40 g used for incubation (hydrolysis negative control) and an aliquot for direct extraction (determination of free volatiles). After incubation (24 h, 40°C with stirring at 120 rpm), salt was added to the hydrolysed aliquots to have the same salting-out effect during extraction of volatile compounds by accelerated solvent extraction.

2.4.2 Accelerated solvent extraction (ASE)

Hydromatrix™ (13 g) and 4-nonanol (11.24 µg diluted in 100 µL of methanol), as internal standard, were mixed with 7.5 g of mashed fruit to obtain a homogeneous powder while inactivating enzymes. The powder was rapidly transferred to a 33 mL pressurized extraction cell for immediate extraction. The extractor was an ASE 200 system (Dionex, Sunnyvale, CA). Extraction conditions were as follows: pentane and ethyl acetate (1:2, mL:mL) as solvent, 10⁷ Pa, 40 °C, 5 min preheating then 5 min static incubation. The extract was concentrated to 1 mL by distillation under vacuum (300 Pa, 25°C, using a Multivapor R12, Büchi, Rungis, France) then by nitrogen flux prior to gas chromatography.

2.4.3 GC–MS

Samples (1 µL) were injected into a GC–MS system (Trace1300-ISQ LT; Thermo scientific, USA) equipped with a TG-WAXMS capillary column [30 m, 0.25 mm i.d., 0.5 µm film thickness] (Thermo scientific).

Injection was in splitless mode at 250 °C. The carrier gas was helium with a constant flow of 1.2 mL.min⁻¹. Oven temperature program was 35 °C for 2 min ramped at 5°C.min⁻¹ to 230°C

then held for 5 min. Mass spectra were obtained by electron ionization at 70 eV, with scanning from m/z 35 to 250 at 2 scans. s^{-1} .

Volatile levels were expressed in micrograms per kilograms of juice (ppb) in 4-nonanol equivalent. Aromatic potential was calculated by difference between hydrolysed samples and incubated control samples. This aromatic potential was compared to the initial free volatile profile to observe molecules increase or decrease. Aromatic potential corresponds to glycoconjugates released by AR2000 and the evolution of these molecules and original free volatiles under action of endogenous enzymes.

Data were collected with GC-MS Solution software Chromeleon 7.2 and, when it was possible, the major compounds were identified by their retention index and their mass spectra using the mass spectral database NIST 14 (US National Institute of Standards and Technology (NIST), Gaithersburg, MD, USA)].

2.5 Statistics

For each batch, three samples of twelve strawberries each were analyzed. Principal Component Analysis (PCA) was performed with XLSTAT software (Addinsoft, France) to compare the 14 strawberry varieties with quantified volatiles as variables. Correlation factor was calculated by Pearson's method.

3. Results

3.1 General characteristics of the strawberries

Global characteristics of the strawberry are summarized in Table 1. The average pH was 3.5 and the average Brix degree was between 8~9. This is coherent with previous studies (Schwieterman et al., 2014). The main acid was citric acid but with a high proportion (~30%) of malic acid. Acid concentrations were similar among all strawberries (fresh market or for

processing). Glucose, fructose and sucrose are present in similar proportions (around 30% for glucose, 45% for fructose and 25% for sucrose) among the strawberries for processing while table strawberries had no sucrose but had higher fructose and glucose concentrations. In strawberries, sucrose does not vary during ripening (Sturm, Koron, & Stampar, 2003) so this difference can not be explained by a simple difference of maturity. Interestingly, the three Clery samples had similar pH but different Brix degree (7.8 for Clery; 6.9 for Clery PC; 8.6 for Clery HS) with an absence of sucrose for Clery PC and Clery HS from CIREF.

3.2 Optimisation of volatile extraction

The extraction solvent was optimized for furaneol, being a key molecule for strawberry aroma. In a pre-test, a synthetic solution of furaneol (0.5 g/L) was extracted by n-pentane/dichloromethane (2:1 mL:mL), dichloromethane or n-pentane/ethyl acetate (1:2 mL:mL). One millilitre of this standard solution was extracted with 1 mL of solvent three times and extracts were injected after concentration. Extraction yields were of 27% with n-pentane/dichloromethane, 65% with dichloromethane and 90% with n-pentane/ethyl acetate, which was therefore retained as solvent for all subsequent extractions. This comparison was completed by ASE on strawberry purees and out of 34 quantified compounds (Figure 2), 30 compounds had a better extraction yield with n-pentane/ethyl acetate (1:2 mL:mL) compared to 4 compounds (isobutyl acetate, 3-penten-2-ol, acetoin and 4-methylpentan-2-ol) for dichloromethane. N-pentane/dichloromethane (2:1 mL:mL) seemed to be the worst solvent for strawberry volatile extraction. N-pentane/ethyl acetate was thus selected to extract volatiles from strawberries for this study.

3.3 Identification of volatiles

More than 60 volatile compounds were detected and identified (Table 2). Most representative compounds of a strawberry aroma were found: furaneol, mesifuran, ethyl butanoate, gamma-decalactone, β -linalool, trans-nerolidol, butyl acetate, ethyl hexanoate, butanoic acid, isobutanoic acid and hexanoic acid (Larsen et al., 1992 ; Ménager, Jost, & Aubert, 2004 ; Schwieterman et al., 2014 ; Ubeda, Callejón, Troncoso, Morales, & Garcia-Parrilla, 2014 ; Lambert, Demazeau, Largeteau, & Bouvier, 1999 ; Darbellay et al., 2004). Some molecules, like 2-phenylethyl alcohol (rose aroma) or 1-terpineol (citrus aroma), were only detected after glycoconjugate release. Cinnamic acid, another characteristic molecule of strawberry, was also found but it had a variability too large to be considered in interpreting the results. In the same triplicate, it can be absent or present in a quantity higher than 10 mg/kg.

3.4 Free volatiles profile

Free volatiles profiles (Suppl. data) were analyzed by PCA (Figure 3) and their repartition by chemical class is presented in Table 3. The sample map (Figure 3.a.) showed two groups of samples. The first principal component clearly separated strawberries for processing from fresh market strawberries while the second principal component differentiated 'Darselect' from all others. Each initial batch was distinct from the others except 'Fraise19' and 'FCM0', which were co-located. These two "varieties" come from a mix used by food industry, which may explain the similitudes between them. The correlation circle (Figure 3.b.) indicated that strawberries for processing are differentiated by volatiles like 2,3-butanediol, pantolactone, acetoin, 3-phenylpropyl alcohol or butanoic acid. Molecules like benzyl alcohol, caprolactones, octanoic acid and 5-(hydroxymethyl)-2-furfural with, respectively, floral, fruity, acidic and caramel notes, seemed responsible for the 'Darselect' differentiation. Fresh market strawberries were differentiated by carbonyls and esters like

hexanal, 3-methylbutyl acetate, isobutyl acetate, 2-hexanone. Again the three 'Clery' batches were distinct, indicating that their volatile profile would not depend only on genetics but also on cultivation method and pedoclimatic conditions.

The variety 'Gariguette' was distinguished by methyl hexanoate, 2-methylbutanoic acid and isobutyric acid.

Fresh market strawberries (Table 3) had on average three times as much volatile compounds as strawberries for processing, and esters as the most abundant molecules, whereas strawberries for processing were rich in acids, except for 'Darselect,' which was rich in esters.

3.5 Glycoconjugated volatile profile

Glycoconjugated volatile profiles (Suppl. data) were analyzed by PCA (Figure 4) and their repartition by chemical class is presented in Table 4.

Alcohols and acids were preponderantly present as glycoconjugates rather than free volatiles. They represent on average 50% of the glycosides. 'Senga Sangana' (Bulgaria) had the highest glycoside concentration and also the highest concentration of glycoconjugated ketones. The highest concentration of glycoconjugated furans, of which furaneol glycoside was the most abundant, was found in 'Siabel' (45% of total glycoconjugates). 'Darselect' was also different from other varieties in glycoconjugates, as it had a very low concentration of coumaran (in "other compounds") and a high level of lactones (25% of total glycoconjugates).

The PCA sample map (Figure 4.a) revealed another trend. All strawberries, except 'Darselect', 'Senga Sangana' (Bulgaria) and 'Gariguette' (but this last with very high variability), were clustered in the center of the PCA sample map, indicating that their glycoconjugated pools were qualitatively similar. 'Darselect' differed in having higher concentrations of 1-terpineol, gamma-caprolactone and ethyl 3-hydroxybutanoate than the

others. ‘Senga Sangana’ (Bulgaria) was differentiated by 2- and 3-hexenoic acid, 1,2-cyclopentanedione and phenol. Finally, 2-hexanone, propanoic acid, acetoin, 2-hydroxyfuraneol and isobutanoic acid were glycoconjugates specific to ‘Gariguette’.

Although strawberries for processing and fresh market strawberries had generally similar glycoconjugate profiles, the ratio “glycoconjugates/volatile compounds” was higher in strawberries for processing. Indeed, strawberries for fresh market had an average of 6% of potential volatile increase, except for “Gariguette” (36%), as opposed to strawberries for processing, which had an average volatile increase of 50%. The highest increase (87 %) was found for “Senga Sangana” (Bulgaria). So strawberries for processing seem to have a higher aromatic potential.

4 Discussion

4.1 Methodology

Glycosides are generally quantified using a SPE pre-treatment but this separation technique, by affinity, may not be optimal to assess quantitatively free volatiles and glycosides. By this method, Roscher, Koch, Herderich, Schreier, & Schwab (1997) estimated the glycosylated furaneol to be between 66% and 750% of the free furaneol in strawberries, whereas in this study it represented on average 14% of free furaneol (except in ‘Gariguette’ where bound furaneol was more than twice the amount of free furaneol). Hampel et al. (2014) confirmed obtaining differing results depending on the method with many molecules, like for 1-nonanol in ‘Cabernet Sauvignon’, which is multiplied by approximately 7 after direct hydrolysis and by more than 22 using the SPE method. The differences are also qualitative. They identified 95 new compounds using direct hydrolysis by glycosidases, to be compared to only 67 using the SPE method in the Chardonnay grapes. Furthermore, they observed that 24 volatiles are significantly increased and 5 decreased using direct hydrolysis as opposed to

SPE, which showed an increase for 17 compounds and a decrease for 13 compounds. Hampel et al. (2014) assumed that glycosides of terpenes are more retained on SPE columns, which is confirmed by Baek & Cadwallader (1999). Hampel et al. (2014) have a similar result on melon. Voirin's thesis (1990) confirms these observations on grape. He obtained an SPE yield of 15% for glycosides with phenyl-type aglycone and 70% to 100% for glycosides of terpene aglycones.

4.2 Volatile molecules of strawberries

Although furaneol is the key molecule for strawberry aroma, it was not qualitatively discriminant to compare volatile profile, as expected because it is ubiquitous in strawberries. This molecule is already found when strawberry volatiles are analyzed by liquid extraction (Larsen et al., 1992; Lambert et al., 1999) and sometimes found with solid phase microextraction (SPME)/headspace (Darbellay et al., 2004) which confirms this conclusion. The comparisons have to rely on other molecules.

In strawberries, the 2-methylbutanoate and 2-methylbutyl esters are present in all reference articles but only 3-methylbutyl acetate and 2-methylbutanoic acid were found here (as identified by their retention index and MS spectra). These molecules come from two different amino acids, L-leucine (for 3-methylbutyl/butanoate) and L-isoleucine (for 2-methylbutyl/butanoate) by the same pathway (Pérez, Olías, Luaces, & Sanz, 2002). The nature of amino acid supplies could explain the absence or presence of 2-methyl and 3-methyl butyl compounds. The C6 carbonyls 3-hexenal and 2-hexenal with their green odor types, other important molecules of strawberry aroma, were absent. This was expected in the aliquots which were salted during mashing as this treatment aims to inhibit the endogenous enzymes of the lipoxygenase (LOX) pathway. This absence could also be explained by their further degradation to hexenol by action of alcohol deshydrogenase (ADH) (Speirs et al.,

1998) or hexenoic acid by oxidation in AR2000-treated samples. Indeed, although 2- and 3-hexenal were not found, 2- and 3-hexenoic acid (quantifiable) and 3-hexenol (not quantifiable) were present. The identified volatiles had mainly fruity or floral odor type. These differences could arise from extraction with ethyl acetate/n-pentane (2:1), which is more polar than dichloromethane (the usual solvent) and the long incubation time (overnight) allowing chemical degradation. Between control samples with and without incubation, decreases of molecules like 2-methylbutanoic acid and hexanoic acid and increases of molecules like 2-hexenoic acid and 2-hydroxy-4-pyranone were observed and some molecules became undetectable like ethyl cinnamate and ethyl hexanoate.

4.3 Comparison of industrial and fresh-market strawberries

Free volatiles

The differentiation between industrial and fresh market strawberry aroma profiles may explain their different uses, resulting from distinct selection goals. Strawberries for processing need to keep their texture and color after treatment especially in dairy products. Fresh market strawberries, which are produced to be eaten as-is, need to have a good instant aromatic quality to attract consumers. This is reflected in the present results by generally higher contents of free volatiles in the fresh-market strawberries. Zorrilla-Fontanesi et al., (2012) studied the natural genetic variability of aroma in strawberry. They compared genomes of plants stem from 2 parents and associated them to aroma differences. This genetic approach probably could be interesting to complete this comparison between strawberries for process and for fresh market. An interesting point was that the 3 studied 'Clery' have different volatiles profiles. These differences can not be explained only by genetics especially considering that the two samples from CIREF come from the same cultivar and were grown at the same location. The only difference was that one sample was in full-field and the other in

hydroponics. The cultivation method also appeared to have a non-negligible effect on aroma profile.

Glycoconjugated volatiles

However limited differences were observed between fresh-market and industrial strawberries in the case of glycoconjugated volatiles. Indeed previous studies (Ubeda et al., 2012; Groyne et al., 1999) found as main glycoconjugated volatiles in strawberries furaneol, benzyl alcohol, 2-phenylethyl alcohol, 4-vinylphenol, 4-vinylguaiacol and hexanoic acid. Results are consistent with literature for ‘Gariguettes’ where Lambert et al. (1999) found also methyl hexanoate, 2-methylbutanoic acid as well as hexanoic acid as main glycoconjugated volatiles. Except for 4-vinylphenol, all were also found in this study. The importance of glycoconjugates of benzyl alcohol and 2-phenylethyl alcohol in strawberries was confirmed by the low discriminant effect of these molecules in the glycoconjugates profiles. Among released aglycones some molecules are interesting for flavor thanks their low thresholds, like 2-phenylethyl alcohol (rose note) with a threshold of 4 µg/kg (4 parts per billion (ppb) (Fenaroli’s Handbook, 2010), furaneol (caramellic note) with a threshold of 0.03 µg/kg or γ-decalactone (peach note) with a threshold of 11 µg/kg. These increases show that deglycosylation could be a way to enrich natural aroma extracts.

5. Conclusion

Direct enzymatic hydrolysis allowed to reexamine the free to glycoconjugated volatile ratios in strawberries. Free volatiles were more abundant than glycoconjugated volatiles, especially in fresh market strawberries, and that the glycosylated/free volatiles ratio was higher in strawberries for processing. All tested strawberries had similar glycoconjugates profiles (except for “Darselect”, “Senga Sangana” (Bulgaria) and “Gariguettes”). Indeed, the

release of glycoconjugated volatiles would enhance aroma in a similar manner for all varieties. It can therefore be an interesting way to enhance the aromatic power of strawberry preparations in industry. Besides fresh market strawberries had a lower aromatic reserve, which indicates that strawberries for processing would be better candidates for future experiments on glycoside hydrolysis during process.

Acknowledgments

This work was supported by funds obtained from BPI (Banque pour l'Investissement) région Bourgogne-Franche Comté, région Provence Alpes Côte d'Azur, France, Europe, FEDER (fonds européen de développement régional), Grand-Dijon, and Grand-Avignon, in the form of Fond Unique Interministériel. The project name is Natarome+. The project is co-labeled by Terralia and Vitagora.

References

- Aharoni, A. (2004). Gain and Loss of Fruit Flavor Compounds Produced by Wild and Cultivated Strawberry Species. *The Plant Cell Online*, 16(11), 3110–3131. <https://doi.org/10.1105/tpc.104.023895>
- Baek, H. H., & Cadwallader, K. R. (1999). Contribution of Free and Glycosidically Bound Volatile Compounds to the Aroma of Muscadine Grape Juice. *Journal of Food Science*, 64(3), 441–444. <https://doi.org/10.1111/j.1365-2621.1999.tb15059.x>
- Bhat, R., Geppert, J., Funken, E., & Stamminger, R. (2015). Consumers Perceptions and Preference for Strawberries—A Case Study from Germany. *International Journal of Fruit Science*, 15(4), 405–424. <https://doi.org/10.1080/15538362.2015.1021408>
- Bianchi, G., Lovazzano, A., Lanubile, A., & Marocco, A. (2014). Aroma Quality of Fruits of Wild and Cultivated Strawberry (FRAGARIA SPP.) in Relation to the Flavour-Related Gene Expression. *Journal of Horticultural Research*, 22(1). <https://doi.org/10.2478/johr-2014-0009>
- Burdock, G. A., & Fenaroli, G. (2010). *Fenaroli's handbook of flavor ingredients* (6th ed). Boca Raton: CRC Press/Taylor & Francis Group.
- Darbellay, C., Luisier, J.-L., Villettaz, J.-C., Amadö, R., & Azodanlou, R. (2004). Changes in flavour and texture during the ripening of strawberries. *European Food Research and Technology*, 218(2), 167–172. <https://doi.org/10.1007/s00217-003-0822-0>
- Forney, C. F., Kalt, W., & Jordan, M. A. (2000). The composition of strawberry aroma is influenced by cultivar, maturity, and storage. *HortScience*, 35(6), 1022–1026.
- Garcia, C., & Renard, C. (2014). Validation des dosages enzymatiques des sucres (glucose, fructose, saccharose) et acides (acide citrique et malique) par un spectrophotomètre avec lecteur de microplaques. *Cahier des Techniques de l'INRA*, 81(1), 18.

- 399 Groyne, J., Lognay, G., & Marlier, M. (1999). Accumulation of glycosidically bound
400 compounds in *Fragaria* ananassa* cv. Elsanta fruits at various developmental stages.
401 *Biotechnologie, Agronomie, Société et Environnement= Biotechnology, Agronomy,*
402 *Society and Environment [= BASE], 3(1), 5–9.*
- 403 Gunata, Y. Z., Bayonove, C. L., Baumes, R. L., & Cordonnier, R. E. (1985). The aroma of
404 grapes I. Extraction and determination of free and glycosidically bound fractions of
405 some grape aroma components. *Journal of Chromatography A, 331, 83–90.*
406 [https://doi.org/10.1016/0021-9673\(85\)80009-1](https://doi.org/10.1016/0021-9673(85)80009-1)
- 407 Hampel, D., Robinson, A. L., Johnson, A. J., & Ebeler, S. E. (2014). Direct hydrolysis and
408 analysis of glycosidically bound aroma compounds in grapes and wines: comparison
409 of hydrolysis conditions and sample preparation methods: Glycoside analysis.
410 *Australian Journal of Grape and Wine Research, 20(3), 361–377.*
411 <https://doi.org/10.1111/ajgw.12087>
- 412 Lambert, Y., Demazeau, G., Largeteau, A., & Bouvier, J.-M. (1999). Changes in aromatic
413 volatile composition of strawberry after high pressure treatment. *Food Chemistry,*
414 *67(1), 7–16.* [https://doi.org/10.1016/S0308-8146\(99\)00084-9](https://doi.org/10.1016/S0308-8146(99)00084-9)
- 415 Larsen, M., Poll, L., & Olsen, C. E. (1992). Evaluation of the aroma composition of some
416 strawberry (*Fragaria ananassa* Duch) cultivars by use of odour threshold values.
417 *Zeitschrift für Lebensmittel-Untersuchung und -Forschung, 195(6), 536–539.*
418 <https://doi.org/10.1007/BF01204558>
- 419 López-Aranda, J. M., Soria, C., Santos, B. M., Miranda, L., Domínguez, P., & Medina-
420 Mínguez, J. J. (2011). Strawberry Production in Mild Climates of the World: A
421 Review of Current Cultivar Use. *International Journal of Fruit Science, 11(3), 232–*
422 *244.* <https://doi.org/10.1080/15538362.2011.608294>

- 423 Ménager, I., Jost, M., & Aubert, C. (2004). Changes in Physicochemical Characteristics and
 424 Volatile Constituents of Strawberry (Cv. Cigaline) during Maturation. *Journal of*
 425 *Agricultural and Food Chemistry*, 52(5), 1248–1254.
 426 <https://doi.org/10.1021/jf0350919>
- 427 Miszczak, A., Forney, C. F., & Prange, R. K. (1995). Development of Aroma volatiles and
 428 color during postharvest ripening of “Kent” strawberries. *Journal of the American*
 429 *Society for Horticultural Science*, 120(4), 650–655.
- 430 Nielsen, T., & Leufven, A. (2008). The effect of modified atmosphere packaging on the
 431 quality of Honeoye and Korona strawberries. *Food Chemistry*, 107(3), 1053–1063.
 432 <https://doi.org/10.1016/j.foodchem.2007.09.025>
- 433 Nuzzi, M., Lo Scalzo, R., Testoni, A., & Rizzolo, A. (2008). Evaluation of Fruit Aroma
 434 Quality: Comparison Between Gas Chromatography–Olfactometry (GC–O) and
 435 Odour Activity Value (OAV) Aroma Patterns of Strawberries. *Food Analytical*
 436 *Methods*, 1(4), 270–282. <https://doi.org/10.1007/s12161-008-9039-y>
- 437 Pérez, A. G., Olías, R., Luaces, P., & Sanz, C. (2002). Biosynthesis of Strawberry Aroma
 438 Compounds through Amino Acid Metabolism. *Journal of Agricultural and Food*
 439 *Chemistry*, 50(14), 4037–4042. <https://doi.org/10.1021/jf011465r>
- 440 Perez, A. G., Rios, J. J., Sanz, C., & Olías, J. M. (1992). Aroma components and free amino
 441 acids in strawberry variety Chandler during ripening. *Journal of Agricultural and*
 442 *Food Chemistry*, 40(11), 2232–2235. <https://doi.org/10.1021/jf00023a036>
- 443 Pérez, A. G., Sanz, C., Olías, R., & Olías, J. M. (1999). Lipoxygenase and Hydroperoxide
 444 Lyase Activities in Ripening Strawberry Fruits. *Journal of Agricultural and Food*
 445 *Chemistry*, 47(1), 249–253. <https://doi.org/10.1021/jf9807519>

- 446 Raab, T. (2006). FaQR, Required for the Biosynthesis of the Strawberry Flavor Compound 4-
 447 Hydroxy-2,5-Dimethyl-3(2H)-Furanone, Encodes an Enone Oxidoreductase. *THE*
 448 *PLANT CELL ONLINE*, 18(4), 1023–1037. <https://doi.org/10.1105/tpc.105.039784>
- 449 Roscher, R., Koch, H., Herderich, M., Schreier, P., & Schwab, W. (1997). Identification of
 450 2,5-dimethyl-4-hydroxy-3[2H]-furanone β -D-glucuronide as the major metabolite of a
 451 strawberry flavour constituent in humans. *Food and Chemical Toxicology*, 35(8), 777–
 452 782. [https://doi.org/10.1016/S0278-6915\(97\)00055-0](https://doi.org/10.1016/S0278-6915(97)00055-0)
- 453 Roscher, René, Herderich, M., Steffen, J.-P., Schreier, P., & Schwab, W. (1996). 2,5-
 454 Dimethyl-4-hydroxy-3[2H]-furanone 6'O-malonyl- β -D-glucopyranoside in strawberry
 455 fruits. *Phytochemistry*, 43(1), 155–159. [https://doi.org/10.1016/0031-9422\(96\)00216-6](https://doi.org/10.1016/0031-9422(96)00216-6)
- 456 Sarry, J., & Gunata, Z. (2004). Plant and microbial glycoside hydrolases: Volatile release
 457 from glycosidic aroma precursors. *Food Chemistry*, 87(4), 509–521.
 458 <https://doi.org/10.1016/j.foodchem.2004.01.003>
- 459 Schwieterman, M. L., Colquhoun, T. A., Jaworski, E. A., Bartoshuk, L. M., Gilbert, J. L.,
 460 Tieman, D. M., ... Clark, D. G. (2014). Strawberry Flavor: Diverse Chemical
 461 Compositions, a Seasonal Influence, and Effects on Sensory Perception. *PLoS ONE*,
 462 9(2), e88446. <https://doi.org/10.1371/journal.pone.0088446>
- 463 Speirs, J., Lee, E., Holt, K., Yong-Duk, K., Steele Scott, N., Loveys, B., & Schuch, W.
 464 (1998). Genetic Manipulation of Alcohol Dehydrogenase Levels in Ripening Tomato
 465 Fruit Affects the Balance of Some Flavor Aldehydes and Alcohols. *Plant Physiology*,
 466 117(3), 1047–1058. <https://doi.org/10.1104/pp.117.3.1047>
- 467 Stahl-Biskup, E., Intert, F., Holthuijzen, J., Stengele, M., & Schulz, G. (1993). Glycosidically
 468 bound volatiles—a review 1986–1991. *Flavour and Fragrance Journal*, 8(2), 61–80.
 469 <https://doi.org/10.1002/ffj.2730080202>

- 470 Sturm, K., Koron, D., & Stampar, F. (2003). The composition of fruit of different strawberry
 471 varieties depending on maturity stage. *Food Chemistry*, 83(3), 417–422.
 472 [https://doi.org/10.1016/S0308-8146\(03\)00124-9](https://doi.org/10.1016/S0308-8146(03)00124-9)
- 473 Ubeda, C., Callejón, R. M., Troncoso, A. M., Morales, M. L., & Garcia-Parrilla, M. C.
 474 (2014). Influence of the production process of strawberry industrial purees on free and
 475 glycosidically bound aroma compounds. *Innovative Food Science & Emerging*
 476 *Technologies*, 26, 381–388. <https://doi.org/10.1016/j.ifset.2014.02.015>
- 477 Ubeda, Cristina, San-Juan, F., Concejero, B., Callejón, R. M., Troncoso, A. M., Morales, M.
 478 L., ... Hernández-Orte, P. (2012). Glycosidically Bound Aroma Compounds and
 479 Impact Odorants of Four Strawberry Varieties. *Journal of Agricultural and Food*
 480 *Chemistry*, 60(24), 6095–6102. <https://doi.org/10.1021/jf301141f>
- 481 Ulrich, D., Hoberg, E., Rapp, A., & Kecke, S. (1997). Analysis of strawberry flavour -
 482 discrimination of aroma types by quantification of volatile compounds. *Zeitschrift für*
 483 *Lebensmitteluntersuchung und -Forschung A*, 205(3), 218–223.
 484 <https://doi.org/10.1007/s002170050154>
- 485 Voirin, S. (1990, April 23). *Connaissance de l'arôme du raisin : analyses et synthèses de*
 486 *précurseurs hétérosidiques* (Thèse en doctorat de Biochimie. Biologie moléculaire et
 487 cellulaire). Montpellier 2, Montpellier. Retrieved from
 488 <http://www.theses.fr/1990MON20074>
- 489 Wein, M., Lavid, N., Lunkenbein, S., Lewinsohn, E., Schwab, W., & Kaldenhoff, R. (2002).
 490 Isolation, cloning and expression of a multifunctional O-methyltransferase capable of
 491 forming 2,5-dimethyl-4-methoxy-3(2H)-furanone, one of the key aroma compounds in
 492 strawberry fruits. *The Plant Journal*, 31(6), 755–765. [https://doi.org/10.1046/j.1365-](https://doi.org/10.1046/j.1365-313X.2002.01396.x)
 493 [313X.2002.01396.x](https://doi.org/10.1046/j.1365-313X.2002.01396.x)

- 494 Young, H., & Paterson, V. J. (1995). Characterisation of bound flavour components in
495 kiwifruit. *Journal of the Science of Food and Agriculture*, 68(2), 257–260.
- 496 Zorrilla-Fontanesi, Y., Rambla, J.-L., Cabeza, A., Medina, J. J., Sanchez-Sevilla, J. F.,
497 Valpuesta, V., ... Amaya, I. (2012). Genetic Analysis of Strawberry Fruit Aroma and
498 Identification of O-Methyltransferase FaOMT as the Locus Controlling Natural
499 Variation in Methylfuran Content. *Plant Physiology*, 159(2), 851–870.
500 <https://doi.org/10.1104/pp.111.188318>
501

Table 1: Physico-chemical characteristics and origin of strawberries material

	<i>Origin</i>	<i>Harvest year</i>	<i>Glucose^a</i>	<i>Fructose^a</i>	<i>Sucrose^a</i>	<i>Citric acid^a</i>	<i>Malic acid^a</i>	<i>Brix</i>	<i>pH</i>
FCMO060	Spain	2015	1.59±0.11	2.26±0.08	1.94±0.37	0.50±0.05	0.28±0.01	6.1±0.2	3.6±0.2
Siabel	Bulgaria	2015	1.75±0.25	2.49±0.08	1.41±0.35	0.70±0.08	0.31±0.03	ND	3.5±0.2
Clery	?	2015	2.12±0.31	3.19±0.46	1.88±0.32	0.40±0.01	0.18±0.01	7.8±0.2	3.7±0.2
Honeoye	Bulgaria	2015	2.30±0.28	2.92±0.33	1.91±0.40	0.62±0.09	0.27±0.03	8.6±0.3	3.8±0.2
Honey	China	2015	1.37±0.28	2.20±0.26	0.96±0.02	0.46±0.02	0.18±0.03	6.2±0.2	3.7±0.2
Senga sangana	Bulgaria	2015	2.49±0.31	2.83±0.34	2.32±0.27	0.70±0.20	0.37±0.01	9.7±0.3	3.5±0.2
Darselect	Germany	2015	2.36±0.62	2.84±0.17	1.43±0.30	0.66±0.21	0.18±0.01	8.4±0.3	3.5±0.2
Senga sangana	Poland	2015	1.59±0.48	2.40±0.37	1.17±0.13	0.50±0.02	0.25±0.05	7.7±0.3	3.4±0.2
Fraise 19	Morocco	2016	1.42±0.17	1.97±0.35	0.73±0.38	0.36±0.22	0.19±0.01	6.4±0.2	3.6±0.2
CIR121	France (CIREF)	2016	3.00±0.04	2.94±0.12	0.00	0.81±0.02	0.29±0.00	9.0±0.3	3.7±0.2
Clery PC	France (CIREF)	2016	2.39±0.03	2.39±0.11	0.00	0.59±0.03	0.19±0.00	6.9±0.2	3.5±0.2
Clery HS	France (CIREF)	2016	2.91±0.07	2.83±0.35	0.00	0.94±0.03	0.25±0.01	8.6±0.3	3.5±0.2
Charlotte	France (CIREF)	2016	3.60±0.04	3.67±0.19	0.00	0.73±0.02	0.21±0.00	11.1±0.3	3.9±0.2
Gariguette	France (CIREF)	2016	2.80±0.15	2.76±0.26	0.00	0.96±0.04	0.35±0.01	9.0±0.3	3.4±0.2

^a in g per 100 g of fresh material ; PC : Full-field ; HS : Hydroponis

Comment citer ce document :

Gaborieau, S. (Auteur de correspondance), Cendrès, Page, D., Ginies, C., Renard, C. (2018). Variability of free and glycosylated volatiles from strawberries destined for the fresh market and for processing, assessed using direct enzymatic hydrolysis. LWT - Food Science and Technology, 98, 187-196. . DOI : 10.1016/i.lwt.2018.08.026

Table 2 : Identified^a volatile compound in GC-MS analyses of strawberries samples

	Calculated RI ^b	Perception threshold in water (µg/L)	Odortype	Ref ^{c,d}
<i>Acids</i>				
(E)-2-Hexenoic acid	1980	NA	fruity	7
2-Methylbutanoic acid	1662	180	acidic	1;2;3;6;7
3-Hexenoic acid	1954	NA	cheesy	
3-Methylhexanoic acid	1916	NA	-	7
Butanoic acid	1625	240	cheesy	1;2;5;7
Citraconic / itaconic acid*	1040	NA	-	
Formic acid	1503	NA	acetic	7
Hexanoic acid	1846	3000	fatty	1;2;5;7
Isobutanoic acid	1570	8100	acidic	1;2;5;7
Nonanoic acid	2171	3000	fatty	5;7
Octanoic acid	2060	3000	waxy	2;5;7
Propanoic acid	1535	20000	acidic	1;7
Succinic anhydride*	2103	NA	-	
<i>Alcohols</i>				
1-Butanol	1142	500	fermented	4;7
1-Hexanol	1355	800	herbal	1;2;3;5;7
2,3-Butanediol	1543	4500	creamy	
3-Phenylpropyl alcohol	2039	NA	balsamic	7
4-Vinylguaiacol	2188	3	spicy	4;7
Benzyl alcohol	1870	10000	floral	2;4;7
Chavicol	2334	NA	phenolic	
4-Ethylphenol	2187	NA	smoky	
Phenol	2000	5900	phenolic	
5-methyl-3-hexen-2-ol	1274	NA	green	
3-hexen-1-ol	1398	70	green	2;7
2-Phenylethyl alcohol	1906	4	floral	7
<i>Aldehydes</i>				
Hexanal	1078	NA	-	7
<i>Esters</i>				
3-Methylbutyl acetate	1122	2	fruity	3;7
Butyl acetate	1074	66	ethereal	1;2;3;5;6;7
Ethyl 3-hydroxybutanoate	1515	NA	fruity	7
Ethyl butanoate	1035	1	fruity	1;3;5;6;7
Ethyl (E)-cinnamate	2127	NA	balsamic	7
Ethyl hexanoate	1233	1	fruity	1;3;5;6;7
Ethyl isobutanoate	961	0.1	fruity	7
Isobutyl acetate	1012	66	fruity	7
Methyl 3-hydroxybutanoate	1461	NA	apple	7

Comment citer ce document :

Gaborieau, S. (Auteur de correspondance), Cendrès, Page, D., Ginies, C., Renard, C. (2018). Variability of free and glycosylated volatiles from strawberries destined for the fresh market and for processing, assessed using direct enzymatic hydrolysis. LWT - Food Science and Technology, 98, 187-196. DOI : 10.1016/j.lwt.2018.08.026

Furans

2-Hydroxyfuraneol	1518	NA	-	
5-(Hydroxymethyl)-2-furfural	1660	NA	fatty	
Furaneol	2031	0.03	caramellic	1;2;3;4;5;6;7
Mesifurane	1580	0.03	musty	1;2;3;4;5;6;7

Ketones

1,2-Cyclopentanedione*	1742	NA	caramellic	
2-Hexanone	1083	NA	fruity	3;7
3-Methyl 2-penten-4-one	1187	NA	vegetable	
3-Penten-2-one	1128	1.5	fruity	7
Acetoin	1284	800	buttery	7
2-hydroxy-4-pyranone	1990	NA	-	

Lactones

delta-Caprolactone	1791	NA	herbal	7
gamma-Caprolactone	1694	1600	tonka	2;5;7
gamma-Decalactone	2138	11	fruity	1;2;3;4;5;7
gamma-Dodecalactone	2374	7	fruity	2;7
Pantolactone	2029	NA	caramellic	4

Terpenes

alpha-(E,E)-Farnesene	1746	NA	woody	
beta-Linalool	1547	6	floral	1;2;3;5;6;7
trans-Linalool oxide (pyranosid)	1721	320	floral	1;2;5;7
alpha-Murolene	1726	NA	woody	
1-Terpineol	1576	300	citrus	4;7
trans-Nerolidol	2042	300	floral	1;2;3;5;7

Others

Coumaran	2389	NA	green tea	
Ethyl methyl benzene*	1225	NA	-	
Ethylene glycol diacetate	1535	NA	-	
1,1-diethoxybutane	990	NA	-	7
unknown	1995	NA	green	

^a Identified by NIST library (version 2.2)

^b On polar column (TG Wax)

^c References where identified volatiles were also found.

^d 1: Larsen et al. (1992); 2: Menager et al. (2004); 3: Schwieterman et al. (2014); 4: Ubeda et al. (2014); 5: Lambert et al. (1999); 6: Azodanlou et al. (2003); 7: VCF Volatile Compounds in Food database – Strawberry fruit (2007-Feb-08 version 9.1)

*only identified by MS

Table 3: Number of molecules and quantitative repartition for chemical classes in strawberries free-volatiles

	TOTAL ($\mu\text{g/kg}^a$)	<i>Acid</i>		<i>Alcohol</i>		<i>Aldehyde</i>		<i>Ester</i>		<i>Furan</i>		<i>Ketone</i>		<i>Lactone</i>		<i>Terpene</i>		<i>Other</i>	
		Number	Ratio	Number	Ratio	Number	Ratio	Number	Ratio	Number	Ratio	Number	Ratio	Number	Ratio	Number	Ratio	Number	Ratio
Clery	16608 $\pm$ 1099	6	44%	3	7%	0	0%	4	15%	1	13%	4	6%	4	4%	2	3%	2	8%
Darselect	13187 $\pm$ 2627	2	38%	1	1%	0	0%	2	19%	3	26%	2	5%	3	10%	1	0%	0	0%
FCMO060	9130 $\pm$ 877	3	27%	1	7%	0	0%	1	15%	1	16%	1	4%	2	24%	1	1%	1	7%
Fraise19	9769 $\pm$ 1415	4	33%	1	5%	0	0%	1	13%	2	13%	1	16%	2	14%	1	1%	1	6%
Honeoye	34176 $\pm$ 1785	6	48%	1	2%	0	0%	2	12%	2	8%	4	4%	4	17%	2	2%	2	6%
Honey	30505 $\pm$ 1274	7	28%	5	21%	0	0%	3	10%	2	8%	4	12%	3	14%	2	1%	2	6%
Senga sangana (Bulgaria)	21154 $\pm$ 2672	7	38%	2	5%	0	0%	4	11%	2	19%	3	3%	3	5%	2	2%	1	19%
Senga sangana (Poland)	19822 $\pm$ 2682	5	43%	2	7%	0	0%	1	10%	3	16%	3	4%	3	15%	2	2%	2	2%
Siabel	22984 $\pm$ 2484	8	20%	2	1%	0	0%	2	24%	2	35%	3	5%	3	9%	1	2%	2	4%
Clery PC	41746 $\pm$ 29012	7	3%	0	0%	0	0%	4	90%	1	1%	3	1%	3	0%	2	1%	2	4%
Gariguette	47980 $\pm$ 7387	10	26%	1	0%	1	1%	8	58%	3	4%	2	1%	1	0%	3	5%	3	4%
Clery HS	95633 $\pm$ 9752	8	6%	0	0%	1	1%	5	79%	2	4%	2	1%	2	0%	2	4%	2	4%
CIR121	115250 $\pm$ 22173	6	5%	0	0%	1	1%	4	84%	2	2%	2	1%	2	4%	2	2%	2	1%
Charlotte	58259 $\pm$ 7228	8	6%	0	0%	1	1%	6	79%	2	5%	2	1%	1	2%	3	2%	2	5%

^a in equivalent 4-nonanol

Comment citer ce document :

Gaborieau, S. (Auteur de correspondance), Cendrès, Page, D., Ginies, C., Renard, C. (2018).
 Variability of free and glycosylated volatiles from strawberries destined for the fresh market and
 for processing, assessed using direct enzymatic hydrolysis. LWT - Food Science and Technology,
 98, 187-196. . DOI : 10.1016/i.lwt.2018.08.026

Table 4: number of molecules and quantitative repartition of chemical classes in strawberries glycoconjugates

	TOTAL ($\mu\text{g/kg}^a$)	<i>Acid</i>		<i>Alcohol</i>		<i>Aldehyde</i>		<i>Ester</i>		<i>Furan</i>		<i>Ketone</i>		<i>Lactone</i>		<i>Terpene</i>		<i>Other</i>	
		Number	Ratio	Number	Ratio	Number	Ratio	Number	Ratio	Number	Ratio	Number	Ratio	Number	Ratio	Number	Ratio	Number	Ratio
Clery	10867 $\pm$ 869	6	39%	4	20%	0	0%	1	1%	1	2%	1	3%	2	2%	1	1%	1	33%
Darselect	7705 $\pm$ 2774	3	43%	1	5%	0	0%	0	0%	1	15%	1	2%	2	26%	2	2%	2	7%
FCMO060	4848 $\pm$ 1697	2	13%	2	19%	1	6%	1	3%	0	0%	1	8%	1	13%	0	0%	1	38%
Fraise19	4655 $\pm$ 745	4	30%	2	17%	0	0%	0	0%	0	0%	1	7%	1	10%	0	0%	1	38%
Honeoye	14044 $\pm$ 5056	5	44%	3	11%	0	0%	0	0%	1	5%	1	5%	2	12%	1	1%	1	23%
Honey	9982 $\pm$ 998	6	47%	4	19%	0	0%	0	0%	0	0%	0	0%	1	7%	1	1%	1	27%
Senga sangana (Bulgaria)	20166 $\pm$ 1613	8	51%	5	9%	0	0%	1	1%	1	0%	1	4%	1	0%	0	0%	1	34%
Senga sangana (Poland)	7350 $\pm$ 2352	4	37%	3	23%	0	0%	1	3%	0	0%	1	2%	2	12%	1	1%	1	25%
Siabel	9268 $\pm$ 1112	3	13%	3	9%	0	0%	0	0%	1	54%	0	0%	2	5%	0	0%	1	19%
Clery PC	2524 $\pm$ 606	4	11%	4	22%	1	0%	0	0%	2	0%	1	12%	1	3%	0	0%	2	52%
Gariguette	18074 $\pm$ 9398	10	54%	2	2%	1	2%	0	0%	2	13%	3	5%	1	0%	2	3%	2	23%
Clery HS	5258 $\pm$ 578	6	19%	3	12%	1	0%	0	0%	2	16%	2	11%	0	0%	1	1%	2	40%
CIR121	7052 $\pm$ 3103	8	20%	3	10%	1	1%	0	0%	1	2%	2	8%	2	14%	1	10%	2	43%
Charlotte	3657 $\pm$ 1060	6	10%	2	2%	1	2%	0	0%	1	4%	2	11%	0	0%	0	0%	2	71%

^a in equivalent 4-nonanol

Comment citer ce document :

Gaborieau, S. (Auteur de correspondance), Cendrès, Page, D., Ginies, C., Renard, C. (2018).
 Variability of free and glycosylated volatiles from strawberries destined for the fresh market and
 for processing, assessed using direct enzymatic hydrolysis. LWT - Food Science and Technology,
 98, 187-196. . DOI : 10.1016/j.lwt.2018.08.026

Figure 1: Scheme of strawberries preparation for assisted-solvent extraction (ASE) of volatiles and sampling point for analysis of free- and bound-volatiles

Figure 2: Concentration of volatiles after extraction by ASE method with: dichloromethane/pentane (1:2) (blue), dichloromethane (red), ethyl acetate/pentane (2:1) (green)

Figure 3: Principal component analysis of free-volatiles in strawberries. A. Sample map; B. Variable correlation map

Figure 4: Principal component analysis of glycoconjugates in strawberries. A. Sample map ; B. Variable correlation map

Highlights:

Glycosylated volatiles can be reliably analyzed by direct enzymatic hydrolysis of the fruit.

Free-volatile profiles split strawberries for fresh market and for processing.

Bound-volatile profiles were similar among the majority of strawberry varieties.

Strawberries for processing were the best candidates for aroma enhancement.

ACCEPTED MANUSCRIPT

Comment citer ce document :

Gaborieau, S. (Auteur de correspondance), Cendrès, Page, D., Ginies, C., Renard, C. (2018). Variability of free and glycosylated volatiles from strawberries destined for the fresh market and for processing, assessed using direct enzymatic hydrolysis. LWT - Food Science and Technology, 98, 187-196. . DOI : 10.1016/j.lwt.2018.08.026