



**HAL**  
open science

## Connecting the dots between computational tools to analyse soil-root water relations.

Sixtine Passot, Valentin Couvreur, Félicien Meunier, Xavier Draye, Mathieu Javaux, Daniel Leitner, Loïc Pagès, Andrea Schnepf, Jan Vanderborght, Guillaume Lobet

► **To cite this version:**

Sixtine Passot, Valentin Couvreur, Félicien Meunier, Xavier Draye, Mathieu Javaux, et al.. Connecting the dots between computational tools to analyse soil-root water relations.. *Journal of Experimental Botany*, 2019, 70 (9), pp.2345 - 2357. 10.1093/jxb/ery361 . hal-02623964

**HAL Id: hal-02623964**

**<https://hal.inrae.fr/hal-02623964v1>**

Submitted on 26 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Connecting the dots between computational tools to analyse soil-root water relations

Sixtine Passot<sup>+1</sup>, Valentin Couvreur<sup>+1</sup>, Félicien Meunier<sup>+1,2</sup>, Xavier Draye<sup>1</sup>, Mathieu Javaux<sup>1,3</sup>, Daniel Leitner<sup>4</sup>, Loïc Pagès<sup>5</sup>, Andrea Schnepf<sup>3</sup>, Jan Vanderborght<sup>3</sup> and Guillaume Lobet<sup>1,3,◦</sup>

<sup>1</sup> Earth and Life Institute, Université catholique de Louvain, Louvain-la-Neuve, Belgium

<sup>2</sup> Computational and Applied Vegetation Ecology lab, Ghent University, Ghent, Belgium

<sup>3</sup> Agrosphere, IBG3, Forschungszentrum Jülich, Jülich, Germany

<sup>4</sup> Simulationswerkstatt, Ortmayrstrasse 20, A-4060 Leonding, Austria

<sup>5</sup> INRA, Centre PACA, UR 1115 PSH, 84914 Avignon cedex 9, France

<sup>+</sup> These authors contributed equally to this work.

<sup>◦</sup> Corresponding author. [g.lobet@fz-juelich.de](mailto:g.lobet@fz-juelich.de)

© The Author(s) 2018. Published by Oxford University Press on behalf of the Society for Experimental Biology. All rights reserved. For permissions, please email: [journals.permissions@oup.com](mailto:journals.permissions@oup.com)

Comment citer ce document :

Passot, S., Couvreur, V., Meunier, F., Draye, X., Javaux, M., Leitner, D., Pagès, L., Schnepf, A., Vanderborght, J., Lobet, G (2019). Connecting the dots between computational tools to analyse soil-root water relations. *Journal of Experimental Botany*, 70 (9), 2345 - 2357. , DOI : 10.1093/jxb/ery361

# Abstract

In the recent years, many computational tools, such as image analysis, data management, process-based simulation and upscaling tools, were developed to help quantify and understand water flow in the soil-root system, at multiple scales (tissue, organ, plant and population). Several of these tools work together or, at least, are compatible. However, for the un-informed researcher, they might seem disconnected, forming a unclear and disorganised succession of tools.

In this article, we present how different pieces of work can be further developed by connecting them to analyse soil-root-water relations in a comprehensive and structured network. This “explicit network of soil-root computational tools” informs the reader about existing tools and help them understand how their data (past and future) might fit within the network. We also demonstrate the novel possibilities of scale-consistent parameterizations made possible by the network with a set of case studies from the literature. Finally, we discuss existing gaps in the network and how we can move forward to fill them.

## Highlights

Many computational tools exist to quantify water flow in the soil-root system. These tools can be arranged in a comprehensive network that can be leveraged to better interpret experimental data.

## Keywords

Computational tools, image analysis, simulation, network, root, soil, water

## Authors informations

Sixtine Passot	sixtine.passot@uclouvain.be	0000-0002-6191-723X
Valentin Couvreur	valentin.couvreur@uclouvain.be	0000-0002-1087-3978
Félicien Meunier	felicien.meunier@uclouvain.be	0000-0003-2486-309X
Xavier Draye	xavier.draye@uclouvain.be	0000-0002-3637-3330
Mathieu Javaux	mathieu.javaux@uclouvain.be	0000-0002-6168-5467
Daniel Leitner	daniel.leitner@simwerk.at	
Loïc Pagès	loic.pages@inra.fr	0000-0002-2476-6401
Andrea Schnepf	a.schnepf@fz-juelich.de	0000-0003-2203-4466
Jan Vanderborght	j.vanderborght@fz-juelich.de	0000-0001-7381-3211
Guillaume Lobet	g.lobet@fz-juelich.de	0000-0002-5883-4572

## Glossary

Term	Definition	Reference
Standard Uptake Fraction	Relative distribution of root water uptake between root segments when water is equally available in space (units: %)	(Couvreur <i>et al.</i> , 2012)
High pressure flow meter	Device designed to measure the root system conductance by perfusing pressurized water into a root system opposite from the natural direction of the transpiration stream	(Tsuda and Tyree, 2000b)
Root pressure probe	Device designed to measure the hydraulic conductance of a single root through variations of water pressure and flow at the cut end of a root	(Steudle and Jeschke, 1983)
Cell pressure probe	Device designed to measure the hydraulic conductivity of the membranes of a single plant cell by observing the relaxation time of water pressure pulses applied to the cell	(Steudle, 1980)
RSML	Root System Markup Language: File format for the storage of root system data	(Lobet <i>et al.</i> , 2015)
MTG	Multi-Tree Graph: File format for the storage of multidimensional tree information, designed for plant models.	(Godin <i>et al.</i> , 1999)
FSPM	Functional-Structural Plant Model: computer model of plant that combine a detailed representation of the plant 3D architecture with functional properties (radial hydraulic conductivity, solute permeability, etc.)	(Godin and Sinoquet, 2005)
Parameter	Fixed input of the models that characterizes a specific property of the system, within the scope of study. For instance, branching rates are parameters in root architectural models	
State variable	Variable that characterizes the state of the system at any moment of the simulation. For instance, the water potential within the plant is a state variable in water FSPM	
Boundary conditions	Variables constraining the model at its external boundaries for the entire duration of the simulation. For instance, in model of water flow within the root system, evaporative demand or soil water potentials are the plant boundary conditions.	
Upscaled property	System property that is an output of the model, at a higher scale	

---

than the input parameters. For instance, the root radial conductivity is an upscaled property of models of root organ water flow.

---

## Introduction

Water deficit is one of the most dramatic abiotic stresses in agriculture (Cattivelli *et al.*, 2008). It occurs when leaf water supply is limited by either the low potential of soil water, and/or by the high hydraulic resistance of the soil-plant system (Sperry *et al.*, 2002). At this point, the atmospheric demand for water is hardly met, leading to a decrease in water potential within the leaf tissues. As a result, stomata will close, reducing the plant transpiration and photosynthesis. To investigate when such limitation occurs, the complex plant-soil-atmosphere system is often conceptualised as a multidimensional hydraulic network, in which both soil and root hydraulic properties may substantially control shoot water supply (Draye *et al.*, 2010; Schoppach *et al.*, 2013; Lobet *et al.*, 2014a).

The **structural properties** of the roots compose the first dimension of the soil-root hydraulic network. Structural properties refer to the physical position and arrangement of the objects of interest. They can be conceptualised at the tissue/organ (transverse anatomy, fig. 1A), plant (root architecture, fig. 1B), or population scale (rooting density profile, fig. 1C).

A second dimension, overlaying structural properties, encompasses the system **functional properties**. When studying water movement, functional properties often refer to hydraulic conductivities or reflection coefficients. Like in the structural layer, these properties can be defined at different scales. Local radial and axial hydraulic conductivities can be defined at the organ scale (fig. 1D) while the entire root system of a single plant can be characterized by its conductance (fig. 1E) and would relate to plant water status (Alsina *et al.*, 2011). An extension of this property to the population scale is the plant population hydraulic conductance per unit horizontal area (fig. 1F), common in canopy models (Cox *et al.*, 1998), and recently integrated in root models (Cai *et al.*, 2017).

Finally, a third dimension describes the plant **environment**. In this contribution, we focus on the soil compartment, which includes the rhizosphere and the bulk soil (fig. 1G). Their respective spatial domains are concentric around individual roots, and their properties differ substantially, so that the rhizosphere is often considered to critically affect plant water availability under water deficit (Carminati *et al.*, 2011; Ahmed *et al.*, 2014). The bulk soil and rhizosphere hydraulic

properties may be described by their water retention and hydraulic conductivity curves. The former defines the pressure needed to extract water from the porous media, and the latter the relation between water flux and water potential gradient in space (van Genuchten, 1980). The water potential that defines the energy level of water is a critical environmental variable, driving the flow of water in the soil-plant system. Similarly to the plant, the soil could be divided into functional and structural components and described according to the studied scale. However, we did not explicit this separation in the following as we rather focus on the plant property description in this study.

Each element of the network is dynamic and heterogeneous. Root systems grow, develop and take up water, while soil water content continuously changes in response to root water uptake and climatic conditions, potentially resulting in complex system behaviour. In addition, some key variables and parameters are hard (if not impossible) to quantify experimentally. As a result, the whole system is difficult to apprehend, and novel approaches might prove useful to study it.

In the recent years, many computational tools (image analysis, data analysis, process based modelling and upscaling tools) were developed to help quantify and understand water dynamics in the soil-plant system. Some of these tools were developed to work together, or at least be compatible. However, for the uninformed researcher, they might seem disconnected, forming a collection of tools with, at best, a common target (plant-water relation exploration) but unrelated to each other.

The overall objective of this paper is to draw and discuss the role of a functional landscape of interconnected experiments and models for the study of soil-plant water relations. It is articulated as 3 sub-objectives: (i) to inform readers about existing procedures and tools used for the quantification of water flow at the organ, root system and plant population scales, as well as their interconnections forming a comprehensive, though non-exhaustive, network, (ii) to provide examples of studies combining experiments, analytical and modelling tools in this network, motivating the use of such approaches to enhance interpretations of available and future data, and (iii) to identify gaps in the network and argue for a better integration of future tools in this workflow with appropriate experiment and model design. A web interface was developed to help researchers use the network: it is available at [https://plantmodelling.shinyapps.io/water\\_network/](https://plantmodelling.shinyapps.io/water_network/).

Through four examples, we illustrate how the dots, consisting in apparently scattered data and tools, can be connected together in a comprehensive network. These examples span over the different scales mentioned above (organ, root system and population) and for all of them, we

present data that can be obtained experimentally, technical limitations that need to be overcome and computational tools readily available. In all these examples, we focus on the soil-root water relation specificities at different scales, except for one where an architectural root growth model is introduced.

## Water flow at the root cross section scale

Different tools and techniques exist to quantify root structural properties at the organ scale. Histology and microscopy techniques enable precise observation of root anatomical structures (the interconnected network of cells). For instance, staining or fluorescence microscopy can be used to acquire images of the organization of different cell types within roots and the nature of cell walls (Costa and Plazenet, 2016). Different image analysis tools are then available to extract quantitative information out of these images. On the one hand, CellSet (Pound *et al.*, 2012) is currently the only tool that enables a complete digitization of the entire cell network. As an output, each single cell is represented by a set of connected edges and nodes. Unfortunately, depending on the image quality, the unautomated part of the procedure can be time-consuming. On the other hand, RootScan (Burton *et al.*, 2012), PHIV-RootCell (Lartaud *et al.*, 2014) and RootAnalyzer (Chopin *et al.*, 2015) are fully automated tools that can quantify anatomical properties (such as the number of cells or the mean size of each cell type) but do not provide a digitized cell network.

As a part of the functional layer, cell hydraulic properties are hard to estimate as water fluxes are difficult to measure at this scale. The cell pressure probe enables this estimation from measurements of water pressure relaxation times of individual cells, at a high time cost (Steudle, 1980; Hose *et al.*, 2000). Osmotic pressure can be measured using nanoliter osmometer (Malone *et al.*, 1989) or scanning electron microscopy (Enns, 1998). However, the latter is expensive and generally not part of the standard equipment of a plant physiology laboratory. At the organ scale, the root pressure probe enables the measurement of axial and radial conductivities of root segments (Steudle and Jeschke, 1983) and junctions to the stem (Meunier *et al.*, 2018b). Some properties of the system can hardly be determined experimentally such as the partitioning of water pathways across cell layers (apoplastic or cell-to-cell) (Barzana *et al.*, 2012).

Detailed root cross-section anatomical descriptions and a minimal set of empirical cell hydraulic properties (e.g. permeability of cell walls and membranes) enables the simulation of water flow across root cross sections. Like at other scales, water flow in the system is solved using transfer equation with boundary water pressures and conductances as input parameters. Such a model can estimate the equivalent hydraulic conductivity of the root cylinder as well as the partitioning of water flow between apoplastic and symplastic compartments of the system. For instance, by combining measurements of cell and root permeability with a hydraulic model, (Bramley *et al.*, 2009) demonstrated that water flow is primarily apoplastic in lupin roots.



A recent study took advantage of these computational tools to estimate the contribution of pearl millet root types to water uptake. Five types were identified based on cross-section anatomy: primary roots, crown roots and 3 types of lateral roots (Passot *et al.*, 2016). A cross-section was thoroughly digitized for each type of root using CellSet (Pound *et al.*, 2012) (fig. 2A). Root axial hydraulic conductances were estimated using the simplified model of Hagen-Poiseuille (fig. 2B), based on measured xylem vessels dimensions. The digitized root anatomical network served as input for a mechanistic model of radial water flow in roots, namely MECHA (Couvreur *et al.*, 2017) (fig. 2C). The model was used to estimate the radial conductivity of a typical segment of each root type. In this example, different tools (image analysis and modelling) were combined to estimate radial and axial conductivities, based on easy-to-acquire experimental data (cross section images). While complementary measurements of root hydraulic properties will always remain an asset (e.g. in order to cross-validate the estimated properties), this method opens the way to high-throughput estimations of root hydraulic properties.

## Root system architecture

Unlike plant cells, the root system of annual crops has a convenient macroscopic scale and all elements (roots) are visible to the human eye. However, the main difficulties faced when retrieving the root system architecture are the hidden nature of this part of the plant, the large number of elements that can possibly overlay and the fragility of the smallest roots, making the full excavation of a complete intact root system particularly difficult. Direct manual methods exist to measure single root architectural traits such as the angle of crown roots with a protractor (Trachsel *et al.*, 2010) or with the basket method (Oyanagi *et al.*, 1993; Uga *et al.*, 2011), the length of individual roots with a ruler (Pritchard *et al.*, 1990; Trachsel *et al.*, 2010) or a combination of several root architectural traits (Trachsel *et al.*, 2010). However, these manual methods do not give access to the full root architecture.

Several digital tools have been developed and are now widely used to access root architectural traits, mostly from images of root system grown in specific experimental setups (see (Paez-Garcia *et al.*, 2015) for a review of existing root phenotyping strategies). These image analysis tools are listed in [www.plant-image-analysis.org](http://www.plant-image-analysis.org) and will not be detailed here (Lobet *et al.* 2013, Lobet 2017). The only point to underline is that each tool generally corresponds to a specific growth medium and image capture technique (eg: RooTrak applies to root system growing in 3D and imaged with X-ray Micro Computed Tomography (Mairhofer *et al.*, 2012)). While some of these tools have been designed to retrace a full root system architecture (often with an important manual input), many of them only extract some root architectural traits (eg. mean lateral root length, number of seminal roots, crown root emergence angle...). Furthermore, even with the use of specifically designed image analysis tools, whole root system digitization

becomes time consuming as soon as the plants are several weeks old. Therefore, subsequent tools are needed to reconstruct full root system architectures from extracted root traits.

Root architecture models, such as SimRoot (Lynch *et al.*, 1997), RootBox (Leitner *et al.*, 2010), RootTyp (Pagès *et al.*, 2013), ArchiSimple (Pagès *et al.*, 2014), OpenSimRoot (Postma *et al.*, 2017b) or CRootBox (Schnepf *et al.*, 2018), are designed to simulate root systems from a limited number of traits, given as input parameters. The major interest of root architectural models is to generate a large number of contrasted root system architectures. Root system modeling enables the exploration of several variants for the same mean traits and the simulation of contrasted architectures, even from synthetic datasets. These contrasting architectures can then be tested in different scenarios, to identify traits that would be beneficial in challenging environments.

An example that illustrates how root architecture models can be applied to interpret experimental data of other root zone processes is given by Schnepf *et al.* (2016). Those authors developed a 3D model of the development of mycorrhizal root systems. The model was designed to simulate primary and secondary root infection with arbuscular mycorrhizal (AM) fungi as well as growth of external fungal hyphae in soil. It was calibrated using root architectural data obtained from pot experiments of *Medicago truncatula*, with and without mycorrhizal inoculum of the AM fungus *Rhizophagus irregularis* BEG 158. In those pots, AM root colonization was determined under a compound microscope and the abundance of *R. irregularis* hyphal biomass was determined using real-time PCR. The root system architecture, however, could not be parameterised from those pot experiments. The authors re-used published images from a previous study (Bourion *et al.*, 2014) (fig. 3A) and re-analysed them with the image analysis tool RootSystemAnalyzer (Leitner *et al.*, 2014a) (fig. 3B). The traits extracted with RootSystemAnalyzer served as parameters for the RootBox model (fig. 3C) (Leitner *et al.*, 2010) which was used to simulate the root system development of mature plants, together with the arbuscular mycorrhizal fungi. This example highlights how published data can be reused to obtain input parameters for modelling. The current literature is filled with similar resources, opening up numerous opportunities. It also highlights the importance of sharing raw experimental data (in this case the images).

## Water flow in the root system

At the root system scale, understanding which root traits positively influence plant water uptake dynamics for a given pedo-climatic situation remains an important research question. Ideotypes have been proposed, but are always tied to a specific environment (Dencic, 1994; Guswa, 2010; Pagès, 2011; Wasson *et al.*, 2012; Lynch, 2013). Different traits, either functional or structural, have been found to maximize the final crop yield depending on the environment (Tardieu, 2012). Ultimately, we need more than single traits or final yield to have a better understanding of plant-environment interactions. We need to understand how water flow within the plant is

dynamically regulated, both spatially and temporally. Unfortunately, accurately measuring water flow is often the limiting step of the experimental pipeline. Several techniques exist to dynamically measure changes in soil water content, such as X-ray computed tomography (Hainsworth and Aylmore, 1983), Electrical Resistivity Tomography (Garré *et al.*, 2013), neutron tomography (Carminati *et al.*, 2010; Esser *et al.*, 2010; Zarebanadkouki *et al.*, 2014a; Tötze *et al.*, 2017), light transmission imaging (Garrigues *et al.*, 2006) or Magnetic Resonance Imaging (Jahnke *et al.*, 2009; Pohlmeier, 2010; Rascher *et al.*, 2011). These techniques can be deployed for a relatively high number of plants. However, due to water capillary flow within the soil domain, observed changes in soil water content are rarely (if never) a direct indication of the location of root water uptake. Water uptake rate itself can be estimated using more advanced but time-consuming lab techniques that use tracers, such as deuterated water that is monitored using neutron radiography (Warren *et al.*, 2013; Zarebanadkouki *et al.*, 2014b).

Functional-structural plant models (FSPM) are often used to decipher plant-environment relationships (Godin and Sinoquet, 2005). FSPMs couple a complete representation of the root system architecture (or whole plant or shoot) with functional properties. Their input parameters are both functional and structural. For FSPMs simulating soil-root interactions, hydraulic parameters can be obtained using a root pressure probe (Stedle *et al.*, 1987) or the outputs of organ scale models but, as stated earlier, are generally difficult to acquire. Thus they are frequently adapted from the available literature. Rhizosphere hydraulic properties can also be coupled to FSPM (Schneider *et al.*, 2009) and would constitute a central component of plant water availability (Carminati *et al.*, 2011). Rhizosphere properties are however difficult to parametrize, and would display complex temporal dynamics (Carminati and Vetterlein, 2013). The FSPM structural input consists of an explicit representation of the root architecture (see Root system architecture section and the related previous case study). Together with the root system geometry, hydraulic properties define the root system hydraulic architecture (Lobet *et al.*, 2014a) and are critical for water stress determination (Leitner *et al.*, 2014b; Vadez, 2014).

Water-related FSPMs provide an exhaustive description of the root water relations (uptake rates, water potentials, etc.) in both space and time. Thus, they constitute an important way to integrate different types of information about properties of the root system and soil state variables in the root zone, which can be obtained experimentally, and to translate this information into a distributed pattern of water flows and local state variables (e.g. water potentials at the soil-root interface) within the root zone. The latter type of information is, as of today, hardly accessible experimentally. An exhaustive review of FSPMs related to water flow can be found in Ndour *et al.* (Ndour *et al.*, 2017).

FSPMs can also be used in so-called *inverse modelling* studies. In such case, the output of the model is known and the model is used to estimate one of the input parameters. For instance, the most likely distribution of root hydraulic properties (that are usually assumed to be age- and order-dependent) can be estimated using a soil-root water flow model and laboratory measurements (Zarebanadkouki *et al.*, 2016). In this study, measurements of local water fluxes were obtained from neutron radiography at different locations in the root system (Zarebanadkouki *et al.*, 2014a). As the experiment took place in a rhizotron, the root system could be fully digitized using an appropriate image analysis tool, which provided accurate information on the root system topology and positions in space (fig. 4A and see also Root system architecture section). Water uptake patterns and axial flows within the root system could then be modelled by applying existing water flow equation resolution algorithms (Doussan *et al.*, 1998; Meunier *et al.*, 2017b) to the segmented root system (fig. 4B). The water flow model requires boundary conditions that need to be estimated or measured. In this case, the water potential at the root collar was measured using a pressure probe and root-soil interface water potentials were estimated from water content distribution.

Such a coupling allowed the authors to estimate the parameters of the root hydraulic conductivity function that best fitted the water flow measurements (fig. 4C and D). These parameters then allowed for novel predictions including water uptake and axial flow distributions everywhere in the root system and not only at observed segments, in homogeneous and heterogeneous soil conditions or under various evaporative demands.

## Water flow at the population scale

The population level is a pivotal scale. It interfaces with general circulation models that represent, among others, the circulation of the atmosphere and its interaction with land surface for climate forecasting (Shrestha *et al.*, 2014). It also introduces variables of critical agronomic interest like crop yield per acre (Holzworth *et al.*, 2018). Like at other scales, robust predictions of the system behaviour require the ability to quantify system properties and a proper validation, here involving field scale observations of water fluxes. These fluxes can be estimated with heavy instrumentation and data analytics, for instance using eddy covariance flux towers (McElrone *et al.*, 2013) or soil moisture sensors grids (Famiglietti *et al.*, 2008).

Structural root information can be obtained using either destructive sampling, such as core sampling (Wasson *et al.*, 2014), monolith excavation (Kuchenbuch *et al.*, 2009), trenches (Vepraskas and Hoyt, 1988) or root crown excavation (Colombi *et al.*, 2015), or non-destructive ones, such as minirhizotrons (Rewald and Ephrath, 2013). None of these techniques allows for direct reconstruction of the root

system, but rather extract synthetic metrics such as a root length density profile, or root crown data (angles, numbers etc.). Some root architectural traits can be derived from data obtained with these techniques using root architecture models and inverse modeling, as stated above (Garré *et al.*, 2012; Vansteenkiste *et al.*, 2013). Functional plant properties, such as root system and stomatal conductances, which coordinate shoot water supply (Alsina *et al.*, 2011) and underground water uptake distribution (Meunier *et al.*, 2018a), can be characterized on individual plants with low-throughput instruments, such as the high-pressure flowmeter (Tsuda and Tyree, 2000a) or the porometer (Rodrigues *et al.*, 2008), then scaled to the population level using the planting density. The main limitation of plant measurement in the field is often the limited sample size, that might not reflect the general behaviour of the system. The same critique can be made about soil hydraulic properties estimated on small and (un)disturbed samples as they may not be representative of the hydrological behaviour at the population level (Hopmans *et al.*, 2002; Vrugt *et al.*, 2004).

These limitations motivate the use of effective descriptions of population water relations, tailored for this specific scale, such as the transpiration correction for “soil water stress” (Verhoef and Egea, 2014) or one-dimensional soil water and nutrient transfer principles (Baram *et al.*, 2016). Two major methodologies address the parametrization of effective field water relations. First, the artificial neural network approach takes advantage of the availability of large amounts of data to train a model. It was used to predict canopy water fluxes from state variables such as the vapor pressure deficit and soil moisture (Whitley *et al.*, 2012; Couvreur *et al.*, 2016). Second, the inverse modelling approach (as described previously) builds on state-of-the-art models to simulate spatio-temporal series of the system state. The model parameter values producing simulated series that best match field observations are considered optimal and representative of the system behaviour. This approach was used to connect models of soil and plant water flow to observations of soil moisture and transpiration in an almond orchard, in order to estimate soil and plant properties, as well as the hardly measurable leaching of water below the root zone (Vrugt *et al.*, 2001). Numerous variables can be used for inverse modeling, such as soil water content, isotopes distributions or root length density profiles.

Going one step further, simplistic macroscale models can be derived from equations of water flow at a lower scale, offering an interesting trade-off between functional simplicity and realism. This type of model involves scale-consistent properties and processes. A cross-validation is thus possible between parameter values estimated directly at the macroscale of interest (e.g. plant population hydraulic conductance per surface area) and derived from the lower scale (e.g. upscaled values derived from root architectural and hydraulic properties). In order to parametrize such a macroscale model of water dynamics in the soil-wheat system, Cai *et al.* (Cai *et al.*, 2017) combined one-dimensional process-based models of water flow (i) in soil (Hydrus-1D, (Simunek and Van Genuchten, 1995)) (fig. 5A), (ii) in roots (Couvreur *et al.*, 2014a), and (iii) in leaves with an isohydric constraint on transpiration (Couvreur *et al.*, 2014b) (fig. 5B).



Regarding soil properties, soil water retention curves were fitted on simultaneous soil water content and pressure head measurements (Cai *et al.*, 2016) with the software RETC (van Genuchten *et al.*, 1991) (fig. 5C). However, the parameters of the soil hydraulic conductivity curve were not experimentally determined (fig. 5D). The vertical distribution of roots in the field (root length density profiles over time) was extracted from *in situ* rhizotube pictures, with the software Rootfly (Zeng *et al.*, 2008) (fig. 5E). Its relative distribution is typically used as proxy for the water uptake distribution in uniformly wet conditions (Feddes *et al.*, 1978; Šimůnek and Hopmans, 2009) (fig. 5F). Plant hydraulic properties could not be observed *in situ* for the wheat population (fig. 5F-G).

An inverse modelling strategy was therefore used to find the “optimal” soil and plant hydraulic properties (fig. 5H) that best fitted the observed soil water status dynamics. The optimized plant hydraulic parameters were cross-validated with properties at the individual plant scale. Conductance parameters obtained for winter wheat at the same stage of maturity using the hydraulic architecture approach turned out to be consistent with the inversely modeled properties at the population scale (Cai *et al.*, 2017).

In order to limit the number of parameters, this approach requires the assumption that system properties are time invariant (e.g. soil hydraulic conductivity curve). Because root system conductance tends to scale with root length, the root conductance per unit root length was assumed invariant in order to accommodate for root growth. Such a constraint also matters when accounting for the spatial heterogeneity of root development under different soil/microclimate environments in macroscale simulations (Cai *et al.*, 2018).

## Discussions and perspectives

Many computational tools exist to better understand water dynamics in soil-plant systems. These tools span different scales (organ, plant and population), types (image analysis, data storage, simulation models) and computational languages (Python, Fortran, C++, C#, Java, ...). For the average user, this multitude might seem overly complex and hard to understand. Yet, most of the tools could work together and form a continuous network. Using this network, experimental data can be transferred from scale to scale and generate new insights (fig. 6, and case studies developed above). Modelling tools currently present in the network are listed in table 1. This list is non-exhaustive as the objective of this paper is less to review existing tools than to encourage their integration in order to enhance our understanding soil-plant water relations. For image analysis tools, we refer the reader to the [www.plant-image-analysis.org](http://www.plant-image-analysis.org) database (Lobet *et al.*, 2013; Lobet, 2017).

We created an interactive online visualization of the network, which contains links to the different tools and makes the understanding of Figure 6 easier. We also added a submission form, such as the community could update the network with new (or missing) tools. The online visualisation tool (open-source) is available at [https://plantmodelling.shinyapps.io/water\\_network/](https://plantmodelling.shinyapps.io/water_network/).

## Identifying gaps in the tool network

Analogies between tool connectivity patterns at each scale in figure 6 reveal the existence of network gaps (represented by the dashed lines in fig.6). Yet, these gaps should not be filled just for the sake of symmetry. Here, we analyze what the function of filling these gaps could be.

The plant “structure” row in Fig. 6 has the most striking pattern, with imaging techniques systematically feeding image-analysis tools. These tools extract two types of data: (i) explicit spatial structures (e.g. RSML, MTG), and (ii) structural pattern properties (e.g. growth rates, branching rates). At the population and plant scales, root development models (Dupuy *et al.*, 2010; Pagès *et al.*, 2013) offer the possibility to convert root pattern properties into predicted root structures. While root anatomical patterns can be automatically characterized by image analysis tools such as PHIV-RootCell (Lartaud *et al.*, 2014), no root anatomy development model exists at that scale. In the perspective of generating a mechanistic model of a whole plant from the cell scale (Band *et al.*, 2014), a root development model would become essential. It would fulfill two main functions: (i) conducting predictions and test hypotheses related to root anatomical development, and (ii) allowing the spatial and temporal interpolation of root anatomies between experimental observations.

Models using explicit root anatomical structures to test hypotheses about hormone signalling (Stoma *et al.*, 2008), tropisms (Robbins and Dinneny, 2018) or radial water flow (Couvreur *et al.*, 2017) have emerged lately. However, models of axial water flow remain largely underexplored. In the broadly used Poiseuille-Hagen model, only the quantity and diameter of xylem vessels are accounted for. Yet, multiple studies have demonstrated that xylem porous plates, pit membranes and persistent primary cross-walls limit root axial conductivity (Sanderson *et al.*, 1988; Shane *et al.*, 2000; Choat *et al.*, 2008, 2012; Brodersen *et al.*, 2013; Tixier *et al.*, 2014) and affect the profile of water uptake along a single root (Meunier *et al.*, 2017a) or its partitioning among root types (Meunier *et al.*, 2018b; Ahmed *et al.*, 2018). Furthermore, cavitation may substantially alter xylem conductivity, particularly in leaves and stems (Rodriguez-Dominguez *et al.*, 2018). X-ray micro-computed tomography now enables the study of this phenomenon *in vivo* in all parts of the plant (root, stem and leaf) (Skelton *et al.*, 2017) and also the *in vivo* embolism removal by root pressure (Brodersen *et al.*, 2018). This process is already simulated by models working at the tissue scale, with experimental functions of percentage loss of conductivity (Mirfenderesgi *et al.*, 2016; Couvreur *et al.*, 2018). However, modelling tools are still missing at the vessel scale and

their development may reveal a complexity that is neglected at higher scales. Such numerical representations of xylem vessel structure and hydraulic properties in the axial direction would allow the use of alternatives to Hagen-Poiseuille law (Giordano *et al.*, 1978; Lewis and Boose, 1995). Indeed, Hagen-Poiseuille law describes water flow at equilibrium state but cavitation is a process happening fundamentally out of equilibrium. Lewis and Boose point out that “Ideally, the exact solutions should be used to calculate [volume flow rate] in xylem conduits, but the equations are difficult to solve without the aid of computer” (Lewis and Boose, 1995). Computer availability is no longer an issue and explicit models of xylem flow are now emerging (Mrad *et al.*, 2018). We expect that filling this gap will shed light on the role of cross-walls in the generation of root hydraulic types, and in root-leaf preferential connectivity (David *et al.*, 2012).

Similarly, at the soil-root interface, imaging tools are now available to precisely observe processes at the scale of the soil particle and root hair (Keyes *et al.*, 2013). Connecting such soil-root interface geometrical descriptions to root hydraulic anatomical models would open new avenues to understand how root hairs enhance plant water availability in dry soils (Carminati *et al.*, 2017).

Soil water fluxes were only explicitly considered in the last case study (population scale). In other case studies, soil was either neglected (organ and plant scales) or included as static boundary conditions. However, in all cases, a model of water flow in the soil domain can be coupled to the plant water flow. Such analyses were for instance carried out at the plant level (Javaux *et al.*, 2008; Huber *et al.*, 2014, 2015; Postma *et al.*, 2017b) or the population level (Gijsman *et al.*, 2002; Hack-ten Broeke *et al.*, 2016). Such models may incorporate multiple soil characteristics such as macropores (Landl *et al.*, 2016), solute convection-dispersion (Schroeder *et al.*, 2012) or specific rhizosphere properties (Schwartz *et al.*, 2016; Roose *et al.*, 2016). For an extensive review of existing soil models, we refer the reader to Vereecken *et al.* (Vereecken *et al.*, 2016).

## Limitations and future developments

Simulating water fluxes in roots with this collection of tools can either help understanding plant water relations as a main goal or be a tool for further application. These tools could also be used as a side usage of a dataset obtained for other purposes. The advent of imaging in plant sciences



and the huge progress made in image analysis allow generating high quality quantitative data of plant structure, suitable for model parameterization. Many models exist at different scales and we highlighted the many possibilities to combine these tools. This set of tools greatly increases the potential of interpretation of experimental data. Yet many authors still publish rich datasets without using modelling tools to interpret them. Using models is not trivial for a large part of the plant science community. Coupling several tools and models together is still rarely applied. Several requirements seem essential to facilitate the use of this pipeline whenever it may add value to the data.

A lot of image analysis tools and models flourish in the water transport domain. Thanks to the wide breadth of scientific literature available (scientific papers, reviews, websites...), developers are usually aware of already existing tools and keep in mind to justify the interest of their tool in this landscape. However, we suggest that further efforts should be made to render the tools compatible with existing ones. In this context, the existence of several tools at the same place of the network (eg. RootTyp and CRootBox for root architecture simulation) is not conflicting. Each one can best suit one scale or specific situation. In our opinion, special attention should be paid to the data format. Indeed, the output data format of upstream tools must be compatible with the input format required by downstream ones. If this is not the case, easy-to-handle tools must exist to convert these data. The multiplication of formats and the need to convert data from one type to another may discourage the use of some of the models. In this respect, the existence of standard formats, such as the Root System Markup Language (Lobet *et al.*, 2015) for root architecture, smooths the interconnection between tools.

When a new tool is created, the documentation of its potential connections with existing ones (e.g. in the user guide) would benefit the whole network. It is indeed expected that the knowledge and the use of all modelling tools will increase. It also underlines the need to keep the models and their documentation updated. Pioneering tools sometimes get outdated by new ones that do similar tasks but that are more user-friendly, faster, use the latest formalisms, or are better connected with newly existing tools. Therefore, either the interconnection between tools needs to be part of a huge maintenance effort for already existing tools, or the acceptance that pioneering tools are doomed to sink into oblivion.

Making the different tools freely available to the community is also a key aspect in their long term maintenance (Lobet, 2017). Many different repositories and licences exist so that everyone should be able to find a combination that suits their (and their institution's) needs. Free access to the tools' source codes would indeed greatly facilitate their evolution, reproducibility of the in silico experiments and allow future developers to interconnect them more easily.

## Acknowledgments

The authors would like to thank Jennifer Brophy, Heike Lindner and Thérèse LaRue (Carnegie Institution for Science, Stanford, CA) for their useful comment on the first version of manuscript. This work was also supported by the Belgian French community ARC 16/21-075 project. VC was also supported by the Interuniversity Attraction Poles Programme-Belgian Science Policy (grant IAP7/29). During the preparation of this manuscript, FM was supported by the “Fonds National de la Recherche Scientifique” (FNRS) of Belgium as a Research Fellow and is grateful to this organization for its financial support.

## Author contributions

	SP	VC	FM	XD	MJ	DL	LP	AS	JV	GL
Conceptualization										
Visualization										
Writing - draft										
Writing - review / editing										

## References

- Ahmed MA, Kroener E, Holz M, Zarebanadkouki M, Carminati A.** 2014. Mucilage exudation facilitates root water uptake in dry soils. *Functional plant biology: FPB* **41**, 1129–1137.
- Ahmed MA, Zarebanadkouki M, Meunier F, Javaux M, Kaestner A, Carminati A.** 2018. Root type matters: measurement of water uptake by seminal, crown, and lateral roots in maize. *Journal of experimental botany* **69**, 1199–1206.
- Alsina MM, Smart DR, Bauerle T, de Herralde F, Biel C, Stockert C, Negron C, Save R.** 2011. Seasonal changes of whole root system conductance by a drought-tolerant grape root system. *Journal of experimental botany* **62**, 99–109.
- Band LR, Wells DM, Fozard JA, et al.** 2014. Systems Analysis of Auxin Transport in the Arabidopsis Root Apex. *The Plant cell*.
- Baram S, Couvreur V, Harter T, Read M, Brown PH, Kandelous M, Smart DR, Hopmans JW.** 2016. Estimating Nitrate Leaching to Groundwater from Orchards: Comparing Crop Nitrogen Excess, Deep Vadose Zone Data-Driven Estimates, and HYDRUS Modeling. *Vadose Zone Journal* **15**, 0.
- Barczy J-F, Rey H, Griffon S, Jourdan C.** 2018. DigR: a generic model and its open source simulation software to mimic three-dimensional root-system architecture diversity. *Annals of botany*.
- Bourion V, Martin C, de Larambergue H, et al.** 2014. Unexpectedly low nitrogen acquisition and absence of root architecture adaptation to nitrate supply in a *Medicago truncatula* highly branched root mutant. *Journal of experimental botany* **65**, 2365–2380.
- Bramley H, Turner NC, Turner DW, Tyerman SD.** 2009. Roles of morphology, anatomy, and aquaporins in determining contrasting hydraulic behavior of roots. *Plant physiology* **150**, 348–364.
- Brodersen CR, Knipfer T, McElrone AJ.** 2018. In vivo visualization of the final stages of xylem vessel refilling in grapevine (*Vitis vinifera*) stems. *The New phytologist* **217**, 117–126.
- Brodersen CR, McElrone AJ, Choat B, Lee EF, Shackel KA, Matthews MA.** 2013. In vivo visualizations of drought-induced embolism spread in *Vitis vinifera*. *Plant physiology* **161**, 1820–1829.
- Burton AL, Williams M, Lynch JP, Brown KM.** 2012. RootScan: Software for high-throughput analysis of root anatomical traits. *Plant and soil* **357**, 189–203.
- Cai G, Vanderborght J, Couvreur V, Mboh CM, Vereecken H.** 2017. Parameterization of Root Water Uptake Models Considering Dynamic Root Distributions and Water Uptake Compensation. *Vadose Zone Journal* **0**, 0.
- Cai G, Vanderborght J, Klotzsche A, van der Kruk J, Neumann J, Hermes N, Vereecken H.** 2016. Construction of Minirhizotron Facilities for Investigating Root Zone Processes. *Vadose Zone Journal* **15**, 1–13.
- Cai G, Vanderborght J, Langensiepen M, Schnepf A, Hüging H, Vereecken H.** 2018. Root growth, water uptake, and sap flow of winter wheat in response to different soil water conditions. *Hydrology and Earth System Sciences* **22**, 2449.
- Carminati A, Moradi AB, Vetterlein D, Vontobel P, Lehmann E, Weller U, Vogel H-J, Oswald SE.** 2010. Dynamics of soil water content in the rhizosphere. *Plant and soil* **332**, 163–176.
- Carminati A, Passioura JB, Zarebanadkouki M, Ahmed MA, Ryan PR, Watt M, Delhaize E.** 2017. Root hairs enable high transpiration rates in drying soils. *The New phytologist*.
- Carminati A, Schneider CL, Moradi AB, Zarebanadkouki M, Vetterlein D, Vogel H-J, Hildebrandt A, Weller U, Schüler L, Oswald SE.** 2011. How the rhizosphere may favor water availability to roots. *Vadose Zone Journal* **10**, 988–998.
- Carminati A, Vetterlein D.** 2013. Plasticity of rhizosphere hydraulic properties as a key for efficient utilization of scarce resources. *Annals of botany* **112**, 277–290.

- Cattivelli L, Rizza F, Badeck F-W, Mazzucotelli E, Mastrangelo AM, Francia E, Marè C, Tondelli A, Stanca AM.** 2008. Drought tolerance improvement in crop plants: An integrated view from breeding to genomics. *Field crops research* **105**, 1–14.
- Choat B, Cobb AR, Jansen S.** 2008. Structure and function of bordered pits: new discoveries and impacts on whole-plant hydraulic function. *The New phytologist* **177**, 608–625.
- Choat B, Jansen S, Brodribb TJ, et al.** 2012. Global convergence in the vulnerability of forests to drought. *Nature* **491**, 752–755.
- Chopin J, Laga H, Huang CY, Heuer S, Miklavcic SJ.** 2015. RootAnalyzer: A Cross-Section Image Analysis Tool for Automated Characterization of Root Cells and Tissues. *PLoS one* **10**, e0137655.
- Colombi T, Kirchgessner N, Le Marié CA, York LM, Lynch JP, Hund A.** 2015. Next generation shovelomics: set up a tent and REST. *Plant and soil*.
- Costa G, Plazanet I.** 2016. Plant Cell Wall, a Challenge for Its Characterisation. *Advances in biological chemistry* **06**, 70–105.
- Couvreur V, Faget M, Lobet G, Javaux M, Chaumont F, Draye X.** 2017. Novel multiscale insights into the composite nature of water transport in roots. *bioRxiv*, 147314.
- Couvreur V, Kandelous MM, Sanden BL, Lampinen BD, Hopmans JW.** 2016. Downscaling transpiration rate from field to tree scale. *Agricultural and Forest Meteorology* **221**, 71–77.
- Couvreur V, Ledder G, Manzoni S, Way DA, Muller EB, Russo SE.** 2018. Water transport through tall trees: A vertically-explicit, analytical model of xylem hydraulic conductance in stems. *Plant, cell & environment*.
- Couvreur V, Vanderborght J, Beff L, Javaux M.** 2014a. Horizontal soil water potential heterogeneity: simplifying approaches for crop water dynamics models. *Hydrology and Earth System Sciences Discussions* **11**, 1203–1252.
- Couvreur V, Vanderborght J, Draye X, Javaux M.** 2014b. Dynamic aspects of soil water availability for isohydric plants: Focus on root hydraulic resistances. *Water resources research* **50**, 8891–8906.
- Couvreur V, Vanderborght J, Javaux M.** 2012. A simple three-dimensional macroscopic root water uptake model based on the hydraulic architecture approach. *Hydrology and Earth System Sciences* **16**, 2957–2971.
- Cox PM, Huntingford C, Harding RJ.** 1998. A canopy conductance and photosynthesis model for use in a GCM land surface scheme. *Journal of Hydrology* **212-213**, 79–94.
- David TS, David JS, Pinto CA, Cermak J, Nadezhdin V, Nadezhdina N.** 2012. Hydraulic connectivity from roots to branches depicted through sap flow: analysis on a *Quercus suber* tree. *Functional plant biology: FPB* **39**, 103–115.
- Dencic S.** 1994. Designing a Wheat Ideotype with Increased Sink Capacity. *Plant breeding = Zeitschrift für Pflanzenzucht* **112**, 311–317.
- Doussan C, Pagès L, Vercambre G.** 1998. Modelling of the hydraulic architecture of root systems: An integrated approach to water absorption - Model description. *Annals of botany* **81**, 213–223.
- Doussan C, Pierret A, Garrigues E, Pagès L.** 2006. Water Uptake by Plant Roots: II – Modelling of Water Transfer in the Soil Root-system with Explicit Account of Flow within the Root System – Comparison with Experiments. *Plant and soil* **283**, 99–117.
- Draye X, Kim Y, Lobet G, Javaux M.** 2010. Model-assisted integration of physiological and environmental constraints affecting the dynamic and spatial patterns of root water uptake from soils. *Journal of experimental botany* **61**, 2145–2155.
- Dupuy L, Fourcaud T, Stokes A, Danjon F.** 2005. A density-based approach for the modelling of root architecture: application to Maritime pine (*Pinus pinaster* Ait.) root systems. *Journal of theoretical biology* **236**, 323–334.
- Dupuy L, Gregory PJ, Bengough AG.** 2010. Root growth models: towards a new generation of continuous approaches. *Journal of experimental botany* **61**, 2131–2143.

**Enns L.** 1998. Solute concentrations in xylem sap along vessels of maize primary roots at high root pressure. *Journal of experimental botany* **49**, 1539–1544.

**Esser HG, Carminati A, Vontobel P, Lehmann EH, Oswald SE.** 2010. Neutron radiography and tomography of water distribution in the root zone. *Journal of Plant Nutrition and Soil Science* **173**, 757–764.

**Famiglietti JS, Ryu D, Berg AA, Rodell M, Jackson TJ.** 2008. Field observations of soil moisture variability across scales. *Water resources research* **44**.

**Feddes RA, Kowalik PJ, Zaradny H.** 1978. *Simulation of Field Water Use and Crop Yield*. Wiley.

**Garré S, Coteur I, Wongleecharoen C, Hussain K, Omsunrarn W, Kongkaew T, Hilger T, Diels J, Vanderborght J.** 2013. Can We Use Electrical Resistivity Tomography to Measure Root Zone Dynamics in Fields with Multiple Crops? *Procedia Environmental Sciences* **19**, 403–410.

**Garré S, Pagès L, Laloy E, Javaux M, Vanderborght\* J, Vereecken H.** 2012. Parameterizing a Dynamic Architectural Model of the Root System of Spring Barley from Minirhizotron Data. *Vadose Zone Journal* **11**, 0.

**Garrigues E, Doussan C, Pierret A.** 2006. Water Uptake by Plant Roots: I – Formation and Propagation of a Water Extraction Front in Mature Root Systems as Evidenced by 2D Light Transmission Imaging. *Plant and soil* **283**, 83–98.

**van Genuchten MT.** 1980. A closed-form equation for predicting the hydraulic conductivity of unsaturated soils. *Soil Science Society of America journal*. *Soil Science Society of America* **44**, 891–898.

**van Genuchten M, Leij FJ, Yates. MS.** 1991. *The RETC Code for Quantifying the Hydraulic Functions of Unsaturated Soils*. EPA Report 600/2-91/065, U.S. Salinity Laboratory, USDA, ARS, Riverside, California.

**Gijsman AJ, Jagtap SS, Jones JW.** 2002. Wading through a swamp of complete confusion: how to choose a method for estimating soil water retention parameters for crop models. *European journal of agronomy: the journal of the European Society for Agronomy* **18**, 77–106.

**Giordano R, Salleo A, Salleo S, Wanderlingh F.** 1978. Flow in xylem vessels and Poiseuille's law. *Canadian journal of botany. Journal canadien de botanique* **56**, 333–338.

**Godin C, Costes E, Sinoquet H.** 1999. A Method for Describing Plant Architecture which Integrates Topology and Geometry. *Annals of botany* **84**, 343–357.

**Godin C, Sinoquet H.** 2005. Functional-structural plant modelling. *The New phytologist* **166**, 705–708.

**Guswa AJ.** 2010. Effect of plant uptake strategy on the water–optimal root depth. *Water resources research* **46**.

**Hack-ten Broeke MJD, Kroes JG, Bartholomeus RP, van Dam JC, de Wit AJW, Supit I, Walvoort DJJ, van Bakel PJT, Ruijtenberg R.** 2016. Quantification of the impact of hydrology on agricultural production as a result of too dry, too wet or too saline conditions.

**Hainsworth JM, Aylmore L.** 1983. The use of computer assisted tomography to determine spatial distribution of soil water content. *Australian Journal of Soil Research* **21**, 435.

**Holzworth D, Huth NI, Fainges J, Brown H, Zurcher E, Cichota R, Verrall S, Herrmann NI, Zheng B, Snow V.** 2018. APSIM Next Generation: Overcoming challenges in modernising a farming systems model. *Environmental Modelling & Software* **103**, 43–51.

**Hopmans JW, Nielsen DR, Bristow KL.** 2002. How useful are small-scale soil hydraulic property measurements for large-scale vadose zone modeling? *Geophysical Monograph Series*. 247–258.

**Hose E, Steudle E, Hartung W.** 2000. Abscisic acid and hydraulic conductivity of maize roots: a study using cell- and root-pressure probes. *Planta* **211**, 874–882.

**Huber K, Vanderborght J, Javaux M, Schröder N, Dodd IC, Vereecken H.** 2014. Modelling the impact of heterogeneous rootzone water distribution on the regulation of transpiration by hormone transport and/or hydraulic pressures. *Plant and soil*.

**Huber K, Vanderborght J, Javaux M, Vereecken H.** 2015. Simulating transpiration and leaf water relations in



response to heterogeneous soil moisture and different stomatal control mechanisms. *Plant and soil* **394**, 109–126.

**Jahnke S, Menzel MI, van Dusschoten D, et al.** 2009. Combined MRI-PET dissects dynamic changes in plant structures and functions. *The Plant journal: for cell and molecular biology* **59**, 634–644.

**Javaux M, Schroeder T, Vanderborght J, Vereecken H.** 2008. Use of a three-dimensional detailed modeling approach for predicting root water uptake. *Vadose Zone Journal* **7**, 1079–1088.

**Keyes SD, Daly KR, Gostling NJ, Jones DL, Talboys P, Pinzer BR, Boardman R, Sinclair I, Marchant A, Roose T.** 2013. High resolution synchrotron imaging of wheat root hairs growing in soil and image based modelling of phosphate uptake. *The New phytologist*.

**Kuchenbuch RO, Horst GH, Buczko U.** 2009. Spatial distribution of maize roots by complete 3D soil monolith sampling. *Plant and soil* **315**, 297–314.

**Landl M, Huber K, Schnepf A, Vanderborght J, Javaux M, Glyn Bengough A, Vereecken H.** 2016. A new model for root growth in soil with macropores. *Plant and soil*, 1–18.

**Lartaud M, Perin C, Courtois B, et al.** 2014. PHIV-RootCell: a supervised image analysis tool for rice root anatomical parameter quantification. *Frontiers in plant science* **5**, 790.

**Leitner D, Felderer B, Vontobel P, Schnepf A.** 2014a. Recovering root system traits using image analysis exemplified by two-dimensional neutron radiography images of lupine. *Plant physiology* **164**, 24–35.

**Leitner D, Klepsch S, Bodner G, Schnepf A.** 2010. A dynamic root system growth model based on L-Systems. *Plant and soil* **332**, 117–192.

**Leitner D, Meunier F, Bodner G, Javaux M, Schnepf A.** 2014b. Impact of contrasted maize root traits at flowering on water stress tolerance – A simulation study. *Field crops research* **165**, 125–137.

**Lewis AM, Boose ER.** 1995. Estimating Volume Flow Rates Through Xylem Conduits. *American journal of botany* **82**, 1112.

**Lobet G.** 2017. Image Analysis in Plant Sciences: Publish Then Perish. *Trends in plant science* **22**, 559–566.

**Lobet G, Couvreur V, Meunier F, Javaux M, Draye X.** 2014a. Plant water uptake in drying soils. *Plant physiology* **164**, 1619–1627.

**Lobet G, Draye X, Périlleux C.** 2013. An online database for plant image analysis software tools. *Plant methods* **9**, 38.

**Lobet G, Pagès L, Draye X.** 2014b. A modeling approach to determine the importance of dynamic regulation of plant hydraulic conductivities on the water uptake dynamics in the soil-plant-atmosphere system. *Ecological modelling* **290**, 65–75.

**Lobet G, Pound MP, Diener J, et al.** 2015. Root System Markup Language: Toward a Unified Root Architecture Description Language. *Plant physiology* **167**, 617–627.

**Lynch JP.** 2013. Steep, cheap and deep: an ideotype to optimize water and N acquisition by maize root systems. *Annals of botany* **112**, 347–357.

**Lynch JP, Nielsen KL, Davis RD, Jablowski AG.** 1997. SimRoot: Modelling and visualization of root systems. *Plant and soil* **188**, 139–151.

**Mairhofer S, Zappala S, Tracy SR, Sturrock C, Bennett M, Mooney SJ, Pridmore T.** 2012. RooTrak: automated recovery of three-dimensional plant root architecture in soil from x-ray microcomputed tomography images using visual tracking. *Plant physiology* **158**, 561–569.

**Malone M, Leigh RA, Tomos AD.** 1989. Extraction and analysis of sap from individual wheat leaf cells: the effect of sampling speed on the osmotic pressure of extracted sap. *Plant, cell & environment* **12**, 919–926.

**McElrone AJ, Shapland TM, Calderon A, Fitzmaurice L, Paw U KT, Snyder RL.** 2013. Surface renewal: an advanced micrometeorological method for measuring and processing field-scale energy flux density data. *Journal of visualized experiments: JoVE*, e50666.

- Meunier F, Couvreur V, Draye X, Vanderborcht J, Javaux M.** 2017a. Towards quantitative root hydraulic phenotyping: novel mathematical functions to calculate plant-scale hydraulic parameters from root system functional and structural traits. *Journal of mathematical biology* **75**, 1133–1170.
- Meunier F, Draye X, Vanderborcht J, Javaux M, Couvreur V.** 2017b. A hybrid analytical-numerical method for solving water flow equations in root hydraulic architectures. *Applied mathematical modelling* **52**, 648–663.
- Meunier F, Rothfuss Y, Bariac T, Biron P, Richard P, Durand J-L, Couvreur V, Vanderborcht J, Javaux M.** 2018a. Measuring and Modeling Hydraulic Lift of Using Stable Water Isotopes. *Vadose Zone Journal* **17**, 0.
- Meunier F, Zarebanadkouki M, Ahmed MA, Carminati A, Couvreur V, Javaux M.** 2018b. Hydraulic conductivity of soil-grown lupine and maize unbranched roots and maize root-shoot junctions. *Journal of plant physiology*.
- Mirfenderesgi G, Bohrer G, Matheny AM, Fatichi S, Frasson M, Prata R, Schäfer KVR.** 2016. Tree level hydrodynamic approach for resolving aboveground water storage and stomatal conductance and modeling the effects of tree hydraulic strategy. *Journal of Geophysical Research: Biogeosciences* **121**, 1792–1813.
- Mrad A, Domec J-C, Huang C-W, Lens F, Katul G.** 2018. A network model links wood anatomy to xylem tissue hydraulic behavior and vulnerability to cavitation. *Plant, cell & environment*.
- Ndour A, Vadez V, Pradal C, Lucas M.** 2017. Virtual Plants Need Water Too: Functional-Structural Root System Models in the Context of Drought Tolerance Breeding. *Frontiers in plant science* **8**, 1577.
- Oyanagi A, Nakamoto T, Morita S.** 1993. The gravitropic response of roots and the shaping of the root system in cereal plants. *Environmental and experimental botany* **33**, 141–158.
- Paez-Garcia A, Motes CM, Scheible W-R, Chen R, Blancaflor EB, Monteros MJ.** 2015. Root Traits and Phenotyping Strategies for Plant Improvement. *Plants* **4**, 334–355.
- Pagès L.** 2011. Links between root developmental traits and foraging performance. *Plant, cell & environment* **34**, 1749–1760.
- Pagès L, Bécel C, Boukcim H, Moreau D, Nguyen C, Voisin A-S.** 2013. Calibration and evaluation of ArchiSimple, a simple model of root system architecture. *Ecological modelling* **290**, 76–84.
- Pagès L, Bécel C, Boukcim H, Moreau D, Nguyen C, Voisin A-S.** 2014. Calibration and evaluation of ArchiSimple, a simple model of root system architecture. *Ecological modelling* **290**, 76–84.
- Pagès L, Vercambre G, Drouet J-L, Lecompte F, Collet C, LeBot J.** 2004. RootTyp: a generic model to depict and analyse the root system architecture. *Plant and soil* **258**, 103–119.
- Passot S, Gnacko F, Moukouanga D, et al.** 2016. Characterization of Pearl Millet Root Architecture and Anatomy Reveals Three Types of Lateral Roots. *Frontiers in plant science* **7**, 829.
- Pohlmeier A.** 2010. MRI in Soils: Determination of Water Content Changes Due to Root Water Uptake by Means of a Multi-Slice-Multi-Echo Sequence (MSME). *The Open Magnetic Resonance Journal* **3**, 69–74.
- Postma JA, Kuppe C, Owen MR, Mellor N, Griffiths M, Bennett MJ, Lynch JP, Watt M.** 2017a. OpenSimRoot: widening the scope and application of root architectural models. *The New phytologist*.
- Postma JA, Kuppe C, Owen MR, Mellor N, Griffiths M, Bennett MJ, Lynch JP, Watt M.** 2017b. OpenSimRoot: widening the scope and application of root architectural models. *The New phytologist* **215**, 1274–1286.
- Pound MP, French AP, Wells DM, Bennet JM, Pridmore TP.** 2012. CellSeT: Novel Software to Extract and Analyze Structured Networks of Plant Cells from Confocal Images. *The Plant cell*.
- Pritchard J, Barlow PW, Adam JS, Tomos AD.** 1990. Biophysics of the inhibition of the growth of maize roots by lowered temperature. *Plant physiology* **93**, 222–230.
- Rascher U, Blossfeld S, Fiorani F, et al.** 2011. Non-invasive approaches for phenotyping of enhanced performance traits in bean. *Functional plant biology: FPB* **38**, 968.
- Rewald B, Ephrath J.** 2013. Minirhizotron Techniques. In: Beeckman T, ed. *Plant Roots*. CRC Press, 42–1–42–16.

- Robbins NE 2nd, Dinneny JR.** 2018. Growth is required for perception of water availability to pattern root branches in plants. *Proceedings of the National Academy of Sciences of the United States of America*.
- Rodrigues ML, Lucília Rodrigues M, Santos TP, Rodrigues AP, de Souza CR, Lopes CM, Maroco JP, Pereira JS, Manuela Chaves M.** 2008. Hydraulic and chemical signalling in the regulation of stomatal conductance and plant water use in field grapevines growing under deficit irrigation. *Functional plant biology: FPB* **35**, 565.
- Rodriguez-Dominguez CM, Carins Murphy MR, Lucani C, Brodribb TJ.** 2018. Mapping xylem failure in disparate organs of whole plants reveals extreme resistance in olive roots. *The New phytologist* **218**, 1025–1035.
- Roose T, Keyes SD, Daly KR, Carminati A, Otten W, Vetterlein D, Peth S.** 2016. Challenges in imaging and predictive modeling of rhizosphere processes. *Plant and soil*, 1–30.
- Sanderson J, Whitbread FC, Clarkson DT.** 1988. Persistent xylem cross-walls reduce the axial hydraulic conductivity in the apical 20 cm of barley seminal root axes: implications for the driving force for water movement. *Plant, cell & environment* **11**, 247–256.
- Schneider CL, Attinger S, Delfs J-O, Hildebrandt A.** 2009. Implementing small scale processes at the soil-plant interface – the role of root architectures for calculating root water uptake profiles. *Hydrology and Earth System Sciences Discussions* **6**, 4233–4264.
- Schnepf A, Leitner D, Landl M, Lobet G, Mai TH, Morandage S, Sheng C, Zoerner M, Vanderborght J, Vereecken H.** 2017. CRooTBox: A Structural-Functional Modelling Framework For Root Systems. *bioRxiv*, 139980.
- Schnepf A, Leitner D, Landl M, Lobet G, Mai TH, Morandage S, Sheng C, Zörner M, Vanderborght J, Vereecken H.** 2018. CRooTBox: a structural–functional modelling framework for root systems. *Annals of botany*.
- Schoppach R my, Wauthélet D, Jeanguenin L, Sadok W.** 2013. Conservative water use under high evaporative demand associated with smaller root metaxylem and limited trans-membrane water transport in wheat. *Functional plant biology: FPB*.
- Schroeder N, Javaux M, Vanderborght J, Steffen B, Vereecken H.** 2012. Effect of Root Water and Solute Uptake on Apparent Soil Dispersivity: A Simulation Study. *Vadoze Zone Journal*.
- Schwartz N, Carminati A, Javaux M.** 2016. The impact of mucilage on root water uptake—A numerical study. *Water resources research* **52**, 264–277.
- Shane MW, McCully ME, Canny MJ.** 2000. Architecture of Branch-root Junctions in Maize: Structure of the Connecting Xylem and the Porosity of Pit Membranes. *Annals of botany* **85**, 613–624.
- Shrestha P, Sulis M, Masbou M, Kollet S, Simmer C.** 2014. A Scale-Consistent Terrestrial Systems Modeling Platform Based on COSMO, CLM, and ParFlow. *Monthly Weather Review* **142**, 3466–3483.
- Šimůnek J, Hopmans JW.** 2009. Modeling compensated root water and nutrient uptake. *Ecological modelling* **220**, 505–521.
- Simunek J, Van Genuchten M.** 1995. Numerical model for simulating multiple solute transport in variably-saturated soils. *Ground water*.
- Skelton RP, Brodribb TJ, Choat B.** 2017. Casting light on xylem vulnerability in an herbaceous species reveals a lack of segmentation. *The New phytologist* **214**, 561–569.
- Sperry JS, Hacke UG, Oren R, Comstock JP.** 2002. Water deficits and hydraulic limits to leaf water supply. *Plant, cell & environment* **25**, 251–263.
- Stedle E.** 1980. Water-relation Parameters of Individual Mesophyll Cells of the Crassulacean Acid Metabolism Plant *Kalanchoë daigremontiana*. *Plant physiology* **66**, 1155–1163.
- Stedle E, Jeschke WD.** 1983. Water transport in barley roots : Measurements of root pressure and hydraulic conductivity of roots in parallel with turgor and hydraulic conductivity of root cells. *Planta* **158**, 237–248.
- Stedle E, Rr O, Schulze E.** 1987. Water transport in maize roots. Measurement of hydraulic conductivity, solute



permeability and of ....

**Stoma S, Lucas M, Chopard J, Schaedel M, Traas J, Godin C.** 2008. Flux-based transport enhancement as a plausible unifying mechanism for auxin transport in meristem development. *PLoS computational biology* **4**, e1000207.

**Tardieu F.** 2012. Any trait or trait-related allele can confer drought tolerance: just design the right drought scenario. *Journal of experimental botany* **63**, 25–31.

**Tixier A, Herbette S, Jansen S, Capron M, Tordjeman P, Cochard H, Badel E.** 2014. Modelling the mechanical behaviour of pit membranes in bordered pits with respect to cavitation resistance in angiosperms. *Annals of botany* **114**, 325–334.

**Tötzke C, Kardjilov N, Manke I, Oswald SE.** 2017. Capturing 3D Water Flow in Rooted Soil by Ultra-fast Neutron Tomography. *Scientific reports* **7**, 6192.

**Trachsel S, Kaeppler SM, Brown KM, Lynch JP.** 2010. Shovelomics: high throughput phenotyping of maize (*Zea mays* L.) root architecture in the field. *Plant and soil* **341**, 75–87.

**Tsuda M, Tyree MT.** 2000a. Plant hydraulic conductance measured by the high pressure flow meter in crop plants. *Journal of experimental botany* **51**, 823–828.

**Tsuda M, Tyree MT.** 2000b. Plant hydraulic conductance measured by the high pressure flow meter in crop plants. *Journal of experimental botany* **51**, 823.

**Uga Y, Okuno K, Yano M.** 2011. Drr1, a major QTL involved in deep rooting of rice under upland field conditions. *Journal of experimental botany* **62**, 2485–2494.

**Vadez V.** 2014. Root hydraulics: The forgotten side of roots in drought adaptation. *Field crops research*, 1–10.

**Vansteenkiste J, Van Loon J, Garré S, Pagès L, Schrevens E, Diels J.** 2013. Estimating the parameters of a 3-D root distribution function from root observations with the trench profile method: case study with simulated and field-observed root data. *Plant and soil* **375**, 75–88.

**Vepraskas MJ, Hoyt GD.** 1988. Comparison of the trench-profile and core methods for evaluating root distributions in tillage studies. *Agronomy journal* **80**, 166–172.

**Vereecken H, Schnepf A, Hopmans JW, et al.** 2016. Modeling Soil Processes: Review, Key Challenges, and New Perspectives. *Vadose Zone Journal* **15**, 0.

**Verhoef A, Egea G.** 2014. Modeling plant transpiration under limited soil water: Comparison of different plant and soil hydraulic parameterizations and preliminary implications for their use in land surface models. *Agricultural and Forest Meteorology* **191**, 22–32.

**Vrugt JA, Schoups G, Hopmans JW, Young C, Wallender WW, Harter T, Bouten W.** 2004. Inverse modeling of large-scale spatially distributed vadose zone properties using global optimization. *Water resources research* **40**.

**Vrugt JA, van Wijk MT, Hopmans JW, Šimunek J.** 2001. One-, two-, and three-dimensional root water uptake functions for transient modeling. *Water resources research* **37**, 2457–2470.

**Warren JM, Bilheux H, Kang M, Voisin S, Cheng C-L, Horita J, Perfect E.** 2013. Neutron imaging reveals internal plant water dynamics. *Plant and soil* **366**, 683–693.

**Wasson AP, Rebetzke GJ, Kirkegaard JA, Christopher J, Richards RA, Watt M.** 2014. Soil coring at multiple field environments can directly quantify variation in deep root traits to select wheat genotypes for breeding. *Journal of experimental botany* **65**, 6231–6249.

**Wasson AP, Richards RA, Chatrath R, Misra SC, Prasad SVS, Rebetzke GJ, Kirkegaard JA, Christopher J, Watt M.** 2012. Traits and selection strategies to improve root systems and water uptake in water-limited wheat crops. *Journal of experimental botany* **63**, 3485–3498.

**Whitley R, Taylor D, Macinnis-Ng C, Zeppel M, Yunusa I, O'Grady A, Froend R, Medlyn B, Eamus D.** 2012. Developing an empirical model of canopy water flux describing the common response of transpiration to solar

radiation and VPD across five contrasting woodlands and forests. *Hydrological processes* **27**, 1133–1146.

**Zarebanadkouki M, Kroener E, Kaestner A, Carminati A.** 2014*a*. Visualization of root water uptake: quantification of deuterated water transport in roots using neutron radiography and numerical modeling. *Plant physiology* **166**, 487–499.

**Zarebanadkouki M, Kroener E, Kaestner A, Carminati A.** 2014*b*. Visualization of root water uptake: quantification of deuterated water transport in roots using neutron radiography and numerical modeling. *Plant physiology* **166**, 487–499.

**Zarebanadkouki M, Meunier F, Couvreur V, Cesar J, Javaux M, Carminati A.** 2016. Estimation of the hydraulic conductivities of lupine roots by inverse modelling of high-resolution measurements of root water uptake. *Annals of botany*.

**Zeng G, Birchfield ST, Wells CE.** 2008. Automatic discrimination of fine roots in minirhizotron images. *The New phytologist* **117**, 549–557.

Accepted Manuscript

## Figure legends

### Figure 1

**Quantifying water relations in the soil plant system.** Tools, properties and state variables used to quantify: (A) the structure of root organ; (B) the structure of root system; (C) the structure of root profiles; (D) the water flow at the organ scale (root section); (E) the water flow at the plant scale (root system); (F) the water flow at the population scale; (G) the water flow in the soil. Without appropriate tools, variables of interest, scales and even plants and their environment seem disconnected.

### Figure 2

**Details of the connected dots to compute the hydraulic properties of the different root types of pearl millet.** Colored parts are the tools, models, properties and state variables used in the approach. Specific tools names were added where relevant. See text for details. A: CellSet, B: Hagen-Poiseuille, C: MECHA, D: Output of the different models

### Figure 3

**Details of the connected dots for root system architecture generation.** Colored parts are the tools, models and properties used in the approach. Specific tools names were added where relevant. See text for details. A: Published data, B: Root System Analyzer, C: RootBox

## Figure 4

**Details of the connected dots for estimating root conductivities from experimental observations through inverse modelling.** Colored parts are the tools, models and inputs used in the approach. Specific tools names were added where relevant. The red arrow indicate the optimisation step used in the inverse modelling. The red boxes highlight the variables evaluated using the inverse modelling. See text for details. A: SmartRoot, B: Hybrid water flow model, C: Optimisation, D: Root hydraulic properties

## Figure 5

**Details of the connected dots for estimate soil and plant-scale conductivities through inverse modelling.** Colored parts are the tools, models and inputs used in the approach. Specific tools names were added where relevant. The red arrow indicate the optimisation step used in the inverse modelling. The red boxes highlight the variables evaluated using the inverse modelling. See text for references to letters A-H.

## Figure 6

**Full network of tools and data used to quantify water flow in the soil plant system.** The network connects experimental procedures, computational tools and data related to water flow in the soil-plant system. It is organised by scales (organ, plant and population) and by the types of information (structural or functional, see text for details). A. Tools to quantify the water flow at the organ scale (root section). B. Tools to quantify the structure of root organ. C. Tools to quantify the water flow at the plant scale (root system). D. Tools to quantify the structure of root system. E. Tools to quantify the water flow at the population scale. F. Tools to quantify the structure of root profiles. G. Tools to quantify water flow in the soil.

## Tables

**Table 1: List of modelling tools fitting into the network.** For image analysis tools, we refer the reader to the [www.plant-image-analysis.org](http://www.plant-image-analysis.org) website

Type	Scale	Name	Reference
Functional	Radial water flow	MECHA	(Couvreur <i>et al.</i> , 2017)
Functional	Root system water flow	R-SWMS	(Javaux <i>et al.</i> , 2008)
Functional - structural	Root system water flow	PlaNet-Maize	(Lobet <i>et al.</i> , 2014b)
Functional - structural	Root system water flow	OpenSimRoot	(Postma <i>et al.</i> , 2017a)
Functional	Root system water flow		(Doussan <i>et al.</i> , 2006)
Functional	Soil water flow	HYDRUS	(Simunek and Van Genuchten, 1995)
Functional	Soil water flow	RSWMS	(Javaux <i>et al.</i> , 2008)
Structural	Root system architecture	CRootBox	(Schnepf <i>et al.</i> , 2017)
Structural	Root system architecture	ArchiSimple	(Pagès <i>et al.</i> , 2013)
Structural	Root system architecture	RootTyp	(Pagès <i>et al.</i> , 2004)
Structural	Root system architecture	DigR	(Barczy <i>et al.</i> , 2018)
Structural	Root system density		(Dupuy <i>et al.</i> , 2005)






# ORGAN SCALE





# PLANT SCALE

# POPULATION SCALE

## CELL



## STRUCTURE



## FUNCTION


## PLANT

### LEGEND

- Experimental procedure
- Image analysis tool
- Modelling tool
- Parameter
- Upscaled property
- Measured property
- Boundary condition
- Predicted state variable
- Measured state variable
- Hypothetical tool or connection


Properties

State variables




## RHIZOSPHERE

## BULK SOIL




Downloaded from https://academic.oup.com/jxb/advance-article-abstract/doi/10.1093/jxb/erz391/5133494 by INRA (Institut National de la Recherche Agronomique) user on 30 October 2018

### WHOLE NETWORK OVERVIEW



### LEGEND



**A**

Published data

**B**



Root System Analyser

Root architecture parameters


Full tracing

Root architecture


RootBox

**C****WHOLE NETWORK OVERVIEW****LEGEND**







**WHOLE NETWORK OVERVIEW**




**LEGEND**





Downloaded from https://academic.oup.com/jxb/advance-article-abstract/doi/10.1093/jxb/ryz381/5533494 by INRA (Institut National de la Recherche Agronomique) user on 30 October 2018

## WHOLE NETWORK OVERVIEW



## LEGEND

