



HAL
open science

Tree aging does not affect the ranking for water use efficiency recorded from $\delta^{13}\text{C}$ in three *Populus deltoides* x *P. nigra* genotypes

Fahad Rasheed, Erwin Dreyer, D. Le Thiec, Zikria Zafar, Sylvain Delagrangé

► To cite this version:

Fahad Rasheed, Erwin Dreyer, D. Le Thiec, Zikria Zafar, Sylvain Delagrangé. Tree aging does not affect the ranking for water use efficiency recorded from $\delta^{13}\text{C}$ in three *Populus deltoides* x *P. nigra* genotypes. *iForest: Biogeosciences and Forestry*, 2019, 12 (3), pp.272-278. 10.3832/ifor2896-012 . hal-02624170

HAL Id: hal-02624170

<https://hal.inrae.fr/hal-02624170v1>

Submitted on 26 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Tree aging does not affect the ranking for water use efficiency recorded from $\delta^{13}\text{C}$ in three *Populus deltoides* × *P. nigra* genotypes

Fahad Rasheed⁽¹⁻²⁾,
Erwin Dreyer⁽¹⁻³⁾,
Didier Le Thiec⁽¹⁻³⁾,
Zikria Zafar⁽²⁾,
Sylvain Delagrangé⁽⁴⁾

A large variability of water use efficiency (assessed from the carbon isotopic discrimination in leaves and leaf soluble sugars) has been detected among poplar genotypes. Checking whether such differences detected in young trees (1-2 years old) remain stable with tree age is a prerequisite to use this trait with confidence for breeding purposes. In this study, a synchronic approach was used to test the age-related stability of the genotypic ranking of carbon isotopic discrimination in wood ($\Delta^{13}\text{C}$) until tree maturity. We sampled 376 trees between 4 and 20 years from three *Populus deltoides* × *P. nigra* genotypes growing in 41 common-garden trials across France. Carbon and nitrogen percentages along with $\delta^{13}\text{C}$ was measured in the bulk wood of the year 2009 and used to compute the $\Delta^{13}\text{C}$. Basal area increment between 2008 and 2009 was also measured. Results showed that $\Delta^{13}\text{C}$ increased (*i.e.*, water use efficiency decreased) between ages 4 to 6 and remained stable later on. Significant differences among genotypes were found but the ranking among genotypes remained stable with age during the assessed life span. Furthermore, basal area increment and $\Delta^{13}\text{C}$ were positively correlated interannually. This large-scale survey shows that despite crossing over in the temporal trend, water use efficiency remained stable with age across 3 poplar genotypes. However, further studies with a large number of genotypes are required to confirm whether this trait can be used to maintain or even improve productivity of poplar plantations, while lowering water consumption.

Keywords: Water Use Efficiency, Age, Wood, Tree Ring, *Populus* × *euramericana*, Basal Area Increment, Synchronic Approach

Introduction

In the context of climate change and fluctuating precipitation patterns (IPCC 2013), water use efficiency (WUE, ratio between biomass accumulated and water transpired) has become an interesting trait for breeding new genotypes (Condon et al. 2004, Richards et al. 2010, Vadez et al. 2014). Isotopic discrimination against ^{13}C ($\Delta^{13}\text{C}$) during CO_2 diffusion through stomata and photosynthesis is an indirect indicator of intrinsic water use efficiency (IWUE, ratio between net CO_2 assimilation rate, A , and stomatal conductance to water vapour, g_s – Farquhar & Richards 1984, Farquhar et al. 1989, Ehleringer et al. 1993). Indeed, this trait was found to be under tight genetic control in several tree species like poplar, pines, chestnut and oaks among others (Lauteri et al. 1997, Brendel et al. 2002, 2008, Rasheed et al. 2015). Furthermore, given the lack of correlation observed between WUE and productivity or, in some instances, the positive correlation between both traits (Vadez et al. 2014), selecting tree species or genotypes based on their ability to express a high WUE is desirable. Such a strategy has been explored in some crops in a context of improving water management (Condon et al. 2004, Richards et al. 2010) and might thus be particularly suitable for sustaining wood production under the uncertainty of water availability.

Poplar genotypes are known to display large differences in growth performance and biomass production (Rasheed et al. 2013). A large genotypic variability of $\Delta^{13}\text{C}$ was found among *Populus deltoides* × *P. nigra* genotypes in controlled environment (Marron et al. 2005) and open field experiments (Monclus et al. 2005, Dillen et al. 2008). Using direct measurements, this variability in $\Delta^{13}\text{C}$ is also found tightly related to variability in the whole plant water use efficiency in *Populus deltoides* × *P. nigra* and in *Populus nigra* (Rasheed et al. 2013, 2015). Interestingly, no correlation was found between productivity and $\Delta^{13}\text{C}$ (Monclus et al. 2005, Dillen et al. 2008), giving opportunity to breed for enhanced WUE without compromising productivity in poplar genotypes. Nevertheless, previous studies on poplar genotypes were conducted on seedlings of 1-2 years and there are few studies investigating the stability of genotype ranking for WUE at tree scale.

Previous studies has demonstrated that $\Delta^{13}\text{C}$ (recorded from tree ring cellulose and bulk wood) is severely affected by age, usually decreasing by a 1-3 per mil with increasing tree age (Francey & Farquhar 1982, Bert et al. 1997, Duquesnay et al. 1998, McCarroll & Pawellek 2001, Brienen et al. 2017). Such an age-related effect may be related to: (i) assimilation of respired CO_2 near the forest floor which is already depleted in ^{13}C (Schleser & Jayasekera

□ (1) INRA, UMR1137 Ecologie et Ecophysiologie Forestières, IFR 110 EFABA, F-54280 Champenoux (France); (2) Department of Forestry & Range Management, University of Agriculture Faisalabad, P.O. Box 38000 (Pakistan); (3) Université de Lorraine, UMR1137 Ecologie et Ecophysiologie Forestières, IFR 110 EFABA, F-54500 Vandoeuvre-lès-Nancy (France); (4) Department of Natural Sciences, Institute of Temperate Forest Sciences (ISFORT), University of Quebec in Outaouais (UQO), 58 Main St, Ripon, Quebec J0V 1V0 (Canada)

@ Fahad Rasheed
(fahad.rasheed@uaf.edu.pk)

Received: Jun 15, 2018 - Accepted: Mar 18, 2019

Citation: Rasheed F, Dreyer E, Le Thiec D, Zafar Z, Delagrangé S (2019). Tree aging does not affect the ranking for water use efficiency recorded from $\delta^{13}\text{C}$ in three *Populus deltoides* × *P. nigra* genotypes. *iForest* 12: 272-278. - doi: 10.3832/ifor2896-012 [online 2019-05-21]

Communicated by: Rossella Guerrieri

1985); (ii) re-assimilation of respired CO₂ retained within the plant canopy (Francey & Farquhar 1982); (iii) decreasing hydraulic conductivity of xylem conduits with age (Monserud & Marshall 2001); (iv) reduced total soil-to-leaf hydraulic conductance with tree size (McCarroll & Loader 2004); or (v) tree height and light availability (Brienen et al. 2017). Records of $\Delta^{13}\text{C}$ in leaves and phloem sap of poplars with different ages revealed a rather stable ranking of the genotypic differences with tree age (Bonhomme et al. 2008). More recently, using a diachronic approach, several genotypes of *Populus deltoides* × *P. nigra* growing in common gardens were compared and the effects of ageing (from 5 to 20 years) on the genotype ranking were studied using the $\Delta^{13}\text{C}$ signals in tree rings as a surrogate for WUE (Rasheed et al. 2011). In the latter study, small changes in $\Delta^{13}\text{C}$ with tree age and no significant interaction between genotype and tree age were found, reinforcing the pertinence of ranking genotypes using $\Delta^{13}\text{C}$. However, in a such diachronic study, the influence of long-term environmental variability and annual effects related to rainfall on $\Delta^{13}\text{C}$ is assumed similar as the sampled trees were grown in common gardens that experienced the same environmental conditions.

In the present study we used a synchron-

ic approach where $\Delta^{13}\text{C}$ was recorded and compared in tree rings with same formation date, but from tree of different ages and contrasting environmental conditions. This sampling technique was used to minimise long-term environment effect and maximise age effect. Three different genotypes of *Populus deltoides* × *P. nigra* were used and carbon isotopic composition ($\delta^{13}\text{C}$) of the bulk wood from the year 2009 was measured and $\Delta^{13}\text{C}$ was computed. Since cellulose and lignin are the major components of bulk wood, and due to different biochemical pathways involved in their formation, the carbon isotopic signals of these two components are also different, i.e., lignin is ~3% depleted than cellulose (Loader et al. 2003, McCarroll & Loader 2004). Any variation in cellulose:lignin ratio can potentially affect the overall $\delta^{13}\text{C}$ signals of whole wood. Therefore, the variation in carbon percentage in the whole wood was investigated, that potentially elucidates the variation of lignin in the bulk wood (Lamlom & Savidge 2003). Moreover, nitrogen percentage in the bulk wood that can give insight regarding the tree response to the variation of available soil nitrogen resource (Guerrieri et al. 2011, Billings et al. 2016) was also measured.

The main difficulty of the synchronic approach is the lack of large-scale common

garden grouping a sufficient number of genotypes with different ages. We therefore used 41 small common gardens (Fig. 1) maintained in different locations in France by the *Institut du Développement Forestier* (IDF-CNPF) and gathered samples for three genotypes of *Populus deltoides* × *P. nigra*. Using this approach, we tested the hypotheses that: (i) water use efficiency estimated from $\Delta^{13}\text{C}$ in wood is affected by tree age; and (ii) genotype ranking based on $\Delta^{13}\text{C}$ does not change with tree age.

Material and methods

The aim of this study was to evidence the effect of age on stability of genotype ranking for $\Delta^{13}\text{C}$. However, studying long-term physiological response in trees can be tricky because of age and environmental signals overlapping each other. Therefore, maximising age-related signals through appropriate sampling technique becomes a prerequisite. In this regard a synchronic technique, where carbon isotope discrimination was compared among tree rings with same formation date but from trees of different ages and environments, thereby maximising the age signal. A major constraint while applying such technique is to find a large number of poplar genotypes of different ages in a common garden. To obtain a satisfying sample of trees of different ages in different genotypes and environments, we sampled trees across a large number of plantations in France (Fig. 1). These small plantations were established by the *Centre National de la Propriété Forestière – Institut pour le Développement Forestier* (CNPF-IDF) to compare the local performance of genotypes.

Study sites

The genotypes were planted in plots of 25 trees at a distance of 7 × 7 m, i.e., at a density of 204 stems ha⁻¹. Forty-one sites were selected (Fig. 1) which were located all over France (see Tab. S1 in Supplementary material for site information and spatial coordinates). Sites covered a large range of climates with mean annual temperature from 10 to 14 °C and precipitation from 600 to 1200 mm. Sampling sites were subsequently categorised as water available (WA, annual rainfall from 1000 to 1200 mm) or water limiting (WL, annual rainfall from 600 to 800 mm; precipitation record maintained by the *Centre National de la Propriété Forestière – Institut pour le Développement Forestier*, CNPF-IDF).

Sample collection and preparation

We focused on 3 *Populus deltoides* × *P. nigra* genotypes: *Koster*, *I-214* and *Dorskamp*, as these genotypes are known to display contrasting $\Delta^{13}\text{C}$ at leaf level (Monclus et al. 2005). Moreover, they were present in a large number of the surveyed plantations and covered the different age classes. The last annual ring corresponding to year 2009 was sampled in February 2010 with an increment borer (0.5 cm²) on ~5 trees site⁻¹




Fig. 1 - Location of 41 sampling sites across France. Each site is represented by a closed star.

Tab. 1 - Carbon and nitrogen percentage, and $\Delta^{13}\text{C}$ measured on the bulk wood from the year 2009 from three genotypes of *Populus deltoides* × *P. nigra* aged from 4 to 18 years. Data was analyzed using linear mixed model for the effects of genotype (G), age (A) and interaction (G × A). Table shows the mean values (\pm SE) combining all age classes. Different letters indicate significant genotype differences after multi-comparison test (Tukey HSD, $P < 0.05$).

Variable	Clone			Effect (Prob.)		
	Dorskamp	I-214	Koster	G	A	G × A
C %	46.8 \pm 0.28 ^a	47.7 \pm 0.097 ^b	47.7 \pm 0.10 ^b	<0.001	0.051	0.305
N %	0.154 \pm 0.004	0.14 \pm 0.007	0.168 \pm 0.007	0.248	<0.001	0.734
$\Delta^{13}\text{C}$ (‰)	20.3 \pm 0.076 ^a	19.9 \pm 0.053 ^b	19.4 \pm 0.078 ^c	<0.001	<0.001	<0.001
BAI ($\text{cm}^2 \text{yr}^{-1}$)	88.4 \pm 9.54	103 \pm 11.0	107.3 \pm 13.4	0.467	0.023	0.167

genotype¹, yielding 376 samples. All cores were divided into seven age classes, i.e., 4, 7, 9, 11, 13, 15 and 18 years. In order to minimise the environmental effect on $\Delta^{13}\text{C}$: (i) date effect was discarded as all cores corresponded to year 2009; and (ii) site effect was averaged in each age class by ensuring the representation of trees from both site types, i.e., WA and WL sites.

The ring formed in 2009 was carefully separated from the adjacent one and from bark with a sharp razor blade. After drying at 70 °C for 48h, each ring was ground separately into fine homogeneous powder using a ring grinder (SODEMI, CEP Industries Department, Cergy-Pontoise, France). One mg of the resulting wood powder was weighed in tin capsules for $\delta^{13}\text{C}$ analysis.

Carbon and nitrogen percentage and carbon isotope analysis

Wood powder was combusted at 1050 °C in sealed evacuated quartz tubes containing cobalt oxide and chromium oxide as catalyst, and an amount of pure oxygen. The gases produced during combustion, CO_2 , H_2O and NO_x were passed through a reduction tube where N_2 was produced and excess of oxygen was removed. Water was trapped by using anhydrous magnesium perchlorate and after reversible absorption, carbon and nitrogen percentages were measured in an elemental analyser (NA 1500-NC[®], Carlo Erba, Milan, Italy – Rasheed et al. 2011). Finally, combusted products were separated by gas chromatography and the CO_2 was delivered to an isotope ratio mass spectrometer (Delta-S[®], Finnigan, Bremen, Germany). Carbon isotope composition was expressed as $\delta^{13}\text{C}$ (eqn. 1):

$$\delta^{13}\text{C} = (R_{\text{sample}} / R_{\text{standard}} - 1) \cdot 1000 \quad (1)$$

where R_{sample} and R_{standard} are the $^{13}\text{C}/^{12}\text{C}$ ratios in a sample and the standard (Vienna-Pee Dee Belemnite) respectively. Accuracy of the measurements was $\pm 0.1\%$. Carbon isotope discrimination between atmosphere and wood was calculated as (eqn. 2):

$$\Delta^{13}\text{C} = \frac{\delta^{13}\text{C}_{\text{air}} - \delta^{13}\text{C}_{\text{wood}}}{1 + \delta^{13}\text{C}_{\text{wood}}/1000} \quad (2)$$

where $\delta^{13}\text{C}_{\text{air}}$ is the carbon isotope composition of CO_2 in the atmosphere and $\delta^{13}\text{C}_{\text{wood}}$ is the carbon isotope composition of the

wood powder. Given that $\delta^{13}\text{C}_{\text{air}}$ was -8.07% during 2003 and an annual decrease of 0.0281% (McCarroll & Loader 2004), $\delta^{13}\text{C}_{\text{air}}$ was estimated at -8.24% for 2009. We assumed that the mean value of $\delta^{13}\text{C}_{\text{air}}$ was similar across sites and did only marginally change with the season.

Basal area increment

The data corresponding to annual circumference (cm) of each individual tree was acquired through the annual growth records from the Centre National de la Propriété Forestière-Insitut pour le Développement Forestier. Assuming trunks were circular in section, circumference was used to calculate the radius and eventually basal area for the years 2008 and 2009. BAI ($\text{cm}^2 \text{y}^{-1}$) for year 2009 was calculated as the difference between the total area corresponding to years 2009 and 2008.

Statistical analysis

Normality and homoscedasticity of data were checked graphically with residual vs. predicted and normal quantile-to-quantile plots. The data set covering the seven age classes was analysed using linear mixed models fitted with genotype, age and their interaction as fixed effects and site as a random effect. Multiple comparison tests (post-hoc Tukey HSD test) were used to evaluate pairwise differences between age classes within each genotype. Further-

more, each age class was tested for genotype effect and post-hoc Tukey HSD test were used to evaluate pair-wise differences between genotypes within each age class. All tests were performed with R (R Core Team 2012) and R packages “nlme” (Pinheiro et al. 2012) and “multcomp” (Hothorn et al. 2008). All tests and correlations were declared significant at $P < 0.05$.

Results

Carbon and nitrogen percentage

Variance between sites was smaller than residual variance (intercept = 0.731; residuals = 1.84) and model R^2 resulted at 0.547 for carbon percentage. No genotype-age interaction was detected in the carbon percentage of bulk wood. It differed significantly among genotypes ($P < 0.001$ – for means, see Tab. 1). A non-significant trend with age was detected due to the slightly lower carbon percentage values at the age of 13 yrs. Nevertheless, carbon percentage did not display any significant trend related to age (Fig. 2). Variance between sites was smaller than residual variance (intercept = 0.009; residuals = 0.071) and model R^2 resulted at 0.525 for nitrogen percentage. For the nitrogen percentage, no interaction between genotype and age was found. Nitrogen percentage did not differ among genotypes but declined significantly with age ($P < 0.001$); it dropped from




Fig. 2 - Age-related variation of Carbon percentage in the bulk wood of the year 2009, sampled during winter 2010 in three *Populus deltoides* × *P. nigra* genotypes over seven age classes. Each point represents the mean of several sites. Error bars represent \pm SE.

Fig. 3 - Age-related variation of mean nitrogen percentage in the bulk wood of the year 2009, sampled during winter 2010. The curve represents the means of three *Populus deltoides* × *P. nigra* genotypes over seven age classes. Error bars represent ± SE and different letters indicate significant differences between age classes after Tukey HSD test ($P < 0.05$).





Fig. 4 - Age-related variation in carbon isotope discrimination ($\Delta^{13}C$) in the 2009 year-ring for three *Populus deltoides* × *P. nigra* genotypes (a) *Dorskamp*, (b) *I-214* and (c) *Koster*. Each point represents a mean value from several sites. Error bars represent ± SE and different letters indicate significant differences between age classes for each genotype after Tukey HSD test ($P < 0.05$).



0.19 down to 0.11% over the tested period (Fig. 3).

Age and genotype effects for $\Delta^{13}C$

Variance between sites was smaller than residual variance (intercept = 0.38; residuals = 0.51) and model R^2 resulted at 0.99.

$\Delta^{13}C$ variation was significantly explained by an interaction between genotype and age ($P < 0.001$ – Tab. 1). A detailed *post-hoc* analysis revealed that an age effect was detectable in all three genotypes (Fig. 4). There was a significant increase of $\Delta^{13}C$ by 2% from 4 to 9 yrs in *Dorskamp* and *Koster*

(Fig. 4a, Fig. 4c) and a less visible one in *I-214* (Fig. 4b) where it increased from 4 to 7 yrs and remained almost stable afterwards.

Genotype effect within each age group was found significant and was assessed with *post-hoc* Tukey HSD test (Tab. 2). In the young age classes (4 to 9 yrs), *Koster* displayed the lowest mean $\Delta^{13}C$ followed by *Dorskamp* and *I-214* that displayed similar values. At higher ages, *Koster* and *I-214* displayed the lowest and *Dorskamp* displayed the highest mean $\Delta^{13}C$, with the exception of age class of 13 yrs, where no significant difference could be detected.

Inter annual and genotype correlation between $\Delta^{13}C$ and BAI

The average BAI of the year 2009 was homogenous among the three genotypes over the tested period (Tab. 1), and therefore the genotypic differences in $\Delta^{13}C$ were independent from any difference in radial growth rate. The inter annual variation of BAI due to the age effect was positively correlated to $\Delta^{13}C$ in the three genotypes (Fig. 5)

Discussion

The present research was aimed to assess the stability of genotype differences in water use efficiency (*WUE*) with tree age; we compared three cultivars of *Populus deltoides* × *P. nigra* for this trait in plantation *in situ* across France and used a synchronic approach, based on $\Delta^{13}C$ recorded from the year 2009 of individuals with different ages.

Genotype and age effect on $\Delta^{13}C$ and relationship with BAI

Whole wood consists of complex mixtures of molecules like cellulose, hemicellulose and lignin with variable isotopic signatures. Early studies used whole wood for the isotopic analysis on tree rings. However, the isotopic composition of individual wood components differs largely (Wilson & Grinstead 1977). Thus, we suspected that the observed significant genotypic difference in mean $\Delta^{13}C$ could potentially be due to changes in cellulose vs. lignin ratios with contrasting isotopic signals. Although previous studies have shown that using whole wood is better than inducing extraction error while using cellulose to measure $\delta^{13}C$ (Guy & Holowachuk 2001, Rasheed et al. 2011), in this study we minimised this effect by using the last annual ring (2009, with same cambial age) for isotope analysis.

Our results showed the existence of an age effect on $\Delta^{13}C$, where $\Delta^{13}C$ increased to a larger extent during first age classes of 4-11 years in two genotypes (*Dorskamp* and *Koster*). This positive trend in $\Delta^{13}C$ with age is contrary to several previous studies reporting age-related decreases in $\Delta^{13}C$ using different dendrochronological approaches (Bert et al. 1997, Duquesnay et al. 1998, McCarroll & Pawellek 2001, Penuelas et al. 2008). $\Delta^{13}C$ has been expected to be an estimator of water use efficiency (Farquhar

Tab. 2 - Carbon isotopic discrimination (± SE) between atmosphere and bulk wood ($\Delta^{13}C$) in the year 2009 for three *Populus deltoides* × *P. nigra* genotypes (*Dorskamp*, *I-214* and *Koster*). Each age class was tested for genotype effect (F-value and P-value) and different letters indicate a significant genotype difference for each age class after Tukey HSD test ($P < 0.05$).

Age class (yrs)	df	<i>Dorskamp</i>	<i>I-214</i>	<i>Koster</i>	F-value	P-value
4	43	18.9 ± 0.156 ^a	19.3 ± 0.107 ^a	18.1 ± 0.132 ^b	22.3	<0.001
7	47	20.3 ± 0.179 ^a	20.0 ± 0.172 ^a	18.7 ± 0.179 ^b	18.4	<0.001
9	88	20.1 ± 0.103 ^a	20.3 ± 0.079 ^a	19.5 ± 0.096 ^b	17.9	<0.001
11	80	20.9 ± 0.191 ^a	19.9 ± 0.095 ^b	19.8 ± 0.136 ^b	13.8	<0.001
13	24	20.8 ± 0.095	20.3 ± 0.175	20.3 ± 0.150	3.03	0.069
15	34	20.6 ± 0.161 ^a	20.1 ± 0.077 ^b	19.8 ± 0.103 ^b	9.26	<0.001
18	39	20.8 ± 0.092 ^a	19.7 ± 0.151 ^b	19.6 ± 0.443 ^b	19.6	<0.001

et al. 1989). Several studies have shown a clear negative correlation between $\Delta^{13}\text{C}$ and A/g_s (intrinsic water use efficiency, $iWUE$) and WUE (whole-plant water use efficiency – Ripullone et al. 2004, Cernusak et al. 2008, Rasheed et al. 2013). Thus, increase in $\Delta^{13}\text{C}$ with age in our study reflects a decrease in $iWUE$ with age. This decrease in $iWUE$ (A/g_s) can either be due to decreased CO_2 assimilation rate (A) or increased stomatal conductance (g_s – Farquhar & Richards 1984, Condon et al. 2004). Furthermore, $\Delta^{13}\text{C}$ and tree growth are often correlated depending upon the growth environment. This correlation was found to be positive for some species, i.e., *Eucalyptus globulus* Labill (Osorio & Pereira 1994, Pita et al. 2001), *Fagus sylvatica* L. (Dupouey et al. 1993) and *Pinus radiata* (Rowell et al. 2008). On the contrary, some other conifer species displayed a negative correlation between $\Delta^{13}\text{C}$ and growth, i.e., *Larix occidentalis* Nutt. (Zhang et al. 1994), *Pinus pinaster* (Nguyen-Queyrens et al. 1998, Brendel et al. 2002), *Picea mariana* Mill (Johnsen et al. 1999) and *Pinus caribaea* Morelet (Xu et al. 2000). However, no correlation between $\Delta^{13}\text{C}$ and tree growth was found in *P. × euramericana* in common garden (Monclus et al. 2005) and poplar plantation (Bonhomme et al. 2008). Within each genotype, we found a positive correlation between BAI and $\Delta^{13}\text{C}$ with age, which is in line with previous findings in different *Populus deltoides* × *P. nigra* genotypes (Rasheed et al. 2011). Similar positive correlation between BAI and $\Delta^{13}\text{C}$ has been reported in *Pinus radiata* (Rowell et al. 2008). The positive correlation between $\Delta^{13}\text{C}$ and BAI suggests that the inter-annual variation of $\Delta^{13}\text{C}$ is controlled to a larger extent by stomatal conductance than by photosynthetic capacity (Johnsen et al. 1999, Xu et al. 2000). In poplar, mostly variation in stomatal conductance rather than photosynthetic capacity seems to control carbon gain and growth (Ceulemans et al. 1987, Monclus et al. 2006, Rasheed et al. 2015). Therefore, based on the significant positive correlation between BAI and $\Delta^{13}\text{C}$ across tested genotypes, we may conclude that the observed increase in $\Delta^{13}\text{C}$ with age and inter-annual variation of $\Delta^{13}\text{C}$ was largely controlled by the variation in stomatal conductance rather than photosynthetic capacity.

Stability of the genotypic ranking for $\Delta^{13}\text{C}$

Checking genotype rank stability with age is of central importance for selecting genotypes for higher water use efficiency. Therefore, genotype ranking in *Populus deltoides* × *P. nigra* was tested for $\Delta^{13}\text{C}$ in controlled vs. open field conditions (Monclus et al. 2005) and subsequently, in well irrigated vs. water stress under field conditions (Monclus et al. 2006). In both studies, genotypic ranking remained stable with no correlation found between $\Delta^{13}\text{C}$ and productivity traits. However, these studies were done on young plants. Parallel to




Fig. 5 - Correlation between carbon isotopic discrimination ($\Delta^{13}\text{C}$) and basal area increment (BAI) in three *Populus deltoides* × *P. nigra* genotypes (a) Dorskamp: $\Delta^{13}\text{C} = 0.0161\text{BAI} + 18.9$; (b) I-214: $\Delta^{13}\text{C} = 0.0087\text{BAI} + 19.0$; and (c) Koster: $\Delta^{13}\text{C} = 0.0138\text{BAI} + 18.0$. Each point represents values measured at a given site.

that, many previous studies evidenced an age effect on $\Delta^{13}\text{C}$ and demonstrated that duration and extent of age effect is variable according to many species (Leavitt 2010). In this context, genotype ranking made on young plants were susceptible to change with age. Our results showed that genotype ranking shuffled across first age class (4 years) to seventh age class (18 years), i.e., I-214 displayed highest and Koster the lowest $\Delta^{13}\text{C}$ values in the first age class (4 years), whereas I-214 was ranked second in the seventh age class (18 years). In spite of rank shuffling with age, genotypic ranking for mean $\Delta^{13}\text{C}$ values matched with that found by Marron et al.

(2005) and Monclus et al. (2005), where Koster had the lowest $\Delta^{13}\text{C}$ and Dorskamp had the highest, and with Rasheed et al. (2011) in Begaar for Dorskamp and I-214. Thus, we may conclude that based on a relatively small number of cultivars, a stability of genotype ranking was observed with age in two cultivars, at least during the approx. 20 years from planting to harvest. This conclusion is based on a limited range of cultivars. Unfortunately, extension to a larger number of genotypes was impossible given the lack of suitable even-aged common garden plantation of poplar cultivars at suitable age, not to speak of common gardens with different tree ages.

Conclusion

In this study, $\Delta^{13}\text{C}$ assessed from the year 2009 increased with tree age in the three genotypes, indicating a decrease in water use efficiency with age at whole tree level. Furthermore, inter-annual variation of $\Delta^{13}\text{C}$ was found positively correlated to BAI in all genotypes which shows that $\Delta^{13}\text{C}$ was largely controlled by stomatal conductance rather photosynthetic capacity. Significant genotypic effect was detected for mean $\Delta^{13}\text{C}$ over the tested period. The genotypic ranking of $\Delta^{13}\text{C}$ was: (i) maintained among the three *Populus deltoides* × *P. nigra* genotypes, despite of crossing detected with tree age; (ii) found consistent with the previous ranking for $\Delta^{13}\text{C}$. Finally, Koster genotype was found highly water use efficient and productive with age among the tested genotypes. However, further studies are required to test other genotypes as well.

Acknowledgements

The authors are very grateful to Cyril Buré (INRA, UMR EEF) for helping in arranging the field visits and in collecting samples together with the different technicians of CRPF. Christian Hossann and Claude Bréchet (INRA, UMR EEF), for spending time doing all isotope analyses. Finally, we thank WFJ Parsons (CEF) for English editing. We are thankful to two anonymous reviewers for their suggestions to improve the manuscript. The research was supported by the European Union FP7 Project “Novel Tree Breeding Strategies” Project no. FP7 - 211868. F. Rasheed was supported by a grant from the Higher Education Commission of Pakistan.

References

- Bert D, Leavitt SW, Dupouey JL (1997). Variation of wood $\delta^{13}\text{C}$ and water use efficiency of *Abies alba* during the last century. *Ecology* 78: 588-596. - doi: [10.1890/0012-9658\(1997\)078\[0588:ZMIIAL\]2.0.CO;2](https://doi.org/10.1890/0012-9658(1997)078[0588:ZMIIAL]2.0.CO;2)
- Billings SA, Boone AS, Stephen FM (2016). Tree-ring $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$, leaf $\delta^{13}\text{C}$ and wood and leaf N status demonstrate tree growth strategies and predict susceptibility to disturbance. *Tree Physiology* 36: 576-588. - doi: [10.1093/treephys/tpw010](https://doi.org/10.1093/treephys/tpw010)
- Bonhomme L, Barbaroux C, Monclus R, Morabito D, Berthelot A, Villar M, Dreyer E, Brignolas F (2008). Genetic variation in productivity, leaf traits and carbon isotope discrimination in hybrid poplars cultivated on contrasting sites. *Annals of Forest Science* 65: 503. - doi: [10.1051/forest:2008024](https://doi.org/10.1051/forest:2008024)
- Brendel O, Le Thiec D, Scotti-Saintagne C, Bodénès C, Kremer A, Guehl J (2008). Quantitative trait loci controlling water use efficiency and related traits in *Quercus robur* L. *Tree Genetic and Genomes* 4: 263-278. - doi: [10.1007/s11295-007-0107-z](https://doi.org/10.1007/s11295-007-0107-z)
- Brendel O, Pot D, Plomion C, Rozenberg P, Guehl J (2002). Genetic parameters and QTL analysis of $\delta^{13}\text{C}$ and ring width in maritime pine. *Plant, Cell and Environment* 25: 945-953. - doi: [10.1046/j.1365-3040.2002.00872.x](https://doi.org/10.1046/j.1365-3040.2002.00872.x)
- Brienen RJW, Gloor E, Clerici S, Newton R, Arppe L, Boom A, Bottrell S, Callaghan M, Heaton T, Helama S, Helle G, Leng MJ, Mielikäinen K, Oinonen M, Timonen M (2017). Tree height strongly affects estimates of water-use efficiency responses to climate and CO_2 using isotopes. *Nature Communications* 8: 288. - doi: [10.1038/s41467-017-00225-z](https://doi.org/10.1038/s41467-017-00225-z)
- Cernusak LA, Winter K, Aranda J, Turner BL (2008). Conifers. Angiosperm trees and lianas growth whole plant water and nitrogen use efficiency and stable isotope composition ($\delta^{13}\text{C}$ and $\delta^{18}\text{O}$) of seedlings grown in a tropical environment. *Plant Physiology* 148: 642-659. - doi: [10.1104/pp.108.123521](https://doi.org/10.1104/pp.108.123521)
- Ceulemans R, Impens I, Imler R (1987). Stomatal conductance and stomatal behaviour in *Populus* clones and hybrids. *Canadian Journal of Botany* 66: 1404-1414. - doi: [10.1139/b88-196](https://doi.org/10.1139/b88-196)
- Condon AG, Richards RA, Rebetzke GJ, Farquhar GD (2004). Breeding for high water-use efficiency. *Journal of Experimental Botany* 55: 2447-2460. - doi: [10.1093/jxb/erh277](https://doi.org/10.1093/jxb/erh277)
- Dillen S, Marron N, Koch B, Ceulemans R (2008). Genetic variation of stomatal traits and carbon isotope discrimination in two hybrid poplar families (*Populus deltoides* “S9-2” × *P. nigra* “Ghoy” and *P. deltoides* “S9-2” × *P. trichocarpa* “V24”). *Annals of Botany* 102: 399-407. - doi: [10.1093/aob/mcn107](https://doi.org/10.1093/aob/mcn107)
- Dupouey JL, Leavitt S, Choiselet E, Jourdain S (1993). Modeling carbon isotope fractionation in tree rings based on effective evapotranspiration and soil water status. *Plant, Cell and Environment* 16: 939-947. - doi: [10.1111/j.1365-3040.1993.tb00517.x](https://doi.org/10.1111/j.1365-3040.1993.tb00517.x)
- Duquesnay A, Bréda N, Stievenard M, Dupouey JL (1998). Changes of tree-ring $\delta^{13}\text{C}$ and water use efficiency of beech (*Fagus sylvatica* L.) in north-eastern France during the past century. *Plant Cell and Environment* 21: 565-572. - doi: [10.1046/j.1365-3040.1998.00304.x](https://doi.org/10.1046/j.1365-3040.1998.00304.x)
- Ehleringer JR, Hall AE, Farquhar GD (1993). Water use in relation to productivity. In: “Stable Isotopes and Plant Carbon-Water Relations” (Ehleringer JR, Hall AE, Farquhar GD eds). Academic Press, New York, USA, pp. 3-8.
- Farquhar G, Richards R (1984). Isotopic composition of plant carbon correlates with water use efficiency of wheat genotypes. *Australian Journal of Plant Physiology* 11: 539-552. - doi: [10.1071/PP9840539](https://doi.org/10.1071/PP9840539)
- Farquhar G, Ehleringer J, Hubick K (1989). Carbon isotope discrimination and photosynthesis. *Annual Review of Plant Physiology and Plant Molecular Biology* 40: 503-537. - doi: [10.1146/annurev.pp.40.060189.002443](https://doi.org/10.1146/annurev.pp.40.060189.002443)
- Francey RJ, Farquhar GD (1982). An explanation of $^{13}\text{C}/^{12}\text{C}$ variations in tree rings. *Nature* 297: 28-31. - doi: [10.1038/297028a0](https://doi.org/10.1038/297028a0)
- Guerrieri R, Mencuccini M, Sheppard LJ, Saurer M, Perks MP, Levy P, Sutton MA, Borghetti M, Grace J (2011). The legacy of enhanced N and S deposition as revealed by the combined analysis of $\delta^{13}\text{C}$, $\delta^{18}\text{O}$ and $\delta^{15}\text{N}$ in tree rings. *Global Change Biology* 17: 1946-1962. - doi: [10.1111/j.1365-2486.2010.02362.x](https://doi.org/10.1111/j.1365-2486.2010.02362.x)
- Guy RD, Holowachuk DL (2001). Population differences in stable carbon isotope ratio of *Pinus contorta* Dougl. ex Loud.: relationship to environment, climate of origin, and growth potential. *Canadian Journal of Botany* 79: 274-283. - doi: [10.1139/b01-001](https://doi.org/10.1139/b01-001)
- Hothorn T, Bretz F, Westfall P (2008). Simultaneous inference in general parametric models. *Biometrical journal* 50: 346-363. - doi: [10.1002/bimj.200810425](https://doi.org/10.1002/bimj.200810425)
- IPCC (2013). Climate change 2013: the physical science basis. In “Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change” (Stocker TF, Qin D, Plattner G-K, Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V, Midgley PM eds). Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 1535.
- Johnsen KH, Flanagan LB, Huber DA, Major JE (1999). Genetic variation in growth carbon isotope discrimination and foliar N concentration in *Picea mariana*: analyses from a half diallel mating design using field grown trees. *Canadian Journal of Forestry Research* 29: 1727-1735. - doi: [10.1139/x99-144](https://doi.org/10.1139/x99-144)
- Lauteri M, Scartazza A, Guido MC, Brugnoli E (1997). Genetic variation in photosynthetic capacity, carbon isotope discrimination and mesophyll conductance in provenances of *Castanea sativa* adapted to different environments. *Functional Ecology* 11: 675-683. - doi: [10.1046/j.1365-2435.1997.00140.x](https://doi.org/10.1046/j.1365-2435.1997.00140.x)
- Leavitt SW (2010). Tree-ring C-H-O isotope variability and sampling. *Science of Total Environment* 408: 5244-5253. - doi: [10.1016/j.scitotenv.2010.07.057](https://doi.org/10.1016/j.scitotenv.2010.07.057)
- Loader NJ, Roberston I, McCarroll D (2003). Comparison of stable carbon isotope ratios in the whole wood, cellulose and lignin of oak tree-rings. *Palaeogeography, Palaeoclimatology, Palaeoecology* 196: 395-407. - doi: [10.1016/S0031-0182\(03\)00466-8](https://doi.org/10.1016/S0031-0182(03)00466-8)
- Lamlom SH, Savidge RA (2003). A reassessment of carbon contents in wood: variation within and between 41 North American species. *Biomass and Bioenergy* 25: 381-388. - doi: [10.1016/S0961-9534\(03\)00033-3](https://doi.org/10.1016/S0961-9534(03)00033-3)
- Marron N, Villar M, Dreyer E, Delay D, Boudouresque E, Petit J-M, Delmotte FM, Guehl JM, Brignolas F (2005). Diversity of leaf traits related to productivity in 31 *Populus deltoides* × *Populus nigra* clones. *Tree Physiology* 25: 425-435. - doi: [10.1093/treephys/25.4.425](https://doi.org/10.1093/treephys/25.4.425)
- McCarroll D, Loader NJ (2004). Stable isotopes in tree rings. *Quaternary Science Reviews* 23: 771-801. - doi: [10.1016/j.quascirev.2003.06.017](https://doi.org/10.1016/j.quascirev.2003.06.017)
- McCarroll D, Pawellek F (2001). Stable carbon isotope ratios of *Pinus sylvestris* from northern Finland and the potential for extracting a climate signal from long Fennoscandian chronologies. *The Holocene* 11: 517-526. - doi: [10.1191/095968301680223477](https://doi.org/10.1191/095968301680223477)
- Monclus R, Dreyer E, Delmotte FM, Villar M, Delay D, Boudouresque E, Petit JM, Marron N, Bréchet N, Brignolas F (2005). Productivity leaf traits and carbon isotope discrimination in 29 *Populus deltoides* × *P. nigra* clones. *New Phytologist* 167: 53-62. - doi: [10.1111/j.1469-8137.2005.01407.x](https://doi.org/10.1111/j.1469-8137.2005.01407.x)
- Monclus R, Dreyer E, Villar M, Delmotte FM, Delay D, Petit JM, Barbaroux C, Le Thiec D, Bréchet C, Brignolas F (2006). Impact of drought on productivity and water-use efficiency in 29 genotypes of *Populus deltoides* × *Populus nigra*. *New Phytologist* 169: 765-777. - doi: [10.1111/j.14](https://doi.org/10.1111/j.14)

69-8137.2005.01630.x

- Monserud RA, Marshall JD (2001). Time-series analysis of $\delta^{13}\text{C}$ from tree rings time trends and autocorrelation. *Tree Physiology* 21: 1087-1102. - doi: [10.1093/treephys/21.15.1087](https://doi.org/10.1093/treephys/21.15.1087)
- Nguyen-Queyrens A, Ferhi A, Loustau D, Guehl JM (1998). Within-ring $\delta^{13}\text{C}$ spatial variability and interannual variations in wood cellulose of two contrasting provenances of *Pinus pinaster*. *Canadian Journal of Forestry Research* 28: 766-773. - doi: [10.1139/x98-044](https://doi.org/10.1139/x98-044)
- Osorio J, Pereira JS (1994). Genotypic differences in water use efficiency and ^{13}C discrimination in *Eucalyptus globulus*. *Tree Physiology* 14: 871-882. - doi: [10.1093/treephys/14.7-8-9.871](https://doi.org/10.1093/treephys/14.7-8-9.871)
- Penuelas J, Hunt JM, Ogaya R, Jump A (2008). Twentieth century changes of tree-ring $\delta^{13}\text{C}$ at the southern range-edge of *Fagus sylvatica*: increasing water-use efficiency does not avoid the growth decline induced by warming at low altitudes. *Global Change Biology* 14: 1076-1088. - doi: [10.1111/j.1365-2486.2008.01563.x](https://doi.org/10.1111/j.1365-2486.2008.01563.x)
- Pinheiro J, Bates B, DebRoy S, Sarkar D, the R Development Core Team (2012). nlme: Linear and nonlinear mixed effects models. R package version 3.1-103. [online] URL: <http://cran.r-project.org/web/packages/nlme/index.html>
- Pita P, Soria F, Canas I, Toval G, Pardos JA (2001). Carbon isotope discrimination and its relationship to drought under field conditions in genotypes of *Eucalyptus globulus* Labill. *Forest Ecology and Management* 141: 211-221. - doi: [10.1016/S0378-1127\(00\)00330-3](https://doi.org/10.1016/S0378-1127(00)00330-3)
- R Core Team (2012). R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. [online] URL: <http://www.R-project.org/>
- Rasheed F, Richard B, Le Thiec D, Montpied P, Paillassa E, Brignolas F, Dreyer E (2011). Time course of $\delta^{13}\text{C}$ in poplar wood: genotype ranking remains stable over the life cycle in plantations despite some differences between cellulose and bulk wood. *Tree Physiology* 31: 1183-1193. - doi: [10.1093/treephys/tp1108](https://doi.org/10.1093/treephys/tp1108)
- Rasheed F, Dreyer E, Richard B, Brignolas F, Montpied P, Le Thiec D (2013). Genotypic differences in ^{13}C discrimination between atmosphere and leaf matter match differences in transpiration efficiency at leaf and whole-plant level in hybrid *Populus deltoides* × *nigra*. *Plant, Cell and Environment* 36: 87-102. - doi: [10.1111/j.1365-3040.2012.02556.x](https://doi.org/10.1111/j.1365-3040.2012.02556.x)
- Rasheed F, Dreyer E, Richard B, Brignolas F, Brendel O, Le Thiec D (2015). Vapour pressure deficit during growth has little impact on genotypic differences of transpiration efficiency at leaf and whole-plant level: an example from *Populus nigra* L. *Plant, Cell Environment* 38: 670-684. - doi: [10.1111/pce.12423](https://doi.org/10.1111/pce.12423)
- Richards RA, Rebetzke GJ, Watt M, Condon AG, Spielmeier W, Dolferus R (2010). Breeding for improved water productivity in temperate cereals: phenotyping, quantitative trait loci, markers and the selection environment. *Functional Plant Biology* 37: 85-97. - doi: [10.1071/FP09219](https://doi.org/10.1071/FP09219)
- Richards RA, Rebetzke GJ, Watt M, Condon AG, Rasheed F, Dreyer E, Richard B, Brignolas F, Brendel O, Le Thiec D (2015). Vapour pressure deficit during growth has little impact on genotypic differences of transpiration efficiency at leaf and whole-plant level: an example from *Populus nigra* L. *Plant, Cell Environment* 38: 670-684. - doi: [10.1111/pce.12423](https://doi.org/10.1111/pce.12423)
- Ripullone F, Lauteri M, Grassi G, Amato M, Borghetti M (2004). Variation in nitrogen supply changes water-use efficiency of *Pseudotsuga menziesii* and *Populus* × *euramericana*; a comparison of three approaches to determine water-use efficiency. *Tree Physiology* 24: 671-679. - doi: [10.1093/treephys/24.6.671](https://doi.org/10.1093/treephys/24.6.671)
- Rowell DM, Ades PK, Tausz M, Arndt SK, Adams MA (2008). Lack of genetic variation in tree ring $\delta^{13}\text{C}$ suggests a uniform stomatally-driven response to drought stress across *Pinus radiata* genotypes. *Tree Physiology* 29: 191-198. - doi: [10.1093/treephys/tpn015](https://doi.org/10.1093/treephys/tpn015)
- Schleser GH, Jayasekera R (1985). $\delta^{13}\text{C}$ variations in leaves of a forest as an indication of reassimilated CO_2 from the soil. *Oecologia* 65: 536-542. - doi: [10.1007/BF00379669](https://doi.org/10.1007/BF00379669)
- Vadez V, Kholova J, Medina S, Kakkera A, Anderberg H (2014). Transpiration efficiency: new insights into an old story. *Journal of Experimental Botany* 65: 6141-53. - doi: [10.1093/jxb/eru040](https://doi.org/10.1093/jxb/eru040)
- Wilson AT, Grinsted MJ (1977). $^{12}\text{C}/^{13}\text{C}$ in cellulose and lignin as palaeothermometers. *Nature* 265: 133-135. - doi: [10.1038/265133a0](https://doi.org/10.1038/265133a0)
- Xu ZH, Saffigna PG, Farquhar GD, Simpson JA, Haines RJ, Walker S, Osborne DO, Guinto D (2000). Carbon isotope discrimination and oxygen isotope composition in clones of the F1 hybrid between slash pine and Caribbean pine in relation to tree growth water use efficiency and foliar nutrient concentration. *Tree Physiology* 20: 1209-1217. - doi: [10.1093/treephys/20.18.1209](https://doi.org/10.1093/treephys/20.18.1209)
- Zhang JW, Fins L, Marshall JD (1994). Stable carbon isotope discrimination photosynthetic gas exchange and growth differences among western larch families. *Tree Physiology* 14: 531-539. - doi: [10.1093/treephys/14.5.531](https://doi.org/10.1093/treephys/14.5.531)

Supplementary Material

Tab. S1 - Complete list of 376 tree cores sampled for three genotypes (*Dorskamp*, *I-214* and *Koster*) under each age class along with information of each site condition (Water available, WA and Water limiting, WL) and their GPS coordinates.

Tab. S2 - Coefficients of predictor variable for Carbon, Nitrogen and along with variance explained by sites, which was taken as random factor.

Tab. S3 - Coefficients of predictor variable for Nitrogen along with variance explained by sites, which was taken as random factor.

Tab. S4 - Coefficients of predictor variable for Carbon isotope discrimination along with variance explained by sites which was taken as random factor.

Link: Rasheed_2896@suppl001.pdf