



HAL
open science

An AgMIP framework for improved agricultural representation in integrated assessment models

Alex C Ruane, Cynthia Rosenzweig, Senthold Asseng, Kenneth J Boote, Joshua Elliott, Frank Ewert, James W Jones, Pierre Martre, Sonali P Mcdermid, Christoph Müller, et al.

► To cite this version:

Alex C Ruane, Cynthia Rosenzweig, Senthold Asseng, Kenneth J Boote, Joshua Elliott, et al.. An AgMIP framework for improved agricultural representation in integrated assessment models. *Environmental Research Letters*, 2017, 12, 10.1088/1748-9326/aa8da6 . hal-02624345

HAL Id: hal-02624345

<https://hal.inrae.fr/hal-02624345>

Submitted on 26 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

LETTER • OPEN ACCESS

An AgMIP framework for improved agricultural representation in integrated assessment models

To cite this article: Alex C Ruane *et al* 2017 *Environ. Res. Lett.* **12** 125003

View the [article online](#) for updates and enhancements.

Related content

- [Implications of climate mitigation for future agricultural production](#)
Christoph Müller, Joshua Elliott, James Chryssanthacopoulos *et al.*
- [Assessing inter-sectoral climate change risks: the role of ISIMIP](#)
Cynthia Rosenzweig, Nigel W Arnell, Kristie L Ebi *et al.*
- [Simulated vs. empirical weather responsiveness of crop yields: US evidence and implications for the agricultural impacts of climate change](#)
Malcolm N Mistry, Ian Sue Wing and Enrica De Cian

Environmental Research Letters



LETTER

An AgMIP framework for improved agricultural representation in integrated assessment models

OPEN ACCESS

RECEIVED

3 March 2017

REVISED

6 September 2017

ACCEPTED FOR PUBLICATION

19 September 2017

PUBLISHED

24 November 2017

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Alex C Ruane^{1,11}, Cynthia Rosenzweig¹, Senthold Asseng², Kenneth J Boote², Joshua Elliott³, Frank Ewert^{4,5}, James W Jones², Pierre Martre⁶, Sonali P McDermid⁷, Christoph Müller⁸, Abigail Snyder⁹ and Peter J Thorburn¹⁰

¹ NASA Goddard Institute for Space Studies, New York, NY, United States of America

² University of Florida, Agricultural and Biological Engineering, Gainesville, FL, United States of America

³ University of Chicago, Computation Institute, Chicago, IL, United States of America

⁴ University of Bonn, Bonn, Germany

⁵ Leibniz Center of Agricultural landscape Research (ZALF), Muencheberg, Germany

⁶ UMR LEPSE, INRA, Montpellier SupAgro, Montpellier, France

⁷ New York University, New York, NY, United States of America

⁸ Potsdam Institute for Climate Impact Research, Potsdam, Germany

⁹ Pacific Northwest National Laboratory, College Park, MD, United States of America

¹⁰ Commonwealth Scientific and Industrial Research Organization, Brisbane, Australia

¹¹ Author to whom any correspondence should be addressed.

E-mail: alexander.c.ruane@nasa.gov

Keywords: AgMIP, climate impacts, crop models, integrated assessment models, agricultural impacts, emulators, model development


Abstract

Integrated assessment models (IAMs) hold great potential to assess how future agricultural systems will be shaped by socioeconomic development, technological innovation, and changing climate conditions. By coupling with climate and crop model emulators, IAMs have the potential to resolve important agricultural feedback loops and identify unintended consequences of socioeconomic development for agricultural systems. Here we propose a framework to develop robust representation of agricultural system responses within IAMs, linking downstream applications with model development and the coordinated evaluation of key climate responses from local to global scales. We survey the strengths and weaknesses of protocol-based assessments linked to the Agricultural Model Intercomparison and Improvement Project (AgMIP), each utilizing multiple sites and models to evaluate crop response to core climate changes including shifts in carbon dioxide concentration, temperature, and water availability, with some studies further exploring how climate responses are affected by nitrogen levels and adaptation in farm systems. Site-based studies with carefully calibrated models encompass the largest number of activities; however they are limited in their ability to capture the full range of global agricultural system diversity. Representative site networks provide more targeted response information than broadly-sampled networks, with limitations stemming from difficulties in covering the diversity of farming systems. Global gridded crop models provide comprehensive coverage, although with large challenges for calibration and quality control of inputs. Diversity in climate responses underscores that crop model emulators must distinguish between regions and farming system while recognizing model uncertainty. Finally, to bridge the gap between bottom-up and top-down approaches we recommend the deployment of a hybrid climate response system employing a representative network of sites to bias-correct comprehensive gridded simulations, opening the door to accelerated development and a broad range of applications.

1. Introduction

Integrated assessment models (IAMs) examine the interactions between human systems and the natural

environment. IAMs thus explore how societal changes, such as global policies, population growth, socioeconomic development, greenhouse gas emissions, and technological advances affect land, air, and water




resources, as well as repercussions when these natural resources are strained (Füssel *et al* 2010, Clarke *et al* 2014). Agriculture has long been central to the relationship between society and natural systems, providing vital foods, fiber, and energy while drawing heavily on land and water resources.

IAMs have traditionally represented agricultural sector changes as exogenous yield changes provided via scenarios aggregated to national or regional level production using current harvested area weights (Müller and Robertson 2014, Nelson *et al* 2014, Wiebe *et al* 2015); however these only draw from a small subset of cutting-edge crop model assessments. A more direct coupling of agricultural responses within IAMs is facilitated by the application of crop model emulators, defined here as computationally-efficient representations of crop model results that capture fundamental responses to climate conditions. Crop model emulators may take the form of lookup tables (e.g. based upon response surfaces in Pirttioja *et al* 2015), simplified response functions (Howden and Crimp 2005, Crimp *et al* 2008, Ruane *et al* 2014, Makowski *et al* 2015), or complex statistical models (Blanc 2017, Mistry *et al* 2017, Moore *et al* 2017), each estimating yield as a function of climate variables with varying degrees of non-linearity and detail about the specific crop variety, farm environment, weather extremes, and crop model emulated. As these emulators get more complex the gain in computational efficiency (compared to just using the crop model itself) is reduced, and in the end a crop model emulator is limited by the performance of the crop model or crop model ensemble that it is emulating. Emulators are distinct from statistical crop models, which are trained upon observational data, with one advantage being that

they may use principles of biophysical process response to explore environments that have not been observed (such as future climate and land use change). The exact specifications and desired detail of a crop model emulator depends on the IAM to which it is coupled, the intended applications, and the capabilities and coverage of the underlying crop model assessments.

IAMs have a lot to gain by better incorporating crop responses to changes in carbon dioxide concentration ($[CO_2]$), temperature, water, nitrogen, and adaptation (CTWNA). CTWNA sensitivity simulations can be more useful than projections driven by global climate models (GCMs) as they provide the information basis to construct crop model emulators for use in IAMs in conjunction with climate emulators (e.g. Meinshausen *et al* 2011, Castruccio *et al* 2014, Hartin *et al* 2015). Figure 1 illustrates how this powerful combination improves agricultural sector representation by allowing IAM land use changes and emissions of greenhouse gases and aerosols to influence regional temperature and precipitation changes (using the climate emulator), affecting crop production and requirements (using the crop model emulator) that feed back into the IAM. This also captures agricultural feedback loops, where societal or environmental changes alter the climate and shift agricultural production in a manner that reinforces or diminishes those changes, and unintended consequences when policies in another sector or region impact distant farming systems (potentially through climate responses or through independent mechanisms such as trade).

The Agricultural Model Intercomparison and Improvement Project (AgMIP; Rosenzweig *et al* 2013, 2015) was launched in 2010 to provide a common



framework and systematic approach for analysis of agricultural challenges. AgMIP connects climate, crop, livestock, and economic models at local, regional, and global scales, allowing multi-model, multi-discipline, multi-scale assessments of agricultural development and food security (Rosenzweig *et al* 2016, Antle *et al* 2015). AgMIP mainly utilizes process-based crop models that represent biophysical processes and their responses to genetics, environment, and management over the course of a growing season, with statistical models also included in some efforts. Integrated assessment modelers examining previous crop modeling studies have been challenged to make sense of differing assumptions, methods, and models in addition to the under-representation of agricultural systems beyond the mid-latitude, high-input breadbaskets (White *et al* 2011, Challinor *et al* 2014a). AgMIP facilitates more robust and transferable findings based on common simulation protocols, multi-model ensembles, the tracking of uncertainty, and an emphasis on under-simulated farm systems. Great strides in computational power are opening new doors for agricultural model development and application, raising the ceiling for multi-model analyses, new scales of decision support, and more accurate crop model emulators for IAM applications.

This article takes stock of the methods used by AgMIP to capture the response of agricultural productivity to changing climate conditions, examining the relative strengths and weaknesses of site, network, and gridded modeling approaches to inform IAMs and related crop model emulators. We then provide a framework for coordinated development and application of agricultural responses drawing value from local to global approaches and linking biophysical and integrated assessments. We conclude with recommendations for priority future work and applications.

2. Survey of crop model outputs germane to IAM emulators

Although AgMIP conducts more than 30 activities (Rosenzweig *et al* 2015), here we survey activities that (a) test for sensitivity to some or all of CTWNA factors and utilize (b) multiple agricultural models, (c) multiple sites, and (d) common protocols. These are described in table 1 along with related activities by the MACSUR project (Modelling European Agriculture with Climate Change for Food Security; Ewert *et al* 2015, Webber *et al* 2015). Figure 2 presents the geographic coverage of these site, network, and gridded activities.

2.1. Site-based approaches

The overwhelming majority of studies in the large literature on crop impacts are site-based studies (White *et al* 2011, Challinor *et al* 2014a), but inconsistent protocols, assumptions, geographic sampling, and methods make generalized interpretation of the results difficult. AgMIP's emphasis on model intercomparison and exploration of climate responses drove initial research activities toward species-based assessment at a small number of carefully selected sites. These 'pilot' projects organized around the application of multiple models on high-quality field datasets (Boote *et al* 2015, Kersebaum *et al* 2015) to expose differences in model structure, process responses, data requirements, and input/output formats.

The first crop pilot was organized by the AgMIP Wheat Team, in which 27 modeling groups ran historical simulations and 30 year sensitivity tests for [CO₂], temperature, and nitrogen response at sites in the Netherlands, India, Argentina, and Australia (Asseng *et al* 2013, Martre *et al* 2015). The Wheat Pilot was open to all interested modeling groups as long as their models were published in peer-reviewed articles.

Table 1. Overview of multi-model, multi-site, protocol-based research activities sampling the carbon–temperature–water–nitrogen–adaptation (CTWNA) change space by AgMIP and related projects.

Research activity	Scope	C (ppm)	T (°C)	W (%)	N	A	Notes
AgMIP-Wheat Phase 1	Sites	360 to 720	−3 to +9	—	−50% to +50%	—	27 wheat models at each of four sites. ^a
AgMIP-Wheat Phase 2a	Sites	—	+0 to +16	—	—	—	30 wheat models simulated at two sites.
AgMIP-Wheat Phase 2b	Global network	—	+0 to +4	—	—	—	30 wheat models at 30 well-watered sites.
AgMIP-Wheat Phase 3	Global network	360 to 550	+0 to +4	—	—	—	32 wheat models at 60 sites (water-limited and well-watered sites).
AgMIP-Maize Phase 1	Sites	360 to 720	−3 to +9	—	—	—	23 maize models at each of four sites.
AgMIP-Maize Phase 2	Sites	387 and 550	—	RF and Irr.	—	—	21 maize models for Braunschweig, Germany, FACE site.
AgMIP-Rice Phase 1	Sites	360 to 720	−3 to +9	—	varied N	—	13 rice models at each of four sites. Two sites included N treatments ranging from 30–150 kg N ha ^{−1} .
AgMIP-Rice Phase 2	Sites	360 to 720	—	—	varied N	—	16 rice models at each of two FACE sites (Japan and China). ^b
AgMIP-Potato	Sites	360 to 720	−3 to +9	−30 to +30	—	—	Nine potato models at each of four sites.
AgMIP-Canola	Sites	360 to 720	−3 to +9	−25 to +25	0% to 150% of observed	—	Eight canola models at each of seven sites.
AgMIP-Sugarcane	Sites	350 to 750	−3 to +9	−30 to +30	—	—	Two sugarcane models at seven Brazilian sites.
AgMIP-Livestock and Grasslands Phase 2	Sites	330 to 900	−1 to +8	−50 to +50	—	—	Common protocols for single model tests at 14 sites. Seven models contributed yield and GHG balance results.
AgMIP Regional Integrated Assessments	Sites	360 to 720	−2 to +8	−75 to +100	0 to 210 kg N ha ^{−1}	—	Two models each for ten sites, multiple crops at many of the sites.
C3MP	Global network	330 to 900	−1 to +8	−50 to +50	—	—	1137 simulation sets in 56 countries; 18 crop species, 23 crop models.
MACSUR-IRS Phase 1	Sites	—	−2 to +9	−50 to +50	—	—	26 wheat models at four sites in Europe.
MACSUR-Crop Rotation	Sites	374 and 550	—	—	100% and 50% of observed	—	15 models with and without crop rotations. CN sensitivity tests performed at Braunschweig, Germany, and N sensitivity at Thibie, France.
GGCMI Phase 2	Global grid	360 to 810	−1 to +6	−50 to +30 plus Irr.	10 to 200 kg N ha ^{−1}	Fully reverse accelerated maturity	12 participating models. ^b Includes no water stress test and no nitrogen stress test. Adaptation adjusts cultivars to maintain planting to maturity duration.

Notes: RF = rainfed; Irr. = irrigated; GHG = greenhouse gas; varied N = multiple nitrogen treatments at each site.

^a Nitrogen tests were only performed for 20 wheat models containing nitrogen dynamics.

^b Ongoing project, final participation may change.

Similar multi-model crop pilots were conducted across selected sites by AgMIP Maize (CT responses; Bassu *et al* 2014), AgMIP Rice (CT responses; Li *et al* 2015), AgMIP Potato (CTW responses; Fleisher *et al* 2016), AgMIP Sugarcane (CTW responses; Marin *et al* 2015), and AgMIP Canola (CTWN responses; Wang *et al* 2017). MACSUR also analyzed TW responses at a transect of four European wheat

sites, providing continuous impacts response surfaces that characterize fundamental crop model properties (Pirttioja *et al* 2015, Fronzek *et al* 2017), and examined the CN response of crop rotations (Kollas *et al* 2015, Yin *et al* 2017). AgMIP's Livestock and Grasslands Team used individual models at a number of sites to create CTW response surfaces for yield and greenhouse gas balances (Erhardt 2017). Phase 2

Table 2. AgMIP-Wheat, AgMIP-Maize, and AgMIP-Rice Team Phase descriptions.

Phase (<i>and key references</i>)	Description
AgMIP-Wheat	
Phase 1 (Asseng <i>et al</i> 2013, Martre <i>et al</i> 2015)	Protocol-based multi-model intercomparison at diverse, high-quality sites. Included limited information and full information calibration settings.
Phase 2a (Asseng <i>et al</i> 2015a, 2015b)	Protocol-based multi-model analysis of temperature response at Hot Serial Cereals artificial heating experiment in Arizona and temperature responses in Mexico.
Phase 2b (Asseng <i>et al</i> 2015a, 2015b, Liu <i>et al</i> 2016)	Intercomparison of temperature responses across 30 sites selected as a representative network of well-watered wheat production regions around the world.
Phase 3 (in progress)	Intercomparison of temperature responses across 60 sites selected to represent both well-watered and water-limited wheat production regions around the world.
AgMIP-Maize	
Phase 1 (Bassu <i>et al</i> 2014)	Protocol-based multi-model intercomparison at diverse, high-quality sites. Included limited information and full information calibration settings.
Phase 2 (Durand <i>et al</i> 2017)	Protocol-based multi-model intercomparison at Free-Air Carbon Enrichment (FACE) site in Germany.
AgMIP-Rice	
AgMIP-Rice Phase 1 (Li <i>et al</i> 2015)	Protocol-based multi-model intercomparison at diverse, high-quality sites. Included limited information and full information calibration settings.
AgMIP-Rice Phase 2 (in progress)	Protocol-based multi-model intercomparison at FACE Sites in Japan and China.

studies by AgMIP Wheat, Maize, and Rice teams have challenged models with field experiments that gauge climate sensitivity at test sites, utilizing ‘Hot Serial Cereal’ heat stress experiments for wheat (Asseng *et al* 2015a, Webber *et al* 2017) and Free-Air Carbon Enrichment data to explore [CO₂] response in maize (Durand *et al* 2017) and rice (Hasegawa *et al* 2017) (table 2). CTWN sensitivity experiments also form a key component of AgMIP’s regional integrated assessments at sites across South Asia and Sub-Saharan Africa (Rosenzweig *et al* 2017).

2.1.1. Strengths and weaknesses of site-based approaches

Intensive, multi-model intercomparisons at high-quality pilot field sites are a critical first component of model evaluation, yielding valuable insight into process responses, structural biases, data requirements, and performance across contrasting systems. These analyses are anchored in field data that enable validation of state variables (e.g. leaf-area index; above-ground biomass and N contents; plant-available soil moisture) across a number of phenological stages as well as end-of-season characteristics (e.g. grain yield and protein content, harvest index). This allows evaluation of the mechanisms by which plants respond to environmental changes, highlighting sensitive biophysical processes and growth stages that in turn help focus climate projections on fundamental stresses (e.g. drought in reproductive stages; heat stress at anthesis). Results demonstrate that multi-model ensembles consistently outperform individual models when evaluated across variables and sites (Martre *et al* 2015, Asseng *et al* 2013, Bassu *et al* 2014, Li *et al* 2015, Fleisher *et al* 2016), although at any given site a subset of models may be preferred (Castañeda-Vera *et al* 2015).

Site-based assessments from the initial AgMIP Pilots are limited in their application to IAMs as they cover only a small number of sites and farming systems. As expected, crops responded differently at the selected sites owing to unique soils, weather, cultivars, and farm management. Additional careful sampling of interactions across the broader CTWNA space is needed, as shown by the benefits of elevated [CO₂] for water use efficiency in recent AgMIP crop team activities (Cammarano *et al* 2016, Deryng *et al* 2016, Durand *et al* 2017).

2.2. Network-based approaches

As AgMIP protocols were developed and tested on individual sites, the next step scaled up these approaches through larger networks of sites coordinated to ensure adherence to a common protocol that enables direct comparison.

2.2.1. Wide ad hoc network approach

AgMIP launched the Coordinated Climate-Crop Modeling Project (C3MP; Ruane *et al* 2014, McDerimid *et al* 2015a), to create CTW impact response surfaces at a range of sites around the world. C3MP samples the CTW space projected by GCMs in the 21st century, enabling the fitting of emulators and response surfaces that can be rapidly applied to estimate the agricultural impacts of new climate projections. C3MP created information technology tools and templates to facilitate the process and invited the agricultural modeling community to participate with their own calibrated sites. The resulting archive reflects submissions from 100 crop modelers, with 1137 simulation sets from 55 countries, including results from 19 crop model families and 18 crop species (McDerimid *et al* 2015a).

2.2.1.1. Strengths and weaknesses of wide ad hoc networks

C3MP's open call for crop model participation led to an unprecedented number and diversity of contributed simulation sets but also challenges in analyses. The result is a network of voluntary 'crowd-sourced' responses rather than a designed plan of geographic coverage, representative sites, or multi-model analyses. Nevertheless, C3MP's wide ad hoc network covers most major agricultural lands and features models calibrated with site-specific information (figure 2). Sampling across all submitted results for a given category of system (e.g. rainfed maize) provides CTW response surfaces isolating the common yield response across a broad sampling of sites and systems as well as uncertainty stemming from model, soil, baseline climate, cultivar, and farming system differences (McDermid *et al* 2015a). Recognizing that IAMs typically track major crops (wheat and rice) and commodity groups (e.g. oil seeds, coarse grains, sugar crops, fruits and vegetables), C3MP's relatively large number of crop species also reduces the amount of crop response mapping that is required to represent climate responses across the diversity of agricultural commodities. C3MP is particularly useful in distinguishing responses within a commodity group (for example, differentiating between millet, sorghum, and maize responses for coarse grains).

Aggregation of the C3MP archive to global production responses is challenging given geographic gaps and under-represented systems, and vetting is difficult given its reliance on prior model calibration and a skew toward common crop models (as were also challenges in the Challinor *et al* 2014a, meta-analysis). We recommend that C3MP analyses do not include simulation sets that use antiquated model versions and a small percentage of flagged sites where low historical yields indicate farming systems that are not presently viable. In some cases, these were conducted as tests of land uses that may become viable in wetter and high-[CO₂] futures, but must be considered distinct from broader CTW analyses. C3MP remains an open process, and each new submission increases the robustness of ensemble statistics and analyses.

2.2.2. Representative network approach

AgMIP Wheat Phase 2b created a global network of 30 well-watered sites selected to represent major wheat systems and regional production areas (irrigated and high-rainfall wheat crops contribute ~70% of global production; see figure 2) (Asseng *et al* 2015a). 30 wheat models are configured for simulation of CT responses at each site, allowing robust ensemble projections and uncertainty analyses (Wallach *et al* 2015, 2016).

2.2.2.1. Strengths and weaknesses of representative networks

The AgMIP Wheat Team network is distinct from C3MP's ad hoc network in that its design allows

multi-model assessment on major regional production systems that together generate the large majority of global wheat production (table 2). Simulated relative impacts are applied to recent FAO country production statistics associated with each simulated location to up-scale to global production impacts (Asseng *et al* 2015a, Liu *et al* 2016).

Even with 30 sites, the network is limited in its spatial coverage and individual sites may not reflect conditions in the broader production regions they represent. The network is concentrated in high-production zones and is likely to miss important responses in areas that were not simulated (AgMIP-Wheat Phase 3 will fill some of these gaps for water-stressed systems). As a simple metric of comprehensiveness of coverage, figure 3 shows how the rainfed and irrigated wheat networks from C3MP and AgMIP-Wheat Phase 2 cover wheat-growing climate conditions as compared with the global Monthly Irrigated and Rainfed Crop Area (MIRCA) year 2000 dataset (Portmann *et al* 2010), using AgMERRA climate data (Ruane *et al* 2015a) and growing seasons from the Global Gridded Crop Model Intercomparison (GGCMI; Elliott *et al* 2015). Both networks are most dense in climate zones that are prominent for wheat production; however the larger C3MP network also includes less common climates for rainfed wheat and samples more from the tails of the irrigated wheat distribution than does AgMIP-Wheat. By simulating more of the cool and wet tails it is likely that C3MP captures more farms that potentially benefit from increases in temperature or are less vulnerable to decreases in precipitation.

Regions with high levels of diversity are difficult to capture given limitations in representative site networks. Sentinel crop modeling sites are often calibrated with data from field experiment datasets designed to highlight potential genetic, fertilizer, water, or pest control treatments, and therefore may not be representative of prevailing agricultural systems within that production region. These site networks tend to be more useful when examining the percentage yield response to a given climate change; this metric has proven robust even in the face of persistent bias in mean regional yields (Challinor *et al* 2014b, Asseng *et al* 2015a).

2.3. Global approaches

Advances in high-performance computing have allowed crop models to enter a new phase of development that is nearly unconstrained by computational limitations. While IAMs are typically run on desktop computers or simple clusters, the 18 modeling groups participating in AgMIP's Global Gridded Crop Modeling Intercomparison (GGCMI; table 3) use parallel computing and advanced data processing pipelines to conduct protocol-based simulations on a 0.5° × 0.5° global grid (Rosenzweig *et al* 2014, Elliott *et al* 2015), with higher resolution gridded studies in the works. These outputs therefore form a desirable basis for more computationally-efficient IAM application through

Table 3. Overview of GGCM phases and model participation.

Phase (and key references)	Description [and # of models participating]
Fast Track Rapid Assessment (Rosenzweig <i>et al</i> 2014)	Conducted for AgMIP/ISIMIP using default versions of global gridded crop models, historical period and future scenarios from downscaled GCMs. Simulated maize, wheat, rice and soybean [seven GGCMs].
Phase 1 Historical Intercomparison (Elliott <i>et al</i> 2015, Müller <i>et al</i> 2017)	Default, harmonized, and no nitrogen stress versions of gridded crop models run over historical period using up to nine climate forcing datasets. Simulated maize, wheat, rice and soybean [15 GGCMs].
Phase 2 CTWNA Sensitivity [results submitted 2016–17]	Default simulations for historical period and sensitivity tests for [CO ₂], temperature, water, nitrogen, and adaptation for all grid cells and crops. Simulated maize, spring wheat, winter wheat, rice and soybean [~12 GGCMs].
Phase 3 Future Assessment [planned for 2017–18]	Conducted for AgMIP/ISIMIP to assess future agricultural production under climate change scenarios [~12–20 GGCMs anticipated].

Note: ISIMIP = the Inter-Sectoral Impacts Model Intercomparison Project (Warszawski *et al* 2014).





Figure 3. Distribution of global (a) rainfed and (b) irrigated wheat area displayed according to growing season mean temperature and precipitation rate (from MIRCA observations, Portmann *et al* 2010). Corresponding C3MP and AgMIP-Wheat Phase 2b network sites are presented to show coverage of global wheat-cropping systems. Note that the AgMIP-Wheat 2b network consists entirely of well-watered sites including both irrigated croplands and rainfed areas that rarely experience water stress.



emulators. GGCM Phase 2 performs a systematic analysis of CTWNA sensitivities for rainfed and irrigated maize, rice, wheat, and soybean with consistent climate information and harmonized planting dates. Adaptation is examined by shifting cultivars to maintain the growing period even as warmer temperatures accelerate phenologic development, thus offsetting some yield losses from climate change.

2.3.1. Strengths and weaknesses of global approaches

GGCM's fast-track results (table 3) provide biophysical impacts across emissions scenarios and five GCMs (Rosenzweig *et al* 2014), providing applications with ensemble mean impacts and uncertainty information from seven GGCMs for four crop species (maize, wheat, rice, and soybean) across the global grid (Nelson *et al* 2014, Wiebe *et al* 2015, Villoria *et al* 2016). It is difficult for crop model emulators to disentangle fundamental responses from these outputs, however, given the many types of changing and interacting climate conditions (e.g. mean temperatures and rainfall; sub-seasonal variations; extreme events). Emulation is also complicated by the inclusion of responsive adaptations allowing management to evolve with climate change in some participating models (Rosenzweig *et al* 2014, supplementary).

GGCM Phase 2 findings indicate considerable spatial variation in CTWNA response across different environments and farm systems, exemplified by the response of rainfed maize to higher $[\text{CO}_2]$ and temperature in the parallel-DSSAT crop model (pDSSAT; Elliott *et al* 2014) (figure 4). These results provide a convenient basis for the construction of crop model emulators, and can also be connected to economic and/or resource availability drivers from IAMs to


dynamically characterize the evolution of socioeconomic yield gap factors such as fertilizer use, irrigation, and adaptation potential.

In contrast to the site networks, GGCMs rely on gridded soil, genetic, management, and weather datasets designed to capture spatially-averaged conditions rather than conditions on a particular farm (Elliott *et al* 2015). While the $0.5^\circ \times 0.5^\circ$ spatial resolution used within GGCM is finer than many GCMs, a grid cell on the equator represents $>310\,000$ ha and thus poses a challenge for comprehensive farm system calibration.

GGCM results are often evaluated using regional yield and production reports, with trend adjustment recommended in recognition of technological development and processes that are not explicitly modeled such as pests, diseases, and widespread flooding (Müller *et al* 2017). Analogously, statistical crop response models are occasionally fitted to similar aggregate yield data that may reflect embedded abiotic factors (e.g. Lobell *et al* 2011). Bias-adjustment is recommended for GGCM application in IAMs, similar to common practices for climate model output applications (e.g. Wilby *et al* 2004, Ruane *et al* 2015b). Overall, GGCMs reflect that there are larger uncertainties in developing country and low-input farming systems, and stand to benefit from improved data collection and sharing in application regions (Kihara *et al* 2015, McDermid *et al* 2015b).

2.4. Emergent characteristics and opportunities from CTWNA simulations

AgMIP site, network, and gridded results demonstrate that multi-model ensembles outperform individual models when analyzed across multiple sites and evaluation variables (e.g. Asseng *et al* 2013,




Rosenzweig *et al* 2014, Martre *et al* 2015, Li *et al* 2015, Bassu *et al* 2014, Wallach *et al* 2015, Ruane *et al* 2016, Fleisher *et al* 2016). Liu *et al* (2016) found relative agreement in wheat response to a 1 °C rise in global temperature, with multi-model ensembles in the well-watered AgMIP Wheat network, GGCM's ISIMIP fast-track, and several statistical model approaches finding 4.1%–6.4% declines in global production.

Uncertainties in input data indicate that there is still room for harmonization that will improve consistency, as illustrated by a comparison of growing seasons at the well-watered AgMIP-Wheat network sites and corresponding GGCM grid cells (figure 5). Uncertainty owing to model structure and parameters remains substantial, and differences in CTWNA responses by two modelers using the same DSSAT model within the MACSUR IRS and AgMIP-Wheat Phase 1 also highlights the potential role of modeler uncertainty stemming from assumptions and subjective decisions made in the absence of supporting data

(Pirttioja *et al* 2015, Confalonieri *et al* 2016). We therefore advise applications to recognize the uncertainty in model-based responses through the use of emulators derived from multiple models or an imposed error term scaled to model-based uncertainty. Evidence across AgMIP activities also recommends avoidance of universal yield functions in favor of yield response functions fitted to broad agro-ecological zones and farming systems (e.g. defined by fertilizer and irrigation inputs).

3. AgMIP framework for improved agricultural representation in IAMs

A cascading pathway of development underlies agricultural representation in IAMs, forming a framework that may be used to drive coordinated development of 'simulation levels', here defined as common communities of development including site-based crop models, network and gridded models, crop model emulators,



and eventual IAM applications (figure 6). Close collaboration and regular updates between site, network, and gridded crop modelers, emulation experts, and IAM groups are needed to keep agricultural impact applications on the cutting edge, to facilitate the use of multiple models, to incorporate understanding from multiple modeling groups, and to avoid the propagation of known biases.

Each simulation level in the AgMIP Framework benefits from improved data access and innovations in core methodologies. Investment in research and development is well served by matching the design, capabilities, and development priorities of models and tools at each level in figure 6. In particular, new biophysical process understanding is best developed within site-based models using field experiment data, particularly for under-sampled agro-ecological zones, crop species, and farming systems under various intensifications (Challinor *et al* 2015, Maiorano *et al* 2017).

Networks and gridded models gain from new datasets that allow extensive configuration for many sites and systems, and have tremendous potential to apply advanced bias-correction and aggregation approaches (Challinor *et al* 2014, van Bussel *et al* 2016, Hoffmann *et al* 2015, 2016, Zhao *et al* 2015, 2016). Crop model emulators are progressed with improved statistical efficiencies and the availability of observed agricultural response data for evaluating strengths and weaknesses. In addition to the potential benefit of adding improved crop model emulators, IAM simulations of long-term shifts in agricultural production are furthered by good data on current systems and advanced representation of the implications of agricultural investment and technological development.

The AgMIP framework for improved agricultural representation in IAMs is non-linear as lower simulation levels build upon advances higher up in the framework and high levels also receive critical

feedback from downstream simulation levels. Pathways of upstream improvements include that assessments of improved models on established grids and networks provide vital feedback for site-based model development on diverse sites. Likewise, emulators often spotlight key sensitivities and uncertainties that may spur further site-based model development and the creation of more representative networks. Network and gridded studies examine the biophysical viability of various simulated farm systems to determine land use pressures, but benefit tremendously by incorporating information on economic viability and resource constraints that IAMs can provide. It is also important to note that many of these simulation levels have extensive applications beyond agricultural representation in IAMs, and that the key bottleneck for one applications may differ from another's crucial development priority.

4. Priority future development and applications

Analysis of the multi-model, multi-site climate sensitivity datasets reviewed in this study suggest that IAMs and other large-scale applications would be well served by the creation and systematic development of a hybrid CTWNA response system that blends the strengths of network and gridded approaches (as noted in figure 6). This hybrid response system would be rooted in (1) detailed process understanding across a representative network of well-calibrated field sites (ideally using field data from prevailing management systems) combined with (2) comprehensive CTWNA coverage from gridded models. Baseline responses generated by these gridded models could initially be compared against the corresponding representative network simulations to assess methodological uncertainty and calculate bias-correction factors. Bias-corrected gridded results could then provide an information basis for crop model emulators and IAM applications, characterizing different farming systems using nitrogen and water components of the CTWNA analysis.

Table 1 highlights that progress toward the creation of this hybrid response system is most advanced for wheat, given the AgMIP-Wheat Phase 2b representative network and spring and winter wheat simulations within GGCM Phase 2. In contrast, soybean is simulated in GGCM but has not yet been the focus of site- or network-based CTWNA analysis, and a number of other important commodities merit inclusion. Coordinated and systematic development of the hybrid response system would foster rapid iterative improvements, as research groups improve the hybrid framework by contributing new process understanding, field sites, model runs, regional configuration information, or statistical approaches. An expanded representative network of models and a fully configured high-resolution gridded (or geo-referenced polygon) model will eventually be interchangeable; however this

hybrid response system provides current state-of-the-art responses and a practical roadmap for applications.

Coordination across AgMIP activities supports the development of linked global and regional assessments to address agricultural sector challenges and food security (Rosenzweig *et al* 2016). Inclusion of IAMs would bring these to a new level, although it is critical that these account for lingering model uncertainty and data gaps even as these are addressed through the coordinated development of agricultural response in linked models.

Acknowledgements

We are grateful for the tremendous effort and contributions provided by all modelers and leaders of all projects listed in table 1. This letter was motivated by discussions with members of the Program on Integrated Assessment Model Development, Diagnostics, and Intercomparisons (PIAMDDI) and the Pacific Northwest National Laboratories, and benefitted from support by the National Aeronautics and Space Agency Science Mission Directorate (WBS 281945.02.03.06.79), the United Kingdom Department for International Development (GB-1-202108), and the Department of Energy (DE-AC5-76RL01830 and DE-SC0005171) as well as the Joint Research Centre of the European Commission. We appreciate the assistance of Meridel Phillips and Greg Reppucci in figure creation.

ORCID iDs

Alex C Ruane  <https://orcid.org/0000-0002-5582-9217>

Cynthia Rosenzweig  <https://orcid.org/0000-0002-7885-671X>

Christoph Müller  <https://orcid.org/0000-0002-9491-3550>

References

- Antle J M, Valdivia R O, Boote K, Janssen S, Jones J W, Porter C H, Rosenzweig C, Ruane A C and Thorburn P J 2015 AgMIP's transdisciplinary agricultural systems approach to regional integrated assessment of climate impacts, vulnerability, and adaptation *Handbook of Climate Change and Agroecosystems: The Agricultural Model Intercomparison and Improvement Project (AgMIP) Integrated Crop and Economic Assessments, Part 1 ICP Series on Climate Change Impacts, Adaptation, and Mitigation* ed C Rosenzweig and D Hillel vol 3 (London: Imperial College Press) pp 27–44
- Asseng S *et al* 2013 Uncertainty in simulating wheat yields under climate change *Nat. Clim. Change* **3** 827–32
- Asseng S *et al* 2015a Rising temperatures reduce global wheat production *Nat. Clim. Change* **5** 143–7
- Asseng S *et al* 2015b Benchmark data set for wheat growth models: field experiments and AgMIP multi-model simulations *Open Data J. Agric. Res.* **1** 1–5
- Bassu S *et al* 2014 Do various maize crop models give the same responses to climate change factors? *Glob. Change Biol.* **20** 2301–20

- Blanc E 2017 Statistical emulators of maize, rice, soybean and wheat yields from global gridded crop models *Agric. Forest Meteorol.* **236** 145–61
- Boote K J, Porter C H, Jones J W, Thorburn P J, Kersebaum K C, Hoogenboom G, White J W and Hatfield J L 2015 Sentinel site data for model improvement—definition and characterization *Improving Modeling Tools to Assess Climate Change Effects on Crop Response, Advances in Agricultural Systems Modeling* ed J L Hatfield and D Fleisher (Madison, WI: ASA, CSSA, and SSSA)
- Cammarano D *et al* 2016 Uncertainty of wheat water use: Simulated patterns and sensitivity to temperature and CO₂ *Field Crops Res.* **198** 80–92
- Castañeda-Vera A, Leffelaar P A, Álvaro-Fuentes J, Cantero-Martínez C and Mínguez M I 2015 Selecting crop models for decision making in wheat insurance *Eur. J. Agron.* **68** 97–116
- Castruccio S, McInerney D J, Stein M L, Crouch F L, Jacob R L and Moyer E J 2014 Statistical emulation of climate model projections based on precomputed GCM runs *J. Clim.* **27** 1829–44
- Challinor A J *et al* 2014a A meta-analysis of crop yield under climate change and adaptation *Nat. Clim. Change* **4** 287–91
- Challinor A, Martre P, Asseng S, Thornton P and Ewert F 2014b Making the most of climate impacts ensembles *Nat. Clim. Change* **4** 77–80
- Challinor A, Parkes B and Ramirez-Villegas J 2015 Crop yield response to climate change varies with cropping intensity *Glob. Change Biol.* **21** 1679–88
- Clarke L *et al* 2014 Assessing transformation pathways climate change 2014: mitigation of climate change *Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* ed O Edenhofer *et al* (Cambridge: Cambridge University Press)
- Confalonieri R *et al* 2016 Uncertainty in crop model predictions: what is the role of users? *Environ. Model. Softw.* **81** 165–73
- Crimp S, Howden M, Power B, Wang E and deVoil P 2008 Global climate change impacts on Australia's wheat crops *Garnaut Climate Change report* (Canberra)
- Deryng D *et al* 2016 Regional disparities in the beneficial effects of rising CO₂ concentrations on crop water productivity *Nature Clim. Change* **6** 786–90
- Durand J L *et al* 2017 How accurately do maize crop models simulate the interactions of atmospheric CO₂ concentration levels with limited water supply on water use and yield? *Eur. J. Agron.* accepted (<https://doi.org/10.1016/j.eja.2017.01.002>)
- Elliott J, Kelly D, Chryssanthacopoulos J, Glotter M, Jhunjhnuwala K, Best N, Wilde M and Foster I 2014 The parallel system for integrating impact models and sectors (pSIMS) *Environ. Model. Softw.* **62** 509–16
- Elliott J *et al* 2015 The global gridded crop model intercomparison: data and modeling protocols for phase 1 (v1.0) *Geosci. Model Dev.* **8** 261–77
- Erhardt F 2017 personal communication
- Ewert F *et al* 2015 Crop modelling for integrated assessment of risk to food production from climate change *Environ. Model. Softw.* **72** 287–303
- Fleisher D H *et al* 2016 A potato model intercomparison across varying climates and productivity levels *Glob. Change Biol.* **23** 1258–81
- Fronzek S *et al* 2017 Classifying multi-model wheat yield impact response surfaces showing sensitivity to temperature and precipitation change *Agric. Syst.* accepted (<https://doi.org/10.1016/j.agry.2017.08.004>)
- Füssel H-M 2010 Modeling impacts and adaptation in global IAMs *WIREs Clim. Change* **1** 288–303
- Hartin C A, Patel P L, Schwarber A, Link R P and Bond-Lamberty B 2015 A simple object-oriented and open source model for scientific and policy analyses of the global climate system—Hector v1.0 *Geosci. Model Dev.* **8** 939–55
- Hasegawa *et al* 2017 personal communication
- Hoffmann H *et al* 2015 Variability of effects of spatial climate data aggregation on regional yield simulation by crop models *Clim. Res.* **65** 53–69
- Hoffmann H *et al* 2016 Impact of spatial soil and climate input data aggregation on regional yield simulations *PLoS ONE* **11** e015178
- Howden M and Crimp S 2005 Assessing dangerous climate change impacts on Australia's wheat industry *MODSIM 2005 International Congress on Modeling and Simulation (December 2005)* pp 170–76 (Modelling and Simulation Society of Australia and New Zealand)
- Kersebaum K C *et al* 2015 Analysis and classification of data sets for calibration and validation of agro-ecosystem models *Environ. Model. Softw.* **72** 402–17
- Kihara J *et al* 2015 Perspectives on climate effects on agriculture: the international efforts of AgMIP in sub-saharan Africa *Handbook of Climate Change and Agroecosystems: The Agricultural Model Intercomparison and Improvement Project (AgMIP) Integrated Crop and Economic Assessments, Part 2 ICP Series on Climate Change Impacts, Adaptation, and Mitigation* ed C Rosenzweig and D Hillel vol 3 (London: Imperial College Press) pp 3–24
- Kollas C *et al* 2015 Crop rotation modelling—a European model intercomparison *Eur. J. Agron.* **70** 98–111
- Li T *et al* 2015 Uncertainties in predicting rice yield by current crop models under a wide range of climatic conditions *Glob. Change Biol.* **21** 1328–41
- Liu B *et al* 2016 Similar negative impacts of temperature on global wheat yield estimated by three independent methods *Nature Clim. Change* **6** 1130–6
- Lobell D B, Schlenker W and Costa-Roberts J 2011 Climate trends and global crop production since 1980 *Science* **333** 616–20
- Maiorano A *et al* 2017 Crop model improvement reduces the uncertainty of the response to temperature of multi-model ensembles *Field Crops Res.* **202** 5–20
- Makowski D *et al* 2015 Statistical analysis of large simulated yield datasets for studying climate effects *Handbook of Climate Change and Agroecosystems: The Agricultural Model Intercomparison and Improvement Project (AgMIP) Integrated Crop and Economic Assessments, Part 1 ICP Series on Climate Change Impacts, Adaptation, and Mitigation* ed C Rosenzweig and D Hillel vol 3 (London: Imperial College Press) pp 279–98
- Marin F R, Thorburn P J, Nassif D S P and Costa L G 2015 Sugarcane model intercomparison: structural differences and uncertainties under climate change *Environ. Model. Softw.* **72** 372–86
- Martre P *et al* 2015 Error of multimodel ensembles of wheat growth: more models are better than one *Glob. Change Biol.* **21** 911–25
- McDermid S P *et al* 2015a The agmip coordinated climate-crop modeling project (C3MP): methods and protocols *Handbook of Climate Change and Agroecosystems: The Agricultural Model Intercomparison and Improvement Project (AgMIP) ICP Series on Climate Change Impacts, Adaptation, and Mitigation* ed C Rosenzweig and D Hillel vol 3 (London: Imperial College Press) pp 191–220
- McDermid S P *et al* 2015b Integrated assessments of the impacts of climate change on agriculture: an overview of AgMIP regional research in South Asia *Handbook of Climate Change and Agroecosystems: The Agricultural Model Intercomparison and Improvement Project (AgMIP) Integrated Crop and Economic Assessments, Part 2 ICP Series on Climate Change Impacts, Adaptation, and Mitigation* ed C Rosenzweig and D Hillel vol 3 (London: Imperial College Press) pp 201–18
- Meinshausen M, Raper S C B and Wigley T M L 2011 Emulating coupled atmosphere-ocean and carbon cycle models with a simpler model, MAGICC6—Part 1: model description and calibration *Atmos. Chem. Phys.* **11** 1417–56
- Mistry M N, Sue Wing I and De Cian E 2017 Simulated vs. empirical weather responsiveness of crop yields: US evidence and implications for the agricultural impacts of climate change *Environ. Res. Lett.* **12** 075007
- Monfreda C, Ramankutty N and Foley J A 2008 Farming the planet: 2. Geographic distribution of crop areas, yields, physiological types, and net primary production in the year 2000 *Glob. Biogeochem. Cycles* **22** GB1022

- Moore F, Baldos U L C and Hertel T 2017 Economic impacts of climate change on agriculture: a comparison of process-based and statistical yield models *Environ. Res. Lett.* **12** 065008
- Müller C and Robertson R D 2014 Projecting future crop productivity for global economic modeling *Agric. Econ.* **45** 37–50
- Müller C *et al* 2017 Global Gridded Crop Model evaluation: Benchmarking, skills, deficiencies and implications *Geosci. Model Dev.* **10** 1403–22
- Nelson G C *et al* 2014 Climate change effects on agriculture: economic responses to biophysical shocks *Proc. Natl Acad. Sci.* **111** 3274–9
- Pirttioja N 2015 A crop model ensemble analysis of temperature and precipitation effects on wheat yield across a European transect using impact response surfaces *Clim. Res.* **65** 87–105
- Portmann F T, Siebert S and Döll P 2010 MIRCA2000—Global monthly irrigated and rainfed crop areas around the year 2000: a new high-resolution data set for agricultural and hydrological modeling *Glob. Biogeochem. Cycles* **24** GB1011
- Ray D K, Gerber J S, MacDonald G K and West P C 2015 Climate variation explains a third of global crop yield variability *Nat. Commun.* **6** 5989
- Rosenzweig C *et al* 2013 The agricultural model intercomparison and improvement project (AgMIP): protocols and pilot studies *Agric. Forest Meteorol.* **170** 166–82
- Rosenzweig C *et al* 2014 Assessing agricultural risks of climate change in the 21st century in a global gridded crop model intercomparison *Proc. Natl Acad. Sci.* **111** 3268–73
- Rosenzweig C, Jones J W, Hatfield J L, Antle J M, Ruane A C and Mutter C Z 2015 The agricultural model intercomparison and improvement project: phase I activities by a global community of science *Handbook of Climate Change and Agroecosystems: The Agricultural Model Intercomparison and Improvement Project (AgMIP) ICP Series on Climate Change Impacts, Adaptation, and Mitigation* ed C Rosenzweig and D Hillel vol 3 (London: Imperial College Press) pp 3–24
- Rosenzweig C, Antle J and Elliott J 2016 Assessing impacts of climate change on food security worldwide *Eos* **97** EO047387
- Rosenzweig C *et al* 2017 Protocols for AgMIP Regional Integrated Assessments (www.agmip.org/regional-integrated-assessments-handbook/)
- Ruane A C, McDermid S, Rosenzweig C, Baigorria G A, Jones J W, Romero C C and Cecil L D 2014 Carbon–temperature–water change analysis for peanut production under climate change: a prototype for the AgMIP coordinated climate–crop modeling project (C3MP) *Glob. Change Biol.* **20** 394–407
- Ruane A C, Goldberg R and Chryssanthacopoulos J 2015a Climate forcing datasets for agricultural modeling: Merged products for gap-filling and historical climate series estimation *Agric. Forest Meteorol.* **200** 233–48
- Ruane A C, Winter J M, McDermid S P and Hudson N I 2015b Agmip climate datasets and scenarios for integrated assessment *Handbook of Climate Change and Agroecosystems: The Agricultural Model Intercomparison and Improvement Project (AgMIP) Integrated Crop and Economic Assessments, Part 1 ICP Series on Climate Change Impacts Adaptation Mitigation* ed C Rosenzweig and D Hillel vol 3 (London: Imperial College Press) pp 45–78
- Ruane A C *et al* 2016 Multi-wheat model ensemble responses to interannual climate variability *Environ. Model. Softw.* **81** 86–101
- van Bussel L *et al* 2016 Spatial sampling of weather data for regional crop yield simulations *Agric. Forest Meteorol.* **220** 101–15
- Villoria N *et al* 2016 Rapid aggregation of global gridded crop model outputs to facilitate cross-disciplinary analysis of climate change impacts in agriculture *Environ. Model. Softw.* **75** 193–201
- Wallach D, Mearns L O, Rivington M, Antle J M and Ruane A C 2015 Uncertainty in agricultural impact assessment *Handbook of Climate Change and Agroecosystems: The Agricultural Model Intercomparison and Improvement Project (AgMIP) ICP Series on Climate Change Impacts, Adaptation, and Mitigation* ed C Rosenzweig and D Hillel vol 3 (London: Imperial College Press) pp 223–60
- Wallach D, Mearns L O, Ruane A C, Rötter R P and Asseng S 2016 Lessons from the climate modeling community on the design and use of ensembles for crop modeling *Clim. Change* **139** 551–64
- Wang *et al* 2017 personal communication
- Warszawski L, Frieler K, Huber V, Piontek F, Serdeczny O and Schewe J 2014 The Inter-Sectoral Impact Model Intercomparison Project (ISI-MIP): project framework *Proc. Natl Acad. Sci. USA* **111** 3228–32
- Webber H, Zhao G, Wolf J, Britz W, Vries W D, Gaiser T, Hoffmann H and Ewert F 2015 Climate change impacts on European crop yields: do we need to consider nitrogen limitation? *Eur. J. Agron.* **71** 123–34
- Webber H *et al* 2017 Canopy temperature for simulation of heat stress in irrigated wheat in a semi-arid environment: a multi-model comparison *Field Crops Res.* **202** 21–35
- White J W, Hoogenboom G, Kimball B A and Wall G W 2011 Methodologies for simulating impacts of climate change on crop production *Field Crops Res.* **124** 357–68
- Wiebe K *et al* 2015 Climate change impacts on agriculture in 2050 under a range of plausible socioeconomic and emissions scenarios *Environ. Res. Lett.* **10** 085010
- Wilby R L, Charles S, Zorita E, Timbal B, Whetton P and Mearns L 2004 Guidelines for use of climate scenarios developed from statistical downscaling methods *IPCC Task Group on Data and Scenario Support for Impacts and Climate Analysis* (www.ipcc-data.org/guidelines/dgm_no2_v1_09_2004.pdf) (Accessed: 2 October 2017)
- Yin X *et al* 2017 Multi-model uncertainty analysis in predicting grain N for crop rotations in Europe *Eur. J. Agron.* **84** 152–65
- Zhao G *et al* 2015 Effect of weather data aggregation on regional crop simulation for different crops, production conditions, and response
- Zhao G *et al* 2016 Evaluating the precision of eight spatial sampling schemes in estimating regional means of simulated yield for two crops *Environ. Model. Softw.* **80** 100–12