



# Cytokinins and the CRE1 receptor influence endogenous gibberellin levels in *Medicago truncatula*

Camille Fonouni-Farde, Erin McAdam, David Nichols, Anouck Diet, Eloise Foo & Florian Frugier

To cite this article: Camille Fonouni-Farde, Erin McAdam, David Nichols, Anouck Diet, Eloise Foo & Florian Frugier (2018) Cytokinins and the CRE1 receptor influence endogenous gibberellin levels in *Medicago truncatula*, *Plant Signaling & Behavior*, 13:2, e1428513, DOI: [10.1080/15592324.2018.1428513](https://doi.org/10.1080/15592324.2018.1428513)

To link to this article: <https://doi.org/10.1080/15592324.2018.1428513>



© 2018 The Author(s). Published with license by Taylor & Francis© Camille Fonouni-Farde, Erin McAdam, David Nichols, Anouck Diet, Eloise Foo and Florian Frugier



View supplementary material [↗](#)



Accepted author version posted online: 26 Jan 2018.  
 Published online: 06 Feb 2018.



Submit your article to this journal [↗](#)



Article views: 304



View related articles [↗](#)



View Crossmark data [↗](#)

## Cytokinins and the CRE1 receptor influence endogenous gibberellin levels in *Medicago truncatula*

Camille Fonouni-Farde<sup>a</sup>, Erin McAdam<sup>b</sup>, David Nichols<sup>c</sup>, Anouck Diet<sup>a</sup>, Eloise Foo<sup>b</sup>, and Florian Frugier<sup>a</sup>

<sup>a</sup>Institute of Plant Sciences Paris-Saclay (IP2S), Centre National de la Recherche Scientifique, Univ. Paris-Sud, Univ. Paris-Diderot, Univ. Evry, Institut National de la Recherche Agronomique, Université Paris-Saclay, Gif-sur-Yvette, France; <sup>b</sup>School of Natural Sciences, University of Tasmania, Hobart, Tasmania, Australia; <sup>c</sup>Central Science Laboratory, University of Tasmania, Hobart, Tasmania, Australia

### ABSTRACT

Gibberellins (GAs) and cytokinins (CKs) are hormones that play antagonistic roles in several developmental processes in plants. However, there has been little exploration of their reciprocal interactions. Recent work in *Medicago truncatula* has revealed that GA signalling can regulate CK levels and response in roots. Here, we examine the reciprocal interaction, by assessing how CKs and the CRE1 (Cytokinin Response 1) CK receptor may influence endogenous GA levels. Real-Time RT-PCR analyses revealed that the expression of key GA biosynthesis genes is regulated in response to a short-term CK treatment and requires the CRE1 receptor. Similarly, GA quantifications indicated that a short-term CK treatment decreases the GA<sub>1</sub> pool in wild-type plants and that GA levels are increased in the *cre1* mutant compared to the wild-type. These data suggest that the *M. truncatula* CRE1-dependent CK signaling pathway negatively regulates bioactive GA levels.

### ARTICLE HISTORY

Received 22 November 2017  
Revised 19 December 2017  
Accepted 26 December 2017

### KEYWORDS

Cytokinin; gibberellin;  
legume; *Medicago truncatula*



Gibberellin (GA) and cytokinin (CK) hormones are known to play antagonistic roles in several plant developmental processes, and some studies have reported development-dependent interactions between these two hormonal pathways.<sup>1</sup> In *Arabidopsis thaliana* seedlings, the up-regulation of GA level and/or GA responses was reported to inhibit CK signaling,<sup>2</sup> and similarly, GA signaling negatively regulates CK levels and/or signaling in *A. thaliana* and *Medicago truncatula* roots.<sup>3,4</sup> However, the reciprocal regulation of endogenous GA levels by CKs is less clear. In *A. thaliana* inflorescences, semi-quantitative RNA gel blot analyses suggested that CKs had no effect on the expression of a GA-responsive gene or a GA biosynthesis gene.<sup>2</sup> However, in the *Arabidopsis* shoot apical meristem, a CK-treatment induced the expression of *AtGA2oxidase* (*GA2ox*) gene, which encodes an enzyme that deactivates bioactive GA,<sup>5</sup> and a genome-wide expression analysis reported the downregulation of several GA biosynthesis genes after a short-term CK treatment in whole *Arabidopsis* seedlings.<sup>6</sup> Similarly, transcriptomic studies performed in *M. truncatula* roots revealed a significant change in the expression of GA biosynthesis and response genes after short-term CK treatment,<sup>7</sup> suggesting that CKs may suppress endogenous GA levels. However, no study has examined the actual changes in endogenous GA levels in response to CKs. Here, we explored whether CKs and the CRE1 (Cytokinin Response 1) receptor that mediates CK responses<sup>8</sup> influenced endogenous GA levels in *M. truncatula*.


Based on transcriptomic data gained in *M. truncatula* roots,<sup>7</sup> we selected two genes rapidly regulated by a CK treatment; *GA20ox1* (GA<sub>20</sub>-oxidase) and *GA2ox1* (GA<sub>2</sub>-oxidase). The

*GA20ox1* gene is predicted to encode an enzyme that catalyzes the formation of the inactive GA precursor GA<sub>20</sub>, which can then be converted to bioactive GA<sub>1</sub> by a *GA3ox* enzyme.<sup>9</sup> The *GA2ox1* gene encodes a putative enzyme that catalyzes the deactivation of GA<sub>1</sub> into inactive GA<sub>8</sub>.<sup>9</sup> The expression of these genes was analyzed by real time RT-PCR as previously described in Fonouni-Farde et al.,<sup>4</sup> in wild-type (WT) and *cre1* mutant roots in response to a short-term CK treatment (BenzylAminoPurine [BAP; Sigma]; 3h; 10<sup>-7</sup>M) (Fig. 1A). In WT, the expression of *GA20ox1* was significantly induced by CKs, whereas the expression of *GA20ox1* was conversely repressed. These CK-dependent regulations were abolished in the *cre1* mutant (Fig. 1A), indicating that a subset of GA metabolic genes requires the CRE1 receptor to be transcriptionally regulated by CKs.

Based on the expression patterns observed in response to the BAP treatment (Fig. 1A), we hypothesized that there may be an increase of bioactive GAs in the *cre1* mutant compared to the WT. We therefore quantified GAs in WT and *cre1* mutant roots following a protocol derived from Boden et al.<sup>10</sup> (Supp. Methods). Like in the closely related legume pea,<sup>11</sup> GA<sub>1</sub> appears to be the predominant bioactive form in *M. truncatula*, as we could detect GA<sub>1</sub> but not GA<sub>4</sub>. A small increase in GA<sub>1</sub> was detected in the roots of the *cre1* mutant compared to the WT, although this change was not significant (Fig. 1B).


As GA<sub>1</sub> levels were near the detection limit in *M. truncatula* roots, we also quantified GAs in shoots of WT plants after a short-term CK treatment (BAP; 3 h; 10<sup>-7</sup>M) (Fig. 1C, D) and

**CONTACT** Florian Frugier  [florian.frugier@cnsr.fr](mailto:florian.frugier@cnsr.fr)  Institute of Plant Sciences - Paris-Saclay (IP2S), Bat. 630, rue de Noetzlin, CS80004, 91192 - Gif-sur-Yvette cedex, France.

 Supplemental data for this article can be accessed on the [publisher's website](#).

© 2018 Camille Fonouni-Farde, Erin McAdam, David Nichols, Anouck Diet, Eloise Foo and Florian Frugier. Published with license by Taylor & Francis

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.



**Figure 1.** CKs and CRE1 regulate GA-metabolic genes and endogenous GA levels. (A) Quantification by RT-qPCR of the expression of *GA2ox1* and *GA20ox1* GA-metabolic genes, previously shown to be regulated by CKs (Ariel et al., 2012), in wild-type (WT) or in *cre1* mutant roots after a BenzylAminoPurine (BAP [Sigma]; 3 h;  $10^{-7}$  M) treatment. (B) Quantification (in ng/g of Root Fresh Weigh [RFW]) of the bioactive *GA1* in WT or *cre1* mutant roots. (C-D) Quantification (in ng/g of Shoot Fresh Weigh [SFW]) of the bioactive *GA1* (C) or the precursor *GA20* (D) in WT shoots treated or not with BAP (3 h;  $10^{-7}$  M). (E-F) Quantification (in ng/g of SFW) of *GA1* (E) or *GA20* (F) in WT or *cre1* mutant shoots. In (A), transcript levels are normalized relatively to untreated control roots to show fold changes and the dotted line indicates a ratio of 1. Error bars represent standard deviations. Asterisks indicate significant differences compared to the untreated control, based on a Mann-Whitney test ( $\alpha < 0.05$ ). Results are the mean of three biological replicates, each replicate being a pool of 25 plants. In (B-F), error bars represent standard errors of the mean. Letters indicate significant differences based on a Mann-Whitney test ( $\alpha < 0.05$ ). In (B), the *GA1* quantification was based on four biological replicates, each replicate being a pool of at least two plants. In (C-F), GA quantifications were based on at least five biological replicates, each replicate being a pool of at least two plants.

in the *cre1* mutant (Fig. 1E, F). In the WT, the level of the bioactive *GA1* was significantly reduced in response to the CK treatment, whereas a small but not significant increase was observed for the immediate precursor to *GA1*, *GA20* (Fig. 1C, D). To determine if the CRE1 receptor was involved in the regulation of endogenous GAs level, we quantified *GA1* and *GA20* in WT and *cre1* shoots (Fig. 1E, F). While the content of *GA1* showed no statistical difference between the WT and the *cre1* mutant, the level of *GA20* was significantly higher in the *cre1*

mutant (Fig. 1F). Taken together, these results indicate that CKs and the CRE1 receptor are able to regulate the level of a GA endogenous precursor and/or of a biologically active GA in *M. truncatula*.

In this study, we show that CKs through the action of the CRE1 receptor influence the levels of GAs in *M. truncatula*. Together with previous results reporting that GAs can regulate CK levels and response in *M. truncatula* roots,<sup>4,7</sup> this suggests that there is a bidirectional antagonistic regulation of CK and GA

levels that likely contribute to a dynamic equilibrium between these two hormones. As bioactive GAs suppress the accumulation of DELLA proteins,<sup>12</sup> we can speculate that in various developmental contexts the CRE1-dependent CK suppression of GA content may increase the DELLA protein accumulation and therefore modulate the expression of target genes.<sup>13,14</sup>

## Acknowledgments

The FF laboratory acknowledges the CNRS, the “Agence Nationale de la Recherche” (ANR) Labex “Saclay Plant Science” (SPS), and the Lidex “Plant Phenotyping Pipeline” (3P) for their support. CFF was the recipient of a Paris-Sud University PhD fellowship. The EF laboratory acknowledges the Australian Research Council (ARC) for financial support, Assoc. Prof. Noel W. Davies and Assoc. Prof. John J. Ross for advice with hormone extraction and analysis and Shelley Urquhart for technical assistance.

## Funding

French National Research Agency (ANR) and Australian Research Council (ARC).

## References

- Weiss D, Ori N. Mechanisms of cross talk between gibberellin and other hormones. *Plant Physiol.* 2007;144:1240–1246. doi:10.1104/pp.107.100370.
- Greenboim-Wainberg Y, Maymon I, Borochoy R, Alvarez J, Olszewski N, Ori N, Eshed Y, Weiss D. Cross talk between gibberellin and cytokinin: The arabidopsis ga response inhibitor spindly plays a positive role in cytokinin signaling. *Plant Cell.* 2005;17:92–102. doi:10.1105/tpc.104.028472.
- Moubayidin L, Perilli S, Dello Ioio R, Di Mambro R, Costantino P, Sabatini S. Report the rate of cell differentiation controls the arabidopsis root meristem growth phase. *Curr Biol.* 2010;20:1138–1143. doi:10.1016/j.cub.2010.05.035.
- Fonouni-Farde C, Kisiala A, Brault M, Emery RJN, Diet A, Frugier F. DELLA1-mediated gibberellin signaling regulates cytokinin-dependent symbiotic nodulation. *Plant Physiol.* 2017;175:1–12. doi:10.1104/pp.17.00919.
- Jasinski S, Piazza P, Craft J, Hay A, Woolley L, Rieu I, Phillips A, Hedden P, Tsiantis M. KNOX action in Arabidopsis is mediated by coordinate regulation of cytokinin and gibberellin activities. *Curr Biol.* 2005;15:1560–1565. doi:10.1016/j.cub.2005.07.023.
- Brenner W. G, Romanov G. A, Köllmer I, Bürkle L, Schmölling T. Immediate-early and delayed cytokinin response genes of Arabidopsis thaliana identified by genome-wide expression profiling reveal novel cytokinin-sensitive processes and suggest cytokinin action through transcriptional cascades. *Plant J.* 2005;44:314–333. doi:10.1111/j.1365-313X.2005.02530.x.
- Ariel F, Brault-Hernandez M, Laffont C, Huault E, Brault M, Plet J, Moison M, Blanchet S, Ichanté JL, Chabaud M, Carrere S, Crespi M, Chan RL, Frugier F. Two direct targets of cytokinin signaling regulate symbiotic nodulation in *Medicago truncatula*. *Plant Cell.* 2012;24:3838–52. doi:10.1105/tpc.112.103267.
- Plet J, Wasson A, Ariel F, Le Signor C, Baker D, Mathesius U, Crespi M, Frugier F. MtCRE1-dependent cytokinin signaling integrates bacterial and plant cues to coordinate symbiotic nodule organogenesis in *Medicago truncatula*. *Plant J.* 2011;1:622–633. doi:10.1111/j.1365-313X.2010.04447.x.
- Hedden P, Kamiya Y. GIBBERELLIN BIOSYNTHESIS: Enzymes, Genes and Their Regulation. *Annu Rev of Plant Physiol and Plant Mol Biol.* 1997;48:431–460. doi:10.1146/annurev.arplant.48.1.431.
- Boden SA, Weiss D, Ross JJ, Davies NW, Trevaskis B, Chandler PM, Swain SM. EARLY FLOWERING3 Regulates Flowering in Spring Barley by Mediating Gibberellin Production and FLOWERING LOCUS T Expression. *Plant Cell.* 2014;26:1557–1569. doi:10.1105/tpc.114.123794.
- Ingram TJ, Reid JB, MacMillan J. The quantitative relationship between gibberellin A1 and internode growth in *Pisum sativum* L. *Planta.* 1986;168:414–420. doi:10.1007/BF00392370.
- Silverstone AL, Jung HS, Dill A, Kawaide H, Kamiya Y, Sun TP. Repressing a repressor: gibberellin-induced rapid reduction of the RGA protein in Arabidopsis. *Plant Cell.* 2001;13:1555–66. doi:10.1105/tpc.13.7.1555.
- Davière J-M, Achard P. A pivotal role of dellas in regulating multiple hormone signals. *Mol Plant.* 2016;9:10–20. doi:10.1016/j.molp.2015.09.011.
- Fonouni-Farde C, Diet A, Frugier F. Root development and endosymbioses: DELLAs lead the orchestra. *Trends Plant Sci.* 2016;21:898–900. doi:10.1016/j.tplants.2016.08.012.