



HAL
open science

Whole-genome sequencing confirms the coexistence of different colonizing Group B Streptococcus isolates underscored by CRISPR typing

Clémence Beauruelle, Maxime Branger, Thierry Cochard, Adeline Pastuszka, Franck Biet, Philippe Lanotte

► To cite this version:

Clémence Beauruelle, Maxime Branger, Thierry Cochard, Adeline Pastuszka, Franck Biet, et al.. Whole-genome sequencing confirms the coexistence of different colonizing Group B Streptococcus isolates underscored by CRISPR typing. *Microbiology Resource Announcements*, 2020, 9 (5), 3 p. 10.1128/MRA.01359-19 . hal-02624834

HAL Id: hal-02624834

<https://hal.inrae.fr/hal-02624834>

Submitted on 26 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Whole-Genome Sequencing Confirms the Coexistence of Different Colonizing Group B *Streptococcus* Isolates Underscored by CRISPR Typing

Clémence Beauvue^{a,b,c} Maxime Branger^b Thierry Cochard^b Adeline Pastuszka^{a,b,c} Franck Biet^b Philippe Lanotte^{a,b,c}

^aUniversité de Tours, INRAE, ISP, Tours, France

^bINRAE, Université de Tours, ISP, Nouzilly, France

^cCHRU de Tours, Service de Bactériologie-Virologie, Tours, France

Clémence Beauvue and Maxime Branger contributed equally to this work. Author order was determined both alphabetically and because the article completes previous work initiated by Clémence Beauvue.

ABSTRACT *Streptococcus agalactiae* is a major pathogen and is the leading cause of neonatal infections in industrialized countries. The diversity of strains isolated from two pregnant women was investigated. Here, we present the draft genome sequences of strains W8A2, W8A6, W10E2, and W10F3, obtained in order to ascertain their phylogenetic affiliation.

Streptococcus agalactiae, or group B *Streptococcus* (GBS), is a major pathogen in humans and is the leading cause of neonatal infections in industrialized countries (1).

In a recent study, we explored the diversity of GBS in the vaginal carriage population in pregnant women at the time of screening (i.e., at 35 to 37 weeks' gestation by vaginal swabbing according to French clinical guidelines) using a CRISPR typing approach (2). For two women (women 8 and 10), two types of isolates (W8A2 and W8A6 and W10E2 and W10F3, respectively) were identified with similarities among the CRISPR arrays between the two types of isolates.

GBS isolates grew in Todd-Hewitt (TH) broth (BD Biosciences). The genomic DNA was isolated using phenol-chloroform extraction and precipitated with 1 M sodium chloride and 2 volumes of ice-cold 100% ethanol. Whole-genome sequencing and library construction were performed by GenoScreen (Lille, France). Libraries from purified genomic DNA were prepared for Illumina sequencing with a Nextera XT sample prep kit (Illumina, San Diego, USA) according to the supplier's recommendations. Sequencing was performed using an Illumina HiSeq platform in paired ends of 250 bp. At least 2.8 million reads were obtained for each strain, for a mean coverage of more than 339× for each based on a 2.1-Mb genome size (Table 1).

A quality check of sequencing data was performed using FastQC v0.11.5 (3), and trimming of reads was done using Sickle v1.33 (4) before assembly using SPAdes v3.11.1 (5). The contigs were annotated using the NCBI Prokaryotic Genome Annotation Pipeline v4.8 (6). All software tools were run using default parameters.

All of the strains were assembled into fewer than 80 contigs (range, 50 to 69) with a total assembly size of 2.1 Mb. About 2,000 coding genes were annotated (range, 2,039 to 2,137). The metrics for each sample are shown in Table 1.

Whole-genome sequencing (WGS) analysis confirmed the presence and structure of CRISPR-Cas genes previously described for these isolates (2). Using PHASTER (7), available for free at <http://phaster.ca/>, remnant phages were discovered on the 4 genomes, and an intact prophage sequence (79% identity with *Streptococcus* pro-

Citation Beauvue C, Branger M, Cochard T, Pastuszka A, Biet F, Lanotte P. 2020. Whole-genome sequencing confirms the coexistence of different colonizing group B *Streptococcus* isolates underscored by CRISPR typing. *Microbiol Resour Announc* 9:e01359-19. <https://doi.org/10.1128/MRA.01359-19>.

Editor Steven R. Gill, University of Rochester School of Medicine and Dentistry

Copyright © 2020 Beauvue et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Philippe Lanotte, philippe.lanotte@univ-tours.fr.

Received 29 October 2019

Accepted 24 November 2019

Published 30 January 2020

TABLE 1 Sequencing, assembly, and annotation metrics for each strain

Strain	BioSample no.	GenBank accession no.	No. of reads	Avg coverage (×)	No. of contigs	N_{50} (kb)	Assembly length (bp)	G+C content (%)	No. of protein CDSs ^a
W8A2	SAMN11415932	SSWT00000000	2,851,690	339	50	139,262	2,078,349	35.26	2,059
W8A6	SAMN11415933	SSWU00000000	2,926,588	348	69	99,289	2,086,162	35.29	2,076
W10E2	SAMN11415934	SSWV00000000	2,851,102	339	55	155,056	2,158,522	35.38	2,137
W10F3	SAMN11415935	SSWW00000000	4,173,810	497	55	149,658	2,085,005	35.34	2,097

^aCDSs, coding sequences.

phage 315.2 [GenBank accession no. [NC_004585.1](#)] was found on W10F3 isolates (contig 4).

In GBS, hemolysin production is encoded by the *cyl* operon, containing 12 genes, including the *cylE* gene (8). Transcriptional regulation of the *cyl* operon is controlled mainly by the two-component system CovS/CovR (*csrS* and *csrR* genes) (9). The *cylE* gene was found strictly conserved in the four different isolates. Even though there was no difference in the *csrS* and *csrR* genes between both types of woman 8 isolates, two mutations were observed in *csrS* genes of isolate W10E2, and one was observed in the W10F3 isolate, which may explain the highly hemolytic activity of W10F3.

Regarding resistance to macrolides, in GBS, erythromycin resistance is commonly due to target modification by an rRNA methyltransferase enzyme encoded by the *ermB* gene (10). An *ermB* gene was found only in isolate W8A6, which presented a high level of resistance to both erythromycin and clindamycin.

One of the major outcomes of this study was the degree of relatedness between isolates originating from each woman. The tree based on single nucleotide polymorphism (SNP) analysis (Fig. 1) allowed the exclusion of a phylogenetic affiliation between isolates W10E2 and W10F3, despite them having the same serotype and a common terminal direct repeat and ancestral spacer. The phylogenetic tree with numbering above the branches indicating the similarity coefficient between the strains (Fig. 1) was constructed with concatenation of SNPs using the software BioNumerics v7.6.3 (Applied Maths) with the settings “SNP-based categorical” and “clustering method by UPGMA.” This result confirms the difference in sequence types found (sequence type 4 [ST4] and ST28). Isolates from type W8A2 and type W8A6 were genetically closely related. These results show the *in vivo* coexistence of carriage of 2 types of closely related strains with a probable common ancestor.




FIG 1 Phylogeny of the clinical isolates of *Streptococcus agalactiae* with reference genomes, resolved with SNPs. The unweighted pair group method with arithmetic mean (UPGMA) phylogenetic tree was inferred from the 19,460 SNPs detected in the whole-genome sequences of clinical isolates in comparison to the reference strain, *Streptococcus agalactiae* NEM316. The number of SNPs identified between the strains is indicated along the top.

Data availability. The draft genome sequences of strains W8A2, W8A6, W10E2, and W10F3 have been deposited in DDBJ/ENA/GenBank under the accession no. [SSWT00000000](#) (BioSample no. [SAMN11415932](#)), [SSWU00000000](#) (BioSample no. [SAMN11415933](#)), [SSWV00000000](#) (BioSample no. [SAMN11415934](#)), and [SSWW00000000](#) (BioSample no. [SAMN11415935](#)), respectively.

ACKNOWLEDGMENTS

We are very grateful to Vanessa Rong for DNA preparation and Aileen Bonade Bottino for technical assistance with the genomic analysis.

REFERENCES

1. Stoll BJ, Schuchat A. 1998. Maternal carriage of group B streptococci in developing countries. *Pediatr Infect Dis J* 17:499–503. <https://doi.org/10.1097/00006454-199806000-00013>.
2. Beauruelle C, Pastuszka A, Mereghetti L, Lanotte P, Beauruelle C, Pastuszka A, Mereghetti L, Lanotte P. 2018. Group B *Streptococcus* vaginal carriage in pregnant women as deciphered by clustered regularly interspaced short palindromic repeat analysis. *J Clin Microbiol* 56:e0149-17. <https://doi.org/10.1128/JCM.01949-17>.
3. Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data. <http://www.bioinformatics.babraham.ac.uk/projects/fastqc>.
4. Joshi NA, Fass JN. 2011. Sickle: a sliding-window, adaptive, quality-based trimming tool for FastQ files (version 1.33). <https://github.com/najoshi/sickle>.
5. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
6. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44: 6614–6624. <https://doi.org/10.1093/nar/gkw569>.
7. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. 2016. PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Res* 44:W16–W21. <https://doi.org/10.1093/nar/gkw387>.
8. Whidbey C, Harrell MI, Burnside K, Ngo L, Becraft AK, Iyer LM, Aravind L, Hitti J, Adams Waldorf KM, Rajagopal L. 2013. A hemolytic pigment of group B *Streptococcus* allows bacterial penetration of human placenta. *J Exp Med* 210:1265–1281. <https://doi.org/10.1084/jem.20122753>.
9. Lamy MC, Zouine M, Fert J, Vergassola M, Couve E, Pellegrini E, Glaser P, Kunst F, Msadek T, Trieu-Cuot P, Poyart C. 2004. CovS/CovR of group B *Streptococcus*: a two-component global regulatory system involved in virulence. *Mol Microbiol* 54:1250–1268. <https://doi.org/10.1111/j.1365-2958.2004.04365.x>.
10. Heelan JS, Hasenbein ME, McAdam AJ. 2004. Resistance of group B streptococcus to selected antibiotics, including erythromycin and clindamycin. *J Clin Microbiol* 42:1263–1264. <https://doi.org/10.1128/jcm.42.3.1263-1264.2004>.