



HAL
open science

Comparison of the in vitro anthelmintic effects of *Acacia nilotica* and *Acacia raddiana*.

Geneviève Zabré, Adama Kaboré, Balé Bayala, Luciana M Katiki, Livio Martins Costa-Júnior, Hamidou H Tamboura, Adrien M G Belem, Adibe L Abdalla, Vincent Niderkorn, Herve Hoste, et al.

► To cite this version:

Geneviève Zabré, Adama Kaboré, Balé Bayala, Luciana M Katiki, Livio Martins Costa-Júnior, et al.. Comparison of the in vitro anthelmintic effects of *Acacia nilotica* and *Acacia raddiana*.. *Parasite*, 2017, 24, pp.44. <10.1051/parasite/2017044>. <hal-02625581>

HAL Id: hal-02625581

<https://hal.inrae.fr/hal-02625581v1>

Submitted on 26 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License

Comparison of the *in vitro* anthelmintic effects of *Acacia nilotica* and *Acacia raddiana*

Geneviève Zabré^{a,b}, Adama Kaboré^{a,*}, Balé Bayala^b, Luciana M. Katiki^c, Lívio Martins Costa-Júnior^d, Hamidou H. Tamboura^a, Adrien M.G. Belem^e, Adibe L. Abdalla^f, Vincent Niderkorn^g, Hervé Hoste^{h,*}, and Helder Louvandini^f

^a Laboratoire de Biologie et Santé Animales-DPA/INERA, 04 BP 8645 Ouagadougou 04, Burkina Faso

^b Université de Ouagadougou / UFR-SVT, 03 BP 7021 Ouagadougou 03, Burkina Faso

^c Instituto de Zootecnia (SAA, APTA), Rua Heitor Penteado 56, Nova Odessa, SP, cep 13460-000, Brazil

^d Universidade Federal do Maranhão – UFMA, Campus do Bacanga CEP 65080-805 São Luís- MA, Brazil

^e Université Nazi Boni, Bobo-Dioulasso, 01 BP 3770 Ouagadougou 01, Burkina Faso

^f Universidade de São Paulo, Centro de Energia Nuclear na Agricultura, NAPTSA, CP 96, CEP 13.400-970, Piracicaba, SP, Brazil

^g UMR1213 Herbivores, INRA – Clermont-Ferrand, Vetagro Sup, 63122 Saint-Genès-Champanelle, France

^h UMR IHAP 1225 INRA/ENVIT, 23 Chemin des Capelles, 31076 Toulouse Cedex, France

Received 7 May 2017, Accepted 20 October 2017, Published online 27 November 2017

Abstract – Gastrointestinal nematodes are a major threat to small ruminant rearing in the Sahel area, where farmers traditionally use bioactive plants to control these worms, including *Acacia nilotica* and *Acacia raddiana*. The main aim of this study was to screen the potential anthelmintic properties of aqueous and acetone extracts of leaves of these two plants based on three *in vitro* assays: (1) the egg hatch inhibition assay (EHA); (2) the larvae exsheathment inhibition assay (LEIA) using *Haemonchus contortus* as a model; and (3) an adult mortality test (AMT) applied on *Caenorhabditis elegans*. For the EHA, only *A. raddiana* was effective with $IC_{50} = 1.58$ mg/mL for aqueous extract, and $IC_{50} = 0.58$ mg/mL for acetonic extract. For the LEIA, all extracts inhibited the exsheathment of larvae compared to the controls, and the aqueous extract of *A. nilotica* was more larvicidal with $IC_{50} = 0.195$ mg/mL. In general, all responses to the substances were dose-dependent and were significantly different from the control group ($p < 0.05$). For the AMT, the extracts of the two *Acacia* species were effective but *A. raddiana* showed greater efficacy with 100% mortality at 2.5 mg/mL and $LC_{50} = 0.84$ mg/mL (acetonic extract). The addition of polyvinyl polypyrrolidone (PVPP) to the extracts suggested that tannins were responsible for blocking egg eclosion and inducing adult mortality but were not responsible for exsheathment inhibition. These results suggest that the leaves of these *Acacia* species possess ovicidal and larvicidal activities *in vitro* against *H. contortus*, and adulticidal effects against *C. elegans*.

Keywords: Natural products, tannins, secondary plant metabolites, nematodes, anthelmintics

Résumé – Comparaison des effets anthelminthiques *in vitro* d'*Acacia nilotica* et *Acacia raddiana*.

Les nématodes gastro-intestinaux demeurent une menace majeure de l'élevage des petits ruminants en zone sahélienne, où les éleveurs utilisent traditionnellement des plantes bioactives pour maîtriser ces parasites, notamment *Acacia nilotica* et *Acacia raddiana*. L'objectif de ce travail était de vérifier l'efficacité anthelminthique d'extraits aqueux et acétoniques de feuilles de ces deux plantes sur la base de 3 tests *in vitro*: 1) l'inhibition de l'éclosion des œufs (EHA), 2) l'inhibition du dégagement des larves (LEA) appliquée au modèle *Haemonchus contortus* et 3) la mortalité des *Caenorhabditis elegans* adultes (AMT). Pour EHA, seul *A. raddiana* a été efficace avec $IC_{50} = 1,58$ mg/mL pour l'extrait aqueux et $IC_{50} = 0,58$ mg/mL pour l'extrait acétonique. Pour LEA, tous les extraits ont inhibé le dégagement des larves par rapport aux témoins et l'extrait aqueux d'*A. nilotica* a été plus larvicide avec un $IC_{50} = 0,195$ mg/mL. De façon générale, toutes les concentrations agissent de façon dose-dépendante et sont significativement différentes du groupe témoin ($P < 0,05$). Pour AMT, les extraits des deux *Acacia* ont été efficaces mais *A. raddiana* a été plus efficace avec 100% de mortalité à 2,5 mg/mL avec une $DL_{50} = 0,84$ mg/mL (extrait acétonique). L'addition de

*Corresponding author: ade1_bf@yahoo.fr

polyvinylpyrrolidone (PVPP) aux extraits a montré que les tannins seraient responsables de l'inhibition de l'éclosion des œufs et de la mortalité chez les vers adultes, mais ne seraient pas responsables de l'inhibition du dégagement des larves. Ces résultats suggèrent que les feuilles de ces deux *Acacia* possèdent des activités ovicides et larvicides *in vitro* contre *H. contortus* et des effets sur les *C. elegans* adultes.

Introduction

Small ruminant rearing contributes strongly to human livelihoods in developing countries, especially for rural farmers [65]. In Sahelian countries, such as Burkina Faso, Mali and Niger, livestock play an important role in the economy [5]. In Burkina Faso, the livestock sector alone contributes over USD 51.3 million annually to the national economy, of which 32% is from small ruminants [49]. Despite the importance of this sector, small ruminant rearing faces feeding and health constraints that limit animal production. Among the health constraints, gastrointestinal nematodes (GINs), and *Haemonchus contortus* in particular (a highly prevalent and pathogenic species of GIN [15,56]), lead to low performances and higher animal mortality [35,58,67]. Like in many developing countries, the majority of livestock owners in rural areas do not use synthetic anthelmintic (AH) drugs because they are not affordable [31,55]. Therefore, they usually rely on practices related to ethnoveterinary medicine to control health problems in livestock [26]. In the Sahel, many natural fodder plants such as different species of *Acacia*, are used to control disease in ruminants, especially GINs. The use of such bioactive plants has several advantages for local farmers: (a) they are locally available; (b) they cost less compared to synthetic AH drugs; and (c) they are well accepted and consumed by sheep and goats.

An ethnoveterinary survey performed in this area of the Sahel showed that among *Acacia* species targeted for ruminants, *Acacia nilotica* var *adansonii* (Guill. & Perr.) O. Ktze and *Acacia raddiana* (Savi) are commonly used because of their high nutritional value and their potential AH effects. [70]. All of the breeders investigated used *A. raddiana* essentially as feed for cattle. However, 57% and 43% of the breeders questioned the use of *A. nilotica* to treat and to feed the cattle, respectively. For example, decoctions prepared from the pods and dried leaves of *A. nilotica* are used to treat foot-and-mouth disease (98%), diarrhoea (73%), leg ulcers (51%), loss of appetite (36%) and gingivitis (32%) in small ruminants [70]. In Mauritania, the powdered seed of *A. nilotica* macerated in fresh water is used to treat diarrhoea. The fruit, leaves and bark extracts of *A. nilotica* have been examined in *in vitro* studies [11,12,34,45] and their effects confirmed in *in vivo* studies [11,12,46]. However, information on *A. raddiana* is sparse.

Both *A. nilotica* and *A. raddiana* species are assumed to contain tannins [61]. Many *in vitro* and *in vivo* studies have now provided consistent evidence to support the AH effect of feed containing tannins and other polyphenols against abomasal and intestinal parasitic nematodes [27,29]. These previous investigations have shown that some local plants that contain tannins can impair different key biological processes of the parasitic nematode life

cycle: (i) the establishment of the infective third-stage larvae [18,20,47]; (ii) the excretion of eggs by adult worms [41,47,66] and (iii) the development of nematode eggs into larvae [51], and can therefore be used as nutraceuticals [30].

The objectives of this study were therefore two-fold: (1) to evaluate, based on *in vitro* assays, the anthelmintic effects of both aqueous and acetonetic extracts of the two *Acacia* plants on two different development stages of *H. contortus* and on adult nematode mortality of members of *Caenorhabditis elegans*; (2) to confirm the possible role of tannins and related polyphenols on *in vitro* AH effects using the tannin inhibitor polyvinyl pyrrolidone (PVPP).

Materials and methods

Plant materials

Fresh leaves of *A. nilotica* and *A. raddiana* were collected in December 2014 at Dori (14° and 15° N; 0° and 3° W), located in north-eastern Burkina Faso. The climate of the region, classified as Sahelian, is marked by a long dry season from November to June and a short rainy season from July to September. The specimens were identified by reference to the herbarium of the French National Centre of Scientific and Technological Research in Ouagadougou, Burkina Faso. Harvested leaves were then cleaned with water and dried at room temperature for a week according to the procedure used by traditional healers in the country.

Dried leaves were first transformed into powders using a blade grinder (basic IKA A 11) and the water content was determined according to the Association of Analytical Communities (AOAC) [6]. Two grams of each powder were dried at 105 °C for 3 h and cooled in a desiccator for 30 min before being weighed. The percentage of water content was calculated according to the following formula: $\% \text{ water content} = ((\text{powders before drying (g)} - \text{powders after drying (g)}) / \text{powders before drying (g)}) \times 100$.

Preparation of extracts

Two extracts were prepared for each *Acacia* species: one aqueous and one water/acetonic extract. For aqueous extracts, 150 g of each powdered material were macerated with 750 mL of distilled water for 24 h. Macerated extracts were then filtered over cotton wool and concentrated under reduced pressure in a rotary evaporator at 40-50 °C before being stored at 4 °C. For water/acetonic extracts, 150 g of each powder were macerated with 750 mL of mixed acetone/water (70/30) (v/v) for 72 h. Macerated extracts were filtered over Whatman paper before being concentrated under reduced pressure

in a rotary evaporator at 40–50 °C and stored at 4 °C. Dry extracts were subjected to phytochemical screening in order to identify the main phytochemical groups.

Quantification of tannins and polyphenolic compounds

Total tannins (TT), total phenols (TP) and condensed tannins (CT) were determined according to Makkar [39] and Makkar et al. [38]. TP and TT were determined by adding 250 µL of Folin-Ciocalteu reagents (1N) and 1.25 mL sodium carbonate solution (20% Na₂CO₃) to an aliquot of the supernatant and then taking absorbance readings at 725 nm. To determine TT, a binding tannin agent, 100 mg of insoluble polyvinyl polypyrrolidone (PVPP), was added to the extract and the measurement was repeated. A calibration curve was prepared from aliquots of the solution of tannic acid. The difference between measurements of TP and TP + PVPP extract readings was an estimate of TT. The concentrations of TP and TT were calculated as tannic acid equivalents (eq) and expressed as g/kg DM. CT were expressed as leucocyanidin equivalents (% of DM) and were determined using 3 mL butanol-HCl reagent and 0.1 mL ferric reagent. The absorbance was read at 550 nm.

Materials for *in vitro* assays

The faecal matter of a donor goat experimentally infected with *H. contortus* was collected at the Department of Pathology at the Federal University of Maranhão (UFMA) (Brazil) to obtain the eggs and L₃ larvae of *H. contortus*. *C. elegans* adult nematodes were obtained by culture according to the method of Chitwood and Feldlauer [22]. All procedures were approved by the Ethics Committee for Animal Experimentation of the Federal University of Maranhão under number 23115018061.

In vitro assays

Three different tests were performed: the egg hatching assay (EHA), the L₃ larval exsheathment assay (LEIA) of *H. contortus*, and the *C. elegans* adult mortality assay (AMT).

Egg hatching assay

This assay was performed according to the method described by Coles et al. [23]. The fresh faecal matter was recovered, crushed and filtered four times on 1-mm, 105-µm, 55-µm and 25-µm sieves. Eggs collected from the 25-µm sieve were centrifuged at 450 g /20 °C for 10 min and the supernatant was removed and replaced by an NaCl solution (density: 1.2), mixed and centrifuged at 1012 g /20 °C for 5 min. The recovered supernatant was filtered with a 25-µm sieve, washed and centrifuged three times at 450 g /20 °C for 10 min. The quantity was adjusted to reach a concentration of 100 eggs/mL. Five increasing concentrations of solutions were prepared for each extract:

0.3, 0.6, 1.25, 2.5 and 5.0 mg/mL diluted in PBS (0.1 M phosphate, 0.05 M NaCl, pH 7.2) for the two aqueous extracts, and in 2% methanol for the acetonic extracts. To test the direct effect of the extracts on nematodes, 100 µL of each concentration + 100 µL of eggs were placed in each well (96-well plate). Four replicates were performed per concentration. The plates were covered with Parafilm and incubated at 27 °C for 48 h. After 48 h of incubation, hatching was stopped by adding lugol iodine solution, and the number of L₁ larvae and eggs per well was counted using a reverse microscope (magnification x 10). Thereafter, the percentage of hatched eggs was determined using the following ratio: number of L₁/(number of eggs + number of L₁).

Larval exsheathment inhibition assay

The assay on larval exsheathment inhibition was performed as previously described by Bahuaud et al. [13]. Larvae were obtained by larval culture from goat faecal matter incubated at room temperature for 14 days to obtain L₃ larvae at the concentration of 2000 larvae/mL. Five increasing concentrations were prepared for each of the two extracts: 2.4, 1.2, 0.6, 0.3 and 0.15 mg/mL, all diluted in phosphate-buffered saline (PBS) (0.1 M phosphate, 0.05 M NaCl, pH 7.2) for aqueous extracts and 1% methanol for acetonic extracts. PBS and 1% methanol were used as a negative control. Briefly, 1 mL of L₃ larvae solution was incubated with 1 mL of extract at 20 °C for 3 h, and then washed with PBS and centrifuged three times at 1380 g for 3 min. Then, 200 µL of the larvae were subjected to the artificial exsheathment process by adding 200 µL of a Milton solution (2% w/v sodium hypochlorite and 16.5% sodium chloride) diluted to 1/300 with PBS. Four replicates were performed per concentration. The kinetics of larval exsheathment in the different experimental treatments were then monitored at 0, 20, 40 and 60 min intervals by microscopic observations (40×).

Mortality assay of adult *C. elegans*

Adult *C. elegans* were isolated according to Katiki et al. [36]. The test was performed with young adults and adults with intact cuticles. Five concentrations were prepared for the assay: 0.6, 1.25, 2.5, 5 and 10 mg/mL and two negative controls: salt solution M-9 (1.5 g KH₂PO₄; 3 g Na₂HPO₄; 2.5 g NaCl; 0.5 mL 1M MgSO₄, final volume of 500 mL) for aqueous extracts, and 2% dimethyl sulfoxide (DMSO) for acetonic extracts used as a solvent. The test was performed using 48-well plates and a nematode stock of 50 larvae/100 µL. In each well, nematodes were placed in contact with each concentration to a final volume of 500 µL/well with four replicates. The plates were then incubated at 25 °C for 20 h. After incubation, plates were read using a microscope and all adult worms were counted and determined as dead or alive, according to Skantar et al. [63]. They were considered dead when they did not show any movement and as alive when there were at least

some tails, head or pharyngeal movements (during 10 s of observation). The negative control consistently showed 95-100% live adults at 20 h after incubation.

Verification of the role of tannins by use of PVPP

In order to confirm the role of tannins on the AH effect, another series of EHA, LEIA and AMT was performed with both extracts of each *Acacia* species. However, for EHA, only *A. raddiana* extracts were used for this test because *A. nilotica* was not effective. Thus, the best concentrations of each extract (5 mg/mL + 100 mg of PVPP for EHA, 10 mg/mL + 200 mg of PVPP for AMT and 1.2 mg/mL + 250 mg of PVPP for LEIA) were pre-incubated with PVPP for 2 h, centrifuged at 1012 g / 10 min at 20 °C before being used for the assay. After centrifugation, all supernatants were removed to perform the three assays according to the procedure previously described and the results compared to an assay without the addition of PVPP. The ratio of pre-incubated PVPP was 1/10 for EHA and AMT. The rate for LEIA was 1/40.

Statistical analyses

Data from parasite tests were recorded on Excel 2010 (Microsoft corporation) and transformed to log ($x + 1$) before being subjected to a variance analysis with SAS software 2010, Version 6.20.4. IC_{50} and LC_{50} were calculated by probit analysis. IC_{50} and LC_{50} were considered significantly different when the 95% LC fails to overlap [60]. Comparison of averages (expressed as mean \pm standard error of mean) was performed using the Kruskal-Wallis test at 0.05.

Results

Quantification of tannins and polyphenolic compounds

Table 1 presents the content of total phenol (TP), condensed tannins (CT) and total tannins (TT) for both *Acacia* species. *A. nilotica* was rich in CT compared to *A. raddiana* which was rich in TT and TP.

Egg hatch inhibition assay

Tables 2 and 4 show the percentage of inhibition of egg hatching at different concentrations and 50% inhibition concentration for *H. contortus*. In general, the two controls recorded the lowest percentages of inhibition (4.3 and 9.8%, respectively, for PBS and 2% methanol) and were significantly different for all extracts ($p < 0.05$). The inhibition of egg hatching increased with increasing concentrations. However, *A. nilotica* was not effective and recorded a percentage of inhibition of less than 22% and an $IC_{50} > 5$ mg/mL. The percentage of egg inhibition of *A. raddiana* ranged from 12.67% to 92.76% with an $IC_{50} = 1.58$ mg/mL for the aqueous extract, and from 27.34% to 91.45% for acetonic extract with an $IC_{50} = 0.58$ mg/mL.

Table 1. Quantity of condensed tannin (CT), total tannin (TT) and total polyphenols (TP) in *Acacia* species.

Samples	CT	TT	TP
<i>A. nilotica</i>	53.60	54.23	76.66
<i>A. raddiana</i>	3.41	391.56	401.28

CT: g of leucocyanidin/kg dried mass (DM) TT and TP: g of tannic acid/kg DM.

Larval exsheathment inhibition essay

Both extracts revealed inhibition activity on L₃ larvae exsheathment of *H. contortus*. The aqueous extract of *A. nilotica* showed the lowest concentration of inhibition (IC_{50}) compared to all of the extracts (Table 4).

Mortality of adult *Caenorhabditis elegans*

Tables 3 and 4 present the efficacy of the extracts on adult *C. elegans* mortality and the lethal concentration. In general, the two controls recorded the lowest mortality rate (4.69% and 0.84%, respectively, for M-9 and 2% DMSO). The differences between concentrations and controls were highly significant ($p < 0.001$). Moreover, both extracts induced adult *C. elegans* mortality. When the concentration increased, the percentage of adult mortality increased only for the acetonic extract of *A. raddiana*, which reached 100% mortality at 2.5 mg/mL with the lowest lethal concentrations: $LC_{50} = 0.84$ mg/mL compared to the other extracts.

Using the inhibitor PVPP to assess the role of tannins and polyphenols

The addition of PVPP to the two extracts of each species significantly affected ($p < 0.05$) egg hatching of *H. contortus* (Table 5) and adult *C. elegans* mortality (Table 6). In contrast, for larval exsheathment, the results seem to indicate that incubation of extracts with PVPP did not change the exsheathment process when compared to non-treated extracts ($p > 0.05$) (Fig. 1).

Discussion

The use of tannin-rich plants as an alternative treatment to chemical anthelmintics is one approach that could reduce the development of parasite resistance [32,34,57]. The objective of this study was to evaluate and compare the *in vitro* ovicidal and larvicidal efficacy of aqueous and acetonic leaf extracts of *A. nilotica* and *A. raddiana* against *H. contortus*, because these two *Acacia* species are widely used by Sahelian breeders in ethno-veterinary medicine. According to the literature, many species of *Acacia* have been reported to have anthelmintic activities: the leaves of *Acacia cyanophylla* [1], *Acacia karoo* [34], *Acacia nilotica* [34,45], *Acacia pennatula* [2,3] and *Acacia polyacantha* [42], and the bark of *Acacia mangium* [54] and *Acacia mearnsii* [69].

Table 2. Percentages of egg hatch of *H. contortus* after 48 h incubation with *Acacia nilotica* and/or *Acacia raddiana* extract at different concentrations.

Concentration (mg/mL)	<i>A. nilotica</i>		<i>A. raddiana</i>	
	Aqueous extract	Acetonic extract	Aqueous extract	Acetonic extract
5	21.14 ± 2.34 a	18.72 ± 3.88 a	92.76 ± 1.92 a	91.45 ± 1.43 a
2.5	15.56 ± 3.34 b	17.23 ± 1.61 a	85.78 ± 2.09 a	88.81 ± 2.53 a
1.25	7.51 ± 1.71 c	10.95 ± 1.24 ab	40.83 ± 2.75 bA	79.88 ± 3.02 abB
0.625	6.48 ± 1.97 c	10.33 ± 1.63 ab	14.47 ± 1.52 cA	51.67 ± 2.83 cB
0.3125	3.42 ± 1.94 dA	9.85 ± 3.42 abB	12.67 ± 3.83 cdA	27.34 ± 2.18 dB
PBS	4.29 ± 2.01 d	-	4.29 ± 2.01 e	-
Methanol 2%	-	9.81 ± 9.98 ab	-	9.81 ± 9.94 e
P	0.0024	0.0067	0.0016	0.0013

- : no administered

(a,b,c,d,e) are compared means within the columns and (A,B) indicate differences within the lines (different extracts) for the two *Acacia* species for each plant. Different letters indicate significant differences ($p < 0.05$)

Table 3. Percentage of adult *Caenorhabditis elegans* mortality after 20 h of incubation with *Acacia nilotica* and *Acacia raddiana* extract at different concentrations.

Concentration (mg/mL)	<i>A. nilotica</i>		<i>A. raddiana</i>	
	Aqueous extract	Acetonic extract	Aqueous extract	Acetonic extract
10	94.97 ± 3.9aA	85.89 ± 3.0 aB	100 ± 0aA	100 ± 0aA
5	87.73 ± 2.9 bA	37.52 ± 2.5 bB	96.75 ± 1.6 aA	100 ± 0aA
2.5	81.78 ± 5.2 bA	19.04 ± 2.2 cB	80.32 ± 8.4 bA	100 ± 0aB
1.25	51.42 ± 2.8cA	3.31 ± 1.2 dB	11.25 ± 1.9 cA	81.08 ± 2.3bB
0.625	34.38 ± 5.5 dA	1.19 ± 2.1 dB	11.16 ± 4.1cA	22.53 ± 1.3cB
M9	4.69 ± 1.8 e	-	4.69 ± 1.8d	-
DMSO 2%	-	0.84 ± 0.8 d	-	0.84 ± 0.8 d
p	< 0.001	< 0.001	< 0.001	< 0.001

- : not administered.

(a,b,c,d,e) are compared means in the columns and (AB) in the lines. Different letters indicate significantly values ($p < 0.05$).

Table 4. Inhibitory concentrations (mg/mL) in the EHA and LEIA (mg/mL) with *Haemonchus contortus* (IC₅₀) and lethal concentrations (mg/mL) for the AMT with *Caenorhabditis elegans* (LC₅₀) with respective 95% confidence intervals for the different plant extracts.

Plant	Extract	Assay		
		EHA	LEIA	AMT
<i>A. nilotica</i>	Aqueous	> 5	0.195 (0.180 – 0.211)a	1.01 (0.78 - 1.23)a,b
	Acetonic	> 5	0.224 (0.226 -0. 265)b,c	5.39 (4.24 - 7.42)c
<i>A. raddiana</i>	Aqueous	1.36 (1.16-1.59)b	0.331 (0.252 – 0.408)c	1.43 (1.07 - 1.84)b
	Acetonic	0.68 (0.51-0.87)a	0.207 (0.173 – 0.246)a,b	0.84 (0.79 - 0.89)a

(a,b,c) are compared between lines (different extracts) for the two *Acacia* species for each plant. Different letters indicate significant differences.

In our studies, three *in vitro* assays were performed with two extracts (aqueous and acetonic extracts) at different concentrations. These types of extracts have been commonly used in many *in vitro* tests. Moreover, solvents and protocols used for extraction caused variations in concentrations and the classes of metabolites present in extracts [40].

The quantification of tannins revealed that *A. nilotica* was 18 times richer in CT than *A. raddiana*. However, *A. raddiana* was seven times richer in TT and five times richer in TP than *A. nilotica*. According to Hoste et al. [27], tannin-rich plants could interact with the proteins of the cuticle, oral cavity, oesophagus, cloaca and vulva of nematodes, modifying their chemical and physical prop-

Table 5. Inhibition of egg hatching after 48 h incubation with extracts treated or not treated with polyvinyl polypyrrolidone (PVPP) for *Acacia raddiana* aqueous and acetic extracts at a concentration of 5 mg/mL.

Extracts	Control	without PVPP (%)	with PVPP (%)
aqueous		89.2 ± 6.5 ^a	2.4 ± 2.9 ^b
acetic	6.2 ± 1.7	98.6 ± 2.7 ^a	3.1 ± 3.4 ^b

(a,b): significant difference ($p < 0.001$) between columns.

Table 6. Adult *Caenorhabditis elegans* mortality percentage, after incubation for 20 h with *Acacia raddiana* and *Acacia nilotica* aqueous and acetic extracts previously treated or not with polyvinyl polypyrrolidone (PVPP).

Extract [] 10 mg/mL	Control	Without PVPP (%)	with PVPP (%)
<i>A. raddiana</i> aqueous	0.69 ^a	100.00 ^b	14.91 ^c
<i>A. raddiana</i> acetic	0.96 ^a	100.00 ^b	13.26 ^c
<i>A. nilotica</i> aqueous	0.69 ^a	94.97 ^b	0.00 ^{ac}
<i>A. nilotica</i> acetic	0.96 ^a	85.89 ^b	4.31 ^{ac}

(a,b,c) = significant difference ($p < 0.05$) between values in rows.

erties. The anthelmintic effects of tannins may be attributed to their capacity to bind free protein available in the tubes for larval nutrition, and this reduced nutrient availability could therefore have resulted in larval starvation or decrease in gastrointestinal metabolism directly through the inhibition of oxidative phosphorylation, causing larval death [7].

Our studies conducted *in vitro* with *A. nilotica* and *A. raddiana* extracts showed the inhibition action on egg hatching and larval exsheathment of *H. contortus* and mortality of *C. elegans* adults. Extracts acted in a dose-dependent manner and their efficacy differed depending on the organ. This anthelmintic efficacy may be attributed to an individual or a combined effect of the bioactive compounds.

In the EHA, *A. nilotica* extracts were not effective compared to *A. raddiana*. *A. nilotica* inhibited 22% of egg hatching for the most effective concentration (5 mg/mL) with $IC_{50} > 5$ mg/mL. However, the two extracts of *A. raddiana* presented a high ovicidal activity (more than 90% at 5 mg/mL) with different concentrations of inhibition: $IC_{50} = 0.68$ mg/mL for acetic extract and $IC_{50} = 1.36$ mg/mL for aqueous extract. Acetic extracts were more ovicidal than aqueous extracts, but no significant difference was recorded between the extracts in our study. Thus, the results obtained for *A. nilotica* contrasted with those obtained by Eguale et al. [24] who reported significant egg hatching inhibition for aqueous extracts for the same plant. Likewise, Badar et al. [12] found that crude aqueous methanol extracts of *A. nilotica* bark ($LC_{50} = 0.201$ mg/mL) had higher inhibitory effects compared with leaves ($LC_{50} = 0.769$ mg/mL). Other results have confirmed the anthelmintic effect of different *Acacia* species on EHA. Thus, Oliveira [54] reported

$IC_{50} = 1.35$ mg/mL and 4.66 mg/mL with *A. mangium* extract on two *H. contortus* (White River: a strain resistant to ivermectin, benzimidazole and closantel; and the Juan strain: susceptible to all synthetic anthelmintics, respectively). Secondary plant metabolites (1) might bind the lipoproteins responsible for eggshell membrane permeability [59], and (2) tannins might inactivate enzymes responsible for the hatching process [43].

In the LEIA, both *Acacia* extracts were effective against larval exsheathment. *A. nilotica* revealed the lowest concentration of inhibition ($IC_{50} = 0.195$ mg/mL) for the aqueous extract compared to the aqueous extract of *A. raddiana* ($IC_{50} = 0.331$ mg/mL). However, the acetic extract of *A. nilotica* was numerically higher compared to *A. raddiana*. Moreover, all extracts at 600 and 1200 μ g/mL blocked the exsheathment process. Previous *in vitro* results indicated that extracts of various tannin-rich woody plants inhibited the exsheathment of nematode L₃ [13]. Alonso-Diaz et al. [2,3] tested *Acacia pennatula* and *A. gaureri* extracts and found an inhibition of 51% at 1200 μ g/mL, 93.5% at 600 μ g/mL, and a total exsheathment process after 70 min at 1.2 mg/mL, respectively. The exsheathment process in trichostrongyle nematodes is a crucial step that represents the transition from the free-living to the parasitic stages.

In the AMT, the free-living soil nematode *C. elegans* was used to test the efficacy of the extracts. *C. elegans* is a system to screen products for their potential anthelmintic effect against small ruminant gastrointestinal nematodes, including *H. contortus* [36]. The two *Acacia* extracts tested were effective against adult *C. elegans* and their lethal concentrations were less than $LC_{50} < 1.5$ mg/mL, except for the acetic extract of *A. nilotica* ($LC_{50} = 5.4$ mg/mL). For the same concentration (2.5 mg/mL), the acetic extract of *A. raddiana* revealed 100% adult mortality compared to 19% for *A. nilotica*. However, their two aqueous extracts revealed the same percentage of mortality (80.32 and 81.78%, respectively). In general, *A. raddiana* was more larvicidal than *A. nilotica*. *A. raddiana* contained high total tannins (TT). Plant extracts containing hydrolysable tannins (HT) such as gallic and/or ellagic acid or containing prodelphinidin CT (with gallic acid units) had higher levels of anthelmintic activity *in vitro* than the proanthocyanidin CT, which lack gallic acid units [17]. According to Katiki et al. [37] plants containing both CT and HT of the gallotannin and ellagitannin types were more lethal to *C. elegans* than plants containing CT lacking gallic acid units.

The results with PVPP for all assays showed that both extracts of *Acacia* species may have variability in the roles of tannins for EHA, LEIA and AMT. For our results, the addition of PVPP in *Acacia* extracts was associated with a significant decrease (restoration of control values) of EHA and AMT, but, did not seem to influence LEIA.

In the AMT, tannins and the associated metabolites seemed responsible for the mortality of adult *C. elegans* for *A. raddiana* extracts compared to *A. nilotica* extracts. Thus, tannins would be responsible for the efficacy of the extracts for the inhibition of egg hatching, but were not the




Figure 1. Larval exsheathment of *H. contortus* in the presence of *acacia* extracts at a concentration of 1.2 mg/mL, and its combination with polyvinyl pyrrolidone (PVPP).

only metabolites involved in adult mortality for *A. raddiana* extracts. These results suggest that a possible synergistic relationship of tannins and other compounds may be involved in killing the worms. For tannins, including condensed tannins, their anthelmintic activities have been reported by Athanasiadou et al. [8], Paolini et al. [57], Min et al. [48], Hoste et al. [27] and Gertrude et al. [25]. Tannins would appear to be able to bind to proteins and glycoproteins of the adult cuticle (a structure rich in proline and hydroxyproline), to the enzymes secreted by worms and involved in various essential functions [27], or to interact with their digestive epithelium to inhibit nutrition and cause the death of the parasite. Similar deductions were made with aqueous and ethanolic extracts of *Moringa oleifera* [25], aqueous extracts of *Anogeissus leiocarpus* and *Daniellia oliveri* [32], *Leucaena leucocephala* and *Gliricidia sepium* [33]. In our study, the inefficacy of PVPP to restore control values of larval exsheathment suggested that tannins were not the metabolites responsible for anthelmintic activity on *H. contortus*. According to Chan-Pérez et al. [21] and Vargas-Magana [68], tannins were not the sole secondary plant metabolites responsible for the AH effects. Our results contrasted with many previous studies that suspected tannins of being responsive to inhibit exsheathment [2,43,44,52]. Thus, Brunet et al. [18] and Hoste et al. [28] observed a structural lesion on *H. contortus* L₃ larvae (*in vitro* and *in vivo*) in contact with sainfoin extracts, a tannin-rich plant. Oliveira et al. [53], showed the inhibition of larval exsheathment in

H. contortus subject to *A. mangium* extract. Son-de Fernex et al. [64] observed an inhibition effect on exsheathment of *H. contortus* larvae subjected to five plants containing CTs. Consequently, we think that other metabolites besides tannins would be able to affect the exsheathment kinetics of L₃ larvae by binding to proteins and glycoproteins of the sheath to prevent the formation of the indented ring in the anterior part.

The work of Barrau et al. [14] and Ayers et al. [9] showed that flavonoids could play an essential role in the anthelmintic activity of some plants. Thus, the sheath of infective larvae of parasitic nematodes would be the target of flavonoids [10,19] as well as polyphenols [4]. Bizimeyera et al. [16] showed that polyphenols could have anthelmintic activities on nematodes. These chemical groups may be able to inhibit the secretion of proteases and acetyl-cholinesterase fluid by the glandular cells of the larvae to thereby prevent digestion and separation of the cap from the rest of the sheath.

However, other bioactive compounds and secondary plant metabolites could interact with multiple molecular targets on the various developmental stages of the parasite. Nandi et al. [50] showed that saponins have the ability to generate ions and cause lipid peroxidation of egg membranes, as well as to cause damage at the larval cuticle to inhibit parasite development. Other authors suggest that the conjugated unsaturated system of the saponins is involved in producing their damaging effect, probably resulting in free radicals, which induce membrane damage through peroxidation in nematodes [62].

A. nilotica and *A. raddiana* extracts both have *in vitro* anthelmintic activities against *H. contortus* and *C. elegans*. The two species of *Acacia* had completely different tannin contents. *A. nilotica*, rich in CT, was (1) not effective in inhibiting egg hatching; (2) partially effective for adult mortality; and (3) effective in inhibiting larval exsheathment. However, *A. raddiana*, rich in TT and TP, was highly effective in the three assays. Thus, in our study, TT-rich plants showed higher and more diverse anthelmintic activities against *H. contortus* compared to CT-rich plants. HT and CT are both found in TT. It is therefore possible that HT are responsible for the efficacy of *A. raddiana* extracts. Moreover, HT are known to show some toxicity for the ruminants, but the leaves of this plant were consumed by small ruminants in the Sahelian region of Burkina Faso. It is possible that the types and the quantity of HT contained in the leaves were not sufficient to cause ruminal toxicity but could affect the parasites. Future studies are required to ascertain which types or HT (gallotannins or ellagitannins) are contained in *A. raddiana* extract and underlie the anthelmintic activities. In general, the two species of *Acacia* used in our study were effective against *H. contortus*. Therefore, the association of these *Acacia* in the treatment of gastrointestinal parasites in a traditional environment might be a valuable approach. However, it would be necessary to conduct *in vivo* parasitological studies to consider the metabolism of extracts in the digestive tract of ruminants.

Acknowledgements. The authors thank the Tripartite Research Project (Africa – Brazil – France), “Fight against desertification in Africa”, funded by INRA/IRD-AIRD and CNPq 457559/2012-8, for supporting the study, as well as the translator of the manuscript.

Conflict of interest

The authors declare that they have no competing interests.

References

- Akkari H, Ben Salem H, Gharbi M, Abidi S, Darghouth MA. 2008a. Feeding *Acacia cyanophylla* Lindl. foliage to Barbarine lambs with or without PEG: Effect on the excretion of gastro-intestinal nematode eggs. *Animal Feed Science and Technology*, 147, 182-192.
- Alonso-Díaz MA, Torres-Acosta JFJ, Sandoval-Castro CA, Aguilar-Caballero AJ, Hoste H. 2008a. *In vitro* larval migration and kinetics of exsheathment of *Haemonchus contortus* exposed to four tropical tanniniferous plants extracts. *Veterinary Parasitology*, 153, 313-319.
- Alonso-Díaz MA, Torres-Acosta JFJ, Sandoval-Castro CA, Capetillo-Leal C, Brunet S, Hoste H. 2008b. Effects of four tropical tanniniferous plants on the inhibition of larval migration and the exsheathment process of *Trichostrongylus colubriformis* infective stage. *Veterinary Parasitology*, 153, 187-192.
- Alonso-Díaz MA, Torres-Acosta JFJ, Sandoval-Castro CA, Hoste H, Aguilar-Caballero AJ, Capetillo-Leal CM. 2008c. Is goats' preference of forage trees affected by their tannin or fiber content when offered in cafeteria experiments. *Animal Feed Science and Technology*, 141, 36-48.
- Association pour la promotion de l'élevage en Savane et au Sahel (APSS). 2015. Le paradoxe de l'élevage au Sahel: forts enjeux, faibles soutiens. Inter-Réseaux Développement rural / Bulletin de synthèse souveraineté alimentaire, n° 16.
- Association of Official Analytical Chemists (AOAC). 1990. *Official Methods of Analysis*, vol. II, 15th ed. Arlington, VA, USA.
- Athanasiadou S, Kyriazakis I, Jackson F, Coop RL. 2001. The effects of condensed tannins supplementation of foods with different protein content on parasitism, food intake and performance of sheep infected with *Trichostrongylus colubriformis*. *British Journal of Nutrition*, 86, 697-706.
- Athanasiadou S, Kyriazakis I, Jackson F, Coop RL. 2001. Direct anthelmintic effects of condensed tannins towards different gastrointestinal species: *in vitro* and *in vivo* studies. *Veterinary Parasitology*, 99, 205-219.
- Ayers S, Zink DL, Mohn K, Powell JS, Brown CM, Murphy T, Brand R, Pretorius S, Stevenson D, Thompson D, Singh SB. 2008. Flavones from *Struthiola argentea* with anthelmintic activity *in vitro*. *Phytochemistry*, 69, 541-545.
- Azando EVB, Hounzangbé-Adoté MS, Olounladé PA, Brunet S, Fabre N, Valentin A, Hoste H. 2011b. Involvement of tannins and flavonoids in the *in vitro* effects of *Newbouldia laevis* and *Zanthoxylum zanthoxyloides* extracts on the exsheathment of third-stage infective larvae of gastrointestinal nematodes. *Veterinary Parasitology*, 180, 292-297.
- Bachaya HA, Iqbal Z, Khan MN, Sindhu ZUD, Jabbar A. 2009. Anthelmintic activity of *Ziziphium nummularia* (bark) and *Acacia nilotica* (fruit) against *Trichostrongylid nematodes* of sheep. *Journal of Ethnopharmacology*, 123, 325-329.
- Badar N, Iqbal Z, Khan MN, Akhtar M.S. 2011. *In vitro* and *in vivo* anthelmintic activity of *Acacia nilotica* (L.) willd. ex delile bark and leaves. *Pakistan Veterinary Journal*, 31, 185-191.
- Bahuaud D, Martinez-Ortiz MC, Chauveau S, Prevot F, Torres-Acosta F, Fouraste I, Hoste H. 2006. Effects of four tanniferous plant extracts on the *in vitro* exsheathment of third-stage larvae of parasitic nematodes. *Parasitology*, 132, 545-554.
- Barrau E, Fabre N, Fouraste I, Hoste H. 2005. Effect of bioactive compounds from sainfoin (*Onobrychis vicifolia* Scop.) on the *in vitro* larval migration of *Haemonchus contortus*: role of tannins and flavonol glycosides. *Parasitology*, 131, 531-538.
- Belem AMG, Kaboré A, Bessin R. 2005. Gastrointestinal helminths of sheep in the central, eastern and northern parts of Burkina Faso. *Bulletin of Animal Health and Production in Africa*, 53, 13-23.
- Bizimenyera ES, Swan GE, Chikoto H, Eloff JN. 2005. Rationale for using *Peltophorum africanum* (Fabaceae) extracts in veterinary medicine. *Journal of South African Veterinary Association*, 76, 54-58.
- Brunet S, Hoste H. 2006. Monomers of condensed tannins affect the larval exsheathment of parasitic nematodes of ruminants. *Journal of Agricultural and Food Chemistry*, 54, 7481-7487.
- Brunet S, Aufrère J, El-babli F, Fourasté I, Hoste H. 2007. The kinetics of exsheathment of infective nematode larvae is disturbed in the presence of a tannin-rich plant extract (sainfoin) both *in vitro* and *in vivo*. *Parasitology*, 134, 1253-1262.

19. Brunet S. 2008. Analyse des mécanismes d'action antiparasitaire de plantes riches en substances polyphénoliques sur les nématodes du tube digestifs des ruminants. Thèse de Doctorat, Université Paul Sabatier de Toulouse. 246 pp.
20. Brunet S, Martínez-Ortiz de Montellano C, Torres-Acosta JFJ, Sandoval-Castro CA, Aguilar-Caballero AJ, Capetillo-Leal C, Hoste H. 2008. Effect of the consumption of *Lysiloma latisiliquum* on the larval establishment of gastrointestinal nematodes in goats. *Veterinary Parasitology*, 157, 81-88.
21. Chan-Pérez JI, Torres-Acosta JFJ, Sandoval-Castro CA, Hoste H, Castañeda-Ramírez GS, Vilarem G, Mathieu C. 2016. *In vitro* susceptibility of ten *Haemonchus contortus* isolates from different geographical origins towards acetone: water extracts of two tannin rich plants. *Veterinary Parasitology*, 217, 53-60.
22. Chitwood DJ, Feldlaufer MFE. 1990. Ecdysteroids in axenically propagated *Caenorhabditis elegans* and culture medium. *Journal of Nematology*, 22, 598-607.
23. Coles GC, Bauer C, Borgsteede FHM, Geerts S, Klei TR, Taylor MA, Waller PJ. 1992. World Association for the Advancement of Veterinary Parasitology (W.A.A.V.P.). Methods for the detection of anthelmintic resistance in nematodes of veterinary importance. *Veterinary Parasitology*, 44, 35-44.
24. Eguale T, Tadesse D, Giday M. 2011. *In vitro* anthelmintic activity of crude extracts of five medicinal plants against egg-hatching and larval development of *Haemonchus contortus*. *Journal of Ethnopharmacology*, 137, 108-113.
25. Gertrude MT, Josué WP, Marie-Claire K, Jeannette Y, Alidou MN, Mpoame M. 2014. Anthelmintic activity of *Moringa oleifera* leaf extracts evaluated *in vitro* on four developmental stages of *Haemonchus contortus* from Goats. *American Journal of Plant Sciences*, 5, 1702-1710.
26. Hilou A, Rappez F, Duez P. 2014. Ethnoveterinary management of cattle helminthiasis among the Fulani and the Mossi (Central Burkina Faso): plants used and modes of use. *International Journal of Biological and Chemical Sciences*, 8, 2207-2221.
27. Hoste H, Jackson F, Athanasiadou S, Thamsborg SM, Hoskin SO. 2006. The effects of tannin-rich plants on parasitic nematodes in ruminants. *Trends in Parasitology*, 22, 253-261.
28. Hoste H, Torres-Acosta JF, Alonso-Diaz MA, Brunet S, Sandoval-Castro C, Hounzangbé-Adoté S. 2008. Identification and validation of bioactive plants for the control of gastrointestinal nematodes in small ruminants. *Tropical Biomedicine*, 25, 56-72.
29. Hoste H, Martínez-Ortiz-De-Montellano C, Manolaraki F, Brunet S, Ojeda-Robertos N, Fourquaux I, Torres-Acosta JFJ, Sandoval-Castro CA. 2012. Direct and indirect effects of bioactive tannin-rich tropical and temperate legumes against nematode infections. *Veterinary Parasitology*, 186, 18-27.
30. Hoste H, Torres-Acosta JFJ, Sandoval-Castro CA, Mueller-Harvey I, Sotiraki S, Louvandini H, Thamsborg SM, Terrill TH. 2015. Tannin containing legumes as a model for nutraceuticals against digestive parasites in livestock. *Veterinary Parasitology*, 212, 5-17.
31. Kaboré A, Tamboura HH, Belem AMG, Traoré A. 2007. Traitements ethno-vétérinaires des parasitoses digestives des petits ruminants dans le plateau central du Burkina Faso. *International Journal of Biological and Chemical Sciences*, 1, 297-304.
32. Kaboré A, Belem AMG, Tamboura HH, Traoré A, Sawadogo L. 2009. *In vitro* anthelmintic effect of two medicinal plants (*Anogeissus leiocarpus* and *Daniellia oliveri*) on *Haemonchus contortus*, an abomasal nematode of sheep in Burkina Faso. *African Journal of Biotechnology*, 8, 4690-4695.
33. Kaboré A, Traoré A, Nignan M, Gnanda BI, Bamogo V, Tamboura HH, Bélem AMG. 2012. *In vitro* anthelmintic activity of *Leuceana leucocephala* (Lam.) De Wit. (Mimosaceae) and *Gliricidia sepium* (Jacq.) Kunth ex Steud (Fabaceae) leave extracts on *Haemonchus contortus* ova and larvae. *Journal of Chemical and Pharmaceutical Research*, 4, 303-309.
34. Kahiya C, Mukaratirwa S, Thamsborg SM. 2003. Effects of *Acacia nilotica* and *Acacia karoo* diets on *Haemonchus contortus* infection in goats. *Veterinary Parasitology*, 115, 265-274.
35. Kamaraj C, Abdul Rahuman A. 2011. Efficacy of anthelmintic properties of medicinal plant extracts against *Haemonchus contortus*. *Research in Veterinary Science*, 91, 400-404.
36. Katiki LM, Ferreirara JFS, Zajac AM, Masler C, Lindsay DS, Chagas ACS, Amarante AFT. 2011. *Caenorhabditis elegans* as a model to screen plant extracts and compounds as natural anthelmintics for veterinary use. *Veterinary Parasitology*, 182, 264-268.
37. Katiki LM, Ferreirara JFS, Gonzalez JM, Zajac AM, Lindsay DS, Chagas ACS, Amarante AFT. 2013. Anthelmintic effect of plant extracts containing condensed and hydrolyzable tannins on *Caenorhabditis elegans*, and their antioxidant capacity. *Veterinary Parasitology*, 192, 218-227.
38. Makkar HPS, Blummel M, Borowy NK, Becker K. 1993. Gravimetric determination of tannins and their correlations with chemical and protein precipitation methods. *Journal of the Science of Food and Agriculture*, 61, 161-165.
39. Makkar HP. 2003. Quantification of tannins in tree and shrub foliage: a laboratory manual. Animal Production and health Section. Joint FAO/IAEA Division of nuclear Technique in Food and Agriculture. International Atomic Energy Agency, Vienna, Austria, 49-53.
40. Marie-Magdeleine C, Hoste H, Mahieu M, Varo H, Archimede H. 2009. *In vitro* effects of *Cucurbita moschata* seed extracts on *Haemonchus contortus*. *Veterinary Parasitology*, 161, 99-105.
41. Martínez-Ortiz-De-Montellano C, Vargas-Magana JJ, Canul-Ku L, Miranda-Soberanis R, Capetillo-Leal C, Sandoval-Castro CA, Hoste H, Torres-Acosta JFJ. 2010. Effect of a tropical tannin-rich plant, *Lysiloma latisiliquum* on adult populations of *Haemonchus contortus* in sheep. *Veterinary Parasitology*, 172, 283-290.
42. Max RA, Kimambo AE, Kassuku AA, Mtenga LA, Buttery PJ. 2007. Effects of tanniferous browse meal on nematode faecal egg counts and internal parasite burdens in sheep and goats. *South African Journal of Animal Science*, 37, 97-106.
43. Molan AL, Waghorn GC, McNabb WC. 2002. Effect of condensed tannins on egg hatching and larval development of *Trichostrongylus colubriformis* *in vitro*. *Veterinary Record*, 150, 65-69.
44. Molan AL, Faraj AM. 2010. The effects of condensed tannins extracted from different plant species on egg hatching and larval development of *Teladorsagia circumcincta* (Nematoda: Trichostrongylidae). *Folia Parasitologica*, 57, 62-68.
45. Moreno FC, Gordon IJ, Wright AD, Benvenuti MA, Saumell CA. 2010. *In vitro* anthelmintic effect of plant extracts against infective larvae of ruminants gastrointestinal nematode parasites. *Archivos de Medicina Veterinaria*, 42, 155-163.

46. Moreno FC, Gordon IJ, Knox MR, Summer PM, Skerrat LF, Benvenuti MA, Saumell CA. 2012. Anthelmintic efficacy of five tropical native Australian plants against *Haemonchus contortus* and *Trichostrongylus colubriformis* in experimentally infected goats (*Capra hircus*). *Veterinary Parasitology*, 187, 237-243.
47. Moreno-Gonzalo J, Osoro K, García U, Frutos P, Celaya R, Ferreira LMM, Ortega-Mora LM, Ferre I. 2013. Effect of the consumption of heather on incoming larvae and established population of *Teladorsagia circumcincta* in experimentally infected Cashmere goats. *Veterinary Parasitology*, 196, 124-129.
48. Min BR, Barry TN, Attwood GT, McNabb WC. 2003. The effect of condensed tannins on the nutrition and health of ruminants fed fresh temperate forages: a review. *Animal Feed Science and Technology*, 106, 3-19.
49. Ministère des Ressources Animales (MRA), 2012. Statistique du secteur de l'élevage. Annuaire 2011. Ouagadougou. 151 p.
50. Nandi B, Roy S, Bhattacharya S, Babu SPB. 2004. Free radicals mediated membrane damage by the saponins *Acacia* side A and *Acacia* side B. *Phytotherapy Research*, 18, 191-194.
51. Niezen JH, Waghorn GC, Graham T, Carter JL, Leathwick DM. 2002. The effect of diet fed to lambs on subsequent development of *Trichostrongylus colubriformis* larvae *in vitro* and on pasture. *Veterinary Parasitology*, 105, 269-283.
52. Olivera LMB, Bevilaqua CML, Macedo ITF, Morais SM, Monteiro MVB, Campello CC, Ribeiro WLC, Batista EKF. 2011a. Effects of six tropical plant extracts on larval exsheathment of *Haemonchus contortus*. *Revista Brasileira de Parasitologia Veterinária, Jaboticabal*, 20, 155-160.
53. Olivera LMB, Bevilaqua CML, Macedo ITF, Morais SM, Machado LKA, Campello CC, Mesquita MA. 2011b. Effects of *Myracrodrum urundeuva* extracts on egg hatching and larval exsheathment of *Haemonchus contortus*. *Parasitology Research*, 109, 893-898.
54. Oliveira GC. 2014. Uso de *Clitoria fairchildiana* Ra Howard Fabaceae e *Acacia mangium* Willd Mimosaceae sobre trichostrongilídeos de caprinos. 84f. Dissertação (Mestrado em Ciências da Saúde) – Programa de Pós-Graduação em Ciências da Saúde – Universidade Federal do Maranhão, São Luís.
55. Olounladé AP. 2005. Effets anthelminthiques des feuilles de *Newbouldia laevis* testées *in vivo* sur les nématodes gastro-intestinaux (*Haemonchus contortus* et *Trichostrongylus colubriformis*) chez les moutons Djallonké. DEA, Université de Lomé, Togo, p. 65.
56. Ouattara L, Dorchies PH. 2001. Helminthes gastro-intestinaux des moutons et chèvres en zones sub-humide et sahélienne du Burkina Faso. *Revue de Médecine Vétérinaire*, 152, 165-170.
57. Paolini V, Bergeaud JP, Grisez C, Prevot F, Dorchies P, Hoste H. 2003. Effects of condensed tannins on goats experimentally infected with *Haemonchus contortus*. *Veterinary Parasitology*, 113, 253-261.
58. Periasamy K, Pichler R, Poli M, Cristel S, Cetrá B, Medus D, Basar M, Thiruvenkadan AK, Ramasamy S, Ellahi MB, Mohammed F, Teneva A, Shamsuddin M, Podesta MG, Diallo A. 2014. Candidate gene approach for parasite resistance in sheep variation in immune pathway genes and association with fecal egg count. *PLoS One* 9, e88337
59. Perry RN. 2002. Hatching. In: Donald L. Lee (Ed.), *The Biology of Nematodes*. Taylor and Francis, London and New York, p. 147-170.
60. Roditakis E, Roditakis NE, Tsagkarakou A. 2005. Insecticide resistance in *Bemisia tabaci* (Homoptera: Aleyrodidae) populations from Crete. *Pest Management Science*, 61, 577-582.
61. Rubanza CDK, Shem MN, Otsyina R, Bakengesa SS, Ichinohe T, Fujihara T. 2005. Polyphenolics and tannins effect on *in vitro* digestibility of selected *Acacia* species leaves. *Animal Feed Science and Technology*, 119, 129-142
62. Sinha Babu SP, Sarkar D, Ghosh NK, Saha A, Sukul NC, Bhattacharya SH. 1997. Enhancement of membrane damage by saponins isolated from *Acacia auriculiformis*. *Japanese Journal of Pharmacology*, 75, 451-454.
63. Skantar AM, Agama K, Meyer SLF, Carta LK, Vinyard BT. 2005. Effects of geldanamycin on hatching and juvenile motility in *Caenorhabditis elegans* and *Heterodera glycines*. *Journal of Chemical Ecology*, 31, 2481-2491.
64. Son-De Fernex EV, Alonso-Diaz MA, Valles-De La Mora B, Capetillo-Leal CM. 2012. *In vitro* anthelmintic of five tropical legumes on the exsheathment and motility of *Haemonchus contortus* infective larvae. *Experimental Parasitology*, 131, 413-418.
65. Sujon MA, Mostoja M, Jahan MS, Das AR, Rob S. 2008. Studies on medicinal plants against gastrointestinal nematodes of goats. *Bangladesh Journal of Veterinary Medicine*, 6, 179-183.
66. Terril TH, Dykes GS, Shaik SA, Miller JE, Kouakou B, Kannan G, Burke JM, Mosjidis JA. 2009. Efficacy of sericea lespedeza hay as a natural dewormer in goats: Dose titration study. *Veterinary Parasitology*, 163, 52-56.
67. Torres-Acosta JFJ, Hoste H. 2008. Alternative or improved methods to limit gastro-intestinal parasitism in grazing sheep and goats. *Small Ruminant Research*, 77, 159-173.
68. Vargas-Magana JJ, Torres-Acosta JFJ, Aguilar-Caballero AJ, Sandoval-Castro CA, Hoste H, Chan-Pérez JI. 2014. Anthelmintic activity of acetone-water extracts against *Haemonchus contortus* eggs: Interactions between tannins and other plant secondary compounds. *Veterinary Parasitology*, 206, 322-327.
69. Yoshihara E, Minho AP, Cardim ST, Tabacow VBD, Yamamura M.H. 2014. *In vitro* ovicidal and larvicidal activity of condensed tannins on gastrointestinal nematode infestations in sheep (*Ovis aries*). *Semina: Ciências Agrárias*, 35, 3173-3180.
70. Zabré G, Kaboré A, Bayala B, Tamboura HH, Belem AMG, Niderkorn V, Costa Junior ML, Louvandini H, Hoste H. 2017. Botanical and ethnoveterinary surveys of two acacias (*Acacia raddiana* and *Acacia nilotica*) exploited in small ruminant rearing in Sahelian area of Burkina Faso. *Animal and Veterinary Sciences*, 5, 63-68.

Cite this article as: Zabré G, Kaboré A, Bayala B, Katiki LM, Costa-Júnior LM, Tamboura HH, Belem AMG, Abdalla AL, Niderkorn V, Hoste H, Louvandini H. 2017. Comparison of the *in vitro* anthelmintic effects of *Acacia nilotica* and *Acacia raddiana*. *Parasite* 24, 44



An international open-access, peer-reviewed, online journal publishing high quality papers on all aspects of human and animal parasitology

Reviews, articles and short notes may be submitted. Fields include, but are not limited to: general, medical and veterinary parasitology; morphology, including ultrastructure; parasite systematics, including entomology, acarology, helminthology and protistology, and molecular analyses; molecular biology and biochemistry; immunology of parasitic diseases; host-parasite relationships; ecology and life history of parasites; epidemiology; therapeutics; new diagnostic tools.

All papers in Parasite are published in English. Manuscripts should have a broad interest and must not have been published or submitted elsewhere. No limit is imposed on the length of manuscripts.

Parasite (open-access) continues **Parasite** (print and online editions, 1994-2012) and **Annales de Parasitologie Humaine et Comparée** (1923-1993) and is the official journal of the Société Française de Parasitologie.

Editor-in-Chief:
Jean-Lou Justine, Paris

Submit your manuscript at
<https://parasite.edmgr.com/>