Tick-pathogen interactions: The metabolic perspective

Alejandro Cabezas Cruz, Pedro Espinosa, Pilar Alberdi, José de La Fuente

► To cite this version:

Alejandro Cabezas Cruz, Pedro Espinosa, Pilar Alberdi, José de La Fuente. Tick-pathogen interactions: The metabolic perspective. Trends in Parasitology, 2019, 35 (4), pp.316-328. 10.1016/j.pt.2019.01.006 . hal-02626522

HAL Id: hal-02626522

<https://hal.inrae.fr/hal-02626522>

Submitted on 22 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution - NonCommercial| 4.0 International License

1 **Tick-pathogen interactions: the metabolic perspective**

2 Alejandro Cabezas-Cruz^{1*}, Pedro Espinosa², Pilar Alberdi², José de la Fuente^{2,3*}

3

4 ¹ UMR BIPAR, INRA, Ecole Nationale Vétérinaire d'Alfort, ANSES, Université Paris-Est, Maisons-Alfort,
5 France.

6 ² SaBio, Instituto de Investigación en Recursos Cinegéticos IREC (CSIC-UCLM-JCCM), Ciudad Real,
7 Spain.

8 ³ Department of Veterinary Pathobiology, Center for Veterinary Health Sciences, Oklahoma State
9 University, Stillwater, OK, United States.

10

11 *Correspondence: jose_delafuente@yahoo.com (J. de la Fuente) and
12 cabezasalejandrocruz@gmail.com (A. Cabezas-Cruz).

13

14 **Abstract**

15 The first tick genome published in 2016 provided an invaluable tool to study the molecular
16 basis of tick-pathogen interactions. Metabolism is a key element in host-pathogen
17 interactions. However, our knowledge of tick-pathogen metabolic interactions is very
18 limited. Recently, a Systems Biology approach using omics datasets has revealed that tick-
19 borne pathogen infection induces transcriptional reprogramming affecting several metabolic
20 pathways in ticks to facilitate infection, multiplication and transmission. Results suggest that
21 the response of tick cells to tick-borne pathogens is associated with tolerance to infection.
22 Here we review our current understanding of tick metabolism modulation by tick-borne
23 pathogens with focus in the model intracellular bacterium *Anaplasma phagocytophilum*.

24

25 **Keywords:** *Ixodes*; *Anaplasma phagocytophilum*; *Borrelia* spp.; metabolism; Tick-pathogen
26 interactions

27

28 **Tick-pathogen metabolic interactions at the center of the infectious storm**

29 Metabolism is key to cellular function [1,2]. In consequence, all major molecular pathways in
30 living cells are interconnected to and regulated by the availability and levels of metabolites
31 [1,2]. Recent studies in mammalian cells showed that interactions between intracellular
32 bacterial pathogens and the host cells can lead to physiological changes in both interacting
33 members [3]. The metabolic adaptations in mammalian cell-pathogen systems promote
34 proliferation or elimination of the pathogen within the host cells [3]. Tick-pathogen
35 interactions, however, involve a more complex array of outcomes which include conflict and
36 cooperation that ultimately benefit both ticks and pathogens [4]. This is not surprising
37 considering that in vector-borne pathogen systems the survival of the vector is essential for
38 the completion of the life cycle of the pathogen. In addition, tick-borne pathogen infection
39 increases tick performance in challenging environmental conditions [5,6]. Therefore, in some
40 cases, vector tolerance to pathogen infection is an advantageous life trait in arthropod
41 vectors [7].

42 Tick metabolism modulation by tick-borne pathogens is a result of coevolution and adaption
43 to a considerable number of tick and reservoir host species [4,8,9]. The *Ixodes scapularis*
44 genome, the first for a medically important chelicerate, is the only tick genome available so
45 far [10]. The sequencing of the *I. scapularis* genome was an essential step towards the
46 understanding of the molecular processes that support the parasitic lifestyle of the tick and
47 its success as a vector of multiple pathogens including bacteria, viruses, protozoa, and
48 helminths which constitute a growing burden for human and animal health worldwide [11].
49 This review focuses on recent research that provided insights on the finely tuned metabolic
50 changes that the model pathogen *Anaplasma phagocytophilum* induces in its tick vector *I.*
51 *scapularis*. For comparative purposes, tick metabolic changes induced by other pathogens
52 including mainly *Borrelia* spp., flaviviruses and fungi will be also included (Table 1). Results
53 suggest that tick-borne pathogens use tick metabolism as a hub to modulate tick physiology
54 and tick cell processes including immunity and apoptosis among others.

55

56 **Nutritional immunity and iron metabolism in ticks**

57 Iron is essential for most organisms because it serves as an electron donor and acceptor in
58 various metabolic processes. Therefore, iron holds a central position in host-pathogen
59 interactions [12]. While host evolved mechanisms to deprive pathogens from iron,
60 pathogens coevolved mechanisms to acquire host iron. Host mechanisms involved in limiting
61 pathogens access to iron were termed 'nutritional immunity' [12]. During feeding, ticks are
62 exposed to the large amounts of iron present in the blood meal. Excess iron can react with
63 H₂O₂ which yields hydroxyl radical, a potent biological oxidant [12]. Therefore, a balanced
64 regulation of iron levels is essential for ticks [13], and potentially, pathogen survival. Two
65 ferritin (FER) proteins, FER1 and FER2, are the primary iron storage and transporter,
66 respectively, in ticks [13]. In addition of controlling the production of reactive oxygen species
67 from iron, like in other model organism [12, 13], ferritins may also contribute to nutritional
68 immunity in ticks by depriving pathogens from essential iron supply. Interestingly,
69 *Escherichia coli* infection induces the expression of a *fer* gene in *Dermacentor variabilis* and
70 silencing of *fer* genes in *Haemaphysalis longicornis* decreases tick survival following a
71 challenge with *E. coli* [13].

72 The tick-borne pathogen *Ehrlichia ruminantium* sense iron levels in mammalian cells and
73 respond to iron starvation by upregulating virulence genes such as those involved in Type IV
74 Secretion System (T4SS), which in turn may participate in iron acquisition by the pathogen
75 [14]. A similar mechanism may be at play during pathogen infection in ticks. An iron-binding
76 protein expressed in the infective stage of *A. phagocytophilum* [15], and hypothesized to be
77 secreted by the T4SS, was found in tick salivary glands and tick cells infected with these
78 bacteria [16]. Bacterial iron-binding proteins may sequester iron to overcome nutritional
79 immunity in ticks. An interesting mechanism to scape nutritional immunity is that of *Borrelia*
80 *burgdorferi* which requires none or very low amounts of iron to survive [17]. In addition,
81 recent research provides evidence that *A. phagocytophilum* modulates tick heme
82 metabolism. Host hemoglobin is cleavage to heme by three sequential set of reactions
83 involving several hemoglobinolytic enzymes in ticks [18]. Due to the high redox potential of
84 heme, this molecule is toxic and in excess can be bacteriostatic [19]. Quantitative proteomics
85 analysis revealed that *A. phagocytophilum* modulates the levels of hemoglobinolytic
86 enzymes which affects the levels of host hemoglobin in tick midguts and salivary glands [18].

87 This finding suggests that *A. phagocytophilum* infection may regulate the levels of heme in a
88 tissue-specific manner with potential effects for pathogen and vector survival.

89

90 **Tick hemocyte metabolism in response to tick-borne pathogen infection**

91 Despite mammalian macrophages possess specialized effector functions to eliminate
92 invading microorganism, many intracellular bacteria infect and replicate inside this immune
93 cells [20]. Globally, macrophages can be divided into inflammatory (M1) and anti-
94 inflammatory (M2) macrophages [20]. Both macrophage types M1 and M2 can be infected
95 by intracellular bacteria; however, the metabolic profiles of these macrophages in response
96 to pathogenic stimulus are substantially different [20]. After stimulation with
97 lipopolysaccharide (LPS), macrophages M1 uptake more glucose via **glucose transporters**
98 (GLUTs, see Glossary) and increase the glycolytic flux and the production of lactate, a process
99 named **aerobic glycolysis**. In contrast, macrophages M2 have the metabolic profile
100 comparable to that of unstimulated cells, but with higher TCA-cycle and oxidative activity
101 [20].

102 Cellular immunity in ticks is mostly mediated by hemocytes, which are involved in the
103 phagocytosis of different microbes [21]. After establishing a primary infection in tick
104 midguts, *A. phagocytophilum* migrates to the salivary glands [11]. A secreted tick protein,
105 P11, enables *A. phagocytophilum* to infect tick hemocytes, which are required for the
106 migration of *A. phagocytophilum* from the gut to the salivary glands [22]. Isolation and *in*
107 *vitro* culture of tick hemocytes has remained elusive. Therefore, the study of tick hemocytes
108 metabolism in response to *A. phagocytophilum* infection has been based on the *I. scapularis*
109 ISE6 cell line. Based on the transcriptional response to *A. phagocytophilum*, ISE6 cells were
110 proposed to constitute a model of tick hemocytes involved in pathogen infection and
111 immune response [23,24]. Comparison of the metabolic profiles of macrophages M1 with
112 that of *A. phagocytophilum*-infected ISE6 cells suggests that the metabolic profile of tick
113 hemocytes in response to pathogen infection is very different to that of inflammatory
114 macrophages M1 (Fig. 1, Key Figure). Two major differences can be summarized: (i) ISE6 cells
115 infected by *A. phagocytophilum* do not display the aerobic glycolysis observed in mammalian

macrophages M1 and (ii) **hypoxia-inducible factor 1-alpha (HIF-1 α)** activation and stabilization in ISE6 cells is not dependent of succinate accumulation and aerobic glycolysis (Fig. 1). This suggests that the metabolic reprogramming associated with macrophage M1 activation does not occurs in tick hemocytes infected with *A. phagocytophilum*. Interestingly, while 2-deoxyglucose (2DG), an inhibitor of **glycolysis**, inhibits HIF-1 α activation in macrophages M1 [25], this compound has no effect on HIF-1 α activation in ISE6 cells [26]. Instead of aerobic glycolysis and succinate accumulation, HIF-1 α activity in *A. phagocytophilum*-infected ISE6 cells appears to be mediated phosphoinositide 3-kinase (PI3K) pathway activation (Fig. 1). In agreement with this, the PI3K inhibitor LY294002 decreased HIF-1 α DNA binding activity in infected ISE6 cells [26]. A metabolic property shared by macrophages M1 and ISE6 cells is, however, an overall decrease in **tricarboxylic acid cycle (TCA cycle)** activity upon bacterial infection [25,26].

The absence of immunogenic molecules such as LPS and diaminopimelic-type peptidoglycans (DAP-PGN) in *A. phagocytophilum* and other bacteria transmitted by *I. scapularis* was previously suggested to contribute to vector tolerance to tick-borne pathogens [7]. The lack of these immune activators in tick-borne pathogens may help these microbes colonize ticks without being eliminated [7]. Interestingly, lipids 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphoglycerol (POPG) and 1-palmitoyl-2-oleoyl diacylglycerol (PODAG) found in *A. phagocytophilum* activate the immune deficiency (IMD) pathway in ticks [27]. Immune priming with POPG and PODAG reduces the *A. phagocytophilum* burden in infected ISE6 cells [27]. This shows that tolerance to infection can be associated with immunity and resistance to tick-borne pathogens in ticks. This is expected considering that unchecked bacterial multiplication within host cells could affect vector, and therefore, tick-borne pathogen survival. Remarkably, recent research shows that IMD is involved in glucose and lipid metabolism regulation in *Drosophila* [28] and constitutive activation of IMD induces a significant reduction in the expression of genes involved in glycolysis, **gluconeogenesis** and TCA cycle in flies [29]. Whether IMD activation by *A. phagocytophilum* regulates tick hemocyte metabolism remains to be tested.

Molecular mechanisms involved in tick metabolism modulation by tick-borne pathogens

146 Tick metabolism modulation by tick-borne pathogens results mainly from the regulation of
147 the levels of metabolic enzymes. Although it remains a challenge to explain how pathogen
148 infection modify the level of thousands of tick genes and hundreds of tick proteins [30], the
149 molecular mechanisms behind this have been partially characterized using the model
150 bacteria *A. phagocytophilum*. To induce transcriptional reprogramming in infected cells, *A.*
151 *phagocytophilum* hijacks the transcriptional regulatory machinery of ticks by inducing the
152 activation of signaling pathways [31] and the recruitment of transcription factors [32],
153 metabolites [33], and histone modifying enzymes [34] (Fig. 2). *A. phagocytophilum* may also
154 induce the degradation of tick proteins by tick proteasome which in turn decreases the
155 levels of targeted metabolic enzymes [23]. In addition, *A. phagocytophilum* infection induces
156 a high proportion of alternatively spliced transcript events (ASEs), which increases the
157 predicted gene ontology processes related to lipids, lipoproteins, and cholesterol/sterol
158 metabolism and other processes in human HL-60 cells infected by this pathogen [35]. The
159 occurrence of ASEs has not been tested in *A. phagocytophilum*-infected ticks or tick cell lines
160 yet. However, there is no reason to rule out the hypothesis that ASEs also occur in ticks in
161 response to *A. phagocytophilum* infection.

162 Activation of tick signaling pathways by *A. phagocytophilum* involves G-protein-coupled-
163 receptor (GPCR) that stimulates *Ixodes* p21-activated kinase (IPAK1) through the activation
164 of PI3K which results in actin-mediated signaling [31]. Activation of IPAK1 results in actin
165 phosphorylation and accumulation of actin in the nucleus which mediates binding of RNA
166 polymerase II (RNAPII) and TATA box-binding protein (TBP) to the promoter of the gene
167 *salp16* inducing its expression [31,36]. PI3K activation was also suggested to induce the
168 expression of the genes *hif-1α*, *hif-1β* and acetyltransferase p300 upon *A. phagocytophilum*
169 infection in tick cells [26] (Fig. 2). Mobilization of HIF-1α, HIF-1β and p300 at HIF target genes
170 may induce HIF-mediated transcriptional activation of glycolytic genes [26].

171 Among the transcription factors (TFs) activated by *A. phagocytophilum* infection is the tick
172 transcriptional activator protein-1 (AP-1) which induces the expression of the genes *iafgp* (*I.*
173 *scapularis* antifreeze glycoprotein) and *kat* (kynurenine aminotransferase) in infected cells
174 [32]. KAT is an enzyme involved in the production of the tryptophan metabolite xanthurenic
175 acid [33]. Results suggests that xanthurenic acid acts as a co-factor of an uncharacterized TF

that regulates the expression of one organic anion transporting polypeptide (*oatp*) gene in *A. phagocytophilum*-infected ISE6 cells [33]. The upregulation of *oatp* and *kat* was proposed to be associated with increased levels of intracellular xanthurenic acid, which in turn favors *A. phagocytophilum* replication in tick salivary glands [33]. Conversely, RNAi-mediated silencing of *oatp* and *kat* expression decrease *A. phagocytophilum* levels in tick cells [33]. A general mechanism is then proposed by which *A. phagocytophilum* nucleomodulins recruit tick metabolites and TFs to activate target genes (Fig. 2).

The capacity of *A. phagocytophilum* to downregulate gene expression in mammalian neutrophils was associated with host histone deacetylase 1 (HDAC1) recruitment to the promoters of target genes by the bacterial nucleomodulin ankyrin repeat protein AnkA [37]. Upon *A. phagocytophilum* infection, AnkA is secreted through T4SS, enters the granulocyte nucleus, binds stretches of AT-rich DNA and recruits host HDAC1 to its binding sites. *A. phagocytophilum* infection also induces HDAC1 binding and deacetylation of histone H3 (H3) which in turn results in downregulation of many host defense genes including *CYBB*, which encodes a component (i.e. NOX2) of the NADPH oxidase [37]. Downregulation of *CYBB* reduces superoxide anion production by NADPH oxidase which is essential for *A. phagocytophilum* survival in neutrophils [37]. Homologs of mammalian HDAC1 and H3 were identified in the *I. scapularis* genome [34]. Whether *A. phagocytophilum* AnkA is secreted and recruits HDAC1 to regulate gene transcription in infected ticks is currently unknown. However, *Ixodes* HDAC1 is overrepresented in *A. phagocytophilum*-infected tick salivary glands and chemical inhibition of this protein decreases *A. phagocytophilum* burden in tick cells [34]. This shows that the expression and activity of HDAC1 is critical for *A. phagocytophilum* survival in both mammalian and tick cells. Notably, in mammalian cells, HDAC1 controls the expression of the metabolic enzyme fructose-1,6-bisphosphatase (FBP), a rate-limiting enzyme in gluconeogenesis [38]. Interestingly, high HDAC1 protein levels [34] were associated with low FBP protein levels in *A. phagocytophilum*-infected tick salivary glands [26]. Conversely, low HDAC1 protein levels [34] correlated with high mRNA and protein levels of *fbp* and FBP, respectively, in *A. phagocytophilum*-infected tick midguts [26]. The negative correlation between the protein levels of HDAC1 and that of *fbp* and FBP strongly suggests that HDAC1 may be recruited by *A. phagocytophilum* nucleomodulins to act as a repressor of *fbp* expression in tick salivary glands (Fig. 2). A decrease in FBP protein

levels may contribute to the inhibition of gluconeogenesis that *A. phagocytophilum* induces in infected tick cells (see below) [23].

Tissue-specific changes of tick carbohydrate metabolism in response to infection

Glucose is the most abundant carbohydrate in the hemolymph of adult ticks and it is required in tick embryonic development [39]. The regulation of glucose metabolic pathways in ticks is essential for *A. phagocytophilum* survival in the vector [23,26]. Notably, *A. phagocytophilum* infection inhibits gluconeogenesis by decreasing the protein levels of **phosphoenolpyruvate carboxykinase**-cytosolic (PEPCK-C) which results in lower levels of glucose in *A. phagocytophilum*-infected ISE6 cells [23,26]. PEPCK-C is also a key regulatory enzyme in **glyceroneogenesis** [40] and thus, low levels of PEPCK-C were associated with reduced glycerol 3-phosphate production in infected tick cells [41]. Low PEPCK-C activity and decreased glucose anabolism inhibits apoptosis which is essential for *A. phagocytophilum* survival in tick cells and tissues [23,30]. Interestingly, glycerol 3-phosphate levels were also reduced in tick nymphs infected by *Borrelia mayonii* (Table 1). Low levels of glycerol 3-phosphate may have a significant impact in fatty acids metabolism, a poorly explored facet in tick-pathogen interactions.

Tissue-specific differential regulation of apoptosis pathways was observed in adult female midguts and salivary glands in response to *A. phagocytophilum* infection [42]. Likewise, the transcriptional reprogramming induced by *A. phagocytophilum* affects the levels of carbohydrate metabolism genes and proteins in a tissue-specific manner [26]. For example, the protein levels of all glycolytic enzymes increase in *A. phagocytophilum*-infected tick midguts, except for the cofactor-independent phosphoglycerate mutase (iPGM) and pyruvate kinase (PK) that despite gene up-regulation, the proteins were under-represented in this tissue [26]. In addition, gene and protein levels of phosphofructokinase (PFK) did not change in tick midguts in response to infection [26]. However, in salivary glands, the glycolysis enzymes fructose-bisphosphate aldolase A (ALDA), PFK and hexokinase (HXK) were under-represented while the rest of the glycolytic enzymes were over-represented. To accurately assess the metabolic impact of the differential protein representation in tick

midguts and salivary glands, a tissue-specific metabolomics approach is required. However, the lower levels of PK and HXK, key regulatory enzymes in glycolysis, in infected tick midguts and salivary glands, respectively, suggest that the production of glycolytic intermediates may be affected in a tissue-specific manner. It would be also important to assess the enzymatic activity of key enzymes which may vary in *A. phagocytophilum*-infected cells and tissues. For example, it was reported that the activity of HXK decreases more than two-folds in *A. phagocytophilum*-infected ISE6 cells [43]. The transcriptional activation of tick glycolytic genes was proposed to be mediated by tick HIF-1 α [26], which contains structural domains conserved in model organisms and the gene *hif-1 α* was significantly upregulated in infected tick midguts [26]. Another important gene targeted by HIF-1 α is pyruvate dehydrogenase kinase 1 (*pdh1*) which encodes the protein PDK1 that controls the enzymatic activity of pyruvate dehydrogenase E1 (PDE1) which transforms pyruvate into **Acetyl-CoA**, the first step in the TCA cycle. PDK1 mRNA and protein levels were higher in *A. phagocytophilum*-infected tick salivary glands suggesting that this pathogen inhibits the TCA cycle in infected salivary glands [26].

Crosstalk between tick amino acid and carbohydrate metabolism during pathogen infection

Several recent studies have shown that amino acids are central to host-pathogen metabolic interaction [3,23,41,44]. When digested, the protein-rich diet of ticks generates large amounts of amino acids. Therefore, amino acid metabolism is of great interest in ticks. For example, tyrosine-associated toxicity is lethal for blood-sucking arthropods which points out the significance of tyrosine degradation pathways. Inhibition of tyrosine aminotransferase (TAT) and 4-hydroxyphenylpyruvate dioxygenase (HPPD), the first two enzymes of the phenylalanine/tyrosine degradation pathway, caused the death of mosquitoes, kissing bugs and ticks after a blood meal [45].

Infection by *A. phagocytophilum* rewires the network of tick vector cell processes and changes the relative importance of some biological pathways including those involved in metabolism [46]. This is expected considering that *A. phagocytophilum* has a small genome

and cannot actively carry out several metabolic processes, and therefore requires and hijacks tick cell resources for survival within the vector [47]. For example, out of twenty amino acids, the *A. phagocytophilum* genome encodes only for the enzymes responsible of the biosynthesis of four amino acids (proline, glutamine, glycine and aspartate) and the glycolysis enzymes are reduced to those that produce glyceraldehyde-3-phosphate and dihydroxyacetone phosphate (DHAP) from **phosphoenolpyruvate** (PEP) [47]. In consequence, *A. phagocytophilum* changes the levels of amino acid metabolism enzymes [41], which in turn affects the intracellular levels of several amino acids and metabolic intermediates [23,33]. Interestingly, tick cells use tyrosine as a fuel to synthesize PEP during early *A. phagocytophilum* infection [41]. This process is achieved by activating the tyrosine/**oxaloacetate** (OAA)/PEPCK-M/PEP node by which *A. phagocytophilum* infection may decrease the tyrosine pool, which in turn may protect ticks against tyrosine-induced toxicity *in vivo* [41].

Tick-borne flaviviruses are known to affect protein processing and amino acid metabolism in tick cells [48,49]. Particularly, infection by Langat virus (LGTV) increased protein levels of acetyl-CoA acetyltransferase 1 (ACAT1), aldehyde dehydrogenase (ALDH5), glutamate dehydrogenase (GLUD1) and fumarylacetoacetase (FAH) in ISE6 cells [49]. Knockdown of the gene *acat1* was associated with decreased LGTV genome replication and LGTV particle release, while knockdown of the gene *fah* decreased only LGTV particle release in ISE6 cells [48]. Interestingly, FAH catalyzes the last step in tyrosine degradation suggesting that flaviviruses infection may, similarly to *A. phagocytophilum*, contribute to tyrosine detoxification in ticks.

Tick lipid metabolism is exploited by different tick-associated microorganisms

Starvation and desiccating conditions that ticks have to endure during off-host periods affect the energy reserves of the tick [50]. For example, starvation of *Dermacentor variabilis* ticks in optimal humidity conditions is known to result in up to 60% losses in lipid reserves [50]. The parietal fat body, the main lipid reserve organ for triacylglycerol and cholesterol, is gaining

increasing interest for the study of infections in insects and ticks [51,52]. However, lipid metabolism signaling pathways are still rarely studied in ticks.

Fatty acids play important roles as energy sources for several microorganisms including pathogenic bacteria [53]. If a pathogen can prevent the use of energy reserves by the host, this will affect its immune system, therefore promoting pathogen infection and persistence. *B. burgdorferi* is not able to synthesize fatty acids and cholesterol, thus relying on the vertebrate host cell or the blood meal of the tick for lipids supply [54]. Similarly, host cholesterol and other lipids are required by *A. phagocytophilum* for infection and multiplication in human cells [55]. Perilipin (PLIN), a major adipocyte lipid droplet-associated phosphoprotein that plays a central role during lipolysis and cholesterol synthesis, is differentially expressed in HL-60 cells in response to infection with *A. phagocytophilum* [56]. The bacteria are able to alter PLIN levels in infected cells, thus modifying the lipid metabolism of the host to facilitate bacterial infection [56]. This phosphoprotein is likely to play a similar role in the metabolism of the tick although its role has not been characterized so far.

Other obligate intracellular pathogens such as viruses need to hijack the cellular machinery for the synthesis of lipids to complete their replication. Flavivirus life cycle is very closely associated to host cell lipids, and they can rearrange intracellular membranes and reorganize the host cell lipid metabolism of infected cells to ensure viral replication [57]. Dengue virus infection can alter metabolic pathways in humans such as fatty acid biosynthesis and **β -oxidation** [58]. Exosomes are small extracellular vesicles that appear to play a central role in cell-to-cell communication [58]. Exosomes transport bioactive molecules, such as messenger RNA (mRNAs) and microRNA (miRNAs) that can be transferred in active form to adjacent cells or to distant organs thus being important vehicles for metabolic cross-talk [59]. Recent research has shown that Flaviviruses use arthropod-derived exosomes secreted by tick cells as a mean for viral RNA and metabolite transmission from the vector, and the vertebrate exosomes for dissemination within the host [60]. The knockdown of transcripts for a glycosylphosphatidylinositol (GPI)-anchored protein (VNN) was associated with decreased LGTV genome replication and LGTV release in ISE6 tick cells [48]. VNN is involved in coenzyme-A (CoA) metabolism [61], which is critical for lipid metabolism.

Fungal infections can also modulate the lipid metabolism of the host [62]. Recent research with different isolates of *Metarhizium anisopliae* showed that infection of *Rhipicephalus microplus* ticks with entomopathogenic fungi can modulate the lipid content thus inducing changes in important metabolic pathways [62]. However, lipid metabolism modulation during *M. anisopliae* infection in ticks was independent of AMP-activated protein kinase (AMPK) and extracellular signal-regulated protein kinase (ERK) pathways [62] which are the major pathways involved in the breakdown of triacylglycerol into fatty acids in mammals [63, 64]. This suggested that tick lipid metabolism modulation by *M. anisopliae* occurs through alternative and uncharacterized lipolysis pathway.

Concluding Remarks

Interactions between ticks, hosts and pathogens involve many metabolic pathways such as carbohydrate, protein, lipid and redox that are affected to produce conflict and cooperation between them. These interactions at the metabolic level lead to mutual beneficial effects of the tick-hosts-pathogen molecular interactions. Among them, amino acids are central to the host-pathogen metabolic interaction. Nevertheless, synthesis and degradation of amino acids is narrowly connected to carbohydrate metabolism, and the later with the lipid metabolism.

Despite recent advances in the study of tick-host-pathogen molecular interactions, most of the pathways and molecules involved in these interactions are not fully characterized. Furthermore, the regulatory mechanisms affecting the metabolic pathways affected by these interactions are only partially characterized (see Outstanding Questions). Future directions should include the application of omics technologies together with data integration algorithms [46] to the investigation of tick interactome, regulome, miRNAome, epigenome and metabolome and their interactions to advance the metabolic perspective on tick-pathogen interactions.

The identification of the metabolic adaptations during tick-pathogen interactions that support tick survival, spread and pathogen infection and transmission could provide new therapeutic targets for the prevention and control of tick-borne diseases. Studies of the

interactions between the pathogen and the host lipid metabolism can help to identify potential targets for the development of vaccines or treatments against health threatening pathogens. These strategies could include the inhibition of lipid biosynthesis. However, manipulation of this major metabolic pathway can also be detrimental for the host. For instance, Lovastatin treatment, an inhibitor of cholesterol synthesis, appears to delay infection and increases survival rates in AG129 mice infected with Dengue Virus Serotype 2 while it also tends to increase viremia which could affect disease progression [65]. A better understanding of the interactions of pathogens with the host cell lipid metabolism and other metabolic pathways could provide new therapeutic targets to control infections by tick-borne pathogens. Furthermore, as recently proposed the application of machine learning and big data analysis to the tick-host-pathogen interactions datasets may also lead to the high throughput identification of candidate vaccine protective antigens [66].

Funding

PA was funded by the Junta de Comunidades of Castilla-La Mancha (Spain).

369 Glossary

370 **Tricarboxylic acid cycle (TCA cycle):** a central route for oxidative phosphorylation in cells. In
371 the TCA cycle, acetyl-CoA is oxidized into carbon dioxide and reducing factors (NADH and
372 FADH₂), that contribute to synthesis of ATP during oxidative phosphorylation, are produced.
373 The TCA cycle is a crossroad of cellular metabolism in which carbohydrate, fatty acids and
374 amino acid metabolisms are interconnected.

375 **β-Oxidation:** the major pathway for the degradation of fatty acids and is essential to
376 produce energy from the generation of acetyl-CoA, NADH and FADH₂ when glucose supply is
377 limiting.

378 **Gluconeogenesis:** a metabolic pathway that during a longer-term fast or starvation results in
379 *de novo* glucose synthesis from non-carbohydrate carbon substrates, mainly lactate and
380 glycerol.

381 **Glyceroneogenesis:** a metabolic pathway that is activated during glucose deprivation to
382 synthesize glycerol 3-phosphate, a building block of triglyceride and fatty acid. In this
383 pathway, glycerol 3-phosphate is synthesized from pyruvate, alanine, glutamine and TCA
384 cycle intermediates.

385 **Glycolysis:** a major catabolic pathway in which one molecule of glucose is transformed into
386 two molecules of pyruvate releasing in the process two net molecules of ATP and two of
387 NADH. The expression of glycolytic genes is regulated by HIF-1α.

388 **Aerobic glycolysis:** A term frequently used to name a metabolic profile similar to that of the
389 *Warburg effect* in which a cell, even under normoxic conditions, generates high amount of
390 energy by using the glycolytic pathway, notably with high production of lactate.

391 **Phosphoenolpyruvate carboxykinase (PEPCK):** PEPCK catalyzes the first committed step in
392 gluconeogenesis, in which OAA is converted into PEP. There are two isoforms of the enzyme:
393 mitochondrial (PEPCK-M) and cytosolic (PEPCK-C). PEPCK-C is a key regulatory enzyme of
394 glyceroneogenesis and gluconeogenesis.

Hypoxia-inducible factor 1-alpha (HIF-1 α): a subunit of the heterodimeric transcription factor HIF-1 that together with HIF-1 β regulates the cellular response to hypoxia by inducing the transcriptional activation of genes involve in glycolysis.

Acetyl-CoA: a central metabolic intermediate synthesized in the mitochondria. Acetyl-CoA is the molecule through which glycolytic pyruvate enters the TCA cycle to be oxidized for energy production. It can be derived from carbohydrates, fatty acids and amino acids.

Glucose transporters (GLUT): a wide group of membrane proteins that transport glucose across the cell membrane. The GLUT family is divided into three classes of which class I member GLUT1 is transcriptionally regulated by HIF-1 α .

Oxaloacetate (OAA): a central metabolite produced in the TCA cycle that takes part in gluconeogenesis, amino acid metabolism and fatty acid synthesis among other pathways. In addition, OAA and PEP result from anaplerotic reactions that keep the metabolic flux of glycolysis and gluconeogenesis.

Phosphoenolpyruvate (PEP): a key metabolite that stands at the crossroad between gluconeogenesis, glycolysis and TCA cycle and can be produced by the activity of two enzymes enolase and PEPCK.

References

1. Ganeshan, K. and Chawla, A. (2014) Metabolic regulation of immune responses. *Annu. Rev. Immunol.* 32, 609-634
2. Green, D.R., Galluzzi, L., Kroemer G. (2014) Cell biology. Metabolic control of cell death. *Science.* 345, 1250256
3. Eisenreich, W. *et al.* (2015) Metabolic Adaptations of Intracellular Bacterial Pathogens and their Mammalian Host Cells during Infection ("Pathometabolism"). *Microbiol. Spectr.* 3, 27–58
4. de la Fuente, J. *et al.* (2016) Tick–host–pathogen interactions: conflict and cooperation. *PLoS. Pathog.* 12, e1005488
5. Cabezas-Cruz, A. *et al.* (2017) Tick-pathogen ensembles: do molecular interactions lead ecological innovation? *Front. Cell. Infect. Microbiol.* DOI: 10.3389/fcimb.2017.00074
6. Busby, A.T. *et al.* (2012) Expression of heat shock proteins and subolesin affects stress responses, *Anaplasma phagocytophilum* infection and questing behaviour in the tick, *Ixodes scapularis*. *Med. Vet. Entomol.* 26, 92-102
7. Shaw, D.K. *et al.* (2018) Vector Immunity and Evolutionary Ecology: The Harmonious Dissonance. *Trends Immunol.* 39, 862-873
8. de la Fuente, J. *et al.* (2015) Flying ticks: anciently evolved associations that constitute a risk of infectious disease spread. *Parasit. Vectors.* DOI: 10.1186/s13071-015-1154-1
9. de la Fuente, J. *et al.* (2016) *Anaplasma phagocytophilum* Uses Common Strategies for Infection of Ticks and Vertebrate Hosts. *Trends Microbiol.* 24, 173-180
10. Gulia-Nuss, M. *et al.* (2016) Genomic insights into the *Ixodes scapularis* tick vector of Lyme disease. *Nat. Commun.* 7, 10507
11. de la Fuente, J. *et al.* (2017) Tick-pathogen interactions and vector competence: identification of molecular drivers for tick-borne diseases. *Front. Cell. Infect. Microbiol.* DOI: 10.3389/fcimb.2017.00114
12. Núñez, G., Sakamoto, K., Soares, M.P. (2018) Innate Nutritional Immunity. *J Immunol.* 201, 11-18
13. Galay, R.L. *et al.* (2015) Iron metabolism in hard ticks (Acari: Ixodidae): the antidote

444 to their toxic diet. Parasitol. Int. 64, 182-189

445 14. Moumène, A. *et al.* (2018) Iron starvation conditions upregulate *Ehrlichia*

446 *ruminantium* type IV secretion system, *tr1* transcription factor and *map1* genes

447 family through the master regulatory protein ErxR. Front. Cell. Infect. Microbiol. DOI:

448 10.3389/fcimb.2017.00535

449 15. Kahlon, A. *et al.* (2013) *Anaplasma phagocytophilum* Asp14 is an invasin that

450 interacts with mammalian host cells via its C terminus to facilitate infection. Infect.

451 Immun. 81, 65-79

452 16. Villar, M. *et al.* (2015) Identification and Characterization of *Anaplasma*

453 *phagocytophilum* Proteins Involved in Infection of the Tick Vector, *Ixodes scapularis*.

454 PLoS. One. DOI: 10.1371/journal.pone.0137237

455 17. Troxell, B., Xu, H., Yang, X.F. (2012) *Borrelia burgdorferi*, a pathogen that lacks iron,

456 encodes manganese-dependent superoxide dismutase essential for resistance to

457 streptonigrin. J Biol. Chem. 287, 19284-19293

458 18. Villar, M. *et al.* (2016) The intracellular bacterium *Anaplasma phagocytophilum*

459 selectively manipulates the levels of vertebrate host proteins in the tick vector *Ixodes*

460 *scapularis*. Parasit. Vectors. DOI: 10.1186/s13071-016-1747-3

461 19. Sachla, A.J. *et al.* (2014) The crimson conundrum: heme toxicity and tolerance in GAS.

462 Front. Cell. Infect. Microbiol. DOI: 10.3389/fcimb.2014.00159

463 20. Price, J.V. and Vance, R.E. (2014) The macrophage paradox. Immunity. 41, 685-693

464 21. Hajdušek, O. *et al.* (2013) Interaction of the tick immune system with transmitted

465 pathogens. Front. Cell. Infect. Microbiol. DOI: 10.3389/fcimb.2013.00026.

466 22. Liu, L. *et al.* (2011) *Ixodes scapularis* salivary gland protein P11 facilitates migration of

467 *Anaplasma phagocytophilum* from the tick gut to salivary glands. EMBO Rep. 12,

468 1196-1203

469 23. Villar, M. *et al.* (2015) Integrated Metabolomics, Transcriptomics and Proteomics

470 Identifies Metabolic Pathways Affected by *Anaplasma phagocytophilum* Infection in

471 Tick Cells. Mol. Cell. Proteomics. 14, 3154-3172

472 24. Alberdi, P. *et al.* (2016) Tissue-specific signatures in the transcriptional response to

473 *Anaplasma phagocytophilum* infection of *Ixodes scapularis* and *Ixodes ricinus* tick cell

474 lines. Front. Cell. Infect. Microbiol. DOI: 10.3389/fcimb.2016.00020

25. Nonnenmacher, Y. and Hiller, K. (2018) Biochemistry of proinflammatory macrophage activation. *Cell. Mol. Life. Sci.* 75, 2093-2109
26. Cabezas-Cruz, A. *et al.* (2017) *Anaplasma phagocytophilum* infection subverts carbohydrate metabolic pathways in the tick vector, *Ixodes scapularis*. *Front. Cell. Infect. Microbiol.* DOI: 10.3389/fcimb.2017.00023
27. Shaw, D.K. *et al.* (2017) Infection-derived lipids elicit an immune deficiency circuit in arthropods. *Nat. Commun.* 8, 14401
28. Kamareddine, L. *et al.* (2018). The *Drosophila* immune deficiency pathway modulates enteroendocrine function and host metabolism. *Cell. Metab.* 28, 449-462
29. Davoodi, S. *et al.* (2018) The Immune Deficiency Pathway Regulates Metabolic Homeostasis in *Drosophila*. *BioRxiv* 253542; DOI: 10.1101/253542
30. Ayllón, N. *et al.* (2015) Systems biology of tissue-specific response to *Anaplasma phagocytophilum* reveals differentiated apoptosis in the tick vector *Ixodes scapularis*. *PLoS. Genet.* 11, e1005120
31. Sultana, H. *et al.* (2010) *Anaplasma phagocytophilum* induces actin phosphorylation to selectively regulate gene transcription in *Ixodes scapularis* ticks. *J Exp. Med.* 207, 1727-1743
32. Khanal, S. *et al.* (2018) Arthropod transcriptional activator protein-1 (AP-1) aids tick-rickettsial pathogen survival in the cold. *Sci. Rep.* 8, 11409
33. Taank, V. *et al.* (2017) Human rickettsial pathogen modulates arthropod organic anion transporting polypeptide and tryptophan pathway for its survival in ticks. *Sci. Rep.* 7, 13256
34. Cabezas-Cruz, A. *et al.* (2016) *Anaplasma phagocytophilum* increases the levels of histone modifying enzymes to inhibit cell apoptosis and facilitate pathogen infection in the tick vector *Ixodes scapularis*. *Epigenetics.* 11, 303-319
35. Dumler, J.S., Sinclair, S.H., Shetty, A.C. (2018) Alternative splicing of differentiated myeloid cell transcripts after infection by *Anaplasma phagocytophilum* impacts a selective group of cellular programs. *Front. Cell. Infect. Microbiol.* DOI: 10.3389/fcimb.2018.00014
36. Sukumaran, B. *et al.* (2006) An *Ixodes scapularis* protein required for survival of *Anaplasma phagocytophilum* in tick salivary glands. *J Exp. Med.* 203, 1507-1517

37. Rennoll-bankert, K.E. *et al.* (2015) Chromatin-bound bacterial effector ankyrin A recruits histone deacetylase 1 and modifies host gene expression. *Cell. Microbiol.* 17, 1640–1652
38. Yang, J. *et al.* (2017) Inhibiting histone deacetylases suppresses glucose metabolism and hepatocellular carcinoma growth by restoring FBP1 expression. *Sci. Rep.* 7, 43864
39. Martins, R. *et al.* (2018) The dynamics of energy metabolism in the tick embryo. *Rev. Bras. Parasitol. Vet.* 27, 259-266
40. Saponaro C. *et al.* (2015) The subtle balance between lipolysis and lipogenesis: a critical point in metabolic homeostasis. *Nutrients.* 7, 9453-9474
41. Cabezas-Cruz, A. *et al.* (2017) *Ixodes scapularis* tick cells control *Anaplasma phagocytophilum* infection by increasing the synthesis of phosphoenolpyruvate from tyrosine. *Front. Cell. Infect. Microbiol.* DOI: 10.3389/fcimb.2017.00375
42. Alberdi, P. *et al.* (2016) *Anaplasma phagocytophilum* manipulates host cell apoptosis by different mechanisms to establish infection. *Vet. Sci.* DOI: 10.3390/vetsci3030015
43. Ayllón N. *et al.* (2013) *Anaplasma phagocytophilum* inhibits apoptosis and promotes cytoskeleton rearrangement for infection of tick cells. *Infect. Immun.* 81, 2415-2425
44. Olive, A.J. and Sasseti, C.M. (2016) Metabolic crosstalk between host and pathogen: sensing, adapting and competing. *Nat. Rev. Microbiol.* 14, 221-234
45. Sterkel, M. *et al.* (2016) Tyrosine detoxification is an essential trait in the life history of blood-feeding arthropods. *Curr. Biol.* 26, 2188-2193
46. Estrada-Peña, A. *et al.* (2018) Use of graph theory to characterize human and arthropod vector cell protein response to infection with *Anaplasma phagocytophilum*. *Front. Cell. Infect. Microbiol.* DOI: 10.3389/fcimb.2018.00265
47. Dunning Hotopp, J.C. *et al.* (2006) Comparative genomics of emerging human ehrlichiosis agents. *PLoS. Genet.* 2, e21
48. Grabowski, J.M. *et al.* (2017) RNAi reveals proteins for metabolism and protein processing associated with Langat virus infection in *Ixodes scapularis* (black-legged tick) ISE6 cells. *Parasit. Vectors.* DOI: 10.1186/s13071-016-1944-0
49. Grabowski JM. *et al.* (2016) Changes in the proteome of Langat-infected *Ixodes scapularis* ISE6 cells: metabolic pathways associated with flavivirus infection. *PLoS. Negl. Trop. Dis.* 10, e0004180

50. Rosendale, A.J. *et al.* (2017) Dehydration and starvation yield energetic consequences that affect survival of the American dog tick. *J Insect. Physiol.* 101, 39-46
51. Angelo, I.C. *et al.* (2013) Neutral lipid composition changes in the fat bodies of engorged females *Rhipicephalus microplus* ticks in response to fungal infections. *Parasitol. Res.* 112, 501-509
52. Charroux, B. and Royet, J. (2010) *Drosophila* immune response: From systemic antimicrobial peptide production in fat body cells to local defense in the intestinal tract. *Fly (Austin)*. 4, 40-47
53. Barisch, C. and Soldati, T. (2017) Breaking fat! How mycobacteria and other intracellular pathogens manipulate host lipid droplets. *Biochimie.* 141, 54-61
54. Hoxmeier, J.C. *et al.* (2017) Metabolomics of the tick-*Borrelia* interaction during the nymphal tick blood meal. *Sci. Rep.* 7, 44394
55. Truchan, H.K., Seidman, D., Carlyon, J.A. (2013) Breaking in and grabbing a meal: *Anaplasma phagocytophilum* cellular invasion, nutrient acquisition, and promising tools for their study. *Microbes. Infect.* 15, 1017-1025
56. Manzano-Roman, R. *et al.* (2008) Expression of perilipin in human promyelocytic cells in response to *Anaplasma phagocytophilum* infection results in modified lipid metabolism. *J Med. Microbiol.* 57, 159-163
57. Martín-Acebes, M.A., Vázquez-Calvo, Á., Saiz, J.C. (2016) Lipids and flaviviruses, present and future perspectives for the control of dengue, Zika, and West Nile viruses. *Prog. Lipid. Res.* 64, 123-137
58. Chotiwan, N. *et al.* (2018) Dynamic remodeling of lipids coincides with dengue virus replication in the midgut of *Aedes aegypti* mosquitoes. *PLoS. Pathog.* 14, e1006853
59. Yu, X., Odenthal, M., Fries, J.W. (2016) Exosomes as miRNA Carriers: Formation-Function-Future. *Int. J Mol. Sci.* DOI: 10.3390/ijms17122028
60. Zhou, W. *et al.* (2018) Exosomes serve as novel modes of tick-borne flavivirus transmission from arthropod to human cells and facilitates dissemination of viral RNA and proteins to the vertebrate neuronal cells. *PLoS. Pathog.* 14, e1006764
61. Kinoshita, T and Fujita, M. (2016) Biosynthesis of GPI-anchored proteins: special emphasis on GPI lipid remodeling. *J Lipid. Res.* 57, 6-24
62. Sá, F.A. *et al.* (2018) *Metarhizium anisopliae* s.l. modulation of lipid metabolism

568 during tick infection is independent of AMPK and ERK pathways. *Parasitol. Res.* 117,
569 793-799

570 63. Jiao, P. *et al.* (2013) Hepatic ERK activity plays a role in energy metabolism. *Mol. Cell.*
571 *Endocrinol.* 375, 157-166

572 64. Hardie, DG. (2014) AMPK-sensing energy while talking to other signaling pathways.
573 *Cell. Metab.* 20, 939-952

574 65. Martinez-Gutierrez, M. *et al.* (2014) Lovastatin delays infection and increases survival
575 rates in AG129 mice infected with dengue virus serotype 2. *PLoS. One.* DOI:
576 10.1371/journal.pone.0087412

577 66. de la Fuente J. *et al.* (2018) High throughput discovery and characterization of tick
578 and pathogen vaccine protective antigens using vaccinomics with intelligent Big Data
579 analytic techniques. *Expert. Rev. Vaccines.* 17, 569-576

580 67. Kalil, S.P. *et al.* (2017) Immune-related redox metabolism of embryonic cells of the
581 tick *Rhipicephalus microplus* (BME26) in response to infection with *Anaplasma*
582 *marginalis*. *Parasit. Vectors.* DOI: 10.1186/s13071-017-2575-9

583

584 **Figure legends**

585 **Figure 1. Metabolic profiles of tick hemocytes and mammalian pro-inflammatory**
586 **macrophages M1.** The figure displays the metabolic profile of *A. phagocytophilum* (bacteria
587 without LPS)-infected ISE6 cells, a model of tick hemocytes, compared to that of mammalian
588 macrophages M1 in response to other intracellular bacteria with LPS. High levels of
589 hexokinase (HXX, green) and phosphofructokinase (PFK, green) are associated with high
590 aerobic glycolysis and increased lactate production in mammalian macrophages M1 [25].
591 The levels and activity of tick HXX decreased (red), and the levels of tick PFK did not change
592 (black), in response to infection which was associated with lower levels of lactate in infected
593 ISE6 cells [23,26,43]. HIF-1 α activation and stabilization (green), that enhances the
594 expression of glycolytic genes, is mediated by aerobic glycolysis and succinate accumulation
595 from glutamine in macrophages M1 [25]. In consequence, blocking aerobic glycolysis with 2-
596 deoxyglucose (2DG) inhibits HIF-1 α activation in these macrophages [25]. In contrast, 2DG
597 does not inhibit HIF-1 α activation (green) in ISE6 cells, an effect that is achieved by

LY294002, an inhibitor of the phosphoinositide 3-kinase (PI3K) pathway [26]. This suggests that HIF-1 α activity in tick hemocytes infected with *A. phagocytophilum* is regulated by PI3K and not aerobic glycolysis. Macrophages M1 and ISE6 cells share a low TCA cycle activity [25, 26]. Vertical arrows represent decrease (red) and increase (green) in metabolite levels.

Figure 2. Molecular mechanisms potentially implicated in tick metabolism modulation by the model pathogen *A. phagocytophilum*. Molecular mechanisms proposed to participate in the transcriptional reprogramming and proteome modulation induced by *A. phagocytophilum* infection in tick cells. Firstly, *A. phagocytophilum* activates the phosphoinositide 3-kinase (PI3K) signaling pathway in a G-protein-coupled-receptor (GPCR)-dependent manner to induce the expression of *salp16* [31], and hypoxia-inducible factor (HIF) *hif-1 α* in ticks [26] (sequential steps and tick molecular factors colored in violet). Secondly, *A. phagocytophilum* recruits tick transcription factors (TF), such as transcriptional activator protein-1 (AP-1) [32] and metabolites, such as xanthurenic acid [33], to regulate the expression of the metabolic genes *oatp* [33] and *kat* [32] (sequential steps and tick molecular factors colored in blue). Other tick metabolic genes may be regulated by the same mechanism in which we propose that *A. phagocytophilum* uncharacterized nucleomodulins may participate in the recruitment of tick TFs. Thirdly, the recruitment of histone deacetylase 1 (HDAC1) to the promoters of target genes through the activity of nucleomodulins may contribute to the repression of tick genes, e.g. fructose-1,6-bisphosphatase (*fbp*) (sequential steps and tick molecular factors colored in green). Lastly, although the mechanism is unknown, *A. phagocytophilum* also targets tick proteins to the proteasome which in turn may result in the underrepresentation of targeted metabolic enzymes [23]. Abbreviations are as follow: *Ixodes* p21-activated kinase (IPAK1), RNA polymerase II (RNAPII), TATA box-binding protein (TBP), acetyltransferase p300 (p300). Closed and open chromatin were represented as red and green double-stranded DNA, respectively.

Table 1. Tick metabolic pathways affected by tick-borne pathogens

Metabolic pathways	Experimental system (<i>in vitro</i> vs. <i>in vivo</i>)	Affected by (name of pathogen species)	Effect on tick metabolism*	References
Glycolysis	ISE6 tick cells	<i>Anaplasma phagocytophilum</i>	Decreased levels and activity of HXK	[26,43]
			Increased levels of PEP	[41]
			Decreased levels of glucose and lactate	[23]
	<i>Ixodes scapularis</i> adults midguts		Upregulation of genes involved in glycolysis (except <i>hxx</i> and <i>pfk</i>)	[26]
			Upregulation of genes <i>hif-1α</i> and <i>hif-1β</i>	
			Increased protein levels of glycolytic enzymes (except iPGM, PK and PFK) and glucose transporters GLUT1A and SGLT2	
<i>Ixodes scapularis</i> adults salivary glands	Upregulation of genes involved in glycolysis (except <i>hxx</i> , <i>pfk</i> , <i>alda</i> , <i>pgk1</i> and <i>enol</i>)	[26]		
	Increased protein levels of glycolytic enzymes (except HXK, PFK and ALDA) and glucose transporter GLUT1A			
Gluconeogenesis	ISE6 tick cells	<i>Anaplasma phagocytophilum</i>	Decreased levels of PEPCK-C and glucose	[23]
			Increased protein levels of PEPCK-M	[26]
	<i>Ixodes scapularis</i> adults midguts		Upregulation of genes involved in gluconeogenesis	[26]
			Decreased levels of gluconeogenesis enzymes (except FBP)	

	<i>Ixodes scapularis</i> adults salivary glands		Decreased levels of gluconeogenesis enzymes	
Glyceroneogenesis	ISE6 tick cells	<i>Anaplasma phagocytophilum</i>	Decreased levels of Glycerol 3-phosphate	[26]
	<i>Ixodes scapularis</i> adults midguts		Increased levels of PEP which may inhibit the enzyme TPI	
	<i>Ixodes scapularis</i> adults salivary glands		Upregulation of genes <i>tpi</i> and <i>gpdh</i> Increased and decreased protein levels of TPI and GPDH, respectively	
	<i>Ixodes scapularis</i> adults salivary glands		Upregulation of gene <i>tpi</i> Increased protein levels of TPI and GPDH	
	<i>Ixodes scapularis</i> nymphs**	<i>Borrelia mayonii</i>	Decreased levels of Glycerol 3-phosphate	[54]
Lipids	<i>Rhipicephalus microplus</i> adults fat body	<i>Metarhizium anisopliae</i>	Increased levels of cholesterol ester and triacylglycerol Decreased levels of free cholesterol	[51]
	<i>Ixodes scapularis</i> nymphs**	<i>Borrelia mayonii</i>	Decreased levels of cholesterol, ethanolamine phosphate and ethanolamine	[54]

Amino Acids	ISE6 tick cells	Anaplasma phagocytophilum	<p>Decreased levels of alanine, glycine, glutamine, glutamate and methionine</p> <p>Increased levels of serine, proline, tyrosine and phenylalanine</p> <p>Tyrosine is used as fuel to synthesize PEP</p>	[23,41]
			<p>Upregulation of gene <i>oatp</i></p> <p>Xanthurenic acid acts as a co-factor for unknown transcription factor(s) that control the expression of <i>oatp</i></p> <p>KAT regulates the expression of <i>oatp</i></p>	[33]
	<i>Ixodes scapularis</i> adults midguts and salivary glands		Changes in the levels of genes and proteins involved in amino acid synthesis and degradation	[41]
	<i>Ixodes scapularis</i> nymphs salivary glands		<p>Upregulation of genes <i>oatp</i> and <i>kat</i></p> <p>Xanthurenic acid acts as a co-factor for unknown transcription factor(s) that control the expression of <i>oatp</i></p>	[33]
	<i>Ixodes scapularis</i> nymphs**	Borrelia mayonii	Decreased levels of alanine, isoleucine, leucine, lysine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, valine, glutamate, glutamine and glycine	[54]
		Borrelia burgdorferi	<p>Increased levels of tryptophan and glutamine</p> <p>Decreased levels of alanine, isoleucine, leucine, phenylalanine, proline, threonine, valine and glycine</p>	

Redox	BME26 tick cells	<i>Anaplasma marginale</i>	Downregulation of pro-oxidant genes and upregulation of antioxidant genes	[67]
	<i>Ixodes scapularis</i> adults midguts and salivary glands	<i>Anaplasma phagocytophilum</i>	Significant changes in the levels of genes and proteins of hemoglobinolytic enzymes affecting host hemoglobin levels in tick tissues	[18]

* Full name of enzymes are in the main text except for phosphoglycerate kinase 1 (PGK1), enolase (ENOL), glycerol-3-phosphate dehydrogenase (GPDH), triosephosphate isomerase (TPI). ** Only the levels of tick metabolites in response to *Borrelia* spp. infection at day 4 of feeding were considered [54].

ISE6 cells - tick hemocytes model

Mammalian macrophages M1

A. phagocytophilum no LPS

Bacteria with LPS

A. phagocytophilum

Tick cell

Molecular factors

Bacterial nucleomodulins

Xanthurenic acid

Phosphate

Tick proteins

Molecular processes

Phosphorylation

Recruitment of tick proteins and/or metabolites

Targeting tick proteins to proteasome degradation

GPCR

Proteasome

AnkA

Cytosol

Nucleus

Activation
(*oatp* and *kat*)

Repression
(*fbp*)

Activation
(*salp16* and *hif-1 α*)

Activation
(glycolytic genes)

RNAPII

RNAPII

RNAPII

RNAPII

p300
HIF-1 β
HIF-1 α

PI3k
IPAK1

Actin

Actin^P

TF

TF

Actin^P
TBP

TF

Actin^P

TBP

Actin^P

TBP