



HAL
open science

Characterization of the phospholipid platelet-activating factor as a mediator of inflammation in chickens

Damien Garrido, Nathalie Katy Chanteloup, Angéline Trotereau, Adrien Lion, Geoffrey Bailleul, Evelyne Esnault, Sascha Trapp, Pascale Quéré, Catherine Schouler, Rodrigo Guabiraba

► To cite this version:

Damien Garrido, Nathalie Katy Chanteloup, Angéline Trotereau, Adrien Lion, Geoffrey Bailleul, et al.. Characterization of the phospholipid platelet-activating factor as a mediator of inflammation in chickens. *Frontiers in Veterinary Science*, 2017, 4, 20 p. 10.3389/fvets.2017.00226 . hal-02626837

HAL Id: hal-02626837

<https://hal.inrae.fr/hal-02626837>

Submitted on 26 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/321882526>

Characterization of the Phospholipid Platelet-Activating Factor As a Mediator of Inflammation in Chickens

Article · December 2017

DOI: 10.3389/fvets.2017.00226

CITATIONS

0

READS

31

10 authors, including:



Adrien Lion

French National Institute for Agricultural Res...

9 PUBLICATIONS 7 CITATIONS

SEE PROFILE



Sascha Trapp

French National Institute for Agricultural Res...

33 PUBLICATIONS 833 CITATIONS

SEE PROFILE



Catherine Schouler

French National Institute for Agricultural Res...

64 PUBLICATIONS 1,043 CITATIONS

SEE PROFILE



Rodrigo Guabiraba

French National Institute for Agricultural Res...

37 PUBLICATIONS 882 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



phagotherapy of avian colibacillosis [View project](#)



Dengue infection pathogenesis in immunocompetent mice [View project](#)



Characterization of the Phospholipid Platelet-Activating Factor As a Mediator of Inflammation in Chickens

Damien Garrido, Nathalie K. Chanteloup, Angéline Trotereau, Adrien Lion, Geoffrey Bailleul, Evelyne Esnault, Sascha Trapp, Pascale Quéré, Catherine Schouler and Rodrigo Guabiraba*

ISP, INRA, Université François Rabelais de Tours, Nouzilly, France

OPEN ACCESS

Edited by:

Dirk Werling,
Royal Veterinary College,
United Kingdom

Reviewed by:

Kenneth James Genovese,
Agricultural Research Service
(USDA), United States
Tatiana Kuster,
Royal Veterinary College,
United Kingdom

*Correspondence:

Rodrigo Guabiraba
rodrigo.guabiraba-brito@inra.fr

Specialty section:

This article was submitted to
Veterinary Infectious Diseases,
a section of the journal
Frontiers in Veterinary Science

Received: 19 October 2017

Accepted: 06 December 2017

Published: 18 December 2017

Citation:

Garrido D, Chanteloup NK,
Trotereau A, Lion A, Bailleul G,
Esnault E, Trapp S, Quéré P,
Schouler C and Guabiraba R (2017)
Characterization of the Phospholipid
Platelet-Activating Factor As a
Mediator of Inflammation in Chickens.
Front. Vet. Sci. 4:226.
doi: 10.3389/fvets.2017.00226

Lipid mediators are known to play important roles in the onset and resolution phases of the inflammatory response in mammals. The phospholipid platelet-activating factor (PAF) is a pro-inflammatory lipid mediator which participates in vascular- and innate immunity-associated processes by increasing vascular permeability, by facilitating leukocyte adhesion to the endothelium, and by contributing to phagocyte activation. PAF exerts its function upon binding to its specific receptor, PAF receptor (PAFR), which is abundantly expressed in leukocytes and endothelial cells (ECs). In chickens, lipid mediators and their functions are still poorly characterized, and the role of PAF as an inflammatory mediator has not yet been investigated. In the present study we demonstrate that primary chicken macrophages express PAFR and lysophosphatidylcholine acyltransferase 2 (LPCAT2), the latter being essential to PAF biosynthesis during inflammation. Also, exogenous PAF treatment induces intracellular calcium increase, reactive oxygen species release, and increased phagocytosis by primary chicken macrophages in a PAFR-dependent manner. We also show that PAF contributes to the *Escherichia coli* lipopolysaccharide (LPS)-induced pro-inflammatory response and boosts the macrophage response to *E. coli* LPS via phosphatidylinositol 3-kinase/Akt- and calmodulin kinase II-mediated intracellular signaling pathways. Exogenous PAF treatment also increases avian pathogenic *E. coli* intracellular killing by chicken macrophages, and PAFR and LPCAT2 are upregulated in chicken lungs and liver during experimental pulmonary colibacillosis. Finally, exogenous PAF treatment increases cell permeability and upregulates the expression of genes coding for proteins involved in leukocyte adhesion to the endothelium in primary chicken endothelial cells (chAEC). In addition to these vascular phenomena, PAF boosts the chAEC inflammatory response to bacteria-associated molecular patterns in a PAFR-dependent manner. In conclusion, we identified PAF as an inflammation amplifier in chicken macrophages and ECs, which suggests that PAF could play important roles in the endothelium-innate immunity interface in birds during major bacterial infectious diseases such as colibacillosis.

Keywords: avian pathogenic *Escherichia coli*, chicken, colibacillosis, endothelial cells, inflammation, macrophages, platelet-activating factor, platelet-activating factor receptor

INTRODUCTION

Inflammation is a complex and a tightly coordinated biological process driven by a wide array of cells and chemical mediators (1). Lipid mediators (e.g., eicosanoids and phospholipids) are molecules well known to take part in the onset of the inflammatory response in different vertebrate organisms (2, 3). These molecules are rapidly biosynthesized by myeloid and epithelial cells within seconds to minutes of acute stimulation of different origins (1, 4). Among these molecules, the phospholipid platelet-activating factor (PAF) potent pro-inflammatory lipid mediator. PAF activates a G protein-coupled receptor named PAF receptor (PAFR) (5–7), which results in pleiotropic and potent biological effects including platelet activation, airway constriction, hypotension, and increased vascular permeability *via* signaling pathways such as phosphatidylinositol 3-kinase (PI3K), PKC, phospholipase A2 (PLA₂), and PLC (5, 8–10). PAF is synthesized *via* two distinct pathways, the *de novo* and remodeling pathways (2, 5, 11), the latter of which is regulated by inflammatory extracellular signals and plays a key role in PAF-induced physiopathology (2, 5, 11, 12). PAF biosynthesis occurs through lyso-PAF formation *via* PLA₂ catalytic activity. In a subsequent step, lysophosphatidylcholine acyltransferase (LPCAT, a member of the lysophospholipid acyltransferase family) generates PAF from lyso-PAF. LPCAT2 is known to biosynthesize PAF upon pro-inflammatory stimuli and may localize to the endoplasmic reticulum (ER) and the Golgi (13, 14). Moreover, PAF levels in the extracellular milieu can be controlled through rapid degradation by PAF-acetylhydrolase (PAF-AH) (15). All these physiological or pathological processes have been well characterized in mammals, more notably in rodents and humans.

Platelet-activating factor synthesis upon inflammatory stimuli such as bacterial lipopolysaccharide (LPS) has been reported in murine peritoneal cells and macrophages (16, 17) and in human eosinophils and neutrophils (18, 19). PAF has been shown to be particularly active in mammalian macrophages during inflammation. It can reprogram macrophages such that there is an enhanced cytokine response to subsequent endotoxin stimulation (20). In addition, PAF acts as an autocrine mediator of macrophage activation upon LPS challenge *via* tumor necrosis factor alpha (TNF- α) expression (21). Systemic administration of PAF mimics a septic shock condition characterized by vasodilation with diminished peripheral vascular resistance and tissue oxygen utilization (22, 23). Moreover, TNF- α and PAF act synergistically to increase neutrophil-endothelium adhesion by stimulating endothelial expression of cell adhesion molecules: intercellular adhesion molecule 1 and E-selectin (24).

Phylogenetically, the regulatory role of PAF is presumed to have appeared in protozoans, having been maintained during evolution (25). In addition to mammals, PAF biosynthesis has been found in marine invertebrates and lower vertebrates (e.g., birds and reptiles). In chickens, PAF production has been observed in the chick's retina (26). Also, PAF has been shown to activate chicken thrombocytes in a Ca²⁺-dependent manner (27). Enzymes involved in the metabolism and catabolism of PAF in chickens were poorly studied, although PAF-AH was reported to be absent in chicken serum (28). Nevertheless, catalytic residues

in PAF-AH sequence are conserved in chickens (29). Overall, very few studies specifically addressed the role of inflammatory lipid mediators in chickens, although some data are available for eicosanoids (30–32). However, to date, no study has addressed the putative regulatory roles for PAF in chickens during inflammation or infection.

Poultry flocks are often impacted by several infectious diseases, and avian colibacillosis is the most prevalent bacterial infection, representing important economic losses to the poultry industry worldwide (33). Colibacillosis is caused by avian pathogenic *Escherichia coli* (APEC) strains, commonly called APEC. Antibiotherapy is leading to antimicrobial resistance, and commercially available vaccines confer protection to a limited set of APEC strains. Pulmonary colibacillosis is the most common form of disease, characterized by acute inflammatory response with massive influx of heterophils and macrophages. Although colibacillosis pathogenesis needs to be better characterized, it is believed that macrophages are crucially involved in bacterial killing and amplification of inflammation in APEC infected chicken lungs (33–35).

In line with that, PAF has been implicated in several animal models of lung injury, and treatment with PAF receptor antagonists attenuates endotoxin-induced lung injury (36). During pulmonary Gram-negative bacteria infection, PAF production is important for bacteria clearance (37). However, during experimental influenza A virus infection in mice, PAF plays a deleterious role by amplifying inflammation and tissue damage (38).

Despite a growing interest in characterizing early mediators of inflammation in chickens, and more notably their crosstalk with innate immunity sentinel cells such as macrophages, no previous works have addressed the mechanisms by which lipid mediators may contribute to inflammation or bacterial clearance. In the present work, we aimed to characterize the PAF/PAFR interactions and to shed light on their functions in chicken macrophages and endothelial cells (ECs), especially with respect to how PAF system molecules are modulated during pulmonary colibacillosis.

MATERIALS AND METHODS

Reagents

Platelet-activating factor (synthetic PAF C16:0, β -acetyl- γ -O-hexadecyl-L- α -phosphatidylcholine hydrate, $\geq 98\%$ HPLC purity tested), human PAF-AH, dimethyl sulfoxide (DMSO), LPS (from *E. coli* O55:B5), and 2',7'-dichlorofluorescein diacetate (DCFDA, $\geq 95\%$ HPLC purity tested) were purchased from Sigma-Aldrich, UK. PAFR antagonists PCA 4248 (C₁₉H₂₃NO₄S, $>95\%$ HPLC purity tested) and WEB 2086 (Apafant, C₂₂H₂₂N₅O₂SCl, $>99\%$ HPLC purity tested), PI3K/Akt inhibitor wortmannin (WMN; SL-2052, C₂₃H₂₄O₈), and calcium calmodulin kinase II (CaMK II) inhibitor KN62 (C₃₈H₃₅N₅O₆S₂, $>98\%$ HPLC purity tested) were purchased from Tocris Biotechnique, UK.

Animals

White leghorn chickens from the GB1 Athens inbred line (B13/B13 histocompatible chickens) or from the PA12 outbred line were hatched and raised specific-pathogen-free (SPF) at INRA

(Plate-Forme d'Infectiologie Expérimentale, PFIE, Nouzilly, France). Food and water were provided *ad libitum*. For APEC infection experiments, B13/B13 histocompatible chickens were housed in BSL2 poultry isolator units.

Cell Culture

HD11, an avian myelocytomatosis virus (MC29)-transformed chicken macrophage-like cell line (39), was cultured in RPMI-1640 medium (Gibco, UK) supplemented with 10% heat-inactivated fetal calf serum (FCS, Gibco, UK), 10 mM HEPES, 2 mM glutamine, 100 U/ml penicillin, and 100 µg/ml streptomycin (all from Sigma-Aldrich, UK) and grown routinely in a 75-cm² flask (Corning, NY, USA) at 40°C under 5% CO₂.

Chicken bone marrow-derived macrophages (BMDM) were outgrown from bone marrow cells using recombinant chicken macrophage colony-stimulating factor 1 (CSF-1) (40) derived from COS-7 cells (fibroblast-like cell line derived from monkey kidney tissue, ATCC, USA) transfected with a pTarget vector (Promega, UK) expressing chicken CSF-1 (kindly provided by Prof. Pete Kaiser, The Roslin Institute, UK). Briefly, femurs and tibia of 3–4-week-old B13/B13 histocompatible or PA12 chickens were removed, cut, and the marrow flushed with sterile phosphate-buffered saline (PBS, Gibco, UK). Next, cells were resuspended in RPMI-1640, loaded onto an equal volume of Histopaque 1.077 gradient (Sigma-Aldrich, UK), and centrifuged at 250 g for 30 min without brake. Cells at the interface were collected, washed twice with PBS, and cultured at a final concentration of 1 × 10⁶ cells/ml in sterile 60 mm bacteriological plates (Corning, NY, USA) in prewarmed complete RPMI-1640 medium supplemented with COS-7 supernatant containing chicken CSF-1, 10% FCS, 10 mM HEPES, 2 mM glutamine, 100 U/ml penicillin, and 100 µg/ml streptomycin at 40°C under 5% CO₂. Half of the medium was replaced with fresh, pre-warmed complete medium containing chicken CSF-1 at day 4. At day 7, adherent cells were washed and harvested in cold PBS containing 2 mM ethylenediaminetetraacetic acid (EDTA, Sigma-Aldrich, UK), centrifuged, and resuspended in complete RPMI-1640 medium. All adherent cells presented macrophage-like morphology as evaluated in glass slides prepared in a CytoSpin[®] Cytocentrifuge (Shandon, USA). Also, 97% of these cells were KUL01+ cells (a chicken macrophage marker) as evaluated by Flow Cytometry using a mouse anti-chicken monoclonal antibody (1:200 dilution, Bio-Rad, UK) and a rat-anti mouse phycoerythrin-coupled secondary antibody (1:300 dilution, Biolegend, UK).

Peripheral blood mononuclear cells (PBMC) were isolated from B13/B13 histocompatible chickens as previously described (41). Briefly, peripheral blood collected in EDTA-coated tubes was mixed with 1% methylcellulose (Sigma-Aldrich, UK, 1:1 v/v) and centrifuged at 25 g for 15 min. The upper phase was then diluted with PBS to the original volume of blood/methylcellulose mixture, carefully layered onto a Histopaque 1.077 gradient and centrifuged at 250 g for 30 min without brake. The PBMC-containing layer was collected, washed, and resuspended in complete RPMI-1640 medium for further incubation at 40°C under 5% CO₂. After 24 h, adherent cells presented monocyte/macrophage-like morphology as evaluated in glass slides prepared in a CytoSpin[®] Cytocentrifuge (Shandon, USA).

Primary chicken aortic endothelial cells (chAEC) were prepared from specific-pathogen-free (SPF) 18-day-old PA12 chicken embryos as previously described (42). Primary chAEC were serially passaged (p2–p5) using 0.05% trypsin-EDTA (Gibco, UK) and kept in EC growth medium (EGM-2 BulletKit, LONZA) at 40°C under 5% CO₂.

General *In Vitro* Stimulation Protocol

Twelve-well tissue culture plates (Corning, NY, USA) were seeded at 7 × 10⁵ cells per well of HD-11 macrophage-like cells, BMDM, PBMC, or primary chAEC in complete medium and incubated at 40°C under 5% CO₂ overnight prior to different stimulation protocols. LPS (10 ng/ml) and hPAF-AH (10 µg/ml) were previously diluted in RPMI-1640. PAF (0.1–10 µM), PCA 4248 (10 µM), WEB 2086 (10 µM), WMN (200 nM), and KN62 (1 µM) were previously diluted in DMSO. Dead bacteria [DB; APEC strain BEN2908 at 10 multiplicity of infection (MOI), 15 min incubation at 65°C for killing] were previously diluted in LB medium. Final concentration of DMSO (the vehicle used for the dilution of PAF, PAFR antagonists, and cell signaling inhibitors) in cell culture wells never exceeded 0.1%. Treatment with PAF receptor antagonists or intracellular signaling pathway inhibitors was performed 1 h prior to stimulation. Cell culture supernatants were recovered 6 h after stimulation (unless otherwise indicated) and stocked at –20 or –80°C for subsequent analyses specified below. Cells were washed in sterile PBS and lysed with RNA lysis buffer (Macherey-Nagel, Germany) containing 2-mercaptoethanol (Merck Millipore, Germany), snap frozen in liquid nitrogen, and stocked at –80°C until RNA extraction. For total protein quantification and Western blot (WB) analysis, cells were washed in PBS followed by cell lysis using RIPA buffer containing a cocktail of protease inhibitors, which also included phenylmethylsulfonyl fluoride and sodium orthovanadate (Santa Cruz Biotechnology, USA), and stocked at –20°C.

Gene Expression Analysis

Total RNA was extracted from frozen cell lysates using the NucleoSpin RNA II kit (Macherey-Nagel, Germany) according to the manufacturer's instructions. Tissue samples were first processed using Trizol (TRI Reagent, Sigma-Aldrich, UK), and RNA was further purified as for cell lysates. RNA quality and concentration were determined using a NanoDrop (Thermo Scientific, USA). Total RNA (up to 1 µg per reaction) was reverse transcribed with iScript cDNA synthesis kit (Bio-Rad, USA). Quantitative real-time PCR (qRT-PCR) was performed on a CFX96 machine (Bio-Rad, USA), the reaction mixture consisting of 1 µl of cDNA, 5 µl of iQ SYBR Green Supermix (Bio-Rad, USA), 0,25 µl of each primer pair (250 nM, Eurogentec, Belgium), and 3,5 µl of nuclease-free water (Sigma-Aldrich, UK) in a total reaction volume of 10 µl. qRT-PCR data were analyzed using the CFX Manager software 3.1 (Bio-Rad, USA). Amplicon size of qRT-PCR products was confirmed in a 2% agarose gel. Gene expression for each target gene was normalized to gene expression levels of chicken hypoxanthine-guanine phosphoribosyltransferase and β-2-microglobulin (β-2-M). Relative normalized expression was calculated using the 2^{–ΔΔC_t} method, and data are represented as fold increase as compared to control group

(unless otherwise indicated). qRT-PCR primers (Table 1) were designed and verified using primer BLAST.¹

Intracellular Calcium

The Fluo-4 NW Calcium Assay kit (Molecular Probes, USA) was used to measure changes in HD11 cells' intracellular calcium levels upon different stimuli. Briefly, HD11 cells at 3×10^4 cells per well in a black 96-multiwell plate (Corning, NY, USA) were loaded with 100 μ l of the dye loading solution containing Fluo-4 NW dye and probenecid, according to the manufacturer's instructions. The 96-well plate was incubated at 40°C under 5% CO₂ for 30 min in the dark, and the stimuli of interest were added to the cells at $T = 0$, and the plates were read immediately. The changes in Fluo-4 NW fluorescence were measured at an excitation wavelength of 494 nm and an emission wavelength of 516 nm in a GloMax®-Multi Detection System plate reader (Promega, UK). Calcium mobilization was recorded over time (213 s, with 3 s intervals).

Apoptosis and Cell Viability

To evaluate the pro-apoptotic and cytotoxic effects of different stimuli, cells were cultured as previously mentioned. Apoptotic/late apoptotic cells were identified using annexin V and propidium iodide (PI) double-positive staining strategy. Briefly, following different stimuli, BMDM were washed with EDTA-containing-PBS, harvested, washed in cold PBS, and resuspended in an annexin-binding buffer (BD Biosciences). Subsequently, annexin V-fluorescein isothiocyanate (FITC) was added to the cells for 30 min. After a washing step, the cells were further incubated with PI (BD Biosciences) for 15 min. After two washing steps, stained cells (50,000 acquired events over total gated cells) were analyzed immediately using a BD FACSCalibur™ (BD Biosciences).

In other experiments, cell viability was determined using the colorimetric methylthiazolotetrazolium bromide (MTT) assay (Sigma-Aldrich, UK) that measures metabolic activity as a surrogate for cell viability/cytotoxicity. Briefly, following different stimuli, MTT was added to a final concentration of 5 μ g/ml per well, and cells were incubated for 2 h at 40°C under 5% CO₂. After complete solubilization of the dye using DMSO, plates were read at 550 nm in a Multiskan Ascent plate reader (Thermo Fisher Scientific, USA).

¹<http://www.ncbi.nlm.nih.gov/tools/primer-blast>.

Caspase 3/7 Activity

Caspase 3/7 activity was carried out using the Caspase-Glo 3/7 assay kit (Promega, UK) according to the manufacturer's protocol. Briefly, BMDM at 3×10^4 cells per well in a black 96-multiwell plate (Corning, NY, USA) were incubated with different stimuli for 6 h. Then, 100 μ l of Caspase-Glo 3/7 reagent was added to the wells. Plates were gently shaken and then incubated in the dark at 20°C for 60 min before luciferase activity was recorded using a GloMax®-Multi Detection System plate reader (Promega, UK).

Intracellular Reactive Oxygen Species (ROS) Analysis

The cell-penetrating dye 2',7'-dichlorofluorescein diacetate (H₂DCFDA) is oxidized by ROS to yield the fluorescent molecule 2',7'-dichlorofluorescein (DCFDA), which is retained intracellularly and visible at the 488 nm excitation laser line. To evaluate ROS production upon different stimuli, BMDM were cultured as previously described and incubated with 10 μ M H₂DCFDA (Sigma-Aldrich, UK) in serum-free medium for 20 min. Then medium was changed, and cells receiving different stimuli were kept at 40°C under 5% CO₂ for 2 h. Cells were then collected and washed in ice-cold PBS prior to flow cytometry analysis (50,000 acquired events over total gated cells) using a BD FACSCalibur™ (BD Biosciences, UK).

Flow Cytometry Analysis of PAFR

For the identification of PAFR expression in BMDM, cells were cultured as previously mentioned, exposed or not to different stimuli, and incubated for 2 h at 40°C under 5% CO₂. Then, cells were harvested, washed in FACS buffer (2 mM EDTA and 2% FCS in PBS), centrifuged, resuspended in FACS buffer, and stained with a rabbit polyclonal antibody raised against amino acids 1–300 of PAFR of human origin (Santa Cruz Biotechnology, USA) at 1:200 dilution. After 45 min at 4°C, cells were washed, resuspended in FACS buffer, and stained with a goat anti-rabbit IgG-FITC (Santa Cruz Biotechnology, USA) at 1:500 dilution. After 40 min at 4°C, cells were washed, centrifuged, resuspended in FACS buffer, and analyzed by flow cytometry (100,000 acquired events over total gated cells) using a BD FACSCalibur™ (BD Biosciences, UK).

Immunofluorescence

Bone marrow-derived macrophages or HD11 cells were grown at 2×10^5 cells per well on sterile glass coverslips placed in 24-well

TABLE 1 | Primer pairs used in the present study for quantitative real-time PCR analysis.

Target mRNA	Forward primer 5'–3'	Reverse primer 5'–3'
β -2-M (<i>B2M</i>)	CGTCCTCAACTGCTTCGCG	TCTCGTGCTCCACCTTGC
COX-2 (<i>PTGS2</i>)	CTGCTCCCTCCCATGTCAGA	CACGTGAAGAATCCGGTGT
CXCL12 (<i>IL8L2</i>)	CTGCGGTGCCAGTGCATTAG	AGCACACCTCTCTCCATCC
Hypoxanthine-guanine phosphoribosyltransferase (<i>HPRT</i>)	TGGTGGGGATGACCTCTCAA	GGCCGATATCCACACTTCG
E-selectin (<i>SELE</i>)	AATGCAAAGCTGTGACCTGC	GCCTGGATTGTCTGTGAGA
IL-1 β (<i>IL-1B</i>)	AGGCTCAACATTGCGCTGTA	CTTGATGCCCTTGATGCCA
iNOS (<i>NOS2</i>)	CCACCAGGAGATGTTGAACATGTC	CCAGATGTGTGTTTTCCATGCA
LPCAT2 (<i>LPCAT2</i>)	GTCCGACAAACCATGGCAAC	TCCAATTGCTCCCCCTTTGG
Platelet-activating factor receptor (<i>PTAFR</i>)	GGAGACCTTTGACACCGTGG	TTTGCCCTCTCCAGACATCG
VCAM-1 (<i>VCAM1</i>)	AGCTTGATGTCAAAGTTCCTCCT	AGGTTCCATTGACTGCTGTT

plates containing complete medium and incubated at 40°C under 5% CO₂ overnight. After adding the stimuli of interest for 2 h, cells were washed in PBS and fixed with 4% paraformaldehyde (Sigma-Aldrich, UK) for 20 min at room temperature, permeabilized with 0.5% Triton X-100 (Sigma-Aldrich, UK) for 5 min, and blocked with PBS, 0.1% Triton X-100, and 2% bovine serum albumin (Sigma-Aldrich, UK). Immunostaining was performed with a rabbit polyclonal antibody raised against amino acids 1–300 of PAFR of human origin or rabbit polyclonal IgG as an isotype control (Santa Cruz Biotechnology, USA) at 1:200 dilution. Secondary antibody goat anti-rabbit IgG-FITC (Santa Cruz Biotechnology, USA) was used at 1:500 dilution. Cell nuclei were counterstained with ProLong Gold Antifade Mountant with DAPI (Thermo Fisher, USA). Cells were observed under an Axiovert 200 M inverted epifluorescence microscope equipped with the Apotome imaging system (Zeiss, Germany). Images were captured with an AxioCam MRm camera and analyzed using the Axiovision software (Zeiss, Germany).

Western Blot

Total protein from BMDM cell lysates was quantified using a Quick Start™ Bradford Protein Assay (Bio-Rad, USA). Thirty micrograms of protein from whole cell lysates were mixed with 2× electrophoresis sample buffer (Santa Cruz Biotechnology, USA). After boiling for 5 min, proteins were separated using 4–15% Mini-PROTEAN TGX™ Precast gels (Bio-Rad, USA) and transferred onto polyvinylidene difluoride (PVDF) membranes using a Trans-Blot® Turbo™ Mini PVDF Transfer pack (Bio-Rad, USA) in a Trans-Blot® Turbo™ Transfer system (Bio-Rad, USA). The membranes were blocked in a buffer containing 5% non-fat dry milk in PBS for 1 h. Primary antibody targeting PAFR was the same as utilized for immunofluorescence (IF) staining at 1:500 dilution. To detect Akt, a rabbit anti-Akt IgG polyclonal antibody (detecting endogenous levels of total Akt1, Akt2, and Akt3 proteins—Cell Signaling Technology, USA) was utilized at 1:500 dilution. To detect CaMK II, a rabbit anti-CaMK II IgG polyclonal antibody (epitope corresponding to amino acids 1–300 mapping at the N-terminus of CaMK II α of human origin, Santa Cruz Biotechnology, USA) was utilized at 1:500. Mouse anti-glyceraldehyde-3-phosphate dehydrogenase IgG1 monoclonal antibody (GAPDH, Merck Millipore, Germany) was utilized as internal loading control at 1:1,000 dilution. After blocking, primary antibodies diluted in 5% non-fat dry milk in PBS were added to the membranes for incubation overnight at 4°C under agitation. Following reaction with the primary antibodies, membranes were thoroughly washed in wash buffer containing 0.1% Tween 20 (Sigma-Aldrich, UK) in PBS and labeled with HRP-conjugated secondary antibodies (goat anti-rabbit or rabbit anti-mouse, Life Technologies, USA, at 1:2,000 dilution) for 1 h. After washing, the antigens were visualized using a Clarity™ ECL Western Blotting Substrate (Bio-Rad, USA) in a chemiluminescence FUSION FX apparatus (Vilber Lourmat, France).

Phagocytosis Assay

Bone marrow-derived macrophages were seeded at 2×10^5 cells per well in 24-well tissue culture plates, treated with different stimuli and immediately incubated with 10 μ g/ml of Zymosan

A (*Saccharomyces cerevisiae*) BioParticles™ Alexa Fluor™ 488 conjugate (Thermo Fisher Scientific, USA) in complete medium at 4°C or 40°C under 5% CO₂ for 1 h. After incubation, cells were harvested, washed twice with ice-cold EDTA-containing PBS, centrifuged, resuspended in PBS, and analyzed by flow cytometry (100,000 acquired events over total gated cells) using a BD FACSCalibur™ (BD Biosciences, UK). Data were expressed as the percentage difference (Δ) between fluorescence positive cells and cells incubated at 40°C (active phagocytosis) and 4°C (limited phagocytosis).

Nitrite Dosage

Nitrite (NaNO₂) content was used as an index of nitric oxide (NO) production by chicken macrophages. Nitrite levels were determined in supernatants of BMDM, PBMC, or HD11 cell cultures by spectrophotometry using the Griess reagent system (Promega, UK). Briefly, 50 μ l of the samples were added to 50 μ l of freshly prepared Griess reagent in a 96-well plate. The absorbance was read at 550 nm in a Multiskan Ascent plate reader (Thermo Fisher Scientific, USA). The nitrite concentration was calculated using a sodium nitrite standard curve.

Lipidomics Analysis

Prostaglandin E2 (PGE2) was quantified from BMDM supernatant using liquid chromatography–tandem mass spectrometry (LC-MS/MS). Briefly, 1 ml of BMDM cell culture supernatant was added to 300 μ l of cold methanol and 5 μ l of internal standards (deuterium labeled compounds). Samples were centrifuged at 900 g for 15 min at 4°C, supernatants were diluted in H₂O to 2 ml and submitted to solid phase extraction using a HRX 96-well plate (50 mg/well, Macherey Nagel, Germany). LC-MS/MS analyses were performed on a Agilent 1290 Infinity HPLC system coupled to an ESI-triple quadrupole G6460 mass spectrometer (Agilent Technologies) as previously described (43).

In Vitro Infection Protocol

HD11 cells were seeded in 12-well plates at 5×10^5 cells per well and incubated at 40°C under 5% CO₂ overnight. APEC strains BEN2908 [O2:K1:H5, a nalidixic acid-resistant derivative of strain MT78 which was isolated from the trachea of a chicken with respiratory infection (44)] and BEN3421 (O2:K1:H4, a nalidixic acid- and tetracycline-resistant strain isolated from a chicken presenting colibacillosis in 2009 in France) were utilized for infection. Cell monolayers were infected with mid-log-phase bacteria at an MOI of 10 and incubated for 1 h in cell culture medium without FCS. Cells were washed and remaining extracellular bacteria were killed by incubation in complete medium containing gentamicin (100 μ g/ml) for 1 h 30 min. A group of cells were then washed and lysed with sterile water for 30 min at 4°C to evaluate the percentage of invasion through bacterial enumeration by viable counts on LB agar plates. Percentage of invasion ($T = 0$) was calculated as the number of intracellular bacteria divided by the number of bacteria in the inoculum. For the other groups, culture medium was changed, and gentamicin concentration was reduced to 10 μ g/ml (a concentration known to kill BEN2908 and BEN3421) to avoid putative accumulation within cells during prolonged incubation. PAF (1 or 10 μ M) was

added together with gentamicin. After 6 or 24 h, supernatant was recovered for nitrite dosage and stocked at -20°C . Cells were washed and lysed with RNA lysis buffer or sterile water for 30 min at 4°C . Intracellular bacteria were enumerated by viable counts on LB agar plates. Percentage of growth was calculated as the number of intracellular bacteria at a given time point divided by the number of bacteria at $T = 0$.

In Vivo Infection and Chicken Tissue Samples

Four-week-old B13/B13 histocompatible chickens were infected intratracheally with 10^9 colony-forming units of the highly adhesive/invasive APEC strain BEN2908 in 200 μl of sterile endotoxin-free NaCl 0.9% solution. The inoculum was prepared with bacteria in the mid-log-phase of growth. The control group was inoculated with sterile NaCl 0.9% solution. Lung and liver samples were collected aseptically from each chicken 6 h post bacterial inoculation. Tissue samples for RNA extraction were snap frozen in liquid nitrogen and stocked at -80°C . For bacterial quantification, lung samples were weighed, placed in gentleMACS™ C tubes containing sterile PBS, and homogenized using a gentle MACS™ Dissociator (Miltenyi Biotec, Germany). Dilutions of lung homogenates were plated onto Drigalski agar plates supplemented with nalidixic acid (30 $\mu\text{g}/\text{ml}$) for bacterial quantification, and 1 ml was incubated in brain heart infusion for qualitative detection of *E. coli*.

Cell Permeability Assay

Primary chAEC were seeded at a density of 5×10^4 cells on fibronectin-coated Falcon® Permeable Support (8.0 μm pore) PET membranes (Corning, NY, USA), placed in 24-well plates containing EC medium and cultured for 72 h at 40°C under 5% CO_2 . Then, chAEC monolayers in the upper chamber received different stimuli and, at different time points, EC growth medium containing 1 mg/ml FITC-dextran (40,000 MW) (Sigma-Aldrich, UK) was added followed by further incubation for 30 min at 40°C under 5% CO_2 . Then, 50 μl were collected from the lower chamber, and fluorescence signals relating to FITC-dextran passage from the upper to the lower chamber were measured using a GloMax®-Multi Detection System plate reader (Promega, UK). Cell layer permeability was expressed as fold increase in FITC-dextran fluorescence as compared to the control group (vehicle).

Protein Homology Analysis

Protein multiple sequence alignment for PAFR and LPCAT 2 (LPCAT2) from chicken, mouse, and human were performed using ClustalW2 through Geneious R10 software (Biomatters Ltd., New Zealand). Homology analyses of transmembrane and cytoplasmic domains for PAFR or LPCAT2, including exclusive LPCAT2 enzymatic-related domains, were performed using Uniprot Blast.² Predicted phosphorylation sites were identified using NetPhos 3.1 server.³ Only intracellular predicted sites with a score above 0.650 and conserved in the three analyzed sequences

were annotated. Predicted N-glycosylation sites conserved in the three analyzed sequences were identified using NetNGlyc 1.0 server.⁴ NCBI Reference Sequence for these proteins were as follows: PAFR (*Homo sapiens*: NP_001158195, *Mus musculus*: NP_001074680, and *Gallus gallus*: XP_004947758) and LPCAT2 (*H. sapiens*: NP_060309, *M. musculus*: NP_766602, and *G. gallus*: NP_001025739).

Statistical Analysis

Comparisons between two groups were performed using a two-tailed unpaired Student's *t* test. Multiple groups were compared using a one-way analysis of variance followed by a Newman-Keuls multiple comparison *post hoc* test. Values for all measurements are expressed as mean \pm SEM. $P < 0.05$ was considered statistically significant. Statistical analyses were performed using GraphPad Prism 5.0 (GraphPad Software, USA). Data are representative of at least two independent experiments.

RESULTS

Protein Homology Analysis of PAFR and LPCAT2

Platelet-activating factor receptor is a class A rhodopsin-like GPCR. Homology analysis showed that chicken PAFR possesses protein sequence identity of 60.3 and 58.0% as compared to human or mouse PAFR protein sequences, respectively. Within transmembrane domains (TMD) 2, 3, and 7 sequence identity may reach 80–90%. Multiple sequence alignments of chicken PAFR to human and mouse counterparts revealed that these proteins have seven hydrophobic α -helical domains and conserved amino acids (or amino-acid motifs) commonly found in members of the rhodopsin family GPCRs: an asparagine (N) in the first TMD, aspartate (D) in the second and seventh TMD, a LXXXD motif in the second TMD, tryptophan (W) in the fourth TMD, conserved cysteine (C) residues in the first two extracellular loops (believed to form an S–S bridge and contribute to protein stability), a FXP motif in the sixth TMD (also found as WXP in other GPCRs), and a DPXXY motif in the seventh TMD (also found as NPXXY in other GPCRs) (Figure S1 in Supplementary Material). Importantly, in the chicken PAFR sequence the residues corresponding to the DRY motif are NRE, lacking the conserved tyrosine residue in the second intracellular loop. The DRY motif is a GPCR signature triplet sequence found downstream of the third TMD, which is believed to be involved in G-protein coupling (45–48). Chicken PAFR also contains two conserved predicted sites for N-linked glycosylation and nine threonine (T) or serine (S) residues identified as potential sites for phosphorylation by protein kinase A, G, or C (Figure S1 in Supplementary Material).

Chicken LPCAT2 homology analysis revealed a protein sequence identity of 62.5% as compared to both human and mouse LPCAT2 protein sequences. Multiple sequence analysis also showed that chicken LPCAT2 contains a large conserved cytoplasmic domain, one conserved TMD (different from human

²<http://www.uniprot.org/blast/>.

³<http://www.cbs.dtu.dk/services/NetPhos/>.


⁴<http://www.cbs.dtu.dk/services/NetNGlyc/>.

LPCAT2, which possesses three TMD) and two conserved motifs within an acyltransferase conserved region: HXXXXD and EGTC, believed to be essential for acyltransferase activity (49) (Figure S2 in Supplementary Material). Also, chicken LPCAT2 possesses a conserved putative EF-hand domain, similar to many Ca^{2+} -binding proteins, which contains eight predicted Ca^{2+} binding sites (13, 49). The presence of a conserved C-terminal sequence motif KKXX suggests that LPCAT2 is likely to be localized to the ER in chickens (50) (Figure S2 in Supplementary Material).

PAFR and LPCAT2 Are Expressed in Selected Chicken Tissues and Macrophages

To gain further insights into PAF function in chickens, we started out to evaluate gene expression for PAFR/*PTAFR* and

LPCAT2/*LPCAT2* in selected healthy chicken tissues (liver, lungs, and bone marrow) and unstimulated macrophages/monocytes by qRT-PCR. We found that PAFR is abundantly expressed in chicken tissues as compared to housekeeping genes, with a two-fold higher expression found in the bone marrow as compared to other organs (Figure 1A). In monocytes/macrophages, PAFR is expressed at similar levels, suggesting that a cell line (HD11), BMDM, and PBMC consistently express PAFR (Figure 1A). LPCAT2 gene expression followed the same pattern as for PAFR expression, with LPCAT2 being highly expressed in the bone marrow as compared to other organs (Figure 1B). In monocytes and macrophages, LPCAT2 was found to be consistently expressed, which suggests that chicken macrophages are able to produce PAF (Figure 1B). Interestingly, following chCSF-1 treatment to promote macrophage differentiation, bone marrow mononuclear cells quickly showed downregulation of PAFR



expression (threefold), which returned to expected (BMDM) levels with the cells getting reprogrammed to differentiate into macrophages (Figure 1C).

PAF Induces Intracellular Calcium Increase in Chicken Macrophages

We next investigated whether chicken macrophages were responsive to exogenous PAF (PAF C16). Intracellular calcium mobilization is a *bona fide* parameter to evaluate early events following GPCR binding and activation. HD11 macrophages were stimulated with PAF (0.1–10 μ M), and intracellular calcium mobilization was measured using a fluorescent calcium probe. PAF induced a quick concentration-dependent increase in intracellular calcium signal which returned to levels close to vehicle-treated wells after 213 s (Figure 1D). While not interfering with intracellular calcium signal alone, pretreatment with canonical PAFR antagonists PCA 4248 or WEB 2086 efficiently blocked intracellular calcium mobilization in chicken macrophages upon exposure to PAF (10 μ M) (Figure 1E). This suggests that PAF seems to act in a specific receptor (PAFR)-mediated fashion in chicken macrophages as in mammalian cells.

PAF Binds to PAFR in Chicken Macrophages

To characterize PAFR protein expression and modulation by exogenous PAF in chicken primary macrophages, IF, WB, and flow cytometry (FACS) analyses using a commercially available antibody predicted to cross-react with different species were performed. PAFR was clearly expressed in chicken BMDM at protein level (Figure 2A, upper panels) as was validated with the isotype control. Interestingly, the addition of PAF (10 μ M) for 2 h led to PAFR downregulation, as seen by the loss of fluorescent signal within stained macrophages (Figure 2A, middle panels). Following PAFR activation, cells become rapidly desensitized, and this refractory state is dependent on PAFR phosphorylation, internalization, and downregulation (51). PAFR antagonist PCA 4248 did not promote PAFR downregulation alone, but abrogated PAFR downregulation induced by exogenous PAF (Figure 2A, middle and lower panels). This phenomenon was confirmed by FACS analysis, where around 23% of BMDM expressed PAFR and the addition of exogenous PAF for 2 h led to receptor downregulation as seen through changes in PAFR mean fluorescence intensity (Figure 2B). Also, WB analysis confirmed that PAFR is partially degraded upon PAF binding and that PCA 4248 blocked this phenomenon (Figure 2C). Altogether, these data revealed that PAFR is expressed at the protein level and can be downregulated and partially degraded by PAF in chicken macrophages.

In addition to this physiological phenomenon, we observed that PAFR gene expression is increased upon *E. coli* LPS or DB stimulation, but not upon exogenous PAF treatment, for 6 h as compared to vehicle-treated group (Figure 2D), as was assessed by IF analysis (Figure 2A, lower right panel). LPCAT2 gene expression was also upregulated by these pro-inflammatory bacterial stimuli, but not by exogenous PAF, which suggests that bacteria-associated molecular patterns are positive signals for

the upregulation of PAFR and LPCAT2 genes, the latter being important for the metabolism of PAF during inflammation.

PAF Promotes Increased Phagocytosis and ROS Release by Chicken Macrophages

To further address the functional role of PAF in chicken primary macrophages, we first assessed whether exogenous PAF were cytotoxic to these cells. We showed that PAF (1 and 10 μ M) was not cytotoxic to chicken BMDM after 6 h stimulation as evaluated through annexin V/PI double-staining strategy (Figure 3A) and caspase 3–7 activation assay (Figure 3B), both canonical markers for early and late cell apoptosis. On the other hand, *E. coli* LPS and DB were cytotoxic to chicken BMDM within the same experimental settings.

We next asked whether PAF could enhance BMDM phagocytosis of fluorescent zymosan beads. We observed that PAF (10 μ M) treatment for 1 h significantly enhanced chicken macrophage phagocytosis and that PAFR antagonist PCA 4248 abrogated this phenomenon (Figure 3C).


In another biologically relevant assay, we assessed whether PAF induces ROS release by chicken macrophages. Using a fluorescent probe, we showed that both PAF (10 μ M) and *E. coli* LPS stimulation for 2 h promoted enhanced ROS release by chicken BMDM. However, PAFR antagonist PCA 4248 reverted ROS release only in PAF-treated BMDM, revealing that ROS release is induced by PAF in a PAFR-dependent manner (Figures 3D,E). Altogether these data suggest that PAF promotes two acute macrophage-related inflammatory events in a PAFR-dependent fashion and independent of any cytotoxic effect in chicken macrophages.

PAF Contributes to LPS Pro-inflammatory Response in Chicken Macrophages

A previous study showed that rabbit alveolar macrophages cytokine production in response to endotoxin is PAF dependent (21). We therefore asked whether endogenous PAF production contributes to the LPS-induced pro-inflammatory response in chicken macrophages at 6 h. We observed that PAFR blockade with PCA 4248 or the addition of recombinant human PAF-AH (which degrades extracellular PAF) together with LPS led to significantly reduced NO production and reduced expression of pro-inflammatory genes such as iNOS/NOS2, COX-2/PTGS2, and IL-1 β /IL1B as compared to groups treated with LPS alone (Figures 4A–D). These data suggest that PAF produced by chicken macrophages upon LPS stimulation contributes to the overall pro-inflammatory effects of LPS *via* PAFR.

PAF Amplifies the Inflammatory Response Induced by *E. coli* LPS

We next asked whether PAF could directly induce the production or expression of inflammatory mediators by chicken macrophages. Starting with NO, we showed that exogenous PAF (0.1–10 μ M) alone is not able to induce NO production by chicken BMDM, whereas LPS is a potent NO production inducer (Figure 4E). However, when added together with LPS, PAF (10 μ M) promoted




a significant increase in LPS-induced NO production by chicken macrophages as compared to groups receiving LPS alone (Figure 4E). We also demonstrated this phenomenon in BMDM from another chicken line (PA12) (Figure 4F) and in PBMC of the B13 chicken line (Figure 4G), suggesting that PAF potentiation of LPS-induced NO production is biologically consistent among chicken macrophages/monocytes. In these latter cells, we also observed the aforementioned contribution of endogenous PAF to LPS-induced production of pro-inflammatory mediators

(here NO) through PAFR blockade with PCA 4248 upon LPS-stimulation (Figures 4F,G), thus corroborating our previous findings.

PAF Potentiation Effects Are Mediated by PI3K and CaMK II Signaling Pathways

To gain further insight into the mechanisms through which PAF potentiates the pro-inflammatory effects of LPS, we first




confirmed that exogenous PAF (10 μ M) not only potentiates the production of NO induced by *E. coli* LPS by chicken BMDM but also enhances iNOS/NOS2, IL-1 β /IL1B, and COX-2/PTGS2 gene expression, without being able to induce the expression of these genes alone (Figures 5A–D). In line with COX-2/PTGS2 upregulation, exogenous PAF potentiates PGE2 (which is metabolized by COX-2) production by chicken macrophages upon LPS stimulation, despite its inability to directly induce the production of this eicosanoid (Figure 5E).

Signaling pathways such as those mediated by PI3K, calcium CaMK II, and mitogen-activated protein kinase (MAPK) have been shown to be involved in mediating the effects of PAF in mammalian leukocytes (52, 53). Pretreatment with the PI3K inhibitor WMN (200 nM) or CaMK II inhibitor KN62 (1 μ M) significantly diminished the potentiation effects of exogenous PAF on LPS-induced upregulation of NO production and iNOS/NOS2, IL-1 β /IL1B, and COX-2/PTGS2 gene expression in chicken BMDM (Figures 6A–D). PI3K inhibition, but not CamK II inhibition, also abrogated exogenous PAF potentiation

of zymosan beads phagocytosis by chicken macrophages, leading to a similar level than that observed when PAF signaling is blocked with PAFR antagonist PCA 4248 (Figure 6E). Importantly, these inhibitors showed mild yet statistically significant effects on LPS-induced NO production and upregulation of pro-inflammatory genes expression (Figures 6A–D), in a magnitude that does not appear to interfere with their inhibitory action on PAFR signaling and its potentiation effects as suggested. We did not observe any statistically significant effects in the aforementioned parameters when the p38 MAPK-mediated signaling pathway was blocked using the specific inhibitor SB 203580 (data not shown).

We also confirmed that Akt (which is phosphorylated by PI3K) and CaMK II are expressed in chicken BMDM at protein levels (Figure 6F), although we could not determine the presence of PI3K due to the unavailability of validated commercial antibodies known to recognize this enzyme in chickens. We therefore presume that PAF exerts its pro-phagocytic and potentiating effects over LPS-induced pro-inflammatory responses through



PI3K/Akt- and/or CaMK II-mediated intracellular signaling pathways.

PAF Contributes to Intracellular Bacteria Killing

In addition to its pro-inflammatory effects in chicken macrophages, we assessed whether exogenous PAF could promote increased intracellular bacteria killing in macrophages infected with two phenotypically different strains of APEC. Although both strains belong to the O2 serotype, the highly adhesive/invasive strain BEN2908 was able to better infect HD11 macrophages (1 log difference), as compared to the BEN3421 strain at the same MOI and at the same time points (Figure 7A).

The addition of exogenous PAF following APEC infection for 6 h did not promote increased intracellular APEC killing as compared to infected groups receiving vehicle only. However, at 24 h, PAF (10 μ M) was able to significantly reduce intracellular bacterial load, notably for the highly virulent BEN2908 strain (Figure 7A). This phenomenon was not related to increased PAF-driven macrophage cytotoxicity since infected cells receiving PAF showed the same cell death level as infected cells receiving vehicle at both 6 and 24 h, as was determined by the MTT assay (Figure 7B). As observed before with LPS, the addition of exogenous PAF to APEC infected macrophages led to time-dependent increase in NO production as compared to untreated infected macrophages (Figure 7C), which suggests






FIGURE 5 | Exogenous platelet-activating factor (PAF) potentiates the pro-inflammatory response elicited by lipopolysaccharide (LPS) in chicken primary macrophages in a PAF receptor (PAFR)-dependent manner. After 6 h stimulation with PAF (10 μM) and/or *E. coli* LPS (10 ng/ml) in the presence or absence of PAFR antagonist PCA 4248 (10 μM), B13/B13 histocompatible chicken-derived bone marrow-derived macrophage (BMDM) supernatants were assessed for the presence of (A) nitric oxide, and cell lysates were used to analyze gene expression of (B) iNOS/NOS2, (C) IL-1 β /IL1B, and (D) COX-2/PTGS2 by quantitative real-time PCR. Data are expressed as relative normalized expression (as compared to vehicle control group). (E) Prostaglandin E2 dosage by mass spectrometry in BMDM supernatants following the same aforementioned stimuli for 6 h. Values are mean \pm SEM. ** $P < 0.01$ and *** $P < 0.001$ when compared to negative (vehicle) control groups. * $P < 0.05$ and ## $P < 0.01$ when compared to positive control groups.



that PAF also amplifies the inflammatory response in APEC-infected macrophages. We also confirmed by IF that PAFR is expressed at protein level in HD11 macrophage cell line (Figure 7D). In addition, APEC infected macrophages showed increased upregulation of PAFR/PTAFR and LPCAT2/LPCAT2 gene expression (Figures 7E,F). Overall, we showed that PAF is able to promote increased intracellular bacteria killing together with an increased pro-inflammatory response. Also, we suggest that APEC infection is likely to promote increased PAF signaling and production by chicken macrophages.

PAFR and LPCAT2 Expression Are Upregulated during Pulmonary Colibacillosis

To follow-up on these findings in a more relevant experimental system, we assessed whether PAFR and LPCAT2 genes are overexpressed in chickens during experimental pulmonary colibacillosis. After 6 h following APEC BEN2908 intratracheal instillation (10^9 bacteria/animal), chickens presented around 10^6 bacteria in the lungs (Figure 8A). We next compared



the expression of acute pro-inflammatory mediators with PAFR and *LPCAT2* expression in chicken lungs and liver, two organs where typical colibacillosis lesions are known to develop. We observed that in both the lungs (**Figure 8B**) and liver (**Figure 8C**), the gene expressions of PAFR/*PTAFR* and *LPCAT2*/*LPCAT2* are upregulated together with pro-inflammatory gene expression such as *COX-2*/*PTGS2*, *iNOS*/*NOS2*, and *IL-1 β* /*IL1B* as compared to vehicle-instilled animals. These data suggest that during pulmonary colibacillosis, genes involved in PAF signaling, and metabolism are overexpressed concomitantly to genes related to the acute

inflammatory response, which is in agreement with the data found in APEC-infected macrophages.

PAF Contributes to Inflammation and Cell Permeability in Chicken Endothelial Cells

Platelet-activating factor contribution to vascular physiopathology, such as increased vascular permeability and leukocyte adhesion, is well described in mammals (5, 8). Using a novel chicken endothelial cell line (chAEC), we aimed at addressing PAF contribution to increased EC permeability and inflammatory response in chickens. We first showed that PAFR/*PTAFR*




FIGURE 8 | Overexpression of platelet-activating factor receptor (PAFR) and LPCAT2 is correlated with increased pro-inflammatory gene expression in chicken organs during pulmonary colibacillosis. **(A)** Bacterial load in the lungs of B13/B13 histocompatible white leghorn chickens at 6 h following intratracheal infection with 10⁹ colony forming units of APEC BEN2908 (O2 serotype) strain. **(B)** Lung or **(C)** liver samples from APEC BEN2908-infected chickens were analyzed for PAFR/PTAFR, LPCAT2/LPCAT2, COX-2/PTGS2, iNOS/NOS2, and IL-1 β /IL1B gene expression by quantitative real-time PCR. Data are expressed as relative normalized expression (as compared to vehicle control group). Values are mean \pm SEM. *** P < 0.001 when compared to negative (vehicle) control group. Sample size was six animals per group.

is expressed in chAEC, and its expression is upregulated by LPS or DB (**Figure 9A**). Next, using a transwell system, we demonstrated that exogenous PAF induced increased EC permeability as assessed by FITC-dextran leakage (**Figure 9B**). A quick phenomenon that peaked at 30 min following exposure to PAF (10 μ M) and lasted for up to 4 h (**Figures 9B,C**). Blockade of PAFR with PCA 4248 abrogated FITC-dextran leakage (**Figures 9B,C**). EC permeability was not related to PAF cytotoxic effects on chAEC as assessed by the MTT assay at 6 h (**Figure 9D**). As observed for macrophages, exogenous PAF treatment for 6 h did not induce pro-inflammatory gene expression alone but potentiated the pro-inflammatory effects of *E. coli* LPS or DB as found for IL-1 β /IL1B and CXCL12/IL8L2 (a heterophil-recruiting chemokine) gene expression (**Figures 9E,F**). Blockade of PAFR with PCA 4248 reduced this potentiation to levels seen for LPS or DB alone (**Figures 9E,F**). Interestingly, exogenous PAF treatment alone upregulated E-selectin/SELE and VCAM-1/VCAM1 gene expression in chAEC to levels close to those found for LPS or DB (**Figure 10**). Altogether, these data reveal that PAF is active in chicken endothelial cells and promotes increased cell permeability, potentiates bacterial-derived inflammatory responses, and may indirectly contribute to leukocyte adhesion to the chicken endothelium.


DISCUSSION

The physiological and pathological roles of lipid mediators in livestock animals, such as the chicken, are often neglected and little is known on their presence and mechanisms of action. Here, we provide novel insights into the function of PAF as a pro-inflammatory mediator in chickens and the mechanism through which PAF contributes to chicken macrophage activation and responses to bacterial LPS. In addition, PAF contributes to intracellular bacterial killing and EC dysfunction, two phenomena that may be correlated with the overexpression of PAFR and LPCAT2 in organs of chickens presenting colibacillosis. Overall,

we provided evidence that PAF may be an early orchestrator of the inflammatory response to pathogenic bacteria in chickens.

Protein homology analysis of chicken PAFR revealed that this receptor shares several potentially functional features with its mammalian counterparts (e.g., structure, N-glycosylation, and phosphorylation sites), all important to effective PAFR signaling (8, 51). In addition, we showed that PAF is able to activate chicken macrophage PAFR, and canonical PAFR antagonists, largely tested in mammalian systems (52, 54, 55), were able to abrogate intracellular calcium increase, ROS release, and phagocytosis induced by exogenous PAF. Together with the identification of PAFR at protein level, we provide evidence that PAF is binding to a now identified PAFR in chicken macrophages. It is relevant to mention that PAFR is downregulated (and to some extent degraded) at protein level upon PAF binding in chicken primary macrophages, a well-known phenomenon in mammalian PAFR (51) that supports the present characterization of a chicken PAFR.


In contrast to what was previously observed with mouse peritoneal macrophages (56), PAF is not able to directly induce the expression of certain pro-inflammatory genes (e.g., cytokines) in chicken macrophages, although still contributing to mechanisms potentially independent of gene transcription, such as ROS release. PAF acts as an autocrine mediator of mammalian macrophage activation upon LPS challenge *via* TNF- α expression (21). Also, PAF can be biosynthesized very quickly (5, 8, 11), for example, at the very first moments of host cellular contact with bacterial LPS, which suggests that both recognition systems may operate together at the inflammatory site. Indeed, PAF is able to potentiate the pro-inflammatory response of LPS or bacteria in chicken macrophages. We could not verify whether inflammatory mediators rapidly induced by PAF (e.g., ROS) would contribute to the potentiation of LPS effects, although this hypothesis should not be discarded. Nevertheless, degradation of endogenous PAF with PAF-AH or blockade of PAFR both limit the pro-inflammatory effects of LPS alone, highlighting that LPS inflammatory response is partly dependent on PAF in chicken



macrophages, as was observed in other studies in mammals (21). Therefore, paracrine- or autocrine-produced PAF is likely to play a direct role in the inflammatory response to Gram-negative bacteria-associated molecular patterns in chickens, notably at the toll-like receptor 4 (TLR4) level, responsible for the recognition of LPS. The biological relevance of these findings merits to be further investigated *in vivo*.

While some studies in mammals suggest that PAF exerts its pro-inflammatory effects through activation of p38-MAPK and NF κ B (9, 21), we could not detect signs of a participation of these proteins in our experimental settings (data not shown). However, PI3K/Akt and CaM KII intracellular signaling pathways are involved in the potentiation effects of PAF on LPS stimulation, with PI3K/Akt being also essential to the pro-phagocytic effects of PAF in chicken macrophages. Both

pathways have been shown to participate in different immune responses following PAFR activation in mammals (53, 56–58). CaM KII has been shown to participate in PAF reprogramming of mammalian macrophages, which would provide a stronger inflammatory response to LPS (57). Also, PAFR signaling in peritoneal macrophages requires TLR4, MyD88, and TRIF, in a so-called transactivation phenomenon (56). Although we clearly highlighted the role of CaM KII in PAFR signaling and its contribution to the LPS response, signaling pathways downstream TLR4 in chickens are not fully understood. Indeed, chickens lack a functional LPS-specific TRAM-TRIF signaling pathway, which may explain differences in response to LPS compared with mammalian species (59). Therefore, the mechanism underlying the interactions between PAFR, CaM KII, and TLR4-associated molecules in chickens remains to be



fully elucidated. Of some interest, PI3K/Akt has been shown to be important to neutrophil recruitment in mammals (60). Furthermore, PAF has been shown to be important to neutrophil activation and adhesion to the endothelium (24, 52, 61). In addition to its role in macrophages, as suggested by the data in the present manuscript, the contribution of PAF to heterophil (the avian ortholog of neutrophils) arrest and recruitment *via* PI3K/Akt signaling pathway merits to be investigated.

Platelet-activating factor is directly involved in the physiopathology of cardiovascular diseases of inflammatory origin (5, 8). PAF has also been associated with EC dysfunction (53, 62). The newly established cellular model of primary chicken aortic endothelial cells (chAEC) (42) allowed us to show that PAF may directly induce increased EC monolayer permeability and strongly amplify the chAEC pro-inflammatory response to bacteria-associated molecular patterns. To our knowledge, this is the first study highlighting the role of a lipid mediator in promoting EC dysfunction in chickens. Interestingly, PAF treatment led to the upregulation of genes coding for the leukocyte adhesion molecules VCAM-1 and E-selectin, which would suggest that PAF, produced within the endothelium and adjacent tissues, could favor not only leukocyte activation but also leukocyte arrest and recruitment. Therefore, the cross-talk between PAF, macrophages, and ECs during host-response to Gram-negative bacteria could contribute to local and sustained inflammatory responses in chickens.

The physiopathology of pulmonary avian colibacillosis encompasses different inflammatory mechanisms in which exacerbated vascular leakage, heterophil and monocyte recruitment, and resident macrophage activation are believed to take place and contribute to disease severity in birds unable to control infection

within a safe timeframe (33–35, 63). The contribution of recruited monocytes or tissue-resident macrophages to the host response during pulmonary colibacillosis is still being unveiled, although several studies have suggested that these resident phagocytes are involved in bacteria recognition and killing, amplification of inflammation, and, potentially, heterophil clearance (33–35, 64, 65). Here, we demonstrated that PAF is able to promote increased bacterial clearance by primary chicken macrophages *in vitro*, together with a strong amplification of the inflammatory response. Our gene expression data showed that PAFR and LPCAT2 genes are overexpressed in lungs and liver of APEC-infected chickens at early time points postinfection. While these data reveal that key components of PAF recognition and biosynthesis are locally expressed in parallel to a strong inflammatory response to APEC, we are still unable to define which cell populations are expressing these molecules. However, it is reasonable to assume that PAF is indeed produced and acts within infected sites, thereby amplifying inflammation and/or contributing to bacterial clearance as observed *in vitro*. Both PAFR and LPCAT2 genes are overexpressed in chicken macrophages and chAEC upon LPS stimulation *in vitro*, which suggests that at least these cell types may be involved in the response to colibacillosis *in vivo*.

We could not quantify PAF production in chicken tissues or cell culture supernatants due to several technical issues (i.e., contamination of closely related phospholipids during mass spectrometry analysis). However, LPCAT2 overexpression has been consistently associated with PAF biosynthesis during inflammation, while LPCAT1 activity and expression are independent of inflammatory signals (13, 66). After the activation of TLR4 by LPS, LPCAT2, but not LPCAT1, is activated and upregulated to promote PAF production in mouse macrophages (67). Indeed, LPCAT2 is believed to be the main pro-inflammatory LPCAT in the remodeling pathway for the biosynthesis of PAF in mammals. Our protein homology data suggest that chicken LPCAT2 is closely related to human and mouse counterparts. Within this scenario, we may propose that, upon contact with bacteria or bacteria-associated molecular patterns, resident macrophages, and other unidentified tissue parenchyma cells, may produce PAF. PAF would in turn amplify local inflammation through eicosanoid, cytokine, and chemokine production to facilitate initial bacterial clearance and trigger the inflammatory response. In parallel, adjacent (e.g., parenchymal) and endothelial PAF production would directly or indirectly (e.g., *via* chemokine production) promote increased leukocyte adhesion and recruitment (e.g., heterophils and monocytes), together with increased vascular leakage, therefore, contributing to the orchestration of classical cellular and molecular events that take place at the onset of the inflammatory response. We therefore suggest that PAF might be an important mediator of inflammation in the host response to Gram-negative bacteria causing economically relevant diseases in poultry farms worldwide, such as APEC.

A PAF receptor antagonist, Modipafant, was already used in the past to treat asthma, although its use is currently discontinued (68). There is also an ongoing clinical trial to validate the use of Modipafant as a therapeutic strategy to limit exacerbated inflammation observed in viral hemorrhagic diseases such as Dengue

fever.⁵ Preliminary experiments suggest that PAFR antagonists would be active *in vivo* in chickens (data not shown), and a proof of concept on the use of PAFR antagonists to limit exacerbated inflammation associated with experimental colibacillosis are currently being performed. Although the feasibility of such therapeutic strategy in the poultry industry is still difficult to envisage, these experiments will bring biologically relevant information on how lipid mediators and their receptors could be pharmacologically manipulated in chickens, so as to pave the way to alternative therapies in face of the widespread antibiotic resistance in poultry farms and the lack of efficacious vaccines for certain avian diseases.

In conclusion, we believe that the present study sheds light on the role and function of PAF in bridging endothelium function and macrophage-associated innate immunity in chickens in response to bacteria-derived stimuli, a hitherto unknown mechanism in birds. Also, these activity profiles may be shared by other yet to be identified lipid mediators of inflammation, a line of research that should be of interest to the scientific community in the fields of avian immunology, evolution, and host–pathogen interaction.

ETHICS STATEMENT

All animals used in the avian pathogenic *Escherichia coli* (APEC) infection protocol were treated according to EU recommendations for animal welfare, and the protocol was approved by the French regional ethics committee number 19 (Comité d’Ethique en Expérimentation Animale Val de Loire) under the reference CL2007-44.

⁵<https://clinicaltrials.gov/ct2/show/NCT02569827>.

REFERENCES

- Bennett M, Gilroy DW. Lipid mediators in inflammation. *Microbiol Spectr* (2016) 4(6):1–21. doi:10.1128/microbiolspec.MCHD-0035-2016
- Serhan CN, Haeggstrom JZ, Leslie CC. Lipid mediator networks in cell signaling: update and impact of cytokines. *FASEB J* (1996) 10(10):1147–58.
- Hla T. Genomic insights into mediator lipidomics. *Prostaglandins Other Lipid Mediat* (2005) 77(1–4):197–209. doi:10.1016/j.prostaglandins.2005.06.008
- Serhan CN, Chiang N, Van Dyke TE. Resolving inflammation: dual anti-inflammatory and pro-resolution lipid mediators. *Nat Rev Immunol* (2008) 8(5):349–61. doi:10.1038/nri2294
- Prescott SM, Zimmerman GA, McIntyre TM. Platelet-activating factor. *J Biol Chem* (1990) 265(29):17381–4.
- Ishii S, Kuwaki T, Nagase T, Maki K, Tashiro F, Sunaga S, et al. Impaired anaphylactic responses with intact sensitivity to endotoxin in mice lacking a platelet-activating factor receptor. *J Exp Med* (1998) 187(11):1779–88. doi:10.1084/jem.187.11.1779
- Ishii S, Shimizu T. Platelet-activating factor (PAF) receptor and genetically engineered PAF receptor mutant mice. *Prog Lipid Res* (2000) 39(1):41–82. doi:10.1016/S0163-7827(99)00016-8
- Zimmerman GA, McIntyre TM, Prescott SM, Stafforini DM. The platelet-activating factor signaling system and its regulators in syndromes of inflammation and thrombosis. *Crit Care Med* (2002) 30(5 Suppl):S294–301. doi:10.1097/00003246-200205001-00020
- Chen LW, Lin MW, Hsu CM. Different pathways leading to activation of extracellular signal-regulated kinase and p38 MAP kinase by formyl-methionyl-leucyl-phenylalanine or platelet activating factor in human neutrophils. *J Biomed Sci* (2005) 12(2):311–9. doi:10.1007/s11373-005-1704-1

AUTHOR CONTRIBUTIONS

RG designed and performed the experiments, analyzed data, and wrote the manuscript. DG, NC, AT, and AL performed the experiments and analyzed data. GB and EE performed the experiments. ST, PQ, and CS contributed reagents, materials, intellectual input, and revised the manuscript.

ACKNOWLEDGMENTS

We thank Nathalie Lallier for her technical assistance on the experimental infection model and the personnel from the experimental unit PFIE (Plateforme d’Infectiologie Expérimentale) at the Centre INRA Val de Loire (Nouzilly, France). We also thank Prof. Mauro M. Teixeira (UFMG, Brazil) and Dr. Cristiana C. Garcia (IOC/Fiocruz, Brazil) for valuable discussions on the experimental design setup.

FUNDING

This work was supported by the French National Institute for Agricultural Research (INRA) through a young researcher grant (to RG) from the Animal Health division (Département Santé Animale, SA).

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at <http://www.frontiersin.org/articles/10.3389/fvets.2017.00226/full#supplementary-material>.

- McLaughlin NJ, Banerjee A, Kelher MR, Gamboni-Robertson F, Hamiel C, Sheppard FR, et al. Platelet-activating factor-induced clathrin-mediated endocytosis requires beta-arrestin-1 recruitment and activation of the p38 MAPK signalosome at the plasma membrane for actin bundle formation. *J Immunol* (2006) 176(11):7039–50. doi:10.4049/jimmunol.176.11.7039
- Snyder F. Platelet-activating factor and its analogs: metabolic pathways and related intracellular processes. *Biochim Biophys Acta* (1995) 1254(3):231–49. doi:10.1016/0005-2760(94)00192-2
- Kihara Y, Ishii S, Kita Y, Toda A, Shimada A, Shimizu T. Dual phase regulation of experimental allergic encephalomyelitis by platelet-activating factor. *J Exp Med* (2005) 202(6):853–63. doi:10.1084/jem.20050660
- Shindou H, Hishikawa D, Nakanishi H, Harayama T, Ishii S, Taguchi R, et al. A single enzyme catalyzes both platelet-activating factor production and membrane biogenesis of inflammatory cells. Cloning and characterization of acetyl-CoA:LYSO-PAF acetyltransferase. *J Biol Chem* (2007) 282(9):6532–9. doi:10.1074/jbc.M609641200
- Tarui M, Shindou H, Kumagai K, Morimoto R, Harayama T, Hashidate T, et al. Selective inhibitors of a PAF biosynthetic enzyme lysophosphatidylcholine acyltransferase 2. *J Lipid Res* (2014) 55(7):1386–96. doi:10.1194/jlr.M049205
- Arai H, Hattori M, Adachi H, Hattori K, Tsujimoto M, Inoue K. Paf acetylhydrolase as a novel calcium-independent phospholipase-A2. *J Cell Biochem* (1994) 18(D):23.
- Shindou H, Ishii S, Uozumi N, Shimizu T. Roles of cytosolic phospholipase A(2) and platelet-activating factor receptor in the Ca-induced biosynthesis of PAF. *Biochem Biophys Res Commun* (2000) 271(3):812–7. doi:10.1006/bbrc.2000.2723
- Shindou H, Ishii S, Yamamoto M, Takeda K, Akira S, Shimizu T. Priming effect of lipopolysaccharide on acetyl-coenzyme A:lyso-platelet-activating

- factor acetyltransferase is MyD88 and TRIF independent. *J Immunol* (2005) 175(2):1177–83. doi:10.4049/jimmunol.175.2.1177
18. Lee T, Lenihan DJ, Malone B, Roddy LL, Wasserman SI. Increased biosynthesis of platelet-activating factor in activated human eosinophils. *J Biol Chem* (1984) 259(9):5526–30.
 19. Owen JS, Baker PR, O'Flaherty JT, Thomas MJ, Samuel MP, Wooten RE, et al. Stress-induced platelet-activating factor synthesis in human neutrophils. *Biochim Biophys Acta* (2005) 1733(2–3):120–9. doi:10.1016/j.bbali.2004.12.016
 20. Maier RV, Hahnel GB, Fletcher JR. Platelet-activating factor augments tumor necrosis factor and procoagulant activity. *J Surg Res* (1992) 52(3):258–64. doi:10.1016/0022-4804(92)90083-C
 21. Bulger EM, Arbabi S, Garcia I, Maier RV. The macrophage response to endotoxin requires platelet activating factor. *Shock* (2002) 17(3):173–9. doi:10.1097/00024382-200203000-00003
 22. Caillard CG, Mondot S, Zundel JL, Julou L. Hypotensive activity of PAF-acether in rats. *Agents Actions* (1982) 12(5–6):725–30. doi:10.1007/BF01965093
 23. Bessin P, Bonnet J, Apffel D, Soulard C, Desgroux L, Pelas I, et al. Acute circulatory collapse caused by platelet-activating factor (PAF-acether) in dogs. *Eur J Pharmacol* (1983) 86(3–4):403–13. doi:10.1016/0014-2999(83)90190-5
 24. Sterner-Kock A, Braun RK, Schrenzel MD, Hyde DM. Recombinant tumour necrosis factor-alpha and platelet-activating factor synergistically increase intercellular adhesion molecule-1 and E-selectin-dependent neutrophil adherence to endothelium in vitro. *Immunology* (1996) 87(3):454–60.
 25. Kulikov VI, Muzya GI. Ether lipids and platelet-activating factor: evolution and cellular function. *Biochemistry (Moscow)* (1997) 62(10):1103–8.
 26. Bussolino F, Torelli S, Stefanini E, Gremo F. Platelet-activating factor production occurs through stimulation of cholinergic and dopaminergic receptors in the chick retina. *J Lipid Mediat* (1989) 1(5):283–8.
 27. Cox CP. Activation of washed chicken thrombocytes by 1-O-hexadecyl/octadecyl-2-acetyl-sn-glycero-3-phosphorylcholine (platelet-activating factor). *Comp Biochem Physiol A Comp Physiol* (1985) 82(1):145–51. doi:10.1016/0300-9629(85)90719-4
 28. Cabot MC, Faulkner LA, Lackey RJ, Snyder F. Vertebrate class distribution of 1-alkyl-2-acetyl-Sn-glycero-3-phosphocholine acetylhydrolase in serum. *Comp Biochem Physiol B* (1984) 78(1):37–40. doi:10.1016/0305-0491(84)90141-X
 29. Tjoelker LW, Eberhardt C, Unger J, Trong HL, Zimmerman GA, McIntyre TM, et al. Plasma platelet-activating factor acetylhydrolase is a secreted phospholipase A2 with a catalytic triad. *J Biol Chem* (1995) 270(43):25481–7. doi:10.1074/jbc.270.43.25481
 30. Whelan J, Golemboski KA, Broughton KS, Kinsella JE, Dietert RR. Characterization of leukotriene production in vivo and in vitro in resident and elicited peritoneal macrophages in chickens and mice. *Prostaglandins Leukot Essent Fatty Acids* (1997) 56(1):41–9. doi:10.1016/S0952-3278(97)90523-8
 31. Kwok AH, Wang Y, Wang CY, Leung FC. Molecular cloning and characterization of chicken prostaglandin E receptor subtypes 2 and 4 (EP2 and EP4). *Gen Comp Endocrinol* (2008) 157(2):99–106. doi:10.1016/j.ygcen.2008.04.001
 32. Kogut MH, He H, Genovese KJ. Bacterial toll-like receptor agonists induce sequential NF-kappaB-mediated leukotriene B4 and prostaglandin E2 production in chicken heterophils. *Vet Immunol Immunopathol* (2012) 145(1–2):159–70. doi:10.1016/j.vetimm.2011.11.003
 33. Guabiraba R, Schouler C. Avian colibacillosis: still many black holes. *FEMS Microbiol Lett* (2015) 362(15):fnv118. doi:10.1093/femsle/fnv118
 34. Matthijs MG, Ariaans MP, Dwars RM, van Eck JH, Bouma A, Stegeman A, et al. Course of infection and immune responses in the respiratory tract of IBV infected broilers after superinfection with *E. coli*. *Vet Immunol Immunopathol* (2009) 127(1–2):77–84. doi:10.1016/j.vetimm.2008.09.016
 35. Horn F, Correa AM, Barbieri NL, Glodde S, Weyrauch KD, Kaspers B, et al. Infections with avian pathogenic and fecal *Escherichia coli* strains display similar lung histopathology and macrophage apoptosis. *PLoS One* (2012) 7(7):e41031. doi:10.1371/journal.pone.0041031
 36. McManus LM, Hanahan DJ, Demopoulos CA, Pinckard RN. Pathobiology of the intravenous infusion of acetyl glyceryl ether phosphorylcholine (AGEPC), a synthetic platelet-activating factor (PAF), in the rabbit. *J Immunol* (1980) 124(6):2919–24.
 37. Soares AC, Pinho VS, Souza DG, Shimizu T, Ishii S, Nicoli JR, et al. Role of the platelet-activating factor (PAF) receptor during pulmonary infection with gram negative bacteria. *Br J Pharmacol* (2002) 137(5):621–8. doi:10.1038/sj.bjp.0704918
 38. Garcia CC, Russo RC, Guabiraba R, Fagundes CT, Polidoro RB, Tavares LP, et al. Platelet-activating factor receptor plays a role in lung injury and death caused by influenza A in mice. *PLoS Pathog* (2010) 6(11):e1001171. doi:10.1371/journal.ppat.1001171
 39. Beug H, von Kirchbach A, Doderlein G, Conscience JF, Graf T. Chicken hematopoietic cells transformed by seven strains of defective avian leukemia viruses display three distinct phenotypes of differentiation. *Cell* (1979) 18(2):375–90. doi:10.1016/0092-8674(79)90057-6
 40. Garceau V, Smith J, Paton IR, Davey M, Fares MA, Sester DP, et al. Pivotal advance: avian colony-stimulating factor 1 (CSF-1), interleukin-34 (IL-34), and CSF-1 receptor genes and gene products. *J Leukoc Biol* (2010) 87(5):753–64. doi:10.1189/jlb.0909624
 41. Kogut MH, McGruder ED, Hargis BM, Corrier DE, Deloach JR. Characterization of the pattern of inflammatory cell influx in chicks following the intraperitoneal administration of live *Salmonella enteritidis* and *Salmonella enteritidis*-immune lymphokines. *Poult Sci* (1995) 74(1):8–17. doi:10.3382/ps.0740008
 42. Lion A, Richard M, Esnault E, Kut E, Soubieux D, Guillory V, et al. Productive replication of avian influenza viruses in chicken endothelial cells is determined by hemagglutinin cleavability and is related to innate immune escape. *Virology* (2017) 513:29–42. doi:10.1016/j.virol.2017.10.007
 43. Le Faouder P, Baillif V, Spreadbury I, Motta JP, Rousset P, Chene G, et al. LC-MS/MS method for rapid and concomitant quantification of pro-inflammatory and pro-resolving polyunsaturated fatty acid metabolites. *J Chromatogr B Analyt Technol Biomed Life Sci* (2013) 932:123–33. doi:10.1016/j.jchromb.2013.06.014
 44. Dho M, Lafont JP. *Escherichia coli* colonization of the trachea in poultry: comparison of virulent and avirulent strains in gnotoxenic chickens. *Avian Dis* (1982) 26(4):787–97. doi:10.2307/1589865
 45. Van Rhee AM, Fischer B, Van Galen PJ, Jacobson KA. Modelling the P2Y purinoceptor using rhodopsin as template. *Drug Des Discov* (1995) 13(2):133–54.
 46. Bockaert J, Pin JP. Molecular tinkering of G protein-coupled receptors: an evolutionary success. *EMBO J* (1999) 18(7):1723–9. doi:10.1093/emboj/18.7.1723
 47. Chung DA, Wade SM, Fowler CB, Woods DD, Abada PB, Mosberg HI, et al. Mutagenesis and peptide analysis of the DRY motif in the alpha2A adrenergic receptor: evidence for alternate mechanisms in G protein-coupled receptors. *Biochem Biophys Res Commun* (2002) 293(4):1233–41. doi:10.1016/S0006-291X(02)00357-1
 48. Boels K, Schaller HC. Identification and characterisation of GPR100 as a novel human G-protein-coupled bradykinin receptor. *Br J Pharmacol* (2003) 140(5):932–8. doi:10.1038/sj.bjp.0705521
 49. Lewin TM, Wang P, Coleman RA. Analysis of amino acid motifs diagnostic for the sn-glycerol-3-phosphate acyltransferase reaction. *Biochemistry* (1999) 38(18):5764–71. doi:10.1021/bi982805d
 50. Shikano S, Li M. Membrane receptor trafficking: evidence of proximal and distal zones conferred by two independent endoplasmic reticulum localization signals. *Proc Natl Acad Sci U S A* (2003) 100(10):5783–8. doi:10.1073/pnas.1031748100
 51. Dupre DJ, Chen Z, Le Gouill C, Theriault C, Parent JL, Rola-Pleszczynski M, et al. Trafficking, ubiquitination, and down-regulation of the human platelet-activating factor receptor. *J Biol Chem* (2003) 278(48):48228–35. doi:10.1074/jbc.M304082200
 52. Hidalgo MA, Ojeda F, Eyre P, LaBranche TP, Smith C, Hancke JL, et al. Platelet-activating factor increases pH(i) in bovine neutrophils through the PI3K-ERK1/2 pathway. *Br J Pharmacol* (2004) 141(2):311–21. doi:10.1038/sj.bjp.0705590
 53. Ko HM, Kang JH, Choi JH, Park SJ, Bai S, Im SY. Platelet-activating factor induces matrix metalloproteinase-9 expression through Ca(2+)- or PI3K-dependent signaling pathway in a human vascular endothelial cell line. *FEBS Lett* (2005) 579(28):6451–8. doi:10.1016/j.febslet.2005.10.027
 54. Stewart AG, Dubbin PN, Harris T, Dusting GJ. Platelet-activating factor may act as a second messenger in the release of icosanoids and superoxide anions from leukocytes and endothelial cells. *Proc Natl Acad Sci U S A* (1990) 87(8):3215–9. doi:10.1073/pnas.87.8.3215
 55. Garcia CC, Guabiraba R, Soriani FM, Teixeira MM. The development of anti-inflammatory drugs for infectious diseases. *Discov Med* (2010) 10(55):479–88.
 56. Agrawal V, Jaiswal MK, Ilievski V, Beaman KD, Jilling T, Hirsch E. Platelet-activating factor: a role in preterm delivery and an essential interaction with

- toll-like receptor signaling in mice. *Biol Reprod* (2014) 91(5):119. doi:10.1095/biolreprod.113.116012
57. Cuschieri J, Bulger E, Garcia I, Jelacic S, Maier RV. Calcium/calmodulin-dependent kinase II is required for platelet-activating factor priming. *Shock* (2005) 23(2):99–106. doi:10.1097/01.shk.0000148075.19190.db
58. Chen J, Lan T, Zhang W, Dong L, Kang N, Zhang S, et al. Platelet-activating factor receptor-mediated PI3K/AKT activation contributes to the malignant development of esophageal squamous cell carcinoma. *Oncogene* (2015) 34(40):5114–27. doi:10.1038/onc.2014.434
59. Keestra AM, van Putten JP. Unique properties of the chicken TLR4/MD-2 complex: selective lipopolysaccharide activation of the MyD88-dependent pathway. *J Immunol* (2008) 181(6):4354–62. doi:10.4049/jimmunol.181.6.4354
60. Pinho V, Russo RC, Amaral FA, de Sousa LP, Barsante MM, de Souza DG, et al. Tissue- and stimulus-dependent role of phosphatidylinositol 3-kinase isoforms for neutrophil recruitment induced by chemoattractants in vivo. *J Immunol* (2007) 179(11):7891–8. doi:10.4049/jimmunol.179.11.7891
61. Ferby IM, Waga I, Sakanaka C, Kume K, Shimizu T. Wortmannin inhibits mitogen-activated protein kinase activation induced by platelet-activating factor in guinea pig neutrophils. *J Biol Chem* (1994) 269(48):30485–8.
62. Humphrey DM, McManus LM, Hanahan DJ, Pinckard RN. Morphologic basis of increased vascular permeability induced by acetyl glyceryl ether phosphorylcholine. *Lab Invest* (1984) 50(1):16–25.
63. Antao EM, Glodde S, Li G, Sharifi R, Homeier T, Laturnus C, et al. The chicken as a natural model for extraintestinal infections caused by avian pathogenic *Escherichia coli* (APEC). *Microb Pathog* (2008) 45(5–6):361–9. doi:10.1016/j.micpath.2008.08.005
64. Reese S, Dalamani G, Kaspers B. The avian lung-associated immune system: a review. *Vet Res* (2006) 37(3):311–24. doi:10.1051/vetres:2006003
65. de Geus ED, Jansen CA, Vervelde L. Uptake of particulate antigens in a nonmammalian lung: phenotypic and functional characterization of avian respiratory phagocytes using bacterial or viral antigens. *J Immunol* (2012) 188(9):4516–26. doi:10.4049/jimmunol.1200092
66. Harayama T, Shindou H, Ogasawara R, Suwabe A, Shimizu T. Identification of a novel noninflammatory biosynthetic pathway of platelet-activating factor. *J Biol Chem* (2008) 283(17):11097–106. doi:10.1074/jbc.M708909200
67. Morimoto R, Shindou H, Oda Y, Shimizu T. Phosphorylation of lysophosphatidylcholine acyltransferase 2 at Ser34 enhances platelet-activating factor production in endotoxin-stimulated macrophages. *J Biol Chem* (2010) 285(39):29857–62. doi:10.1074/jbc.M110.147025
68. Kuitert LM, Angus RM, Barnes NC, Barnes PJ, Bone ME, Chung KF, et al. Effect of a novel potent platelet-activating factor antagonist, modipafant, in clinical asthma. *Am J Respir Crit Care Med* (1995) 151(5):1331–5. doi:10.1164/ajrccm.151.5.7735582

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

The reviewer TK and handling editor declared their shared affiliation.

Copyright © 2017 Garrido, Chanteloup, Trotereau, Lion, Bailleul, Esnault, Trapp, Quéré, Schouler and Guabiraba. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.