



Draft genome sequence and characterization of commensal *Escherichia coli* strain BG1 isolated from bovine gastro-intestinal tract

Audrey Segura, Pauline Auffret, Christophe C. Klopp, Yolande Bertin, Evelyne Forano

► To cite this version:

Audrey Segura, Pauline Auffret, Christophe C. Klopp, Yolande Bertin, Evelyne Forano. Draft genome sequence and characterization of commensal *Escherichia coli* strain BG1 isolated from bovine gastro-intestinal tract. *Standards in Genomic Sciences*, 2017, 12 (1), 10.1186/s40793-017-0272-0 . hal-02626891

HAL Id: hal-02626891

<https://hal.inrae.fr/hal-02626891>

Submitted on 26 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

EXTENDED GENOME REPORT

Open Access



Draft genome sequence and characterization of commensal *Escherichia coli* strain BG1 isolated from bovine gastro-intestinal tract

Audrey Segura^{1*} , Pauline Auffret^{1†}, Christophe Klopp², Yolande Bertin¹ and Evelyne Forano¹

Abstract

Escherichia coli is the most abundant facultative anaerobic bacteria in the gastro-intestinal tract of mammals but can be responsible for intestinal infection due to acquisition of virulence factors. Genomes of pathogenic *E. coli* strains are widely described whereas those of bovine commensal *E. coli* strains are very scarce. Here, we report the genome sequence, annotation, and features of the commensal *E. coli* BG1 isolated from the gastro-intestinal tract of cattle. Whole genome sequencing analysis showed that BG1 has a chromosome of 4,782,107 bp coding for 4465 proteins and 97 RNAs. *E. coli* BG1 belonged to the serotype O159:H21, was classified in the phylogroup B1 and possessed the genetic information encoding “virulence factors” such as adherence systems, iron acquisition and flagella synthesis. A total of 12 adherence systems were detected reflecting the potential ability of BG1 to colonize different segments of the bovine gastro-intestinal tract. *E. coli* BG1 is unable to assimilate ethanolamine that confers a nutritional advantage to some pathogenic *E. coli* in the bovine gastro-intestinal tract. Genome analysis revealed the presence of i) 34 amino acids change due to non-synonymous SNPs among the genes encoding ethanolamine transport and assimilation, and ii) an additional predicted alpha helix inserted in cobalamin adenosyltransferase, a key enzyme required for ethanolamine assimilation. These modifications could explain the incapacity of BG1 to use ethanolamine. The BG1 genome can now be used as a reference (control strain) for subsequent evolution and comparative studies.

Keywords: *Escherichia coli*, Commensal, Bovine, Gastro-intestinal tract, Whole genome sequencing, Virulence factors, Ethanolamine

Introduction

Escherichia coli is a common inhabitant of the gastro-intestinal tract of humans and animals [1]. In particular, *E. coli* is typically the most common facultative anaerobe in the lower intestine of mammals and its presence in the environment is usually considered to reflect fecal contamination [1, 2]. The *E. coli* population is multiclonal and fluctuates in its predominance depending on diet, exposure to antibiotics or interactions with the host endogenous microbiota [1].

The intestinal microbiota predominantly comprises strict anaerobic bacteria, especially in the colon. *E. coli* exists in a symbiotic relationship with strict anaerobes: *E. coli* ferments monosaccharides generated by the degradation of polysaccharides by anaerobes (*E. coli* being unable to synthesize the necessary hydrolase enzymes) and in turn, *E. coli* is able to consume oxygen and therefore to favor the strict anaerobe multiplication by creating a more anaerobic environment [2, 3]. Similarly, the host-*E. coli* relationship is mutualistic: the intestinal environment promotes efficient *E. coli* survival and multiplication and in turn, the *E. coli* population produces vitamins K and B12, which are required by mammalian hosts, and competitively excludes pathogens from the host intestinal tract [2]. *E. coli* strains are able to colonize various locations in the mammalian gastro-

* Correspondence: audrey.segura@inra.fr

†Equal contributors

¹Université Clermont Auvergne, INRA, MEDIS, F-63000 Clermont-Ferrand, France

Full list of author information is available at the end of the article



intestinal tract, but they are mainly found on the mucus layer used by *E. coli* as an essential nutritional source [4]. Successful colonization of the gastro-intestinal tract by *E. coli* depends upon several factors: competition for nutrients with the autochthonous microbiota, production of adhesins to bring the bacteria closer to the epithelia, penetration of the mucus layer, rapid growth and biofilm formation ability [1, 2, 4]. If *E. coli* growth does not exceed the turnover rate of the mucus layer, the bacterial cells are sloughed off into the intestine lumen and then eliminated in the feces [4]. Therefore, *E. coli* must display metabolic flexibility and grow in biofilm in order to succeed in this very competitive biotope [4].

Although considered as commensal in the mammalian gut, *E. coli* also causes a broad range of intestinal or extra-intestinal diseases due to the acquisition of mobile genetic elements encoding virulence factors. Among pathogenic *E. coli*, STEC is the major food-borne pathogen responsible for hemorrhagic colitis and hemolytic uremic syndrome [5]. In particular, a STEC strain subgroup EHEC belonging mostly to the serotype O157:H7 is responsible for serious public health concern and financial burden [5]. STEC strains are mainly transmitted to humans through contaminated meat or unpasteurized milk consumption [6]. It is of interest to note that healthy ruminants, mainly cattle, are the principal reservoir for *E. coli* O157:H7 strains, but cattle lack the Shiga-toxin vascular receptor, which explains why they are Shiga-toxin tolerant [6].

The cost of whole genome sequencing has decreased drastically and it is now possible to sequence a large number of isolates and use bioinformatic approaches to extract strain relatedness and gene carriage data. *E. coli* strains involved in human infections have been extensively studied and many whole genome sequences of *E. coli* associated with human illness are now available, allowing exploration of pathogenicity processes and identification of virulence factors. Due to cattle STEC dissemination, a significant number of whole genomes of *E. coli* O157:H7 strains isolated from bovine have also been sequenced. While previous genome sequencing efforts with commensal intestinal *E. coli* have focused on human strains [7–9], such data are scarce concerning commensal *E. coli* strains isolated from the bovine gastro-intestinal tract. It would be valuable to have recent and reliable genomic data on bovine commensal strains to be used as reference genomes.

In this study, we report the draft genome sequence and preliminary functional annotation of the commensal *E. coli* strain BG1 isolated from the digestive tract of a cow. The strain BG1 has been previously included in studies concerning the adaptation of pathogenic and commensal *E. coli* strains in the bovine gastro-intestinal tract [10, 11]. This study aimed to characterize the

genomic features of the BG1 strain in order to provide information for future genomic scale (whole genome) comparative analyses. The organism is not part of a larger genomic survey project.

Organism information


Classification and features

As described for the genus *Escherichia*, *E. coli* BG1 is a Gram-negative, rod-shaped bacterium belonging to the *Enterobacteriaceae* family (Table 1). *E. coli* is a facultative anaerobe that is motile by means of flagella (Fig. 1). *E. coli* strains are typically able to grow over a wide temperature range (15–48 °C) with optimum growth

Table 1 Classification and general features of *E. coli* BG1 [58]

MIGS ID	Property	Term	Evidence code ^a
	Classification	Domain <i>Bacteria</i>	TAS [59]
		Phylum <i>Proteobacteria</i>	TAS [60]
		Class <i>Gammaproteobacteria</i>	TAS [61, 62]
		Order <i>"Enterobacteriales"</i>	TAS [63]
		Family <i>Enterobacteriaceae</i>	TAS [64, 65]
		Genus <i>Escherichia</i>	TAS [66, 67]
		Species <i>Escherichia coli</i>	TAS [66, 67]
		Gram stain	Negative
	Cell shape	Rod	IDA, TAS [1]
	Motility	Motile	TAS [1]
	Sporulation	None	TAS [1]
	Temperature range	≈ 15–48 °C	TAS [1]
	Optimum temperature	37–42 °C	TAS [1]
pH range; Optimum	5.5–8.0; 7	TAS [1]	
Carbon source	Carbohydrates, amino acids	IDA, TAS [1]	
MIGS-6	Habitat	Bovine digestive tract	IDA
MIGS-6.3	Salinity	Not reported	
MIGS-22	Oxygen requirement	Facultative anaerobe	TAS [1]
MIGS-15	Biotic relationship	Commensalism	IDA
MIGS-14	Pathogenicity	Non-pathogenic	
MIGS-4	Geographic location	France	
MIGS-5	Sample collection	January 14, 2009	
MIGS-4.1	Latitude	Not reported	
MIGS-4.2	Longitude	Not reported	
MIGS-4.4	Altitude	Not reported	

^aEvidence codes – IDA Inferred from Direct Assay; TAS Traceable Author Statement (i.e., a direct report exists in the literature); NAS Non-traceable Author Statement (i.e., not directly observed for the living, isolated sample, but based on a generally accepted property for the species, or anecdotal evidence). These evidence codes are from the Gene Ontology project [68]



from 37 to 42 $^{\circ}\text{C}$ and within a pH range of 5.5–8.0 (the best growth occurs at pH 7) [1] (Table 1). Like typical members of the *E. coli* species, the commensal strain BG1 utilizes D-glucose, D-mannitol, L-rhamnose, D-saccharose, D-melibiose and L-arabinose. Unlike most pathogenic O157:H7 EHEC strains, the strain BG1 is able to use sorbitol as a carbon source. In addition, *E. coli* BG1 is positive for arginine dihydrolase, ornithine decarboxylase, β -galactosidase and indole production.

In silico serotyping using SerotypeFinder (version 1.1) [12] revealed that *E. coli* BG1 belongs to the serotype O159:H21. The whole genome of *E. coli* BG1 lacked all the genes encoding antimicrobial resistance screened using ResFinder (version 2.1) [13]. *E. coli* strains can be divided into different phylogroups (A, B1, B2, D and E) commonly used to investigate the evolution and diversity of *E. coli* strains [14]. Phylogrouping was performed in silico using the quadruplex method described by Clermont et al. [14] and the primersearch program from the EMBOSS open software suite [15]. *E. coli* BG1 belongs to the phylogroup B1, which is commonly distributed among both bovine commensal and human pathogenic *E. coli* strains [16, 17].

A whole genome phylogenetic analysis based on single nucleotide polymorphism (SNP) differences in *E. coli*

BG1, bovine and human commensal *E. coli* strains, bovine pathogenic *E. coli* strains and bovine O157:H7 STEC strains (Additional file 1: Table S1) was conducted using CSI Phylogeny (version 1.4) [18]. Published *E. coli* genomes representing different *E. coli* pathotypes were selected for genomic comparison (Additional file 1: Table S1). In addition, two reference *E. coli* strains, one of which is the *E. coli* type strain (NCTC9001^T), were also included in this study. As shown in Fig. 2, the bacterial strains were clustered according to the phylogroup classification: BG1 was clustered with commensal and pathogenic *E. coli* strains belonging to phylogroup B1 (EHEC, STEC, ETEC, EAEC, APEC and *E. coli* responsible for postpartum metritis in dairy cows). The closest relative strains to BG1 were *E. coli* K71 isolated from the environment of a cow shed and *E. coli* W26 isolated from bovine feces, both of which belong to the phylogroup B1 (Fig. 2). In contrast, BG1 was more distantly clustered to pathogenic bovine and human *E. coli* strains (Fig. 2). However, *E. coli* KCJ852 (phylogroup B1), which is responsible for metritis, was more closely clustered to BG1 than the P4 and VL2732 strains associated with bovine mastitis (phylogroup A) (Fig. 2). It is of interest to note that i) the bovine *E. coli* strains of commensal origin (BG1, K71 and W26) were distantly related to bovine STEC O157:H7 strains (phylogroup E) and ii) the SNP-based phylogeny analysis failed to cluster the commensal *E. coli* strains according to their human or animal origin.

Genome sequencing information

Genome project history

Bovine commensal *E. coli* strains are poorly documented. Therefore, the *E. coli* BG1 strain was selected for genome sequencing to provide valuable genetic information for future genomic scale (whole genome) comparative analysis. *E. coli* BG1 has been used as a reference strain in studies related to carbon and nitrogen nutrition of *E. coli* strains in the bovine gastro-intestinal tract [10, 11]. The strain BG1 was isolated from the small intestine content of a cow at the slaughterhouse in January 2009. The animal was raised and slaughtered in accordance with the guidelines of the local ethics committee and current INRA (National Institute for Agricultural Research) ethical guidelines for animal welfare (Slaughterhouse Permit number: 63,345,001). The bovine intestinal samples were collected after the slaughter of animals required for experiments specifically approved by the “Comité d’éthique en matière d’expérimentation animale en Auvergne” (Permit number: CE22-08) in the experimental slaughterhouse of the “Herbipole”, INRA Saint-Genès-Champanelle, France. The Whole Genome Shotgun project was deposited at DDBJ/ENA/GenBank under the accession MOAH00000000 (Oct 31, 2016). A summary of the sequencing project information is provided in Table 2.




Fig. 2 Phylogenetic tree highlighting the position of *E. coli* BG1 relative to other *E. coli* strains. The whole genome SNP based phylogeny was established with CSI phylogeny 1.4 [28] using the genome of K71 as a reference and standard input parameters. The tree was midpoint rooted and plotted using Seaview (version 4.6.1) [56]. Each strain is identified as H (Human), B (Bovine), A (Avian), F (Food) or K12 (Laboratory strain), and its clinical or non-pathogenic (NP) characteristic is specified.

Growth conditions and genomic DNA preparation

E. coli BG1 was inoculated in Luria-Bertani broth from a single colony and incubated at 37 °C with shaking (200 rpm) to early stationary phase. The bacterial suspension was then centrifuged (10,000 g for 15 min) and the total DNA was extracted from the bacterial pellet using the DNeasy Blood and Tissue Kit following the manufacturer’s recommendations (Qiagen). DNA was

quantified using a Nanodrop spectrophotometer and DNA integrity was electrophoretically verified by ethidium bromide staining.

Genome sequencing and assembly

Whole genome sequencing was performed at the GeT-PlaGe core facility (INRA Toulouse, France). DNA-seq libraries were prepared according to Illumina’s protocols using the Illumina TruSeq Nano DNA LT Library Prep Kit. Briefly, DNA was fragmented by sonication using a Covaris M220 and adapters were ligated to be sequenced. Eight cycles of PCR were applied to amplify libraries. Library quality was assessed using the Agilent Bioanalyzer and libraries were quantified by qPCR using the Kapa Library Quantification Kit. DNA-seq experiments were performed on an Illumina MiSeq using a paired-end read length of 2 × 250 bp with the Illumina MiSeq Reagent Kits v2. The raw reads were stored in ng6 [19] and quality was checked using fastqc [20]. They were assembled with SPAdes (version 3.1.1) [21] using standard parameters.

Genome annotation

The assembled contigs were annotated with Prokka (version 1.10) [22] using standard parameters. Predicted genes were also assigned to functional categories of Clusters of Orthologous Groups (COGs) of proteins

Table 2 Genome sequencing project information for <i>E. coli</i> BG1		
MIGS ID	Property	Term
MIGS 31	Finishing quality	High quality draft
MIGS-28	Libraries used	Paired ends library
MIGS 29	Sequencing platforms	Illumina MiSeq
MIGS 31.2	Fold coverage	127x
MIGS 30	Assemblers	SPAdes version 3.1.1
MIGS 32	Gene calling method	PROKKA version 1.10
	Locus Tag	BLX34
	Genbank ID	MOAH000000000
	GenBank Date of Release	2017-02-24
	GOLD ID	
MIGS 13	BIOPROJECT	PRJNA351833
	Source Material Identifier	BG1
	Project relevance	Commensal <i>E. coli</i> (control strain)

using blastp against the NCBI COG 2014 database [23]. Additional gene features were predicted using TMHMM Server 2.0 [24], SignalP Server (version 4.1) [25], CRISPRfinder (last update 2016–09-01) [26] and ISSaga (version 2.0) [27]. PHASTER [28] was then used to identify prophage regions in the BG1 genome. A prophage region was considered to be intact if the associated completeness score was above 90, questionable if the score was between 70 and 90 and incomplete if the score was less than 70 [28].

Genome properties

The genome of *E. coli* BG1 consists of 4,782,107 bp with no discernible plasmid (no match retrieved with PlasmidFinder version 1.3 [29]), and a G + C content of 50.7%. The genome has been assembled into 84 contigs. Of the 4562 predicted genes, 4465 coded for protein and 97 were RNA-related (including eight 5S rRNA genes, suggesting the presence of 8 rRNA operons, and 86 tRNA genes). In addition, 22 pseudo genes were identified. Among the 4465 protein coding genes, 3831 (85.8%) had an assigned function while the 634 remaining genes (14.2%) encoded proteins annotated as hypothetical or unknown. In addition, the BG1 genome contained 38 predicted insertion sequences (ISs), 4 intact and 1 questionable prophage regions, and 2 CRISPR elements suggesting possible genetic crosstalk, such as horizontal gene transfer among the *E. coli* population. The genome properties are presented in Table 3. The distribution of genes into COGs functional categories is summarized in Table 4.

Table 3 Genome statistics

Attribute	Value	% of Total ^a
Genome size (bp)	4,782,107	100.00
DNA coding (bp)	4,218,785	88.22
DNA G + C (bp)	2,424,397	50.70
DNA scaffolds	84	
Total genes	4562	100.00
Protein coding genes	4465	97.88
RNA genes	97	2.13
Pseudo genes	22	0.48
Genes in internal clusters	1171	25.67
Genes with function prediction	3831	83.98
Genes assigned to COGs	3814	83.60
Genes with Pfam domains	275	6.03
Genes with signal peptides	174	3.81
Genes with transmembrane helices	1080	23.67
CRISPR repeats	2	

^aThe total is based on either the size of the genome in base pairs or the total number of proteins coding genes in the annotated genome
All the information has been obtained from Prokka annotation

Table 4 Number of genes associated with general COGs functional categories

Code	Value	% age ^a	Description
J	250	6.55	Translation, ribosomal structure and biogenesis
A	2	0.05	RNA processing and modification
K	293	7.68	Transcription
L	154	4.04	Replication, recombination and repair
B	0	0.00	Chromatin structure and dynamics
D	41	1.07	Cell cycle control, cell division, chromosome partitioning
V	93	2.44	Defense mechanisms
T	176	4.61	Signal transduction mechanisms
M	271	7.11	Cell wall/membrane/envelope biogenesis
N	156	4.09	Cell motility
U	60	1.57	Intracellular trafficking, secretion, and vesicular transport
O	153	4.01	Post-translational modifications, protein turnover, chaperones
C	282	7.39	Energy production and conversion
G	381	9.99	Carbohydrate transport and metabolism
E	335	8.78	Amino acid transport and metabolism
F	101	2.65	Nucleotide transport and metabolism
H	169	4.43	Coenzyme transport and metabolism
I	119	3.12	Lipid transport and metabolism
P	190	4.98	Inorganic ion transport and metabolism
Q	53	1.39	Secondary metabolites biosynthesis, transport and catabolism
R	211	5.53	General function prediction only
S	238	6.24	Function unknown
–	750	7.43	Not in COGs

^aThe total is based on the total number of proteins coding genes in the annotated genome

Extended insights

Genome repertoire comparison

It is admitted that bacterial genome sequences show significant diversity due to horizontal gene transfers, gene loss and other genomic rearrangements [1]. In this report, characteristics of whole genome datasets of a selection of *E. coli* strains were compared with those of *E. coli* BG1 (Table 5). Our main objective was to compare the genome of BG1 with that of bovine (K71 and W26) and human (SE15 and Nissle) commensal *E. coli* strains, but we also included a bovine pathogenic strain (VL2732) and a human EHEC pathogen (Sakai), as the bovine intestine is the main reservoir of EHEC [10]. A human uropathogenic strain (NCTC9001^T), which is also the *E. coli* type strain, was also included as reference. These strains were assigned to different phylogroups (Additional file 1: Table S1; Fig. 2). As expected, the

Table 5 Characteristics of whole genome datasets of different *E. coli* strains

Strain name	Phylo group	Origin ^a	Plasmid(s)	Genome size (bp) [chromosome + plasmid(s)]	G + C ratio (%)	CDS (nb)	Protein coding regions (nb)	rRNA operons (nb) ^b	tRNA genes (nb)
BG1	B1	Bc	0	4,782,107	50.7	4562	4465	8	86
K71	B1	Bc	0	5,115,070	50.7	5178	4872	4	65
W26	B1	Bc	0	5,118,532	50.6	4925	4852	4	66
Nissle 1917	B2	Hc	0	5,441,200	50.6	5417	4970	10	121
SE15	B2	Hc	1	4,839,683 [4,717,338 + 122,345]	50.7	4763	4572	7	85
NCTC86	A	Hc	0	5,111,920	50.6	5243	4934	7	87
VL2732	A	Bp	0	4,664,032	50.6	4615	4363	4	71
Sakai	E	Hp	2	5,594,477 [5,498,450 + 92,721 + 3306]	50.5	5447	5324	7	103
NCTC9001 ^T	B2	Hp	0	5,038,133	50.6	5154	4859	6	62

^aB: bovine; H: human; c: commensal; p: pathogen^bMinimal number of rRNA operons based on Prokka (BG1) or Genbank (K71, W26, VL2732, NCTC86, NCTC9001^T) annotation or on *rrnDB* (version 5.1) information [69]

greatest difference in genome size was observed between BG1 and the EHEC strain Sakai (the genome size of BG1 is 812,370 bp smaller than the Sakai genome [17.0% of the BG1 genome]). This difference could be explained by the number of mobile genetic elements: the Sakai genome contains 18 prophage regions (at most 5 in the BG1 genome) and 80 insertion sequences (38 in the BG1 genome) [30]. About half of the Sakai-specific sequences are of bacteriophage origin and carry the genes involved in EHEC pathogenesis (bloody diarrhea, hemolytic uremic syndrome) [30]. More surprisingly, the chromosome length of the commensal *E. coli* Nissle 1917 is 659,093 bp larger than the BG1 genome (13.8% of the BG1 genome). *E. coli* Nissle 1917 is a human commensal strain known to be a successful colonizer of the human gut and used as a probiotic for the treatment of various intestinal disorders [31]. It is well documented that the Nissle genome carries at least three genomic islands (GEIs) inserted at different tRNA sites (*serX*, *argW* and *pheV*) probably acquired by horizontal gene transfer [32, 33]. These GEIs contained genes encoding proteins considered as fitness factors (microcins, iron uptake systems, proteases ...) contributing to survival of *E. coli* Nissle and successful colonization of the human body [32, 33]. These GEIs were found in non-pathogenic *E. coli* strains but were also frequently distributed among ExPEC strains [32]. Sequence comparison showed that the genes carried by Nissle 1917 GEIs (*mch*, *mcm*, *iro*, *iuc*, *sat*, *iha*, *ybt*) are absent in the BG1 genome, suggesting the absence of these GEIs in BG1.

In accordance with the differences in genome size, the highest number of tRNA genes, described as common sites for integration of foreign DNA elements (bacteriophages, genomic islands), were detected in the genome

of *E. coli* strains Nissle and Sakai (121 and 103 tRNA genes, respectively while only 86 were identified in the BG1 draft genome (Table 5). The genome of the remaining strains carried 62 (in the type strain NCTC 9001^T) to 85 tRNA-encoding genes (Table 5). These numbers may be slightly different depending on the annotation pipeline used for the draft genome sequences.

Virulence factors

The genes encoding virulence factors in the *E. coli* BG1 genome were analyzed using blastn against the Virulence Factors Database genomic dataset [34]. A total of 164 genes encoding virulence factors were identified in BG1 (Additional file 2: Table S2), while 181 and 202 genes encoding virulence factors were found in the reference strains NCTC86 and NCTC9001^T, respectively. In-depth analysis of the BG1 genome showed that most of these genes are involved in bacterial adherence to the host epithelium, iron acquisition systems (siderophores) and flagella synthesis. As expected, genes coding for toxins produced by pathogenic *E. coli* strains responsible for diarrhea or intestinal damage in mammals (Shiga-toxin, heat stable [ST] toxin, heat-labile [LT] toxin, heat-stable enterotoxin 1 [EAST1], cytotoxic necrotizing factor 1 [CNF1]) are absent in the BG1 genome. The *E. coli* BG1 genome also lacks the genes encoding α -hemolysin and enterohemolysin which are involved in the virulence of pathogenic *E. coli* strains.

Adherence systems

A total of 49 genes coded for the synthesis of organelles involved in adherence of *E. coli* to host intestinal epithelium (Additional file 3: Table S3). Accordingly, the transmission electron micrograph of *E. coli* BG1 showed numerous fimbriae surrounding the bacteria (Fig. 1). Removal of partial genes and incomplete gene clusters

Table 6 Adherence systems encoded by the *E. coli* BG1 genome

Adherence system	Gene or genes cluster	Pathotype ^a	in vitro cell adherence ^b	Receptor
Curli fimbriae	<i>csgDEFG, csgBA</i>	EHEC, ETEC, aEPEC, APEC	T84	Matrix, plasma proteins
EhaA autotransporter	<i>ehaA</i>	EHEC, EAEC, ETEC, AIEC, EPEC	Primary bovine epithelial cells (terminal rectum)	Unknown
EhaB autotransporter	<i>ehaB</i>	EHEC, UPEC, ETEC, EIEC, EAEC	NA ^c	Collagen I, laminin
EhaC autotransporter	<i>ehaC (yfaL)</i>	EHEC, UPEC	Unknown	Unknown
ECP (<i>E. coli</i> Common Pilus)	<i>ecpRABCD</i>	ETEC, EHEC, NMEC, EAEC, aEPEC, septicemia	HT29, Hep-2, HeLa, HTB-4	Arabinosyl residues
ELF (<i>E. coli</i> Laminin-binding Fimbriae)	<i>ycbQRST</i>	EHEC, aEPEC	HT29, Hep-2, MDBK	Laminin
F9 Fimbriae	<i>z2200-z2206</i>	EHEC, UPEC, APEC, AIEC, EAEC, EPEC	EBL	Bovine fibronectin, Galβ1-3GlcNAc
EaeH adhesin	<i>eaeH</i>	UPEC, EHEC, ETEC, NMEC	UM-UC-3, Caco-2, CHO, HeLa, Vero	Unknown
HCP (Hemorrhagic Coli Pilus)	<i>hcpABC (ppdD-hofBC)</i>	ETEC, EHEC, aEPEC, APEC	T84, Caco-2, HeLa, Hep-2, MDBK, cow colon explants	Laminin, fibronectin
Stg fimbriae	<i>stgABCD</i>	APEC, UPEC	UM-UC-3, INT 407	Unknown
T1P (Type I pili)	<i>fimBE, fimAICDFGH</i>	UPEC, aEPEC, EAEC, APEC, STEC	HeLa, REC, colonic and ileal enterocytes	Mannose
UpaG autotransporter	<i>upaG</i>	UPEC	T24	Fibronectin, laminin

^aSee the "Abbreviations" paragraph^bCell lines: T84 (human colonic adenocarcinoma), HT29 (human colorectal adenocarcinoma), Hep-2 (epithelial cells from epidermoid carcinoma of the human larynx), HeLa (human cervix epithelial carcinoma), HTB-4 (human bladder transitional carcinoma), MDBK (Madin-Darby bovine kidney), EBL (embryonic bovine lung), UM-UC-3 (human bladder carcinoma), Caco-2 (human colon carcinoma), CHO (Chinese hamster ovary), Vero (kidney epithelial cells from an African green monkey), INT 407 (HeLa derivative), REC (human B cell lymphoma), T24 (human bladder transitional carcinoma)^cNA: no adherence to the cells lines tested

revealed that BG1 possessed the genetic information required to encode 12 potentially functional full adherence systems (Table 6). All these systems are known to be produced by pathogenic *E. coli* and to adhere in vitro to different cells lines (Table 6) (for reviews see [35–37]). These adherence systems reflect the ability of commensal *E. coli* to colonize distinct niches during its transit through the different compartments of the bovine gastro-intestinal tract. It is also of interest to note that some of these adherence systems possess characteristics corresponding to physiological conditions encountered in the bovine gastro-intestinal tract: i) *eaeH* expression is induced at 39 °C, the internal bovine temperature, but not at 37 °C [38] ii) the pili HCP is involved in adherence of *E. coli* to bovine gut explants [39] and iii) the F9 fimbriae are essential for in vivo colonization of calves [40]. Furthermore, the *stg* and *F9* gene clusters are strongly associated with *E. coli* belonging to phylogenetic group B1 [41, 42]. To broaden these results, in silico analysis of adherence systems carried by additional *E. coli* strains (human and bovine commensal and pathogenic isolates) (Additional file 1: Table S1; Additional file 4: Figure S1) was also performed. A hierarchical clustering based on the presence/absence of 78 distinct adherence systems encoding genes was built using R (version 3.3.1) [43]. As shown in Additional file 4: Figure S1, bovine

and human *E. coli* strains were not separately distributed (the closest relative strains to BG1 were the human *E. coli* strains S11 and IA11 [Additional file 4: Figure S1]) suggesting that the adherence systems are associated with the adaptation of *E. coli* to a specific habitat (i.e. the digestive tract) rather than host specificity. As expected, the uropathogenic strain NCTC9001^T possesses the *pap* ACDE-GHIK genes which are specific to UPEC strains [44].

Some of these adherence systems possess redundant properties: EhaB, ELF, HCP and UpaG are known to bind to laminin and curli, EhaA, EhaB, EhaC, ECP, F9, EaeH, HCP and UpaG are involved in biofilm formation (Table 6). This suggested an important role of both laminin binding and biofilm formation in survival and/or multiplication of commensal *E. coli*. Laminin is an extracellular matrix commonly present in the mammalian intestine which act as an interlinking molecule in connective tissues that promote bacterial adhesion and colonization to the host tissues [45]. Moreover, commensal *E. coli* strains can reside in mixed biofilms in the mucus layer covering the mouse intestine [4, 46]. Because the survival of *E. coli* depends on anaerobes that degrade polysaccharides included in the mucus layer, it has been hypothesized that the anaerobes in the mixed biofilms provide *E.*

coli with monosaccharide locally rather than from a mixed pool available to all species [4, 46]. Therefore, the mixed biofilm formation can result in a more efficient carbon source for commensal *E. coli* strains in the mammalian gut [4, 46].

As discussed above, the adhesion systems encoded by the BG1 genome were associated with *E. coli* strains mostly isolated from clinical cases (Table 6). However, it is important to note that the BG1 genome did not carry the genes encoding the F17, F5 and F41 fimbriae and the afimbrial adhesin CS31A mainly associated with bovine pathogenic *E. coli* strains involved in diarrhea [47]. For example, a recent epidemiological study showed that the F5/F41 fimbriae were prevalent among bovine diarrheagenic *E. coli* isolated in France [48]. The genes encoding F17, F5 and F41 are not detected in the genome of the human and bovine *E. coli* strains included in this study suggesting that these adherence systems are specific to bovine intestinal pathogenic *E. coli*.

Flagella synthesis

A total of 47 genes encoding proteins required for flagella synthesis were present in the BG1 genome. Accordingly, the transmission electron micrograph of *E. coli* BG1 showed peritrichous flagella attached to the bacterial cell surface and clearly distinct from fimbriae (Fig. 1). Flagella are mainly locomotive organelles allowing bacterial movements. However, it is well documented that the flagella (also known as H-antigen) of some pathogenic *E. coli* mediate the adhesion to or invasion of epithelial cells (NMEC, aEPEC, ETEC, EAEC, EHEC, APEC) and contribute to biofilm formation (UPEC, ETEC) (for a review see Zhou et al. [49]). In particular, flagella of aEPEC, ETEC and EHEC strains specifically recognized a receptor located at the microvillus tips of human enterocytes [50]. Interestingly, *E. coli* BG1 possesses the genetic information required to encode the flagella H21, a H antigen type reported to be involved in the invasion of EHEC O113:H21 into HCT-8 colonic epithelial cells [49]. Also, it should be noted that STEC strains with serotype O159:H21 have been isolated from bovine as well as porcine feces [51, 52].

Iron acquisition systems

Complete genetic information required for enterobactin synthesis (*entABCDEFGHI*) and ferric-enterobactin uptake (*fepABCDEFGH*) was present in the genome of *E. coli* BG1 (Additional file 2: Table S2). Siderophores, including enterobactin, are mechanisms secreted by *E. coli* to scavenge iron in order to survive and multiply in hosts or external environments. Siderophores are usually described as crucial for the proliferation of pathogenic *E. coli* in the host and have been classified as virulence

factors. However, enterobactin is frequently produced by commensal *E. coli* isolated from healthy mammals (human and animal isolates) [53]. *ent* and *fep* genes were also found in the genome of the reference strain NCTC86 (data not shown). Accordingly, Pi et al. have demonstrated that enterobactin plays a fundamental role in the colonization of healthy mouse gastro-intestinal tract by non-pathogenic *E. coli* [54].

Ethanolamine utilization

In a previous study, we demonstrated that ethanolamine present in the bovine gut is used by EHEC as a nitrogen source [11]. Furthermore, ethanolamine promotes expression of fimbrial genes and influenced EHEC adherence to epithelial cells [55]. Interestingly, *E. coli* BG1 is unable to degrade ethanolamine present in the bovine intestine, while the EHEC reference strain EDL933 gains a growth competitive advantage by assimilating ethanolamine in bovine intestinal content [11]. Therefore, we performed in-depth analysis of the genes involved in ethanolamine utilization in order to understand the inability of the commensal strain BG1 to use ethanolamine as a nitrogen source.

The degradation and assimilation of ethanolamine by EHEC EDL933 requires exogenous adenosylcobalamin (Ado-Cbl) and are encoded by 17 genes included in the *eut* operon [11]. In this study, we used blastn and SeaView (version 4.6.1) [56] to compare the *eut* genes of *E. coli* BG1 with those of EHEC EDL933. Sequence alignment showed 317 SNPs between the two *eut* operons (97.82% identity) (Additional file 5: Table S4). In addition, no premature stop codon was detected and only 34 amino acid changes due to non-synonymous SNPs were identified among the 17 predicted polypeptides encoded by the *eut* operon of BG1 (Additional file 5: Table S4). Furthermore, the presence of a 72 bp insertion was also identified in the *eutT* gene coding for cobalamin adenosyltransferase in the BG1 genome compared with the EDL933 genome (Additional file 6: Figure S2). It is important to note that ethanolamine ammonia-lyase, the key enzyme in ethanolamine degradation, required the Ado-Cbl cofactor produced by EutT to be active. The 72 bp insertion sequence at position 395 resulted in a modified translated polypeptide with 24 additional amino acids at position 132. The possible EutT conformation illustrated in Fig. 3 was predicted using Phyre (version 2.0) [57] and showed that 18 of the 24 amino acids encoded by the 72 bp sequence were predicted to form an additional alpha helix in the BG1 EutT protein.

In summary, in view of the 34 amino acid changes due to non-synonymous SNPs among the 17 predicted polypeptides encoded by the *eut* operon and the prediction



Fig. 3 Predicted secondary structure modeling of the EutT protein of *E. coli* BG1 obtained with Phyre version 2.0 [57]

of an additional alpha helix in BG1 EutT cobalamin adenosyltransferase, we suspected a reduced or abolished ethanolamine ammoniolyase activity, which could explain the inability of BG1 to assimilate ethanolamine in the bovine digestive tract.

Conclusion

The comparison of whole genomes provides information on gene content and organization, and gives an overview of how organisms are related. The draft genome sequence of *E. coli* BG1 isolated from the bovine intestine is now available and can provide valuable information at the genomic scale to explore the genetic and functional features adapted to the bovine gut. The genome of *E. coli* BG1 can be used as a reference for subsequent evolution and comparative studies (some examples of genome comparative analysis have already been described in this report).

As expected, the BG1 genome does not carry the genetic information encoding toxins responsible for

intestinal damage. More surprisingly, the *E. coli* BG1 strain possesses the genetic information required to encode systems classified as “virulence factors” and produced by pathogenic *E. coli*. This could suggest that genes encoding virulence factors are “in transit” from commensal species that act as genetic depositories with the ability to transmit DNA fragments to pathogenic *E. coli* strains. However, both pathogenic and non-pathogenic *E. coli* strains are able to colonize the gut and seem to use similar factors to adhere to the host epithelial cells. Therefore, it is questionable whether the ability of intestinal *E. coli* to colonize the host gut (resistance to the intestinal flux), excrete siderophores (iron uptake from the surrounding environment) and produce flagella (capacity to move toward nutrient-rich environments) can be considered as “virulence factors”. The terms “virulence”, “fitness” and “colonization” factors appear to overlap for *E. coli* species. In fact, factors contributing to *E. coli* survival in a given environment should be considered as fitness and adaptation factors

enabling successful colonization of the host rather than strict markers of pathogenesis. In contrast, the factors responsible for disease establishment or intestinal damages during infection (e.g. aqueous or hemorrhagic diarrhea), such as toxins or the type III secretion system, appear to be true virulence factors.

Additional files

Additional file 1: Table S1. *E. coli* strains included in this study (XLSX 13 kb)

Additional file 2: Table S2. Genes encoding virulence factors in the *E. coli* BG1 genome (XLSX 31 kb)

Additional file 3: Table S3. Genes encoding adherence systems in *E. coli* BG1 genome (XLSX 11 kb)

Additional file 4: Figure S1. Hierarchical clustering of *E. coli* strains according to adherence systems encoding genes. The dendrogram and associated heatmap are generated on the basis of gene presence/absence considering 78 genes involved in adherence, using binary distance and complete clustering method, R version 3.3.1. [43]. Blue color indicates gene presence, red gene absence. The origin of each strain is identified with B (Bovine) or H (Human). The color of the strain name corresponds to its phylogroup as in Fig. 2. (DOCX 67 kb)

Additional file 5: Table S4. Genes encoding the transport and assimilation of ethanolamine in *E. coli* BG1 genome. (XLSX 12 kb)

Additional file 6: Figure S2. Nucleotide sequence alignment of the *eutT* gene. The sequences of the *eutT* gene and the translated EutT polypeptide were aligned respectively from *E. coli* strains BG1 and EDL933 using Seaview version 4.6.1 [56]. (DOCX 15 kb)

Abbreviations

AIEC: Adherent-invasive *E. coli*; APEC: Avian pathogenic *E. coli*; EAaggEC: Enteropathogenic *E. coli*; EHEC: Enterohemorrhagic *E. coli*; EPEC: Enteropathogenic *E. coli*; ETEC: Enterotoxigenic *E. coli*; ExPEC: Extraintestinal pathogenic *E. coli*; NMEC: Neonatal meningitis *E. coli*; STEC: Shiga-producing *E. coli*; UPEC: Uropathogenic *E. coli*

Acknowledgements

The authors thank Frédérique Chaucheyras-Durand for critical reading of the manuscript, Alexandra Durand and Marine Bertoni for excellent technical assistance, Brigitte Gaillard-Martinie for the transmission electron microscopy and Olivier Bouchez for genome sequencing. The genome sequencing was performed at the GeT core facility, Toulouse, France (<http://get.genotoul.fr>), and was supported by France Génomique National infrastructure, funded as part of the "Investissement d'avenir" program managed by the Agence Nationale pour la Recherche (contract ANR-10-INBS-09). We are also grateful to the Genotoul bioinformatics platform Toulouse Midi-Pyrénées (Genotoul Bioinfo) for providing computing and storage resources.

Authors' contributions

EF and YB designed and coordinated the study; PA and CK performed the bioinformatic analysis; CK performed the genome assembly and annotation; AS performed the laboratory experiments; EF, YB, PA, AS and CK wrote the manuscript. All the authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Université Clermont Auvergne, INRA, MEDIS, F-63000 Clermont-Ferrand, France. ²Plateforme Bioinformatique Toulouse, Midi-Pyrénées UBIA, INRA, Auzerville Castanet-Tolosan, France.

Received: 7 March 2017 Accepted: 21 September 2017

Published online: 10 October 2017

References

- Welch RA. The genus *Escherichia*. In: Dworkin M, Falkow S, Rosenberg E, Schleifer K-H, Stackebrandt E, editors. The Prokaryotes, vol. 6. Third ed. Berlin: Springer; 2006. p. 60–71.
- Blount ZD. The unexhausted potential of *E. coli*. *elife*. 2015;4:e05826.
- Jones SA, Gibson T, Maltby RC, Chowdhury FZ, Stewart V, Cohen PS, Conway T. Anaerobic respiration of *Escherichia coli* in the mouse intestine. *Infect Immun*. 2011;79:4218–26.
- Conway T, Cohen PS. Commensal and pathogenic *Escherichia coli* metabolism in the gut. *Microbiol Spectr*. 2015; doi:10.1128/microbiolspec.MBP-0006-2014.
- Kaper JB, Nataro JP, Mobley HL. Pathogenic *Escherichia coli*. *Nat Rev Microbiol*. 2004;2:123–40.
- Karmali MA, Gannon V, Sargeant JM. Verocytotoxin-producing *Escherichia coli* (VTEC). *Vet Microbiol*. 2010;140:360–70.
- Oshima K, Toh H, Ogura Y, Sasamoto H, Morita H, Park SH, Ooka T, Iyoda S, Taylor TD, Hayashi T, et al. Complete genome sequence and comparative analysis of the wild-type commensal *Escherichia coli* strain SE11 isolated from a healthy adult. *DNA Res*. 2008;15:375–86.
- Rasko DA, Rosovitz MJ, Myers GS, Mongodin EF, Fricke WF, Gajer P, Crabtree J, Sebahia M, Thomson NR, Chaudhuri R, et al. The pangenome structure of *Escherichia coli*: comparative genomic analysis of *E. coli* commensal and pathogenic isolates. *J Bacteriol*. 2008;190:6881–93.
- Toh H, Oshima K, Toyoda A, Ogura Y, Ooka T, Sasamoto H, Park SH, Iyoda S, Kurokawa K, Morita H, et al. Complete genome sequence of the wild-type commensal *Escherichia coli* strain SE15, belonging to phylogenetic group B2. *J Bacteriol*. 2010;192:1165–6.
- Bertin Y, Chaucheyras-Durand F, Robbe-Masselot C, Durand A, de la Foye A, Harel J, Cohen PS, Conway T, Forano E, Martin C. Carbohydrate utilization by enterohaemorrhagic *Escherichia coli* O157:H7 in bovine intestinal content. *Environ Microbiol*. 2013;15:610–22.
- Bertin Y, Girardeau JP, Chaucheyras-Durand F, Lyan B, Pujos-Guillot E, Harel J, Martin C. Enterohaemorrhagic *Escherichia coli* gains a competitive advantage by using ethanolamine as a nitrogen source in the bovine intestinal content. *Environ Microbiol*. 2011;13:365–77.
- Joensen KG, Tetzschner AM, Iguchi A, Aarestrup FM, Scheut F. Rapid and easy in silico serotyping of *Escherichia coli* isolates by use of whole-genome sequencing data. *J Clin Microbiol*. 2015;53:2410–26.
- Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother*. 2012;67:2640–4.
- Clermont O, Christenson JK, Denamur E, Gordon DM. The Clermont *Escherichia coli* phylo-typing method revisited: improvement of specificity and detection of new phylo-groups. *Environ Microbiol Rep*. 2013;5:58–65.
- Rice P, Longden I, Bleasby A. EMBOS: the European Molecular Biology Open Software Suite. *Trends Genet*. 2000;16:276–7.
- Askari Badouei M, Jajarmi M, Mirsalehian A. Virulence profiling and genetic relatedness of Shiga toxin-producing *Escherichia coli* isolated from humans and ruminants. *Comp Immunol Microbiol Infect Dis*. 2015;38:15–20.
- Bok E, Mazurek J, Stosik M, Wojciech M, Baldy-Chudzik K. Prevalence of virulence determinants and antimicrobial resistance among commensal *Escherichia coli* derived from dairy and beef cattle. *Int J Environ Res Public Health*. 2015;12:970–85.
- Kaas RS, Leekitcharoenphon P, Aarestrup FM, Lund O. Solving the problem of comparing whole bacterial genomes across different sequencing platforms. *PLoS One*. 2014;9:e104984.
- Mariette J, Escudie F, Allias N, Salin G, Noirot C, Thomas S, Klopp C. NG6: Integrated next generation sequencing storage and processing environment. *BMC Genomics*. 2012;13:462.
- Andrews S. FastQC: a quality control tool for high throughput sequence data. 2010. <http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>. Accessed 10 Oct 2016.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, et al. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol*. 2012;19:455–77.

22. Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics*. 2014;30:2068–9.
23. Geer LY, Marchler-Bauer A, Geer RC, Han L, He J, He S, Liu C, Shi W, Bryant SH. The NCBI BioSystems database. *Nucleic Acids Res*. 2010;38:D492–6.
24. Krogh A, Larsson B, von Heijne G, Sonnhammer EL. Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *J Mol Biol*. 2001;305:567–80.
25. Petersen TN, Brunak S, von Heijne G, Nielsen H. SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nat Methods*. 2011;8:785–6.
26. Grissa I, Vergnaud G, Pourcel C. CRISPRFinder: a web tool to identify clustered regularly interspaced short palindromic repeats. *Nucleic Acids Res*. 2007;35:W52–7.
27. Varani AM, Siguier P, Goubeyre E, Charneau V, Chandler M. ISSaga is an ensemble of web-based methods for high throughput identification and semi-automatic annotation of insertion sequences in prokaryotic genomes. *Genome Biol*. 2011;12:R30.
28. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Res*. 2016;44:W16–21.
29. Carattoli A, Zankari E, Garcia-Fernandez A, Voldby Larsen M, Lund O, Villa L, Moller Aarestrup F, Hasman H. In silico detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence typing. *Antimicrob Agents Chemother*. 2014;58:3895–903.
30. Hayashi T, Makino K, Ohnishi M, Kurokawa K, Ishii K, Yokoyama K, Han CG, Ohtsubo E, Nakayama K, Murata T, et al. Complete genome sequence of enterohemorrhagic *Escherichia coli* O157:H7 and genomic comparison with a laboratory strain K-12. *DNA Res*. 2001;8:11–22.
31. Lodinova-Zadnikova R, Sonnenborn U. Effect of preventive administration of a nonpathogenic *Escherichia coli* strain on the colonization of the intestine with microbial pathogens in newborn infants. *Biol Neonate*. 1997;71:224–32.
32. Grozdanov L, Raasch C, Schulze J, Sonnenborn U, Gottschalk G, Hacker J, Dobrindt U. Analysis of the genome structure of the nonpathogenic probiotic *Escherichia coli* strain Nissle 1917. *J Bacteriol*. 2004;186:5432–41.
33. Sun J, Gunzer F, Westendorf AM, Buer J, Scharfe M, Jarek M, Gossling F, Blocker H, Zeng AP. Genomic peculiarity of coding sequences and metabolic potential of probiotic *Escherichia coli* strain Nissle 1917 inferred from raw genome data. *J Biotechnol*. 2005;117:147–61.
34. Chen L, Yang J, Yu J, Yao Z, Sun L, Shen Y, Jin Q. VFDB: a reference database for bacterial virulence factors. *Nucleic Acids Res*. 2005;33:D325–8.
35. Antao EM, Wieler LH, Ewers C. Adhesive threads of extraintestinal pathogenic *Escherichia coli*. *Gut Pathog*. 2009;1:22.
36. Croxen MA, Law RJ, Scholz R, Keeney KM, Wlodarska M, Finlay BB. Recent advances in understanding enteric pathogenic *Escherichia coli*. *Clin Microbiol Rev*. 2013;26:822–80.
37. McWilliams BD, Torres AG. Enterohemorrhagic *Escherichia coli* adhesins. *Microbiol Spectrum*. 2014; doi: 10.1128/microbiolspec.EHEC-0003-2013.
38. Easton DM, Allsopp LP, Phan MD, Moriel DG, Goh GK, Beatson SA, Mahony TJ, Cobbold RN, Schembri MA. The intimin-like protein FdeC is regulated by H-NS and temperature in enterohemorrhagic *Escherichia coli*. *Appl Environ Microbiol*. 2014;80:7337–47.
39. Xicohtencatl-Cortes J, Monteiro-Neto V, Ledesma MA, Jordan DM, Francetic O, Kaper JB, Puente JL, Giron JA. Intestinal adherence associated with type IV pili of enterohemorrhagic *Escherichia coli* O157:H7. *J Clin Invest*. 2007;117:3519–29.
40. Dziva F, van Diemen PM, Stevens MP, Smith AJ, Wallis TS. Identification of *Escherichia coli* O157 : H7 genes influencing colonization of the bovine gastrointestinal tract using signature-tagged mutagenesis. *Microbiology*. 2004;150:3631–45.
41. Lymberopoulos MH, Houle S, Daigle F, Leveille S, Bree A, Moulin-Schouleur M, Johnson JR, Dozois CM. Characterization of Stg fimbriae from an avian pathogenic *Escherichia coli* O78K80 strain and assessment of their contribution to colonization of the chicken respiratory tract. *J Bacteriol*. 2006;188:6449–59.
42. Wurpel DJ, Totsika M, Allsopp LP, Hartley-Tassell LE, Day CJ, Peters KM, Sarkar S, Ulett GC, Yang J, Tiralongo J, et al. F9 fimbriae of uropathogenic *Escherichia coli* are expressed at low temperature and recognise Galbeta1-3GlcNAc-containing glycans. *PLoS One*. 2014;9:e93177.
43. Team RC. R: A language and environment for statistical computing. Vienna, Austria: The R Project for Statistical Computing; 2016.
44. Arthur M, Campanelli C, Arbeit RD, Kim C, Steinbach S, Johnson CE, Rubin RH, Goldstein R. Structure and copy number of gene clusters related to the *pap* P-adhesin operon of uropathogenic *Escherichia coli*. *Infect Immun*. 1989;57:314–21.
45. Simon-Assmann P, Spenle C, Lefebvre O, Keding M. The role of the basement membrane as a modulator of intestinal epithelial-mesenchymal interactions. *Prog Mol Biol Transl Sci*. 2010;96:175–206.
46. Leatham-Jensen MP, Fridmott-Moller J, Adediran J, Mokszycki ME, Banner ME, Caughron JE, Krogfelt KA, Conway T, Cohen PS. The streptomycin-treated mouse intestine selects *Escherichia coli* *envZ* missense mutants that interact with dense and diverse intestinal microbiota. *Infect Immun*. 2012;80:1716–27.
47. Nagy B, Fekete PZ. Enterotoxigenic *Escherichia coli* in veterinary medicine. *Int J Med Microbiol*. 2005;295:443–54.
48. Valat C, Forest K, Auvray F, Metayer V, Meheut T, Polizzi C, Gay E, Haenni M, Oswald E, Madec JY. Assessment of adhesins as an indicator of pathovar-associated virulence factors in bovine *Escherichia coli*. *Appl Environ Microbiol*. 2014;80:7230–4.
49. Zhou M, Yang Y, Chen P, Hu H, Hardwidge PR, Zhu G. More than a locomotive organelle: flagella in *Escherichia coli*. *Appl Microbiol Biotechnol*. 2015;99:8883–90.
50. Sampaio SC, Luiz WB, Vieira MA, Ferreira RC, Garcia BG, Sinigaglia-Coimbra R, Sampaio JL, Ferreira LC, Gomes TA. Flagellar cap protein FliD mediates adherence of atypical enteropathogenic *Escherichia coli* to enterocyte microvilli. *Infect Immun*. 2016;84:1112–22.
51. Baranzoni GM, Fratamico PM, Gangiredla J, Patel I, Bagi LK, Delannoy S, Fach P, Boccia F, Anastasio A, Pepe T. Characterization of shiga toxin subtypes and virulence genes in porcine shiga toxin-producing *Escherichia coli*. *Front Microbiol*. 2016;7:574.
52. Pigatto CP, Schocken-turrino RP, Souza EM, Pedrosa FO, Comarella L, Irino K, Boccia MA, Farah SM, Warth JF, Fadel-Picheth CM. Virulence properties and antimicrobial susceptibility of Shiga toxin-producing *Escherichia coli* strains isolated from healthy cattle from Parana State. *Brazil Can J Microbiol*. 2008;54:588–93.
53. Searle LJ, Meric G, Porcelli I, Sheppard SK, Lucchini S. Variation in siderophore biosynthetic gene distribution and production across environmental and faecal populations of *Escherichia coli*. *PLoS One*. 2015;10:e0117906.
54. Pi H, Jones SA, Mercer LE, Meador JP, Caughron JE, Jordan L, Newton SM, Conway T, Klebba PE. Role of catecholate siderophores in gram-negative bacterial colonization of the mouse gut. *PLoS One*. 2012;7:e50020.
55. Garsin DA. Ethanolamine utilization in bacterial pathogens: roles and regulation. *Nat Rev Microbiol*. 2010;8:290–5.
56. Gouy M, Guindon S, Gascuel O. SeaView version 4: A multiplatform graphical user interface for sequence alignment and phylogenetic tree building. *Mol Biol Evol*. 2010;27:221–4.
57. Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJ. The Phyre2 web portal for protein modeling, prediction and analysis. *Nat Protoc*. 2015;10:845–58.
58. Field D, Garrity G, Gray T, Morrison N, Selengut J, Sterk P, Tatusova T, Thomson N, Allen MJ, Angiuoli SV, et al. The minimum information about a genome sequence (MIGS) specification. *Nat Biotechnol*. 2008;26:541–7.
59. Woese CR, Kandler O, Wheelis ML. Towards a natural system of organisms: proposal for the domains Archaea, Bacteria, and Eucarya. *Proc Natl Acad Sci U S A*. 1990;87:4576–9.
60. Garrity GM, Bell JA, Lilburn T. Phylum XIV. Proteobacteria *phyl. nov.* In: Garrity GM, Brenner DJ, Krieg NR, Staley JR, editors. *Bergey's Manual of Systematic Bacteriology. Second edition, Volume 2, Part B*. New York: Springer; 2005. p. 1.
61. Garrity GM, Bell JA, Lilburn T. Class III. Gammaproteobacteria *class. nov.* In: Garrity GM, Brenner DJ, Krieg NR, Staley JR, editors. *Bergey's Manual of Systematic Bacteriology. Second edition, Volume 2, Part B*. New York: Springer; 2005. p. 1.
62. Euzéby J. Validation list no. 106. Validation of publication of new names and new combinations previously effectively published outside the IJSEM. *Int J Syst Evol Microbiol*. 2005;55:2235–8.
63. Garrity GM, Holt JG. Taxonomic outline of the archaea and bacteria. In: Garrity GM, Boone DR, Castenholz RW, editors. *Bergey's Manual of Systematic Bacteriology*, vol. 1. Second ed. New York: Springer; 2001. p. 155–66.
64. Hill LR, Skerman VBD, Sneath PHA. Corrigenda to the approved lists of bacterial names: edited for the international committee on systematic bacteriology. *Int J Syst Bacteriol*. 1984;34:508–11.
65. Rahn O. New principles for the classification of bacteria. *Zentralblatt für Bakteriologie, Parasitenkunde, Infektionskrankheiten und Hygiene*. 1937;96:273–86.
66. Skerman VBD, McGowan V, Sneath PHA. Approved lists of bacterial names. *Int J Syst Bacteriol*. 1980:225–420.
67. Castellani A, Chalmers AJ. Genus *Escherichia* Castellani and Chalmers, 1918. Wood W. And Co. Manual of Tropical Medicine. Third edition. New York; William Wood and Company; 1919. p.941-3.

68. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, et al. Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet.* 2000;25:25–9.
69. Stoddard SF, Smith BJ, Hein R, Roller BR, Schmidt TM. rrnDB: improved tools for interpreting rRNA gene abundance in bacteria and archaea and a new foundation for future development. *Nucleic Acids Res.* 2015;43:D593–8.

Submit your next manuscript to BioMed Central
and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

